



# Gene expression and highly diluted molecules

Marta Marzotto, Debora Oliosio and Paolo Bellavite\*

Section of General Pathology, Department of Pathology and Diagnostics, University of Verona, Verona, Italy

\*Correspondence: paolo.bellavite@univr.it

## Edited by:

Michael Heinrich, University College London, UK

## Reviewed by:

Fang-Rong Chang, Kaohsiung Medical University, Taiwan

Hwee-Ling Koh, National University of Singapore, Singapore

Benedict Green, Agricultural Research Services-United States Department of Agriculture, USA

**Keywords:** *Gelsemium sempervirens*, high dilutions, gene expression regulation, microarray technology, neurocyte cell line, Homeopathy

## A commentary on

### High diluted molecules and gene expression

by Chirumbolo, S. (2014). *Front. Pharmacol.* 5:183. doi: 10.3389/fphar.2014.00183

Plants of the genus *Gelsemium* have measurable effects on anxiety-like symptoms in laboratory models (Magnani et al., 2010; Liu et al., 2013; Meyer et al., 2013; Jin et al., 2014). Searching for a possible mechanism of action, we found that very low doses and homeopathic dilutions of *Gelsemium sempervirens* (*Gelsemium*) modulate the expression of genes involved in neuronal functions (G-protein coupled receptor signaling pathways, calcium homeostasis, inflammatory response and receptors) (Marzotto et al., 2014; Oliosio et al., 2014).

The commentary's first criticism is that "the search for an involvement of neural genes related to anxiety/depression or mood disorders is biased by the expression of human genes having no orthologs/homologs in mice, where the authors reported evidence about *Gelsemium* action on behavioral tests in animal anxiety models." Frankly, we do not see what type of bias can be found in our line of research: *Gelsemium* is traditionally used as a remedy for anxiety-related symptoms in humans; we have shown that it works in mice models (Magnani et al., 2010; Bellavite et al., 2012), and we wanted to study its mechanism of action at the molecular level, using human neurocyte cell lines. This type of experimental procedure, which addresses various knowledge gaps using

both animal studies and *in vitro* models, is very common in pharmacological studies.

Furthermore, the commentary (Chirumbolo, 2014) states that "some genes indicated to be downregulated by *Gelsemium* 2c, should not be expressed by neuronal cells (e.g., CD163, MPO, C8B, LST1, TREM2, notoriously expressed in immune cell)." Actually, it is well known that neurons express genes of cellular pathways also involved in cytokine/chemokine and immune responses. Genes related to immune and inflammatory responses are expressed in SH-SY5Y cells, as reported by many researchers (Gatta et al., 2011; Toyama et al., 2011; Cui et al., 2012; Xu et al., 2013).

Although *Gelsemium* contains several different compounds (Dutt et al., 2010; Jin et al., 2014), the major active alkaloid of this plant is gelsemine. In the cited commentary we read that "concentration of gelsemine was not assessed, as it was solely calculated on previous spectrometry investigations and new preparations, from ethanol draw extracts, were not further quantified by analytical chemistry." This statement is misleading because the concentration of gelsemine which we correctly reported (0.021 g/100 ml, corresponding to  $6.5 \times 10^{-4}$  mol/L) (Marzotto et al., 2014, p. 2) was precisely determined by liquid chromatography (not "spectrometry") in the mother tincture from which the samples were prepared. The concentration of UV-VIS absorbing substances decreased by a factor of 100 at each centesimal dilution step, to become analytically undetectable after a few passages, as we have shown by means of spectrometry in our paper. What we were interested in was to check the

accuracy of the procedures as far as possible, not to determine the gelsemine concentration in all subsequent centesimal dilutions. Perhaps we need to remind readers that normally, when one performs a study of dose-response and the concentration in the highest dose and the dilution factor are known, there is no need to determine the concentration of substances in all successive dilutions.

Another erroneous criticism which forces us to reply is the statement (Chirumbolo, 2014) that our UV-VIS spectra (Marzotto et al., 2014) "showed a peak at 250 nm caused by contaminating millimolar ethanol in *Gelsemium* 2c." This is untrue, and we fail to understand how the writer could have reached such a conclusion, since UV-visible absorption spectra were performed with a double-beam spectrophotometer using drug samples and reference controls, having exactly the same ethanol concentration. Subsequently the commentary claims that "the authors tested a complex mixture of *G. sempervirens* extract, containing at least about 0.154 mM EtOH at 2c, if dilutions were conducted exactly" and that "concentration of EtOH, set at 30% v/v, faded out to 0.003% in tested dilutions but the authors did not clarify how much for each centesimal dilution in the Methods section." The writer supposes that high ethanol concentration could have caused "latent apoptosis." This is confusing and misleading. As clearly indicated in our paper (Marzotto et al., 2014), the final ethanol concentration was 0.03% v/v (p. 2), and "no significant differences in cell viability were observed between cells treated with the ethanol control solution 0.03% (v/v) and untreated cells" (p. 5).


Furthermore, as other researchers have found (Do et al., 2013), cell viability of neurocytes is unaffected by doses of ethanol up to 10 mmol/L (0.06%). Low doses of ethanol may influence gene expression, but would have acted in the same way in the *Gelsemium* and in the control samples. To rule out any misinterpretation of our findings, **Figure 1** gives a brief outline of our microarray experiments.

Finally, the commentary (Chirumbolo, 2014) maintains that “Gene array profile of expression following 24 h incubation with *Gelsemium* 2c, showed down-regulation of 49 genes, namely 87.5% gene array.” It is hard to understand how this 87.5%

of genes was calculated, since the 49 genes represent about 0.1% of the genes on the microarray chip. The observation that “many gene products, listed in the expression profile of 56 genes array, such as LOC154872, KIAA0825, LOC150763, C1orf167, have not been identified” is bizarre. The presence of some as yet unidentified sequences in the human genome database is of course well known.

We hope that these clarifications will be welcomed in the interests of providing correct and truthful information to readers. In summary, we provided reliable evidence that *Gelsemium* exerts a prevalently inhibitory effect on a series of neurocyte genes across a wide dose

range (Marzotto et al., 2014; Oliosio et al., 2014). The effect decreases with decreasing doses, but whole genome expression analysis made it possible to detect statistically significant changes even at extremely low doses and homeopathic dilutions (e.g., 5th and 9th centesimal dilution, corresponding to final gelsemine concentrations of  $6.5 \times 10^{-15}$  mol/L and  $6.5 \times 10^{-23}$  mol/L respectively). More robust conclusions about the role of the genes involved will require determination whether proteins encoded by the affected genes are similarly changed, using proteomic and phosphoproteomic approaches, and/or further studies using purified active plant compounds.



## REFERENCES

- Bellavite, P., Conforti, A., Marzotto, M., Magnani, P., Cristofolletti, M., Olioso, D., et al. (2012). Testing homeopathy in mouse emotional response models: pooled data analysis of two series of studies. *Evid. Based Complement. Alternat. Med.* 2012:954374. doi: 10.1155/2012/954374
- Chirumbolo, S. (2014). High diluted molecules and gene expression. *Front. Pharmacol.* 5:183. doi: 10.3389/fphar.2014.00183
- Cui, W. Y., Wang, J., Wei, J., Cao, J., Chang, S. L., Gu, J., et al. (2012). Modulation of innate immune-related pathways in nicotine-treated SH-SY5Y cells. *Amino Acids* 43, 1157–1169. doi: 10.1007/s00726-011-1171-0
- Do, H., Park, H. J., Sohn, E. H., Kim, B. O., Um, S. H., Kwak, J. H., et al. (2013). Ethanol induces cell cycle arrest and triggers apoptosis via Sp1-dependent p75NTR expression in human neuroblastoma cells. *Cell Biol. Toxicol.* 29, 365–380. doi: 10.1007/s10565-013-9260-3
- Dutt, V., Thakur, S., Dhar, V. J., and Sharma, A. (2010). The genus gelsemium: an update. *Pharmacogn. Rev.* 4, 185–194. PMID:22228960. doi: 10.4103/0973-7847.70916
- Gatta, V., Drago, D., Fincati, K., Valenti, M. T., Dalle, C. L., Sensi, S. L., et al. (2011). Microarray analysis on human neuroblastoma cells exposed to aluminum, beta(1-42)-amyloid or the beta(1-42)-amyloid aluminum complex. *PLoS ONE* 6:e15965. doi: 10.1371/journal.pone.0015965
- Jin, G. L., Su, Y. P., Liu, M., Xu, Y., Yang, J., Liao, K. J., et al. (2014). Medicinal plants of the genus Gelsemium (Gelsemiaceae, Gentianales)-A review of their phytochemistry, pharmacology, toxicology and traditional use. *J. Ethnopharmacol.* 152, 33–52. doi: 10.1016/j.jep.2014.01.003
- Liu, M., Huang, H. H., Yang, J., Su, Y. P., Lin, H. W., Lin, L. Q., et al. (2013). The active alkaloids of Gelsemium elegans Benth. are potent anxiolytics. *Psychopharmacology (Berl.)* 225, 839–851. doi: 10.1007/s00213-012-2867-x
- Magnani, P., Conforti, A., Zanolin, E., Marzotto, M., and Bellavite, P. (2010). Dose-effect study of Gelsemium sempervirens in high dilutions on anxiety-related responses in mice. *Psychopharmacology (Berl.)* 210, 533–545. doi: 10.1007/s00213-010-1855-2
- Marzotto, M., Olioso, D., Brizzi, M., Tononi, P., Cristofolletti, M., and Bellavite, P. (2014). Extreme sensitivity of gene expression in human SH-SY5Y neurocytes to ultra-low doses of Gelsemium sempervirens. *BMC Complement. Altern. Med.* 14:104. doi: 10.1186/1472-6882-14-104
- Meyer, L., Boujedaini, N., Patte-Mensah, C., and Mensah-Nyagan, A. G. (2013). Pharmacological effect of gelsemine on anxiety-like behavior in rat. *Behav. Brain Res.* 253, 90–94. doi: 10.1016/j.bbr.2013.07.010
- Olioso, D., Marzotto, M., Moratti, E., Brizzi, M., and Bellavite, P. (2014). Effects of Gelsemium sempervirens L. on pathway-focused gene expression profiling in neuronal cells. *J. Ethnopharmacol.* 153, 535–539. doi: 10.1016/j.jep.2014.02.048
- Toyama, T., Yoshida, E., Shinkai, Y., and Kumagai, Y. (2011). DNA microarray analysis of human neuroblastoma SH-SY5Y cells exposed to methylmercury. *J. Toxicol. Sci.* 36, 843–845. doi: 10.2131/jts.36.843
- Xu, L. J., Jiang, T., Zhang, F. J., Han, J. F., Liu, J., Zhao, H., et al. (2013). Global transcriptomic analysis of human neuroblastoma cells in response to enterovirus type 71 infection. *PLoS ONE* 8:e65948. doi: 10.1371/journal.pone.0065948

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 31 August 2014; accepted: 17 October 2014; published online: 12 November 2014.

Citation: Marzotto M, Olioso D and Bellavite P (2014) Gene expression and highly diluted molecules. *Front. Pharmacol.* 5:237. doi: 10.3389/fphar.2014.00237

This article was submitted to *Ethnopharmacology*, a section of the journal *Frontiers in Pharmacology*.

Copyright © 2014 Marzotto, Olioso and Bellavite.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.