


RESEARCH ARTICLE

Open Access



The serine-48 residue of nucleolar phosphoprotein nucleophosmin-1 plays critical role in subcellular localization and interaction with porcine circovirus type 3 capsid protein

Jianwei Zhou¹, Juan Li¹, Haimin Li¹, Ying Zhang¹, Weiren Dong¹, Yulan Jin¹, Yan Yan¹, Jinyan Gu^{1*} and Jiyong Zhou^{1,2*} 

Abstract

The transport of circovirus capsid protein into nucleus is essential for viral replication in infected cell. However, the role of nucleolar shuttle proteins during porcine circovirus 3 capsid protein (PCV3 Cap) import is still not understood. Here, we report a previously unidentified nucleolar localization signal (NoLS) of PCV3 Cap, which hijacks the nucleolar phosphoprotein nucleophosmin-1 (NPM1) to facilitate nucleolar localization of PCV3 Cap. The NoLS of PCV3 Cap and serine-48 residue of N-terminal oligomerization domain of NPM1 are essential for PCV3 Cap/NPM1 interaction. In addition, charge property of serine-48 residue of NPM1 is critical for nucleolar localization and interaction with PCV3 Cap. Taken together, our findings demonstrate for the first time that NPM1 interacts with PCV3 Cap and is responsible for its nucleolar localization.

Keywords: porcine circovirus type 3, capsid protein, nucleolar localization signal, nucleolar phosphoprotein nucleophosmin-1, amino acid charge property

Introduction

Viruses of genus *Circovirus*, in the family *Circoviridae*, have been detected in terrestrial, aquatic and avian species, including pigs, ducks, dogs, minks, rats, palm civets, geese, pigeons, canaries, parrots, and others [1–8]. Circoviruses are circular, single-stranded DNA viruses that are the smallest known autonomously replicating viruses in mammals [9]. Two porcine circovirus (PCV) genotypes, PCV1 and PCV2, have been extensively studied [10, 11]. PCV1 is nonpathogenic to pigs and was identified in the porcine kidney cell line PK-15 [12, 13]. PCV2

is pathogenic and causes porcine circovirus-associated diseases (PCVAD) and immunosuppression [12, 14, 15].

In 2015, a novel PCV, designated as PCV3, was first identified by next generation sequence (NGS) analysis as a pathogenic agent in sows that died and displayed acute porcine dermatitis and nephropathy syndrome (PDNS)-like clinical signs, reproductive failure, cardiac pathology, and multisystemic inflammation in the United States [15, 16] and then in China, Poland, South Korea, Brazil, Thailand, Germany, Denmark, Spain, and Italy [17–24]. Retrospective research data have shown that PCV3 possesses high sequence homology to bat circovirus and that the earliest cases of PCV3 infection can be traced back to 1966 in China, suggesting that PCV3 may have been derived early from bats and gradually adapted to pigs [25]. Recently, Fu et al. found that

*Correspondence: gujinyan@zju.edu.cn; jyzhou@zju.edu.cn

¹ MOA Key Laboratory of Animal Virology, Center of Veterinary Sciences, Zhejiang University, 866 Yuhangtang Road, Hangzhou, Zhejiang 310058, PR China

Full list of author information is available at the end of the article



© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

PCV3 has been circulated in swine herds for nearly half a century and may have originated from a bat-associated circovirus [26]. This implicates that in addition to PCV2, the newly discovered PCV3, is also pathogenic to pigs and serves as an etiological agent for PCVAD.

The PCV3 genome is a single-stranded circular DNA of 2.0 kb, which contains three major open reading frames (ORFs) [16]. The ORF2-encoded capsid protein (Cap) is necessary for virion packaging and participates in genome replication by interacting with Rep protein and is also the main viral immunogen and acts as a key regulator of viral replication as well as the virus-host interaction [27–34]. The N-terminus of PCV3 Cap is rich in basic amino acids, and the amino acid residues 1–34 was identified as nuclear localization signal (NLS), which is essential for nuclear transport of PCV3 Cap [35]. Besides, PCV3 Cap interacts with signal transducer and activator of transcription 2 (STAT2) to inhibit type I interferon signaling [36]. Nuclear entry of viral proteins is responsible for genome replication and is mediated by cellular shuttle proteins. The capsid of herpes simplex virus type 1 (HSV-1) and adenovirus 2 (AdV-2), the nucleoprotein of influenza A virus (IAV), and NS5A of hepatitis C virus (HCV) bound to nuclear pore complex (NPC) proteins, nuclear import soluble factors, heat shock protein 70 (HSP70), and histone H1 for nuclear shuttling, which were essential for viral propagation [37–39]. Besides NLS, there may be involvement of host proteins for nuclear entry of PCV3 Cap. For example, hepatitis delta virus (HDV) was studied for nucleolar localization and it was found that interacting with nucleolin (NCL) promoted viral replication [40]. Since PCV3 Cap is a nucleus-located protein, whether host proteins participating in its localization need to be elucidated.

Nucleolar phosphoprotein nucleophosmin-1 (NPM1) is a nuclear shuttle factor and resides mainly in the nucleolus, which bears a N-terminal oligomerization domain (OligoD), a central histone-binding domain (HistonD), and a C-terminal nucleic acid-binding domain (NBD) [41, 42]. NPM1 plays roles in many cellular processes such as ribosome biogenesis, DNA replication and repair, stress response, centrosome duplication, and nucleocytoplasmic transport [43]. NPM1 has been reported to interact with viral proteins, such as human immunodeficiency virus type 1 (HIV-1) Rev protein [44] and adenoviral core proteins [45]. However, whether NPM1 interacts PCV3 Cap remains unclear.

In the present study, we found that knockdown of *NPM1* abolished nucleolar localization of PCV3 Cap, and confirmed the essential role of NoLS of PCV3 Cap for binding to NPM1. We proved that charge property of serine-48 residue within N-terminal OligoD of NPM1 determines its subcellular localization and interaction with PCV3 Cap. Taken together, our findings highlight the critical role of NPM1 in nucleolar entry of PCV3 Cap.

Materials and methods

Cells

The Epithelial PK-15 cell line, free of PCV was provided by the China Institute of Veterinary Drugs Control and was maintained in our laboratory. The cell line was cultured in minimal essential medium (MEM) (Gibco, Carlsbad, CA, USA). HEK293T cells (CRL-11268; ATCC, Manassas, VA, USA) were cultured in Dulbecco's modified Eagle medium (DMEM) (Gibco). Both media were supplemented with 10% fetal bovine serum (FBS) (CCS30010.02; MRC, Australia).

Antibodies and reagents

Rabbit polyclonal antibodies (pAb) against Myc (R1208-1), GFP (SR48-02), Flag (0912-1), and mouse monoclonal antibodies (mAb) against β -actin (M1210-2) and GST (M0807-1) were purchased from Huaan Biological Technology (Hangzhou, China). Mouse anti-Myc (05-419) and anti-Flag (F1804) mAbs were obtained from Sigma-Aldrich (St. Louis, MO, USA). Anti-Flag affinity resin (A2220) for immunoprecipitation was also purchased from Sigma-Aldrich. Rabbit mAb against NPM1 (ab52644) was purchased from Abcam (Cambridge, MA). NP-40 cell lysis buffer (50 mM Tris [pH 7.4], 150 mM NaCl, 1% NP-40) was purchased from Beyotime (P0013F; Shanghai, China). Horseradish peroxidase (HRP)-labeled goat anti-mouse and anti-rabbit IgG were purchased from KPL (Milford, MA, USA).

Plasmid construction and transfection

DNA fragments encoding full-length and truncated PCV3 *Cap* variants were amplified by PCR from full-length genomic DNA of PCV3 [16] (accession no. KT869077.1), and subcloned separately into vectors pCMV-Myc-N (Clontech, Palo Alto, CA, USA), pCMV-Flag-N (Clontech), pCMV-Flag-gst-N (Clontech), pEGFP-C3 (Clontech), and pmCherry-C1 (Clontech) for different purposes. The resultant plasmids were

Flag-PCV3-Cap, Flag-gst-PCV3-Cap, Myc-PCV3-Cap, GFP-PCV3-Cap, GFP-PCV3-Cap(1-34aa), GFP-PCV3-Cap(35-214aa), mCherry-PCV3-Cap, mCherry-PCV3-Cap(1-34aa), mCherry-PCV3-Cap(35-214aa). The nucleotide fragments PCV1-NLS, PCV4-NLS, CanaryCV-NLS, CanineCV-NLS, MinkCV-NLS, DragonflyCV-NLS, CoCV-NLS, BtCV-NLS, DuCV-NLS, GoCV-NLS, and BFDV-NLS were synthesized and inserted into the

vector pEGFP-C3 by Sunya Biotechnology (Zhejiang, China). GFP-PCV2-NLS plasmid was constructed and stored in our laboratory. The nuclear localization signals (NLSs) within capsid protein of circoviruses from different species are listed in Table 1. The full-length cDNA sequences of *NCL* (accession no. XM_021074959.1), *NPM1* (accession no. XM_013990662.2), and truncated *NPM1* variants were amplified from PK-15 cells

Table 1 The nuclear localization signals (NLSs) of circovirus Cap from different species

Circoviruses	Accession No	Sequences of NLSs
PCV1	AY193712.1	MTWPRRRYRRRTRPRSHLGNILRRRYPYLAHPAFNRNRYRWRRK
PCV2	AY188355.1	MTYPRRRYRRRHRPRSHLQILRRRPWLHPRHRYRWRRK
PCV3	KT869077.1	MRHRAIFRRRPRRRRRHRRRYARRRLFIRRP
PCV4	MK986820.1	MPIRSRYSRRRRNRNRQRRRGLWPRANRRRYRWRRKN
CanaryCV	AJ301633.2	MWLTFNQVARRRRLAPRRRRWRRRYWXRRIIPANRR
CanineCV	MK731981.1	MRVRRHARASRRRYRTRPLIRYRRRQNNFKLFHLRLRR
MinkCV	MK561562.1	MPVRSRYSRRRRWRNRNRGPRRYARGGYRWRR
DragonflyCV	HQ638058.1	MVRYRAARRPVRRARRSRVKLFRRRRYYHRR
CoCV	HQ401281.1	MRRRFYRRRAPIRRRRIRRRRLSRMRGHR
BtCV	JF938082.1	MARFRRRVARRRPVRSIRRIIRRRRYGRRR
DuCV	EF370476.1	MRRSTYRRAYARRRRRGLRRRLRRRLRIGRP
GoCV	AF536940.1	MPLYRARPRSLYRRRRAANRRRYYRRRLHIGRIR
BFDV	AY450445.1	MNCACAIQIRRRRYARPYRRRHNRYYRRRYYFRRRR

Table 2 Primers used for cloning PCR

Gene product	Sense primer (5'–3')	Antisense primer (5'–3')
PCV3 Cap(1-214aa)	ATGAGACACAGACTATATTC	TTAGAGAACGGACTTGTAACGAATC
PCV3 Cap(35-214aa)	ATGGCTGGCACATACTACACAAAGAAAT	TTAGAGAACGGACTTGTAACGAATC
PCV3 Cap(1-34aa)	ATGAGACACAGACTATATTC	TTA TGTGGGCTCCTAATGAATA
NPM1(1-294aa)	ATGGAAGATTCGATGGATAT	TTAAAGAGACTTCCTCCACT
NPM1(1-117aa)	ATGGAAGATTCGATGGATAT	TACTAAGTGTGTCCTAATATGC
NPM1(118-188aa)	GCTGTAGAGGAAGATGCAGAGT	TTCCGCTTCTCATCATCAA
NPM1(189-294aa)	GAAAAAGCTCCAGTAAAGAAAT	TTAAAGAGACTTCCTCCACT
NPM1(1-188aa)	ATGGAAGATTCGATGGATAT	TTCCGCTTCTCATCATCAA
NPM1(118-294aa)	GCTGTAGAGGAAGATGCAGAGT	TTAAAGAGACTTCCTCCACT
NPM1(1-117aa + 189-294aa)	ATGGAAGATTCGATGGATAT TAGTGGACAGCCTTAGTAGAAAAAGCTCCAGTAAAG ATGGAAGATTCGATGGATAT	CTTTACTGGAGCTTTTCTACTAAGTGCTGCCACTA TTAAAGAGACTTCCTCCACT TTAAAGAGACTTCCTCCACT
NPM1-S48A	TTATCTTTAAGAACGGTGCATTAGGGGCT	AGCCCCTAATGCGACCGTTCTTAAAGATAA
NPM1-S48E	TTATCTTTAAGAACGGTGAATTAGGGGCTGGCGCAA	TTTGCGCCAGCCCCTAATTGACCGTTCTTAAAGATAA
NPM1-S88A	AACCTTTGAAAATGTCTGTACAGCCGACGGTTGCACTTG	CAAGTGCAACCGTGGCTGTACAGACATTTCAAAGTT
NPM1-S88E	AACCTTTGAAAATGTCTGTACAGCCGACGGTTGAACTTG	CAAGTTCAACCGTGGCTGTACAGACATTTCAAAGTT
NPM1-T95A	TTGGGGGCTTTGAAATAGACCACCTGTGGTCTTA	TAAGACCACAGGTGGTGTATTTCAAAGCCCCCAA
NPM1-T95D	TTGGGGGCTTTGAAATAGATCCACCTGTGGTCTTA	TAAGACCACAGGTGGTGTATTTCAAAGCCCCCAA
NPM1-S48T	TTATCTTTAAGAACGGTGCATTAGGGGCT	AGCCCCTAATGCGACCGTTCTTAAAGATAA
NPM1-S48D	AAAATGAGCACCAGTTATCTTTAAGAACGGTTCGATTTA	TAAATGACCGTTCTTAAAGATAACTGGTGCTCATTTT
NPM1-S48K	TTATCTTTAAGAACGGTCAAATTAGGGGCTGGCGCAA	TTTGCGCCAGCCCCTAATTGACCGTTCTTAAAGATAA
NPM1-S48R	TTATCTTTAAGAACGGTGCAGATTAGGGGCTGGCGCAA	TTTGCGCCAGCCCCTAATCTGACCGTTCTTAAAGATAA

and cloned into vectors pmCherry-C1, pCMV-Flag-N, pEGFP-C3, and pGEX-4T-1 (GE Healthcare Biosciences, Piscataway, NJ, USA) using specific primers. The resultant plasmids were mCherry-NCL, GFP-NCL, mCherry-NPM1, Flag-NPM1, GST-NPM1, GFP-NPM1(1-294aa), GFP-NPM1(1-117aa), GFP-NPM1(118-188aa), GFP-NPM1(189-294aa), GFP-NPM1(1-188aa), GFP-NPM1(118-294aa), and GFP-NPM1(1-117aa + 189-294aa). Mutants were generated by site-specific mutagenesis. The primers used are summarized in Table 2. For transfection, PK-15 and HEK293T cells were seeded onto plates or glass coverslips at a suitable density according to the experimental plan and grown up to 70% to 90% confluency. jetPRIME transfection reagent (Polyplus Transfection, New York, NY, USA) was used for PK-15 cell transfection, and ExFect transfection reagent (T101-01/02; Vazyme Biotechnology, Nanjing, China) was used for 293T cell transfection according to the manufacturers' instructions.

Confocal microscopy

To study the colocalization of NPM1, NCL and full-length PCV3 Cap or its truncated variants, HEK293T or PK-15 cells were cotransfected with indicated vectors fused with GFP and mCherry tags. The cells were fixed using 4% paraformaldehyde (PFA) for 20 min and permeabilized with 0.2% Triton X-100 for 5 min at room temperature. Cellular nuclei were counterstained with 10 µg/mL 4', 6'-diamidino-2-phenylindole (DAPI; 10236276001; Roche, Mannheim, Germany). The cells were then visualized under LSM780 laser scanning confocal microscope (Zeiss, Oberkochen, Germany). GFP signal was detected after excitation at 488 nm with an emission long-band filter at 505–530 nm (green). mCherry fluorescence was detected after excitation at 561 nm with an emission long-pass filter at 550–600 nm (red). DAPI was detected after excitation at 405 nm with an emission long-pass filter at 445–450 nm (blue).

SDS-PAGE and immunoblotting

For western blotting, cells were lysed in lysis buffer after transfection. Samples were collected, and proteins were separated by standard SDS-PAGE and transferred to

nitrocellulose membranes (GE Healthcare) followed by blocking in phosphate-buffered saline (PBS) containing 5% skimmed milk powder and 0.05% Tween 20. The membrane was washed three times with PBS containing 0.05% Tween 20. After that, the membranes were incubated with primary antibody overnight at 4 °C. After three to five washes with PBS containing 0.05% Tween 20, the membranes were incubated with HRP-labeled secondary antibody at room temperature for 1.0 h. Immunoreactive protein bands were then visualized using enhanced chemiluminescence reagent (Amersham Biosciences, Little Chalfont, United Kingdom) and imaged using AI680 Images (GE Healthcare).

Co-IP and GST pull-down assays

For Co-IP assays, HEK293T cells were transfected with the indicated plasmids for 48 h. Cells were lysed in NP-40 cell lysis buffer containing a protease inhibitor cocktail. After centrifugation at $12,000 \times g$ for 10 min, the supernatant was treated with protein A/G plus agarose (sc-2002; Santa Cruz Biotechnology, Santa Cruz, CA, USA) for 1.0 h at 4 °C to eliminate nonspecific binding to the agarose beads, and then immunoprecipitated using anti-Flag beads. The beads were washed three times with NP-40 buffer and then boiled in protein loading buffer before the SDS-PAGE and western blotting. For GST pull-down assays, GST and GST-NPM1 were expressed in *Escherichia coli* BL21 cells and purified using Pierce glutathione agarose beads (21516; Thermo, Rockford, IL, USA). To prepare the bait proteins, purified GST as well as the GST-NPM1 were immobilized on glutathione agarose beads (16100; Thermo Fisher Scientific, USA), while lysates of Flag-PCV3-Cap transfected 293T cells were used as the prey protein. Equal amount of purified GST, GST-NPM1 and Flag-PCV3-Cap proteins were added together and incubated overnight at 4 °C and washed three times with NP-40 buffer. After that, it was boiled in protein loading buffer and subjected SDS-PAGE and western blotting using mouse mAbs against Flag and GST. For the conventional GST pull-down assays, HEK293T cells were transfected with the indicated plasmids for 48 h. Cells were lysed in NP-40 buffer containing a protease inhibitor cocktail. Subsequently, the cell lysates were precleared

and immobilized on glutathione agarose beads and incubated for 4.0 h at 4 °C. The beads were then washed three times with NP-40 buffer and boiled in protein loading buffer. Finally, the protein samples were separated and subjected to western blotting using mouse mAbs against Flag and GST or rabbit pAb against GFP.

***NPM1* knockdown by lentivirus-mediated RNA interference (RNAi)**

NPM1 knockdown was performed as previously described [46] with slight modifications. Briefly, the effective sh*NPM1* (targeting sequence GGATGAGTTGCA CATTGTATT) or shCON (targeting sequence CCGATC GCTACAAATAAG) RNA was co-transfected with the helper lentiviral packaging plasmids psPAX2.0 (12260; Addgene) and pMD2.G (12259; Addgene) into HEK293T cells to produce a lentivirus containing sh*NPM1* or shCON respectively. PK-15 cells were then infected with the resultant lentiviruses and cultured in complete medium for another 24 h and then subjected to puromycin (5 µg/mL; A1113803; Invitrogen) treatment for 7 days







to obtain *NPM1*-silenced cells. Finally, the nuclear DNA content of shRNA-control and *NPM1*-silenced cells were measured using propidium iodide (PI) staining and fluorescence-activated cell sorting (FACS) analysis to analyze cell cycle.

Cell cycle analysis

Flow cytometry was used to analyze cell cycle. In brief, the adherent shCON and sh*NPM1* cells were harvested by trypsin digestion, washed with PBS, and pelleted by centrifugation. Afterwards, the cells were fixed in 70% ethanol overnight at 4 °C and then stained for nuclear DNA content with 50 µg/mL FxCycleTMPI/RNase Staining Solution (2149269; Thermo Fisher Scientific, USA) at room temperature for 30 min. Determination of PI-stained cells was performed using fluorescence-activated cell sorting (FACS) on BD FACSverse. At least 100,000 cells were counted for each sample. Data were analyzed using FlowJo software.

(See figure on next page.)

Figure 1 *NPM1* knockdown abolished the nucleolar localization of PCV3 Cap. **A** *NPM1*-knocked down and shCON-transfected PK-15 cells were respectively transfected with mCherry-PCV3-Cap for 24 h, and cells were fixed and then subjected to confocal microscopy analysis. GFP signals indicate silenced PK-15 cells. Nuclei were stained with 4', 6'-diamidino-2-phenylindole (DAPI). Scale bar, 10 µm. **B**, **C** Images of PK-15 cell line expressing shRNA mediated by lentivirus (**B**), and the knockdown efficiency of *NPM1* in shCON-transfected and *NPM1*-silenced PK-15 cells (**C**). The cell lysates were subjected to immunoblotting using the indicated antibodies. **D** The nuclear DNA content of shRNA-control and *NPM1*-silenced cells were measured using PI staining and FACS analysis to analyze the cell cycle. **E** The histogram was analyzed quantitatively by a curve-fitting program to determine the percentage of cells in the S phase of both cells. Data are presented as means ± SD of three independent biological experiments. *ns* not significant.



Statistical analysis

All results are presented as means \pm standard deviations (SD). Statistical analysis was performed using Student's *t*-test. *p*-values of <0.05 were considered significant.

Results

The NPM1 is essential for the nucleolar localization of PCV3 Cap

To investigate the role of NPM1 in the nucleolar localization of PCV3 viral protein Cap, the *NPM1*-silenced and control PK-15 cells were transfected with mCherry-PCV3-Cap for 24 h. Confocal imaging showed that PCV3 Cap exhibited apparent nucleolar localization in shCON-transfected cells, and distributed to the nucleoplasm in *NPM1*-silenced cells (Figure 1A). The knockdown level of NPM1 in shCON-transfected and *NPM1*-silenced PK-15 cells was apparent as shown in Figure 1B, C. To rule out the effect of cell cycle (S phase) on the nucleolar localization of PCV3 Cap, the nuclear DNA content of shRNA-control and *NPM1*-silenced cells was measured using propidium iodide (PI) staining and fluorescence-activated cell sorting (FACS) analysis. The results indicated that the G0/G1, S, and G2/M phases in *NPM1*-silenced cells were nearly same as


compared to that in the shRNA control cells (Figure 1D). The histogram was further analyzed quantitatively by a curve-fitting program to determine the percentage of cells in the S phase. The percentage was similar in both group of cells (Figure 1E). Taken together, the results indicate that the NPM1 is responsible for supporting the nucleolar localization of PCV3 Cap.

PCV3 Cap colocalizes and interacts with nucleolar protein NPM1

Given that silencing NPM1 expression could repress the nucleolar localization of PCV3 Cap, we investigated the colocalization of PCV3 Cap with NPM1. The results showed that PCV3 Cap colocalized with NPM1 in nucleoli of both HEK293T and PK-15 cells cotransfected with GFP-PCV3-Cap and mCherry-NPM1 (Figure 2A, B). To confirm the interaction between PCV3 Cap and NPM1, lysates of PK-15 cells transfected with Flag-PCV3-Cap were immunoprecipitated with anti-Flag beads and probed for the presence of NPM1 protein with anti-NPM1 monoclonal antibody (mAb), indicating that PCV3 Cap indeed interacted with endogenous NPM1 protein (Figure 2C). Consistently,

(See figure on next page.)

Figure 2 PCV3 Cap colocalizes and interacts with NPM1. **A, B** NPM1 colocalization with PCV3 Cap in transfected cells. HEK293T (**A**) and PK-15 (**B**) cells were cotransfected with GFP-PCV3-Cap and mCherry-NPM1 for 24 h, and cells were fixed and then subjected to confocal microscopy analysis. Nuclei were stained with 4', 6'-diamidino-2-phenylindole (DAPI) (**A, B**). Scale bar, 10 μ m. **C.** PK-15 cells were transfected with empty vector and Flag-PCV3-Cap for 48 h. **D, E** HEK293T cells were cotransfected with Myc-PCV3-Cap and Flag-NPM1 for 48 h. The cell lysates were immunoprecipitated with Flag beads (**C, D**) or anti-Myc purified IgG (**E**). **F** Lysates of Flag-PCV3-Cap transfected 293T cells mixed with the GST, GST-NPM1 proteins were pulled down with glutathione *S*-transferase (GST) beads, and then subjected to GST pull-down assays, followed by immunoblotting using corresponding antibodies.



Flag-tagged NPM1 could interact with Myc-tagged PCV3 Cap in 293T cells (Figure 2D, E). Furthermore, glutathione *S*-transferase (GST) pull-down assays and immunoblotting revealed a direct interaction of NPM1 with PCV3 Cap (Figure 2F).

Taken together, these data indicated that PCV3 Cap interacts directly with NPM1.

The N-terminal residues 1–34 of PCV3 Cap is a nucleolar localization signal and critical for binding to NPM1








To identify the domain in PCV3 Cap required for interaction with NPM1, we constructed two truncated mutants of PCV3 Cap (Figure 3A). Co-IP and GST pull-down assays showed that amino acids (aa) 1–34 (M1), as well as the full-length PCV3 Cap (WT) could interact with NPM1, whereas aa 35–214 (M2) of Cap could not able to interact with NPM1 (Figure 3B, C). These data showed that the N-terminal residues 1–34 of PCV3 Cap that

form an NLS (34) are crucial for Cap binding to NPM1. Further Co-IP experiments showed that the NLSs within capsid protein of porcine circovirus type 1, 2, 3, 4 and circoviruses from terrestrial, aquatic and avian species, including pigs, canaries, canines, minks, dragonflies, pigeons, ducks, bats, geese, and parrots, were also indispensable for the binding to NPM1 (Figure 3D, E), indicating that the interaction is highly conserved in evolution.

As shown in Figure 2A, B, PCV3 Cap is located in nucleolus in the transfected-cells. To investigate whether PCV3 Cap has a nucleolus localization signal (NoLS), we used online tools (Nucleolar location sequence Detector, <http://www.compbio.dundee.ac.uk/www-nod/index.jsp>; and NLS Mapper, http://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_form.cgi). Surprisingly, a potential NoLS was predicted at the N-terminus of PCV3 Cap. To validate this, plasmids PCV3 Cap-WT, -M1, and -M2, were respectively cotransfected into 293T cells along with

(See figure on next page.)

Figure 3 Binding domain identification of PCV3 Cap with NPM1. **A** Schematic representation of the truncation mutants of PCV3 Cap used in this study. **B, C** The N-terminal nuclear localization signal of PCV3 Cap interacted with NPM1. HEK293T cells were cotransfected with plasmids encoding full-length PCV3 Cap or truncation mutants fused with a GFP-, or Flag-gst tag, along with Flag-NPM1; cell lysates were subjected to immunoprecipitation or GST pull-down and immunoblotting using the indicated antibodies. **D, E** The nuclear localization signals (NLSs) within capsid protein of porcine circovirus type 1, 2, 3, 4 and circoviruses from other species were responsible for the binding to NPM1. HEK293T cells were cotransfected with plasmids encoding NLSs of PCV1, 2, 3, 4 (**D**) and circoviruses from terrestrial, aquatic and avian species, including pigs, canaries, canines, minks, dragonflies, pigeons, ducks, bats, geese, and parrots (**E**), along with Flag-NPM1; cell lysates were subjected to immunoprecipitation and immunoblotting using the indicated antibodies. **F** Identification of the nucleolar localization signal in PCV3 Cap. HEK293T cells were cotransfected with GFP-NCL and mCherry-PCV3-Cap-WT, -M1, or -M2 for 24 h. The resultant cells were fixed, stained with DAPI and subjected to confocal microscopy analysis. Scale bar, 10 μ m. **G** Amino acid sequence alignment of the nucleolar localization signal from different PCV3 genotypes.



GFP-nucleolin (NCL). Confocal imaging showed that only PCV3 Cap-WT and -M2 colocalized with NCL (Figure 3F). These data confirmed that the N-terminal residues 1–34 of PCV3 Cap is a NoLS. Aligning the amino acid sequences of NLS from different genotypes (PCV3a, PCV3b) of PCV3 using Jalview software revealed that NLSs of PCV3 Cap are highly conserved (Figure 3G).

Ser48 in OligoD of NPM1 is critical for binding to PCV3 Cap



Various functional domains have been identified within NPM1, including an N-terminal oligomerization domain (OligoD) having chaperone activity, the C-terminal nucleic acid binding domain (NBD), and two central acidic domains for histone binding (HistonD) [42]. To characterize the domain responsible for binding of NPM1 to PCV3 Cap, we constructed a series of NPM1-truncation mutants fused with GFP: OligoD (aa 1–117),

HistonD (aa 118–188), NBD (aa 189–294), OligoD-HistonD (aa 1–188), HistonD-NBD (aa 118–294), OligoD-NBD (aa 1–117 + 189–294) (Figure 4A) and mapped the domains of NPM1 necessary for interaction with PCV3 Cap. Co-IP and GST pull-down assays showed that the constructs OligoD, OligoD-HistonD and OligoD-NBD interacted with PCV3 Cap, whereas HistonD, NBD and HistonD-NBD did not (Figure 4B, C), indicating that the OligoD of NPM1 is required for binding to PCV3 Cap. Binding pattern of the OligoD of NPM1 with PCV3 Cap NLS in silico using PyMOL software also showed that the N-terminal OligoD of NPM1 bound well to the NLS of PCV3 Cap in a plastic and flexible manner (Figure 4D).

Phosphorylation of the oligomerization domain is essential for the regulation of NPM1 structural polymorphism [47]. To explore if these phosphorylation sites played roles in interaction with PCV3 Cap and to

(See figure on next page.)





Figure 4 Serine-48 of NPM1 is critical for binding to PCV3 Cap. **A** Schematic representation of the OligoD, HistonD, and NBD of NPM1 and their truncation mutants used in this study. **B, C** The OligoD of NPM1 (aa 1–117) interacted with PCV3 Cap. HEK293T cells were cotransfected with plasmid GFP-NPM1-WT or serial GFP-NPM1 truncated mutants D1 to D6, along with Flag-PCV3-Cap and Flag-gst-PCV3-Cap expression vectors. The cell lysates were subjected to immunoprecipitation or GST pull-down and immunoblotting using the indicated antibodies. **D** Prediction of binding pattern of N-terminal OligoD from NPM1 with NoLS of PCV3 Cap using PyMOL software. **E** The N-terminal amino acid sequence of the OligoD of NPM1. Serine (S) or threonine (T) residues are marked in red. **F, G** Mapping the crucial amino acids of OligoD responsible for binding to PCV3 Cap. HEK293T cells were cotransfected with NPM1 or NPM1 mutants (-S48A, -S48E, -S88A, -S88E, -T95A, or -T95D), along with Flag-PCV3-Cap and Flag-gst-PCV3-Cap expression vectors, and the cell lysates were subjected to immunoprecipitation or GST pull-down and immunoblotting using the indicated antibodies.



E

```

MEDSMDMDMSPLRPQNYLFGCELKADKDYHF
KVDNDENEHQSLRITVS48LGAGAKDELHIVEA
EAMNYEGSPIKVTLATLKMSVQPTVS88LGGFEI
T95PPVVLRLKCGSGPVHISGQHLV
    
```



identify the key amino acid residues in the OligoD critical for the interaction, we constructed a series of putative phosphorylation site mutants of GFP-NPM1, including GFP-NPM1-S(serine)48A(alanine), -S48E(glutamic acid), -S88A, -S88E, -T(threonine)95A, and -T95D(aspartic acid) (Figure 4E), and subjected to Co-IP and GST pull-down assays with Flag-PCV3-Cap and Flag-gst-PCV3-Cap. The results showed that the non-phosphorylated S48A, S88A and T95A, and mimic-phosphorylated S88E and T95D bound with Flag-PCV3-Cap and Flag-gst-PCV3-Cap nearly as well as the WT. However, the mimic-phosphorylated S48E mutant failed to binding to PCV3 Cap (Figure 4F, G). In summary, these findings indicate that the unphosphorylated serine-48 of NPM1 is responsible for the interaction between NPM1 and PCV3 Cap.

Charge property of the 48th amino acid within NPM1 determines its nucleolar localization and NPM1/Cap interaction


The non-phosphorylated S48A, but not the mimic-phosphorylated S48E interacted with PCV3 Cap, which

compelled us to further characterize the 48th amino acid. Due to the different charge properties of S48A and S48E, we investigated if charge property of 48th amino acid within NPM1 is responsible for binding to PCV3 Cap. Overexpression plasmids of NPM1 with different charge property of 48th amino acid, including GFP-NPM1-WT, -S48E, -S48D, -S48A, -S48T, -S48K(lysine), and -S48R(arginine) were constructed and proceeded to Co-IP and GST pull-down assays with Flag-PCV3-Cap and Flag-gst-PCV3-Cap. The results showed that NPM1 harboring a neutral amino acid in position 48 retained interaction with PCV3 Cap, whereas switching the neutral amino acid to an acidic or basic one abolished this property (Figure 5A, B). Taken together, these data demonstrated that charge property of 48th amino acid in OligoD of NPM1 is critical for interaction with PCV3 Cap.

To further explore whether charge property of 48th amino acid within NPM1 made a difference in its nucleolar localization, we employed GFP fused NPM1 48th amino acid mutants (GFP-NPM1-WT, -S48E, -S48D, -S48A, -S48T, -S48K, -S48R) and nucleolus-located protein mCherry-NCL. Immunofluorescence assays showed that mutants S48A and S48T exhibited nucleolar localization and colocalized with NCL as same as WT, whereas

(See figure on next page.)

Figure 5 Charge property of serine-48 within NPM1 determines its subcellular localization and interaction with PCV3 Cap. A, B Charge property of serine-48 of NPM1 is responsible for binding to PCV3 Cap. HEK293T cells were cotransfected with NPM1 or NPM1 mutants (-S48E, -S48D, -S48A, -S48T, -S48K, or -S48R), along with Flag-PCV3-Cap and Flag-gst-PCV3-Cap expression vectors, and the cell lysates were subjected to immunoprecipitation or GST pull-down and immunoblotting using the indicated antibodies. **C** Nucleolin colocalization with NPM1 and various NPM1 mutants in transfected cells. HEK293T cells were cotransfected with GFP-NPM1 or NPM1 mutants (-S48E, -S48D, -S48A, -S48T, -S48K, or -S48R), along with mCherry-NCL for 24 h, and cells were fixed and then subjected to confocal microscopy analysis. Nuclei were stained with 4', 6'-diamidino-2-phenylindole (DAPI). Scale bar, 10 μ m.



the S48E, S48D, S48K, and S48R variants distributed into nucleoplasm uniformly, and failed to colocalize with NCL (Figure 5C), indicating that charge property of 48th amino acid within NPM1 is responsible for its nucleolar localization. Overall, we concluded that charge property of 48th amino acid within NPM1 plays crucial role in its nucleolar localization and interaction with PCV3 Cap.

Discussion

NPM1 is a constitutively expressed protein which interacts with several host and viral proteins and participates in several steps of viral life cycle such as nuclear entry, viral genome replication, transcription, capsid assembly and egress [48–51]. Different domains of NPM1 enable this protein to shuttle between cytoplasm, nucleoplasm and nucleolus, but still, it primarily resides into the nucleolus. Under certain conditions, NPM1 may be transported to cytoplasm to perform specific functions, mediated by binding of chromosome region maintenance 1 (CRM1) to the nuclear export signal (NES) motifs or some other factors. However, in normal physiologic conditions, nuclear import predominates over export, and NPM1 is particularly resides in nucleolus to perform its functions [52].

As a multifunctional nucleolar phosphoprotein, NPM1 interacts with various viral proteins. Two stretches of negatively charged amino acids in the primary structure of NPM1 are particularly important because these motifs may potentially bind with positively-charged amino acid stretches in viral proteins, such as arginine-rich motif (ARM) and NLS [51]. Viral positively charged amino acid sequences always play important regulatory roles in different stages of the viral replication cycle, such as nucleocytoplasmic transport, viral genome replication and transcription, and viral particle assembly [50, 51]. However, whether PCV3 Cap, harboring a conserved NoLS, could also bind to NPM1 protein has not been reported. Our results showed that knockdown of *NPM1* suppressed the nucleolar localization of PCV3 Cap (Figure 1). PCV3 Cap colocalizes and interacts with NPM1 (Figure 2), and the NLSs within capsid protein of circoviruses from various species were all indispensable for binding to NPM1 (Figure 3), indicating that the interactions between NPM1 and circoviruses Cap NLSs are highly conserved during the evolution. Moreover, we verified that charge property of serine-48 within N-terminal oligomerization domain of NPM1 is vital for its subcellular localization and interaction with PCV3 Cap (Figures 4 and 5). PCV3 Cap may enter the nucleolus at the beginning of infection to support viral transcription, or alter the cell cycle with retention of the S phase and expression of related host proteins, or favor the synthetic replication of viral genome DNA as well [53]. During

the infection cycle, the viral proteins Cap, Rep and Rep' accumulate in the nucleoplasm, implying that DNA replication and encapsidation of the circular closed single-stranded DNA (ssDNA) occur in the nucleus and not in cytoplasmic compartments [53, 54]. The capsid assembly in the nucleus is a prerequisite for translocation to the cytoplasm and subsequent release of mature virions during late infection [55]. It will be interesting to investigate whether PCV3 Cap interacts with cellular factors regulating transcription.

Intriguingly, when Ser48 residue of NPM1 was mutated to neutral amino acids (Ala, Thr et al.), it retained nucleolar localization. But, when it was mutated to acidic amino acids (Glu, Asp) or alkaline amino acids (Lys, Arg, His), it did no longer localize in nucleolus (Figure 5). Phosphorylation of the oligomerization domain is responsible for the thermodynamic stability and spatial conformation of NPM1 and its nucleolar localization [47], we thus speculate that different charge properties of NPM1 affect its subcellular localization and interaction with other proteins. Viral exploitation of nucleolar function may lead to alterations in host cell transcription and translation, or to hijacking of nucleo–cytoplasmic transport and capsid assembly to favor viral replication [56]. Considering that the nucleolar localization signal (NoLS) at the N-terminus of PCV3 Cap functions as an NPM1 binding site and mediates its transport to the nucleolar compartment, we hypothesize that NPM1 promotes intracellular nucleolar trafficking of PCV3 Cap. The NPM1 targets PCV3 Cap to the nucleolus via interaction with its NoLS and facilitates encapsidation viral genome DNA, and assembly of viral particles and hence it is essential for viral replication inside the nucleus of infected cells, which is consistent with the previous study [57]. The NLS of PCV2 Cap can interact with the nuclear membrane receptor (gC1qR) to regulate DNA [58]. Likely, this suggests that the NLS of PCV3 Cap may be also involved in DNA binding. NPM1 may act as an adaptor for transport PCV3 Cap into nucleolus and facilitate its nucleolar localization. Further study is needed to verify whether the nuclear entry of PCV3 virion is also mediated by NPM1.

In summary, our results showed that knockdown of *NPM1* impaired the nucleolar localization of PCV3 Cap and charge property of 48th amino acid within NPM1 plays critical role in its nucleolar localization and interaction with PCV3 Cap. Taken together, this study broadens our understanding of nucleolar entry of PCV3 Cap and mechanism of PCV3 replication, which will be helpful to identify novel potential targets for therapeutic and prophylactic intervention of PCV3 infection.

Abbreviations

NPM1: nucleolar phosphoprotein nucleophosmin-1; PCV: porcine circovirus; PCV1: porcine circovirus type 1; PCV2: porcine circovirus type 2; PCV3: porcine circovirus type 3; Cap: capsid protein; PCVAD: porcine circovirus-associated diseases; NoLS: nucleolar localization signal; NGS: next generation sequence; PDNS: porcine dermatitis and nephropathy syndrome; ORF: open reading frame; NLS: nuclear localization signal; STAT2: signal transducer and activator of transcription 2; HSV-1: herpes simplex virus type 1; AdV-2: adenovirus 2; IAV: influenza A virus; HCV: hepatitis C virus; NPC: nuclear pore complex; HSP70: heat shock protein 70; HDV: hepatitis delta virus; NCL: nucleolin; OligoD: oligomerization domain; HistonD: histone-binding domain; NBD: nucleic acid-binding domain; HIV-1: Human immunodeficiency virus type 1; MEM: minimal essential medium; DMEM: Dulbecco's modified Eagle medium; FBS: fetal bovine serum; pAb: polyclonal antibody; mAb: monoclonal antibody; EGFP: enhanced green fluorescent protein; GST: glutathione S-transferase; HRP: horseradish peroxidase; PCR: polymerase chain reaction; CanaryCV: Canary circovirus; MinkCV: mink circovirus; DragonflyCV: dragonfly cyclovirus; CoCV: Columbid circovirus; BtCV: bat associated cyclovirus; DuCV: duck circovirus; GoCV: goose circovirus; BFDV: Beak and feather disease virus; PFA: paraformaldehyde; DAPI: 4', 6'-Diamidino-2-phenylindole; SDS: sodium dodecyl sulfate; PAGE: polyacrylamide gel electrophoresis; PBS: phosphate-buffered saline; Co-IP: co-immunoprecipitation; RNAi: RNA interference; PI: propidium iodide; FACS: fluorescence-activated cell sorting; aa: amino acid; CRM1: chromosome region maintenance 1; NES: nuclear export signal; ARM: arginine-rich motif; ssDNA: single-stranded DNA; gC1qR: globular heads of complement component C1q receptor.

Acknowledgements

We thank Ms. Yunqin Li (Bio-Ultrastructure Analysis Lab, Center of Agrobiological and Environmental Sciences, Zhejiang University) for technical help with confocal microscopy observations.

Authors' contributions

JwZ and JyZ designed the experiments. JwZ interpreted the data and wrote the manuscript. JwZ performed the experiments with assistance and advice from JL, HmL, YZ, WrD, YJ and YY. JyG and JyZ revised the paper. All authors read and approved the final manuscript.

Funding

This work was supported by the Key Research & Development Program of Zhejiang Province (Grant No. 2020C02011).

Availability of data and materials

All data and materials generated for this study are included in the article.

Ethics approval and consent to participate

Not applicable.

Consent for publication

The authors declare that they agree to publish.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ MOA Key Laboratory of Animal Virology, Center of Veterinary Sciences, Zhejiang University, 866 Yuhangtang Road, Hangzhou, Zhejiang 310058, PR China. ² Collaborative Innovation Center and State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, First Affiliated Hospital, Zhejiang University, Hangzhou, PR China.

Received: 28 August 2020 Accepted: 3 December 2020

Published online: 07 January 2021

References

- Nauwynck HJ, Sanchez R, Meerts P, Lefebvre DJ, Saha D, Huang L, Misinz G (2012) Cell tropism and entry of porcine circovirus 2. *Virus Res* 164(1–2):43–45
- Lorincz M, Csagola A, Farkas SL, Szekeley C, Tuboly T (2011) First detection and analysis of a fish circovirus. *J Gen Virol* 92(Pt 8):1817–1821
- Schoemaker NJ, Dorrestein GM, Latimer KS, Lumeij JT, Kik MJ, van der Hage MH, Campagnoli RP (2000) Severe leukopenia and liver necrosis in young African grey parrots (*Psittacus erithacus*) infected with psittacine circovirus. *Avian Dis* 44(2):470–478
- Phenix KV, Weston JH, Ypelaar I, Lavazza A, Smyth JA, Todd D, Wilcox GE, Raidal SR (2001) Nucleotide sequence analysis of a novel circovirus of canaries and its relationship to other members of the genus *Circovirus* of the family *Circoviridae*. *J Gen Virol* 82(Pt 11):2805–2809
- Todd D, Weston JH, Soike D, Smyth JA (2001) Genome sequence determinations and analyses of novel circoviruses from goose and pigeon. *Virology* 286(2):354–362
- Li L, McGraw S, Zhu K, Leutenegger CM, Marks SL, Kubiski S, Gaffney P, Jr, Dela Cruz FN, Wang C, Delwart E et al (2013) Circovirus in tissues of dogs with vasculitis and hemorrhage. *Emerg Infect Dis* 19(4):534–541
- Tischer I, Gelderblom H, Vettermann W, Koch MA (1982) A very small porcine virus with circular single-stranded-DNA. *Nature* 295(5844):64–66
- Lian H, Liu Y, Li N, Wang Y, Zhang S, Hu R (2014) Novel circovirus from mink, China. *Emerg Infect Dis* 20(9):1548–1550
- Virus taxonomy, 6th report of the International Committee on Taxonomy of Viruses. *Arch Virol Suppl* 1995, 10:1–586.
- Ellis J (2014) Porcine circovirus: a historical perspective. *Vet Pathol* 51(2):315–327
- Meng XJ (2013) Porcine circovirus type 2 (PCV2): pathogenesis and interaction with the immune system. *Annu Rev Anim Biosci* 1:43–64
- Denner J, Mankertz A (2017) Porcine circoviruses and xenotransplantation. *Viruses* 9(4):83
- Huang L, Lu Y, Wei Y, Guo L, Liu C (2012) Identification of three new type-specific antigen epitopes in the capsid protein of porcine circovirus type 1. *Arch Virol* 157(7):1339–1344
- Li LL, Kapoor A, Slikas B, Bamidele OS, Wang CL, Shaikat S, Masroor MA, Wilson ML, Ndjanga JBN, Peeters M et al (2010) Multiple diverse circoviruses infect farm animals and are commonly found in human and chimpanzee feces. *J Virol* 84(4):1674–1682
- Phan TG, Giannitti F, Rossow S, Marthaler D, Knutson TP, Li L, Deng X, Resende T, Vannucci F, Delwart E (2016) Detection of a novel circovirus PCV3 in pigs with cardiac and multi-systemic inflammation. *Virol J* 13(1):184
- Palinski R, Pineyro P, Shang PC, Yuan FF, Guo R, Fang Y, Byers E, Hause BM (2017) A novel porcine circovirus distantly related to known circoviruses is associated with porcine dermatitis and nephropathy syndrome and reproductive failure. *J Virol* 91(1):e01879
- Wen S, Sun W, Li Z, Zhuang X, Zhao G, Xie C, Zheng M, Jing J, Xiao P, Wang M et al (2018) The detection of porcine circovirus 3 in Guangxi, China. *Transbound Emerg Dis* 65(1):27–31
- Tochetto C, Lima DA, Varela APM, Loiko MR, Paim WP, Scheffer CM, Herpich JJ, Cerva C, Schmitz C, Cibulski SP et al (2018) Full-genome sequence of porcine circovirus type 3 recovered from serum of sows with stillbirths in Brazil. *Transbound Emerg Dis* 65(1):5–9
- Kedkovid R, Woonwong Y, Arunorat J, Sirisereewan C, Sangpratum N, Lumyai M, Kesdangakonwut S, Teankum K, Jittimanee S, Thanawongnuwech R (2018) Porcine circovirus type 3 (PCV3) infection in grower pigs from a Thai farm suffering from porcine respiratory disease complex (PRDC). *Vet Microbiol* 215:71–76
- Stadejek T, Wozniak A, Milek D, Biernacka K (2017) First detection of porcine circovirus type 3 on commercial pig farms in Poland. *Transbound Emerg Dis* 64(5):1350–1353
- Kwon T, Yoo SJ, Park CK, Lyoo YS (2017) Prevalence of novel porcine circovirus 3 in Korean pig populations. *Vet Microbiol* 207:178–180
- Faccini S, Barbieri I, Gilioli A, Sala G, Gibelli LR, Moreno A, Sacchi C, Rosignoli C, Franzini G, Nigrelli A (2017) Detection and genetic characterization of Porcine circovirus type 3 in Italy. *Transbound Emerg Dis* 64(6):1661–1664
- Fux R, Sockler C, Link EK, Renken C, Krejci R, Sutter G, Ritzmann M, Eddicks M (2018) Full genome characterization of porcine circovirus type 3 isolates reveals the existence of two distinct groups of virus strains. *Virol J* 15(1):25
- Franzo G, Legnardi M, Hjulsager CK, Klaumann F, Larsen LE, Segales J, Drigo M (2018) Full-genome sequencing of porcine circovirus 3 field strains from Denmark, Italy and Spain demonstrates a high

- within-Europe genetic heterogeneity. *Transbound Emerg Dis* 65(3):602–606
25. Li G, He W, Zhu H, Bi Y, Wang R, Xing G, Zhang C, Zhou J, Yuen KY, Gao GF et al (2018) Origin, genetic diversity, and evolutionary dynamics of novel porcine circovirus 3. *Adv Sci* 5(9):1800275
 26. Fu X, Fang B, Ma J, Liu Y, Bu D, Zhou P, Wang H, Jia K, Zhang G (2018) Insights into the epidemic characteristics and evolutionary history of the novel porcine circovirus type 3 in southern China. *Transbound Emerg Dis* 65(2):e296–e303
 27. Nawagitgul P, Morozov I, Bolin SR, Harms PA, Sorden SD, Paul PS (2000) Open reading frame 2 of porcine circovirus type 2 encodes a major capsid protein. *J Gen Virol* 81(Pt 9):2281–2287
 28. Blanchard P, Mahe D, Cariolet R, Keranflec'h A, Baudouard MA, Cordioli P, Albina E, Jestin A (2003) Protection of swine against post-weaning multisystemic wasting syndrome (PMWS) by porcine circovirus type 2 (PCV2) proteins. *Vaccine* 21(31):4565–4575
 29. Fenaux M, Opriessnig T, Halbur PG, Elvinger F, Meng XJ (2004) Two amino acid mutations in the capsid protein of type 2 porcine circovirus (PCV2) enhanced PCV2 replication in vitro and attenuated the virus in vivo. *J Virol* 78(24):13440–13446
 30. Timmusk S, Fossum C, Berg M (2006) Porcine circovirus type 2 replicase binds the capsid protein and an intermediate filament-like protein. *J Gen Virol* 87(Pt 11):3215–3223
 31. Wiederkehr DD, Sydlar T, Buergi E, Haessig M, Zimmermann D, Pospischil A, Brugnera E, Sidler X (2009) A new emerging genotype subgroup within PCV-2b dominates the PMWS epizooty in Switzerland. *Vet Microbiol* 136(1–2):27–35
 32. Cao J, Lin C, Wang H, Wang L, Zhou N, Jin Y, Liao M, Zhou J (2015) Circovirus transport proceeds via direct interaction of the cytoplasmic dynein IC1 subunit with the viral capsid protein. *J Virol* 89(5):2777–2791
 33. Fermin G, Tennant P (2018) Host-virus interactions: battles between viruses and their hosts. In: *Viruses: molecular biology, host interactions, and applications to biotechnology*. Academic Press, p 245–271. <https://doi.org/10.1016/B978-0-12-811257-1.00010-3>
 34. Wang HJ, Zhang KL, Lin C, Zhou JW, Jin YL, Dong WR, Gu JY, Zhou JY (2019) Conformational changes and nuclear entry of porcine circovirus without disassembly. *J Virol* 93(20):824
 35. Mou C, Wang M, Pan S, Chen Z (2019) Identification of nuclear localization signals in the ORF2 protein of porcine circovirus type 3. *Viruses* 11(12):1086
 36. Shen HQ, Liu XH, Zhang PF, Wang SY, Liu YL, Zhang LY, Song CX (2020) Porcine circovirus 3 Cap inhibits type I interferon signaling through interaction with STAT2. *Virus Res* 275:197804
 37. Hennig T, O'Hare P (2015) Viruses and the nuclear envelope. *Curr Opin Cell Biol* 34:113–121
 38. Sapphire AC, Guan T, Schirmer EC, Nemerow GR, Gerace L (2000) Nuclear import of adenovirus DNA in vitro involves the nuclear protein import pathway and hsc70. *J Biol Chem* 275(6):4298–4304
 39. Trotman LC, Mosberger N, Fornerod M, Stidwill RP, Greber UF (2001) Import of adenovirus DNA involves the nuclear pore complex receptor CAN/Nup214 and histone H1. *Nat Cell Biol* 3(12):1092–1100
 40. Lee CH, Chang SC, Chen CJ, Chang MF (1998) The nucleolin binding activity of hepatitis delta antigen is associated with nucleolus targeting. *J Biol Chem* 273(13):7650–7656
 41. Yun JP, Chew EC, Liew CT, Chan JYH, Jin ML, Ding MX, Fai YH, Li HKR, Liang XM, Wu QL (2003) Nucleophosmin/B23 is a proliferate shuttle protein associated with nuclear matrix. *J Cell Biochem* 90(6):1140–1148
 42. Okuwaki M (2008) The structure and functions of NPM1/Nucleophosmin/B23, a multifunctional nucleolar acidic protein. *J Biochem* 143(4):441–448
 43. Lindstrom MS (2011) NPM1/B23: a multifunctional chaperone in ribosome biogenesis and chromatin remodeling. *Biochem Res Int* 2011:195209
 44. Fankhauser C, Izaurralde E, Adachi Y, Wingfield P, Laemmli UK (1991) Specific complex of human immunodeficiency virus type 1 rev and nucleolar B23 proteins: dissociation by the Rev response element. *Mol Cell Biol* 11(5):2567–2575
 45. Samad MA, Okuwaki M, Haruki H, Nagata K (2007) Physical and functional interaction between a nucleolar protein nucleophosmin/B23 and adenovirus basic core proteins. *FEBS Lett* 581(17):3283–3288
 46. Lin C, Gu J, Wang H, Zhou J, Li J, Wang S, Jin Y, Liu C, Liu J, Yang H et al (2018) Caspase-dependent apoptosis induction via viral protein ORF4 of porcine circovirus 2 binding to mitochondrial adenine nucleotide translocase 3. *J Virol* 92(10):e00238
 47. Mitrea DM, Grace CR, Buljan M, Yun MK, Pytel NJ, Satumba J, Nourse A, Park CG, Madan Babu M, White SW et al (2014) Structural polymorphism in the N-terminal oligomerization domain of NPM1. *Proc Natl Acad Sci USA* 111(12):4466–4471
 48. Jeong H, Cho MH, Park SG, Jung G (2014) Interaction between nucleophosmin and HBV core protein increases HBV capsid assembly. *FEBS Lett* 588(6):851–858
 49. Liu CD, Chen YL, Min YL, Zhao B, Cheng CP, Kang MS, Chiu SJ, Kieff E, Peng CW (2012) The nuclear chaperone nucleophosmin escorts an Epstein-Barr Virus nuclear antigen to establish transcriptional cascades for latent infection in human B cells. *PLoS Pathog* 8(12):e1003084
 50. Day PM, Thompson CD, Pang YY, Lowy DR, Schiller JT (2015) Involvement of nucleophosmin (NPM1/B23) in assembly of infectious HPV16 capsids. *Papillomavirus Res* 1:74–89
 51. Nouri K, Moll JM, Milroy LG, Hain A, Dvorsky R, Amin E, Lenders M, Nagel-Steger L, Howe S, Smits SH et al (2015) Biophysical characterization of nucleophosmin interactions with human immunodeficiency virus rev and herpes simplex virus US11. *PLoS ONE* 10(12):e0143634
 52. Szebeni A, Herrera JE, Olson MO (1995) Interaction of nucleolar protein B23 with peptides related to nuclear localization signals. *Biochemistry* 34(25):8037–8042
 53. Finsterbusch T, Steinfeldt T, Caliskan R, Mankertz A (2005) Analysis of the subcellular localization of the proteins Rep, Rep' and Cap of porcine circovirus type 1. *Virology* 343(1):36–46
 54. Liu Q, Tikoo SK, Babiuk LA (2001) Nuclear localization of the ORF2 protein encoded by porcine circovirus type 2. *Virology* 285(1):91–99
 55. Chung AK, Bolin SR (2002) Kinetics of porcine circovirus type 2 replication. *Arch Virol* 147(1):43–58
 56. Finsterbusch T, Steinfeldt T, Doberstein K, Rodner C, Mankertz A (2009) Interaction of the replication proteins and the capsid protein of porcine circovirus type 1 and 2 with host proteins. *Virology* 386(1):122–131
 57. Duan ZQ, Chen J, Xu HX, Zhu J, Li QH, He L, Liu HM, Hu SL, Liu XF (2014) The nucleolar phosphoprotein B23 targets Newcastle disease virus matrix protein to the nucleoli and facilitates viral replication. *Virology* 452:212–222
 58. Fotso GBK, Bernard C, Bigault L, de Boisseson C, Mankertz A, Jestin A, Grasland B (2016) The expression level of gC1qR is down regulated at the early time of infection with porcine circovirus of type 2 (PCV-2) and gC1qR interacts differently with the Cap proteins of porcine circoviruses. *Virus Res* 220:21–32

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.