

ARTICLE

Received 7 Jul 2014 | Accepted 15 Oct 2014 | Published 27 Nov 2014

DOI: 10.1038/ncomms6576

OPEN

# Long non-coding RNA-mediated transcriptional interference of a permease gene confers drug tolerance in fission yeast

Ryan Ard<sup>1</sup>, Pin Tong<sup>1</sup> & Robin C. Allshire<sup>1</sup>

Most long non-coding RNAs (lncRNAs) encoded by eukaryotic genomes remain uncharacterized. Here we focus on a set of intergenic lncRNAs in fission yeast. Deleting one of these lncRNAs exhibited a clear phenotype: drug sensitivity. Detailed analyses of the affected locus revealed that transcription of the *nc-tgp1* lncRNA regulates drug tolerance by repressing the adjacent phosphate-responsive permease gene *transporter for glycerophosphodiester 1* (*tgp1*<sup>+</sup>). We demonstrate that the act of transcribing *nc-tgp1* over the *tgp1*<sup>+</sup> promoter increases nucleosome density, prevents transcription factor access and thus represses *tgp1*<sup>+</sup> without the need for RNA interference or heterochromatin components. We therefore conclude that *tgp1*<sup>+</sup> is regulated by transcriptional interference. Accordingly, decreased *nc-tgp1* transcription permits *tgp1*<sup>+</sup> expression upon phosphate starvation. Furthermore, *nc-tgp1* loss induces *tgp1*<sup>+</sup> even in repressive conditions. Notably, drug sensitivity results directly from *tgp1*<sup>+</sup> expression in the absence of the *nc-tgp1* RNA. Thus, transcription of an lncRNA governs drug tolerance in fission yeast.

<sup>1</sup> Wellcome Trust Centre for Cell Biology and Institute of Cell Biology, School of Biological Sciences, The University of Edinburgh, Max Born Crescent, Edinburgh EH9 3BF, Scotland, UK. Correspondence and requests for materials should be addressed to R.C.A. (email: robin.allshire@ed.ac.uk).

Eukaryotic genomes are pervasively transcribed. Frequently this transcription generates long non-coding RNAs (lncRNAs), which may be transcribed antisense to protein-coding genes, from within introns, or from intergenic regions of the genome. RNA polymerase II (RNAPII) is responsible for generating both messenger RNAs (mRNAs) and lncRNAs<sup>1</sup>. As with mRNAs, many lncRNAs are processed (that is, capped, spliced, polyadenylated), however, in contrast to protein-coding mRNAs, lncRNAs are predominantly nuclear and many are rapidly degraded by the exosome<sup>2</sup>, the major cellular 3' → 5' RNA degradation machinery<sup>3</sup>. Consequently, the majority of lncRNAs exhibit low steady-state levels compared with mRNAs. This instability coupled with their general lack of primary sequence conservation has led to the suggestion that many lncRNAs might simply result from spurious, inconsequential 'transcriptional noise'<sup>4</sup>. Nonetheless, accumulating evidence indicates that an increasing number of lncRNAs act to regulate gene expression<sup>2,5</sup>.

The mere act of lncRNA transcription, including accompanying chromatin modifications and resulting changes in nucleosome density<sup>6</sup>, can have a profound impact on neighbouring gene expression. In the simplest scenario, lncRNA expression can provide an environment that is either suitable or unsuitable for transcription factor binding. For example, cascading lncRNA transcription upstream of the fission yeast *Schizosaccharomyces pombe* *fbp1*<sup>+</sup> gene is required to induce *fbp1*<sup>+</sup> expression following glucose starvation<sup>7</sup>. In addition, in a process termed 'transcriptional interference', serine-mediated repression of the budding yeast *Saccharomyces cerevisiae* *SER3* gene is brought about by lncRNA transcription into the gene promoter, which increases nucleosome density and prevents transcription factor access<sup>8–10</sup>. These examples illustrate the positive and negative influence that lncRNA transcription can exert on gene regulation in response to environmental changes.

lncRNAs can also be processed into smaller regulatory RNAs (for example, short interfering RNA)<sup>11</sup>. In *S. pombe*, lncRNAs transcribed from centromeric outer repeats are processed by Dicer (Dcr1) into short interfering RNAs, which target the Clr4 H3K9 methyltransferase via Ago1 (within the RNA-induced transcriptional silencing complex) to establish repressive heterochromatin through the methylation of lysine 9 on histone H3 (refs 12–15). In addition, lncRNAs may directly associate with and recruit factors that alter chromatin status, in *cis* or in *trans*, silencing genes or behaving as enhancers<sup>16,17</sup>. For example, lncRNAs aid the response of *S. cerevisiae* cells to specific changes in nutrient availability by recruiting chromatin-modifying complexes (for example, histone deacetylases) to dynamically regulate several genes<sup>18–20</sup>. Related mechanisms have since been reported in multicellular eukaryotes<sup>21,22</sup>. Recent analyses also suggest that patches of transient heterochromatin can form under particular conditions at specific euchromatic loci in *S. pombe*<sup>23–25</sup>. This mechanism involves the RNA-binding protein Mmi1, which recruits the RNA-surveillance machinery to specific determinant of selective removal (DSR) motifs in target transcripts, leading to their exosome-mediated degradation<sup>26</sup>. Mmi1 and its associated factor Red1 are reported to also recruit chromatin-modifying activities via nascent mRNA and lncRNA targets to deposit H3K9 methylation (H3K9me2) at these locations<sup>23,25,27,28</sup>. It is therefore evident that lncRNAs employ a variety of mechanisms to regulate gene expression.

Despite rapid advances in lncRNA identification, only a small number have been characterized in detail. A clear challenge in assigning function has been a lack of lncRNA sequence conservation between even the most closely related species<sup>29</sup>. However, the order of genes flanking the transcription units that encode lncRNAs can be preserved through evolution<sup>30</sup> (that is,

synteny) and provides another criterion by which we can identify potential functionally conserved lncRNAs whose primary sequences might have diverged too much so as not to retain detectable homology.

Only a few of the ~500 annotated intergenic lncRNAs in *S. pombe* are conserved at the sequence level in three divergent *Schizosaccharomyces* species, although many retain synteny with flanking genes in at least one other species<sup>31</sup>. We identified eight discrete intergenic lncRNAs that exhibit synteny in at least three of the four *Schizosaccharomyces* species. Deletion of one of these loci (SPNCRNA.1343 or *ncRNA.1343* for short) exhibited a drug-sensitivity phenotype. We demonstrate that *ncRNA.1343* encodes a bidirectional lncRNA promoter and that its deletion causes loss of expression of the divergent unstable transcript *nc-tgp1*. Our analyses reveal that *nc-tgp1* is targeted for Mmi1-directed exosome degradation and is required to repress a downstream phosphate-responsive gene (SPBC1271.09 designated *transporter for glycerophosphodiester 1* (*tgp1*<sup>+</sup>)). However, rather than involving transient heterochromatin formation as a result of targeted RNA degradation, the regulation of *tgp1*<sup>+</sup> by the *nc-tgp1* RNA appears to be mediated by transcriptional interference. Most importantly, tolerance of *S. pombe* to a broad spectrum of compounds relies on the regulation of *tgp1*<sup>+</sup> by *nc-tgp1*.


## Results

**Deletion of SPNCRNA.1343 causes drug hypersensitivity.** The *S. pombe* genome is predicted to encode ~500 intergenic lncRNAs<sup>32</sup>. Although few of these lncRNAs exhibit detectable sequence conservation, ~100 are conserved in synteny with putative lncRNA orthologues in at least one of the three other known *Schizosaccharomyces* species<sup>31</sup>. For example, the functionally characterized telomerase RNA (*ter1*<sup>+</sup>/SPNCRNA.214) is syntenic despite its lack of sequence conservation (see Supplementary Fig. 1a).

To identify other potential functionally conserved lncRNAs, we selected eight lncRNAs, including *ter1*<sup>+</sup> as a control, where surrounding gene order is retained in *S. pombe* and at least two other *Schizosaccharomyces* species. Each lncRNA gene was deleted by replacement with a loxP-flanked *ura4*<sup>+</sup> marker (Supplementary Fig. 1b). Apart from *ter1Δ*, the selected lncRNAs were not essential for normal cell growth (Supplementary Figs 1c and 2). However, since many characterized lncRNAs regulate gene expression in response to environmental changes and stress<sup>33</sup>, we tested the growth of these lncRNA deletion strains in response to the following stresses: temperature, the microtubule destabilizing drug thiabendazole (TBZ), DNA synthesis inhibitor hydroxyurea (HU), ultraviolet-induced DNA damage, H<sub>2</sub>O<sub>2</sub>-induced oxidative stress and caffeine, an inhibitor of cyclic AMP phosphodiesterase. Cells lacking SPNCRNA.1343 (*ncRNA.1343* for short) displayed a phenotype: hypersensitivity to TBZ, HU and caffeine but not to temperature extremities, ultraviolet-irradiation or oxidative stress (Supplementary Fig. 1c and Supplementary Fig. 2).

### Drug sensitivity of 1343Δ cells is caused by *tgp1*<sup>+</sup> induction.

lncRNAs can act in *cis* to regulate the expression of nearby genes<sup>2</sup>. To determine the cause of drug sensitivity in 1343Δ cells we examined the expression of genes flanking the locus by real-time quantitative reverse transcriptase-PCR (RT-qPCR) in wild-type cells, cells with *ncRNA.1343* replaced by loxP-flanked *ura4*<sup>+</sup> marker (1343Δ::*ura4*<sup>+</sup>) and cells with the *ura4*<sup>+</sup> marker subsequently removed (1343Δ; Fig. 1a). SPBC1271.09 transcript levels increased >50-fold in both 1343Δ::*ura4*<sup>+</sup> and 1343Δ cells (Fig. 1b), while the expression of other neighbouring genes was unaltered. SPBC1271.09 encodes a conserved



**Figure 1 | Drug sensitivity following *ncRNA.1343* deletion is due to increased *tgp1*<sup>+</sup> expression.** (a) Schematic representation of genes flanking *ncRNA.1343*. (b) RT-qPCR experiments measured transcript levels for nearby genes in wild-type cells and following replacement of *ncRNA.1343* with *ura4*<sup>+</sup> (*1343Δ::ura4*<sup>+</sup>) or deletion (*1343Δ*). Error bars represent s.e.m. resulting from at least three independent replicates. (c) Northern analysis of *tgp1*<sup>+</sup> transcript levels in wild-type and *1343Δ* cells grown in the presence of phosphate. (d) Serial dilutions of wild-type, *1343Δ::ura4*<sup>+</sup>, *1343Δ* and *tgp1Δ1343Δ* double mutant spotted on non-selective YES medium or in the presence of TBZ (20 μg ml<sup>-1</sup>), HU (10 mM) or caffeine (15 mM), respectively.

glycerophosphodiester membrane transporter (designated as *tgp1*<sup>+</sup>) orthologous to the *S. cerevisiae* permease *GIT1*. As with *S. cerevisiae* *GIT1*, the *tgp1*<sup>+</sup> gene is repressed when cells are grown in the presence of phosphate and induced upon phosphate starvation<sup>34,35</sup>. Northern analysis confirmed that *tgp1*<sup>+</sup> was indeed highly expressed in *1343Δ* cells but not wild-type cells, both grown in the presence of phosphate (repressed condition; Fig. 1c).

To determine whether the drug sensitivity of *1343Δ* cells is a direct result of increased *tgp1*<sup>+</sup> expression, the *tgp1*<sup>+</sup> gene was deleted from *1343Δ* cells (*tgp1Δ1343Δ*). This manipulation restored TBZ, HU and caffeine tolerance to levels comparable with wild-type cells (Fig. 1d). We conclude that increased *tgp1*<sup>+</sup> expression is directly responsible for the drug-sensitivity phenotype of cells lacking *ncRNA.1343*.


**Bidirectional lncRNA promoter upstream of *tgp1*<sup>+</sup>.** Previous RNA-seq analysis indicates that an lncRNA is transcribed in the sense orientation upstream of *tgp1*<sup>+</sup> (refs 27,31). We identified two divergent transcriptional start sites arising within *ncRNA.1343*: one lncRNA transcribed towards the *tgp1*<sup>+</sup> gene (*nc-tgp1*) and the other in the opposite orientation (*nc-1343*; Fig. 2a; Supplementary Fig. 3). *lacZ* reporter assays demonstrate that the bidirectional promoter drives greater levels of transcription in the *nc-tgp1* direction (Supplementary Fig. 3). This finding is consistent with Rpb1 Chromatin Immunoprecipitation (ChIP) analysis showing that RNAPII is enriched over the *nc-tgp1* transcription unit, while much lower RNAPII levels are detected on *nc-1343* (Fig. 2b).

We next examined the regulation of the *nc-1343* and *nc-tgp1* transcripts produced from this bidirectional promoter. A ~0.9 kb transcript for *nc-1343* was readily detected in wild-type cells. The size and levels of the *nc-1343* transcript increased in exosome defective (*rrp6Δ*) cells, but not cells lacking Mmi1 or Red1 (Fig. 2c,d; Supplementary Fig. 4). The lncRNA corresponding to *nc-tgp1* was previously detected in *rrp6Δ* and *red1Δ* cells<sup>27</sup>. We identified a consensus DSR motif for Mmi1 binding at position +820 nt within the *nc-tgp1* transcript and RNA IP (RIP) experiments confirmed a direct interaction between Mmi1 and the *nc-tgp1* RNA (Supplementary Fig. 5). Northern analysis identified that an ~1.9 kb *nc-tgp1* RNA accumulates in *rrp6Δ*, *mmi1Δ* and *red1Δ*, but not in wild-type cells (Fig. 2e,f; Supplementary Fig. 4). Interestingly, a recent study found that the repressive lncRNA transcribed upstream of the phosphate-responsive *pho1*<sup>+</sup> gene in *S. pombe* also contains a DSR motif and is targeted by Mmi1 for exosome-mediated degradation<sup>28</sup>, indicating that a similar regulatory mechanism might control expression of *tgp1*<sup>+</sup> and *pho1*<sup>+</sup>. In sum, both *nc-1343* and *nc-tgp1* transcripts are processed by the exosome, but only *nc-tgp1* is regulated by Mmi1-mediated recruitment of the nuclear exosome.

A moderate increase in *tgp1*<sup>+</sup> transcript levels has previously been reported in cells lacking Mmi1 (ref. 23). In agreement with this, we detected a similar increase (approximated four-fold) in *tgp1*<sup>+</sup> transcript levels in *mmi1Δ* or exosome (*rrp6Δ* or *dis3-5Δ*) mutant cells by RT-qPCR, however, this increase is significantly less than the >50-fold upregulation of *tgp1*<sup>+</sup> observed in *1343Δ* cells (Fig. 2g,h; Supplementary Fig. 4). Moreover, we failed to detect the *tgp1*<sup>+</sup> transcript in *rrp6Δ* or *mmi1Δ* cells by Northern analysis, indicating that *tgp1*<sup>+</sup> is not induced in the absence of these factors. Thus, Mmi1-mediated exosome degradation is not the predominant mechanism involved in *tgp1*<sup>+</sup> regulation.

***tgp1*<sup>+</sup> is repressed by the *nc-tgp1* lncRNA.** The presence of the unstable *nc-tgp1* RNA upstream of *tgp1*<sup>+</sup> suggests that either *nc-tgp1*, *nc-1343* or both regulate *tgp1*<sup>+</sup> expression. To test the involvement of these lncRNAs in *tgp1*<sup>+</sup> regulation, a series of strategic genetic manipulations were performed (Fig. 3a). Truncations of *nc-1343* (that is, *AΔ* and *BΔ*) that retain its 5' end did not result in the drug-sensitivity phenotype presented by *1343Δ* cells (Fig. 3b) and, similarly, did not induce *tgp1*<sup>+</sup> expression (Fig. 3c). This indicates that full-length *nc-1343* is not required for *tgp1*<sup>+</sup> repression. We next tested if *nc-tgp1* is involved in repressing *tgp1*<sup>+</sup>. Our analyses show that transcription of *nc-tgp1* starts within the encoded *ncRNA.1343* transcription unit (Supplementary Fig. 3). Thus, deletion of the entire locus (*1343Δ*) removes the *nc-tgp1* promoter, and the 5' end of its transcript, resulting in the observed loss of *nc-tgp1* expression (Figs 2f and 3c). The *AΔ* and *BΔ* truncations of *nc-1343*, which retain the *nc-tgp1* promoter, do not affect *nc-tgp1* transcription or relieve repression of *tgp1*<sup>+</sup>. In contrast, interruption of the *nc-tgp1* transcription unit by insertion of the *ura4*<sup>+</sup> marker gene (*nc-tgp1::ura4*<sup>+</sup>) prevented *nc-tgp1* transcription, induced *tgp1*<sup>+</sup> expression to levels observed in *1343Δ* levels and increased sensitivity of these cells to TBZ, HU and caffeine (Fig. 3b,c). These analyses demonstrate that it is *nc-tgp1*, not *nc-1343*, that is critical for repressing *tgp1*<sup>+</sup> in the presence of phosphate.

**Phosphate starvation induces *tgp1*<sup>+</sup> by repressing *nc-tgp1*.** Upon phosphate starvation of fission yeast, several genes involved in the phosphate response are induced (including *tgp1*<sup>+</sup> and *pho1*<sup>+</sup>) (ref. 35). To determine how the transcription of *nc-tgp1* is altered in response to phosphate and how it might influence *tgp1*<sup>+</sup> expression we assessed expression in phosphate-rich




**Figure 2 | Two distinct lncRNAs are transcribed from a bidirectional promoter upstream of *tgp1<sup>+</sup>*.** (a) Previously published strand-specific RNA-Seq analysis (Rhind *et al.*,<sup>31</sup>) upstream of SPBC1271.09/*tgp1<sup>+</sup>*, represented as reads per kilobase per million (RPKM). Location of qPCR primer pairs and probes for Northern analysis are shown below. (b) Rbp1 ChIP-qPCR experiments performed in wild-type cells. (c,e,g) Northern analysis of *nc-1343*, *nc-tgp1* and *tgp1<sup>+</sup>* transcript levels in wild-type, *rrp6Δ*, *mmi1Δ* and *1343Δ*, respectively. (d,f,h) RT-qPCR experiments measured *nc-1343*, *nc-tgp1* and *tgp1<sup>+</sup>* transcript levels in wild-type, *rrp6Δ*, *mmi1Δ* and *1343Δ*, respectively. Error bars represent s.e.m. resulting from at least three independent replicates.

(+PO<sub>4</sub>) and phosphate-deprived (−PO<sub>4</sub>) conditions. As expected, the levels of *tgp1<sup>+</sup>* and the *pho1<sup>+</sup>* control increased upon phosphate starvation (Fig. 4a,b). In contrast, the levels of both *nc-tgp1* and *nc-1343* RNAs decreased significantly in the absence of phosphate (Fig. 4a; Supplementary Fig. 6). The observed reduction in *nc-tgp1* levels is consistent with a situation whereby loss or reduction of *nc-tgp1* transcription permits *tgp1<sup>+</sup>* induction. In agreement with this, significantly less RNAPII associates with the *nc-tgp1* transcription unit in both phosphate-starved wild-type cells and phosphate-replete *1343Δ* cells, which do not transcribe *nc-tgp1* (Fig. 4c). Therefore, preventing *nc-tgp1* transcription, even in phosphate-rich medium, recapitulates the changes in RNAPII occupancy that normally accompany *tgp1<sup>+</sup>* induction upon phosphate deprivation.

**RNAi-directed heterochromatin does not repress *tgp1<sup>+</sup>*.** Cells with defective exosome function (*rrp6Δ*) accumulate non-coding RNAs, some of which have been reported to attract Mmi1-dependent RNA elimination factors, along with RNA interference (RNAi) components and the Clr4 H3K9 methyltransferase, leading to the formation of transiently regulated HOODs (heterochromatin domains)<sup>25</sup>. The regions containing the *tgp1<sup>+</sup>* and *pho1<sup>+</sup>* genes are included in HOOD-17 and HOOD-24, respectively, and both form a region of Mmi1-directed transient heterochromatin in *rrp6Δ* cells<sup>24,27</sup>. The *nc-tgp1* transcript is clearly regulated by Mmi1-directed exosome degradation

(Fig. 2e,f), however, we do not detect methylated H3K9 (H3K9me2) over the *tgp1<sup>+</sup>*, *nc-tgp1* or *nc-1343* genes within HOOD-17 in wild-type cells (Fig. 5a). Likewise, only very low levels of H3K9me2, slightly above background in cells lacking the H3K9 methyltransferase (*clr4Δ*), could be detected on the *pho1<sup>+</sup>* gene and the upstream Mmi1-targeted lncRNA (*nc-pho1*) within HOOD-24. Moreover, this low level of H3K9me2 did not drop appreciably upon induction of *tgp1<sup>+</sup>* and *pho1<sup>+</sup>* (−PO<sub>4</sub>; Fig. 5a). Equivalent background levels of H3K9me2 were detectable on another Mmi1-targeted lncRNA gene (*sme2<sup>+</sup>*) and the highly expressed actin gene (*act1<sup>+</sup>*). In contrast, H3K9me2 was ~100-fold enriched over the centromeric outer repeats (*dg*) in wild-type cells, but reduced to background in *clr4Δ* cells, indicating that H3K9-methylated chromatin had been efficiently immunoprecipitated. In addition, the transcript levels of *tgp1<sup>+</sup>*, *nc-tgp1*, *nc-1343*, *pho1<sup>+</sup>* and *nc-pho1* were unaffected by loss of RNAi (for example, *ago1Δ* or *dcr1Δ*) or heterochromatin components (for example, *clr4Δ* or *swi6Δ*) (Fig. 5b; Supplementary Fig. 7a). Nor were the kinetics of *tgp1<sup>+</sup>* or *pho1<sup>+</sup>* induction following phosphate starvation altered in cells lacking heterochromatin (Supplementary Fig. 7b,c). In contrast, *nc-tgp1*, *nc-pho1* and *sme2<sup>+</sup>* RNA levels were clearly elevated in cells lacking Mmi1-mediated exosome degradation (*mmi1Δ* and *rrp6Δ*). Thus, although H3K9me2 accumulates at particular regions in *rrp6Δ* cells (for example, HOOD-17: *tgp1<sup>+</sup>* and HOOD-24: *pho1<sup>+</sup>*), we conclude that RNAi and heterochromatin play no appreciable role in regulating these




**Figure 3 | *nc-tgp1*, not *nc-1343*, represses *tgp1*<sup>+</sup> to confer drug tolerance.** (a) Schematic diagram indicating strategic manipulations of lncRNAs upstream of *tgp1*<sup>+</sup> including *1343Δ*, shorter deletions of *ncRNA.1343* (*AΔ* and *BΔ*) and *ura4<sup>+</sup>* integration within the *nc-tgp1* lncRNA locus (*nc-tgp1:ura4<sup>+</sup>*) in wild-type background. (b) Serial dilutions of wild-type, *1343Δ*, *AΔ*, *BΔ* and *nc-tgp1:ura4<sup>+</sup>* were spotted on non-selective YES medium or in the presence of TBZ (20 μg ml<sup>-1</sup>), HU (10 mM) or caffeine (15 mM), respectively. (c) RT-qPCR experiments measured *tgp1*<sup>+</sup>, *nc-tgp1* and *nc-1343* transcript levels in wild-type, *1343Δ*, *AΔ*, *BΔ* and *nc-tgp1:ura4<sup>+</sup>* cells. Error bars represent s.e.m. resulting from three independent replicates.

genes under normal physiologically repressive conditions or during their induction.

***nc-tgp1* prevents Pho7 transcription factor binding.** The above analyses indicate that *nc-tgp1* is transcribed into the *tgp1*<sup>+</sup> promoter and suggest that production of this upstream lncRNA represses *tgp1*<sup>+</sup> expression. We therefore investigated if transcription of *nc-tgp1* interferes with the induction mechanism of *tgp1*<sup>+</sup> in response to phosphate starvation. The Pho7 transcription factor has previously been shown to engage phosphate-response gene promoters in phosphate-starved cells<sup>35,36</sup>. Our ChIP analyses confirmed that Pho7-green fluorescent protein (Pho7-GFP) accumulates on the *pho1*<sup>+</sup> promoter in phosphate-depleted cells (Supplementary Fig. 8). In addition, Pho7-GFP levels accumulate over the region upstream of *tgp1*<sup>+</sup> when activated (Fig. 6a). However, in cells unable to transcribe *nc-tgp1* (*1343Δ*), higher levels of Pho7-GFP associate with the region upstream of *tgp1*<sup>+</sup> even in repressive conditions (that is, +PO<sub>4</sub>). We conclude that loss of *nc-tgp1* expression due to phosphate starvation or by preventing production of this lncRNA (for example, *1343Δ*) allows Pho7 binding and subsequent *tgp1*<sup>+</sup> induction.


Active RNAPII promoters display reduced nucleosome density<sup>37</sup>. lncRNA transcription over promoters can increase nucleosome density and prevent gene induction<sup>8,10,20</sup>. We




**Figure 4 | Phosphate starvation induces *tgp1*<sup>+</sup> and reduces lncRNA transcription.** (a) RT-qPCR experiments measured *tgp1*<sup>+</sup>, *nc-tgp1* and *nc-1343* transcript levels in wild-type cells grown in phosphate-rich medium (+PO<sub>4</sub>) or in the absence of phosphate (-PO<sub>4</sub>). *pho1*<sup>+</sup> is a positive control for phosphate starvation. (b) Northern analysis of *tgp1*<sup>+</sup> in wild-type cells grown in the presence or absence of phosphate and *1343Δ* grown in the presence of phosphate. (c) Rpb1 ChIP-qPCR experiments performed in wild-type cells grown in the presence or absence of phosphate and *1343Δ* grown in the presence of phosphate. Error bars represent s.e.m. resulting from three independent replicates.

found that histone H3 levels were greater over the *tgp1*<sup>+</sup> gene and upstream region when it is repressed (+PO<sub>4</sub>) compared with when it is expressed (-PO<sub>4</sub>; Fig. 6b). In contrast, H3 levels over control loci (*act1*<sup>+</sup>, *sme2*<sup>+</sup> and *dg* repeats) were unaffected by phosphate availability. Thus, upstream transcription appears to alter nucleosome density over the *tgp1*<sup>+</sup> promoter and thereby occlude Pho7 binding. Likewise, a considerable drop in H3 levels was observed on the *pho1*<sup>+</sup> gene and *nc-pho1* lncRNA region upstream in phosphate-poor conditions, implying a similar mechanism may also operate to regulate the expression of *pho1*<sup>+</sup>. We conclude that transcription of the upstream lncRNA inhibits expression of *tgp1*<sup>+</sup> by a transcriptional interference mechanism that alters the chromatin landscape, preventing access to the key phosphate-responsive transcription factor Pho7.


To directly test if transcriptional interference of *tgp1*<sup>+</sup> by *nc-tgp1* is responsible for *tgp1*<sup>+</sup> repression, we replaced the *nc-tgp1* promoter with the strong, thiamine-regulated *nmt1* promoter (*nmt1-nc-tgp1*) (Fig. 7a). Transcription of *nc-tgp1* from the *nmt1* promoter is rendered unresponsive to phosphate. Instead, *nc-tgp1* is repressed or derepressed in the presence or absence of thiamine, respectively. When *nc-tgp1* was transcribed from the *nmt1* promoter, *tgp1*<sup>+</sup> remained repressed regardless of phosphate availability (Fig. 7b). In contrast, repression of *nmt1*-driven *nc-tgp1* by thiamine resulted in the induction of *tgp1*<sup>+</sup> expression in phosphate-rich media and consequently caused drug sensitivity (Fig. 7b,c). In addition, H3 levels over the region upstream of *tgp1*<sup>+</sup> were high when *nc-tgp1* was transcribed and



**Figure 5 | *tgp1+* is not regulated by RNAi/heterochromatin.** (a) H3K9me2 ChIP-qPCR experiments performed in the presence or absence of phosphate. *clr4Δ* was used as a negative control. The euchromatic actin gene (*act1+*) and centromeric *dg* repeats (*dg*) are positive and negative controls for heterochromatin. *pho1+* is a phosphate-regulated gene repressed by *nc-pho1*, a lncRNA target of Mmi1. *sme2+* is another lncRNA target of Mmi1. H3K9me2 to bulk H3 ratio has not been presented due to background methyl H3K9 levels detected at these loci. (b) RT-qPCR experiments measured *tgp1+*, *nc-tgp1* and *nc-1343* transcript levels in wild-type cells and cells lacking factors involved in heterochromatin formation and stability, respectively. Error bars represent s.e.m. resulting from at least three independent replicates.




**Figure 6 | *nc-tgp1* transcription prevents stable Pho7 binding and increases nucleosome density upstream of *tgp1+*.** (a) GFP ChIP-qPCR experiments were performed in the presence or absence of phosphate in cells with C-terminally GFP-tagged Pho7. An untagged strain was used as a negative control. Primer pair #3 was used to detect Pho7 binding at the *tgp1+* promoter. (b) Nucleosome density was measured by histone H3 ChIP-qPCR experiments in wild-type cells grown in the presence or absence of phosphate. Error bars represent s.e.m. resulting from three independent replicates.



**Figure 7 | *nmt1* controlled *nc-tgp1* alters drug tolerance in response to thiamine.** (a) Schematic diagram of *nc-tgp1* under the control of the strong, thiamine-repressible *nmt1* promoter. (b) RT-qPCR experiments measured *tgp1+* and *nc-tgp1* levels in response to thiamine and phosphate availability using *nmt1-nc-tgp1* cells. (c) Serial dilutions of wild-type, *1343Δ* and *nmt1-nc-tgp1* cells were spotted on non-selective PMG medium or in the presence of TBZ, HU or caffeine, respectively, with or without thiamine as indicated. (d) H3 ChIP-qPCR experiments in *nmt1-nc-tgp1* cells grown in the presence or absence of thiamine. Error bars represent s.e.m. resulting from three independent replicates.

reduced when *nc-tgp1* was repressed by thiamine (Fig. 7d). Lastly, exogenous expression of full-length *nc-tgp1* from a plasmid failed to repress *tgp1+*, ruling out the possibility that *nc-tgp1* operates in *trans* (Supplementary Fig. 9). Collectively, these findings

confirm that it is the transcription of *nc-tgp1* over the *tgp1+* promoter that alters nucleosome density to regulate *tgp1+* induction (see Fig. 8) and, as a consequence, drug tolerance of fission yeast cells.



**Figure 8 | Model for transcriptional interference at  $tgp1^+$ .** The presence of phosphate induces transcription of an unstable lncRNA ( $nc-tgp1$ ). lncRNA transcription increases nucleosome density, occludes Pho7 transcription factor binding and thus represses  $tgp1^+$  expression.  $nc-tgp1$  transcription is reduced following phosphate starvation, decreasing nucleosome density, allowing Pho7 to stably engage the  $tgp1^+$  promoter and induce  $tgp1^+$  expression.

## Discussion

An increasing number of lncRNAs have been shown to tightly regulate eukaryotic gene expression following intra-/extra-cellular environment changes that require rapid, integrated responses at the level of transcription<sup>2</sup>. In *S. pombe*, for example, the balance of antisense lncRNAs and sense transcription controls various stress-response pathways<sup>33,38</sup>. However, little is known about the majority of *S. pombe* intergenic lncRNAs. Here we selected and deleted eight stable, discrete lncRNAs in *S. pombe* that show conserved synteny in at least two of the three other known *Schizosaccharomyces* species. Excluding the *ter1+* control, only deletion of *ncRNA.1343* exhibited a definitive phenotype: sensitivity to various compounds due to induction of a nearby phosphate-responsive permease gene ( $tgp1^+$ ). Closer inspection revealed that the *ncRNA.1343* promoter is bidirectional. Furthermore, transcription from this bidirectional promoter favours the production of a previously unannotated and unstable lncRNA ( $nc-tgp1$ ) towards the  $tgp1^+$  gene under repressive conditions.

Recent studies in fission yeast have implicated lncRNAs in directing repression of specific genes by a mechanism involving transient RNAi-dependent heterochromatin formation<sup>27</sup>. For example, the Mmi1-targeted lncRNA upstream of  $pho1^+$  has recently been reported to recruit RNAi-directed heterochromatin to repress  $pho1^+$  in response to phosphate availability<sup>28</sup>. However, these findings differ from genome-wide H3K9me2 mapping which show that  $tgp1^+$  and  $pho1^+$ , both of which are regulated by upstream lncRNAs that are targeted for exosome-mediated degradation by Mmi1 (Fig. 2; ref 28), only accumulate RNAi-directed H3K9me2 in mutants with defective RNA processing/degradation (for example, *rrp6Δ*) and not in wild-type cells grown under repressive phosphate-rich conditions<sup>24</sup>. The significance of *rrp6Δ*-dependent heterochromatin at the  $tgp1^+$  and  $pho1^+$  genes is therefore unclear. Cells lacking Rrp6 accumulate aberrant RNAs and exhibit disrupted heterochromatin globally, including significantly decreased H3K9me2 over centromeric repeats<sup>39</sup>. Therefore caution must be exercised when interpreting the analyses of mutants with such severe defects in RNA processing/degradation. Importantly, we do not detect significant levels of H3K9me2 enrichment on the  $tgp1^+$  and  $pho1^+$  promoters/genes in wild-type cells under repressive (phosphate-rich) conditions. We cannot exclude the possibility that distinct assay conditions in a previous report allowed detection of low H3K9me2 levels on the  $pho1^+$  promoter when repressed<sup>28</sup>, however, the consequence of such H3K9me2 remains uncertain given that our analyses show that the expression of  $pho1^+$  or  $tgp1^+$  is unaffected by loss of RNAi/heterochromatin. We note that our findings are in agreement with previous expression profiling analyses, showing unaltered

$tgp1^+$  and  $pho1^+$  levels in *S. pombe* cells lacking RNAi/heterochromatin<sup>40</sup>. In contrast, transcripts arising from *bone fide* heterochromatin in centromeric outer repeats are clearly elevated when RNAi/heterochromatin is defective. Thus, our analyses indicate that the repression of both  $tgp1^+$  and  $pho1^+$  is unlikely to involve regulated heterochromatin in wild-type cells. Instead, we favour a model whereby  $tgp1^+$  and  $pho1^+$  are repressed by a transcriptional interference mechanism.

Transcriptional interference is well-established in many systems. In the bacterium *Escherichia coli*, the gene encoding the *clr* transcriptional activator is repressed in response to nitrogen starvation by the act of lncRNA transcription from an alternate upstream promoter<sup>41</sup>. In the single celled eukaryote *S. cerevisiae*, which lacks RNAi and heterochromatin, transcription of the *SRG1* lncRNA into the *SER3* promoter, or heterologous promoters, was found to alter nucleosome density and interfere with transcription factor binding<sup>8–10</sup>. Similarly, in *S. cerevisiae*, non-coding transcription over the *IME1* (ref. 20), *GAL7* (ref. 42) and *FLO11* (ref. 43) promoters prevent gene induction. Analogous mechanisms have also been reported in multicellular eukaryotes. For example, the *Drosophila Ubx* gene<sup>44</sup>, the human dihydrofolate reductase gene<sup>45</sup> and the imprinted *Igf2r* gene in mammals<sup>46</sup> are repressed independent of RNAi or transient heterochromatin formation by non-coding transcription into their respective promoters. These examples illustrate that transcriptional interference is a simple, conserved mechanism for modulating specific genes without requiring additional *trans*-acting regulatory factors. Our results are consistent with both *nc-tgp1* and *nc-pho1* mediating repression of downstream genes ( $tgp1^+$  and  $pho1^+$ , respectively) by transcriptional interference, not by the formation of transient heterochromatin. We base this conclusion on our findings that: (i)  $tgp1^+$  and  $pho1^+$  expression is unaffected by loss of RNAi/heterochromatin; (ii) H3K9me2 is not associated with  $tgp1^+$  or  $pho1^+$  loci in wild-type cells; (iii) *nc-tgp1* transcription declines when  $tgp1^+$  is induced ( $-PO_4$ ); (iv) loss of the *nc-tgp1* transcript allows induction of  $tgp1^+$  under normally repressive ( $+PO_4$ ) conditions (similarly, loss of lncRNA transcription upstream induces  $pho1^+$  in repressive medium<sup>27,28</sup>); (v) transcription of *nc-tgp1* by a thiamine-repressible promoter brings  $tgp1^+$  under the control of thiamine, rather than phosphate; (vi) RNAPII and nucleosome density is increased over the  $tgp1^+$  promoter region when the repressive *nc-tgp1* RNA is transcribed and (vii) the Pho7 activator binds the  $tgp1^+$  promoter region when *nc-tgp1* transcription is lost.

Genome-wide RNA sequencing has allowed the detection of a large number of lncRNAs in a variety of species. However, it remains unclear how many of these lncRNA are functional transcripts that act to influence gene expression and/or chromatin landscapes. Examples such as Xist RNA in mammals and roX RNAs in *Drosophila* represent functional transcripts that are critical for mediating dosage compensation by altering chromatin status and expression levels from sex chromosomes<sup>47</sup>. However, enthusiasm for lncRNA function has been somewhat dampened by reports showing that the ablation in animal models of some of the best-characterized lncRNAs (for example, HOTAIR, MALAT1, Kcnq1ot1, NEAT1) exhibited less dramatic or undetectable phenotypes<sup>48–53</sup>. Of the discrete stable lncRNAs that we deleted in fission yeast, only one (*ncRNA.1343*) had an obvious phenotype in the growth conditions tested. Detailed analysis was required to reveal that deletion of *ncRNA.1343* actually affected expression of a divergent unstable lncRNA (*nc-tgp1*) transcribed in the opposite orientation as the annotated locus. Only after further manipulation and analyses could we conclude that the expression of *nc-tgp1* interferes with the expression of  $tgp1^+$  downstream. The fact that the unstable

*nc-tgp1* transcript is the functional partner of the apparently non-functional stable *nc-1343* RNA transcribed from the same bidirectional promoter demonstrates the importance of comprehensive analyses of ncRNAs and the consequences of their deletion. Based on our analyses, we surmise that the low level expression of *nc-1343* represents transcriptional noise, resulting as a byproduct of ample *nc-tgp1* transcription. The syntenic conservation of *ncRNA.1343* within the *Schizosaccharomyces* genus<sup>31</sup> hints at the possibility of a conserved regulatory mechanism that involves lncRNA transcription into the promoter region of *tgp1*<sup>+</sup> in related species. Thus, although genome-wide approaches can rapidly catalogue the presence and response of various lncRNAs to different conditions, much more detailed locus-specific analyses is required to pinpoint the function of each individual lncRNA with respect to *cis* regulation of nearby genes or *trans* regulation of genes at distal loci.

## Methods

**Yeast strains, plasmids and standard techniques.** *S. pombe* strains used in this study are listed in Supplementary Table 1. Standard methods were used for fission yeast growth, genetics and manipulations<sup>54</sup>. All strains were grown in Yeast extract plus supplement medium (YES), unless otherwise indicated. For phosphate starvation experiments, cells were grown to mid-log phase in YES medium, washed twice in dH<sub>2</sub>O, and then grown for indicated times in Pombe minimal glutamate (PMG) synthetic medium without Na<sub>2</sub>HPO<sub>4</sub> (–PO<sub>4</sub>). Genetic deletions and protein tagging were carried out by lithium acetate transformation. All genetic modifications were confirmed by colony PCR. Plasmids were transformed by electroporation. Selections were performed on PMG/agar plates with according auxotrophy or on YES/agar plates with appropriate antibiotic(s) and grown at 32 °C. Serial (1:4) dilutions of equal number of cells were spotted onto YES/agar and grown at 32 °C, unless indicated otherwise. For drug-sensitivity experiments, cells were spotted onto YES/agar or PMG/agar with DMSO or TBZ (20 µg ml<sup>–1</sup>), HU (10 mM), caffeine (15 mM) and H<sub>2</sub>O<sub>2</sub> (1 mM). For ultraviolet-sensitivity experiments, spotted cells were ultraviolet-irradiated at 80 J m<sup>–2</sup> with a Stratagene UV Crosslinker and grown in the dark at 25 °C. The plasmids containing *lacZ* under the control of the *nc-tgp1* and *nc-1343* bidirectional promoter were cloned as follows. The non-coding promoter was amplified from *S. pombe* genomic DNA in both orientations (using *lacZ\_1\_F/lacZ\_1\_R* and *lacZ\_2\_F/lacZ\_2\_R* primer pairs; see Supplementary Table 2) and ligated into pREP vector containing *lacZ* using PstI/SalI restriction sites. To test if *nc-tgp1* can repress *tgp1*<sup>+</sup> in *trans*, the *nc-tgp1* transcription unit was amplified from *S. pombe* genomic DNA (using *nc-tgp1\_Sall\_F* and *nc-tgp1\_XmaI\_R* primer pairs, see Supplementary Table 2) and ligated into pREP3x using Sall/XmaI restriction sites.

**Liquid assay for β-galactosidase activity.** Assays for β-galactosidase activity were performed as described<sup>55</sup>. Briefly, yeast containing vectors expressing *lacZ* under the control of various promoters were grown to log phase (OD<sub>595</sub> of ~0.5) in selective media. Cells were permeabilized by SDS/chloroform. Cell extracts were equilibrated at 30 °C for 5 min before the addition of ortho-Nitrophenyl-β-galactoside (ONPG). The reaction was stopped with Na<sub>2</sub>CO<sub>3</sub> once the solution turned yellow and elapsed time was recorded. Cell debris was spun and the OD<sub>420</sub> was measured. Units were calculated as follows: Units/OD = 1000 × (OD<sub>420</sub>/Volume × Time × OD<sub>595</sub>).

**Chromatin and RIP.** Cells were grown to mid-log phase at 32 °C in YES. For phosphate starvation experiments, cells in mid-log phase were washed twice in dH<sub>2</sub>O before being grown in PMG (–PO<sub>4</sub>) for 4 h. ChIP was performed essentially as described<sup>12</sup>. Briefly, cells were fixed with 1% paraformaldehyde for 15 min at room temperature. Cells were lysed by bead beating (Biospec Products) and sonicated using a Bioruptor (Diagenode) sonicator at 5 °C on high for a total of 20 min (30 s ON/OFF cycles). Five microlitres of Rpb1 antibody (#2629; Cell Signaling), 2 µl GFP antibody (G10362; Life Technologies), 2 µl H3 antibody (ab1791; Abcam) and 1 µl of H3K9me2 antibody (m5.1.1; ref. 56) were used for IPs. RIP experiments were performed essentially as described<sup>13</sup>. Hisx6-TEV-Protein A-tagged Mmi1 was captured from cell lysate with IgG Dynabeads (Life Technologies). Mmi1-bound RNA was isolated by phenol-chloroform extraction, DNase treated and reverse transcribed. Quantitative analysis was performed by qPCR.

**RNA analysis.** RNA was isolated from *S. pombe* using RNeasy Mini- or Midi-Kits as per manufacturer's instructions (Qiagen). For RT-qPCR experiments, first strand complementary DNA synthesis was performed on Turbo DNase (Life Technologies) treated RNA using random hexamers and Superscript III (Invitrogen) as per manufacturer's instructions. Negative controls lacking RT were performed alongside all RT-qPCR experiments. Northern analysis of long non-coding

transcripts was performed using UTP-[a<sup>32</sup>P]-labelled RNA probes as described<sup>57</sup>. Transcription start sites were mapped using the SMARTer RACE complementary DNA Amplification Kit as per manufacturer's instructions (Clontech).

**Quantitative real-time PCR.** Primers used in this study are listed in Supplementary Table 2. qPCR was performed using SYBR Green on a Roche Lightcycler. Data was analyzed with LightCycler 480 Software 1.5.0.39. RT-qPCR levels were calculated by normalizing product of interest to an internal reference gene (*act1*<sup>+</sup>). Expression levels were expressed relative to levels detected in wild-type cells. ChIP enrichments were calculated as the ratio of product of interest from IP sample normalized to the corresponding input sample and expressed as %IP. Error bars represent s.e.m., resulting from at least three independent replicates.

## References

- Guttman *et al.* Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* **458**, 223–227 (2009).
- Ponting, C. P., Oliver, P. L. & Reik, W. Evolution and functions of long noncoding RNAs. *Cell* **136**, 629–641 (2009).
- Mitchell, P., Pefalski, E., Shevchenko, A., Mann, M. & Tollervey, D. The exosome: a conserved eukaryotic RNA processing complex containing multiple 3' → 5' exoribonucleases. *Cell* **91**, 457–466 (1997).
- Struhl, K. Transcriptional noise and the fidelity of initiation by RNA polymerase II. *Nat. Struct. Mol. Biol.* **14**, 103–105 (2007).
- Wilusz, J. E., Sunwoo, H. & Spector, D. L. Long noncoding RNAs: functional surprises from the RNA world. *Genes Dev.* **23**, 1494–1504 (2009).
- Li, B., Carey, M. & Workman, J. L. The role of chromatin during transcription. *Cell* **128**, 707–719 (2007).
- Hirota, K. *et al.* Stepwise chromatin remodeling by a cascade of transcription initiation of non-coding RNAs. *Nature* **456**, 130–134 (2008).
- Hainer, S. J., Pruneski, J. A., Michell, R. D., Monteverde, R. M. & Martens, J. A. Intergenic transcription causes repression by directing nucleosome assembly. *Genes Dev.* **25**, 29–40 (2011).
- Martens, J. A., Laprade, L. & Winston, F. Inter Thebault genic transcription is required to repress *Saccharomyces cerevisiae* SER3 gene. *Nature* **429**, 571–574 (2004).
- Thebault, P. *et al.* Transcription regulation by the noncoding RNA SRG1 requires Spt2-dependent chromatin deposition in the wake of RNA polymerase II. *Mol. Cell Biol.* **31**, 1288–1300 (2011).
- Fejes-Toth, K. *et al.* Post-transcriptional processing generates a diversity of 5'-modified long and short RNAs. *Nature* **457**, 1028–1032 (2009).
- Bayne, E. H. *et al.* Stc1: a critical link between RNAi and chromatin modification required for heterochromatin integrity. *Cell* **140**, 666–677 (2010).
- Motamedi, M. R. *et al.* Two RNAi complexes, RITS and RDRC, physically interact and localize to noncoding centromeric RNAs. *Cell* **119**, 789–802 (2004).
- Verdel, A. *et al.* RNAi-mediated targeting of heterochromatin by the RITS complex. *Science* **303**, 672–676 (2004).
- Volpe, T. A. *et al.* Regulation of heterochromatin silencing and histone H3 lysine-9 methylation by RNAi. *Science* **297**, 1833–1837 (2002).
- Rinn, J. L. *et al.* Functional demarcation of active and silent chromatin domains in human HOX loci by non-coding RNAs. *Cell* **129**, 1311–1323 (2007).
- Ørom, U. A. *et al.* Long noncoding RNAs with enhancer-like function in human cells. *Cell* **143**, 46–58 (2010).
- Camblong, J., Iglesias, N., Fickentscher, C., Dieppl, G. & Stutz, F. Antisense RNA stabilization induces transcriptional gene silencing via histone deacetylation in *S. cerevisiae*. *Cell* **131**, 706–717 (2007).
- Houseley, J., Rubbi, L., Grunstein, M., Tollervey, D. & Vogelauer, M. A ncRNA modulates histone modification and mRNA induction in the yeast GAL gene cluster. *Mol. Cell* **32**, 685–695 (2008).
- van Werven, F. J. *et al.* Transcription of two long non-coding RNAs mediates mating-type control of gametogenesis in budding yeast. *Cell* **150**, 1170–1181 (2012).
- Heo, J. B. & Sung, S. Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. *Science* **331**, 76–79 (2011).
- Wang, K. C. *et al.* A long noncoding RNA maintains active chromatin to coordinate homeotic gene expression. *Nature* **472**, 120–124 (2011).
- Hiriart, E. *et al.* Mmi1 RNA surveillance machinery directs RNAi complex RITS to specific meiotic genes in fission yeast. *EMBO J.* **31**, 2296–2308 (2012).
- Yamanaka, S. *et al.* RNAi triggered by specialized machinery silences developmental genes and retrotransposons. *Nature* **493**, 557–560 (2013).
- Zofall, M. *et al.* RNA elimination machinery targeting meiotic mRNAs promotes facultative heterochromatin formation. *Science* **335**, 96–100 (2012).
- Harigaya, Y. *et al.* Selective elimination of messenger RNA prevents an incidence of untimely meiosis. *Nature* **442**, 45–50 (2006).
- Lee, N. N. *et al.* Mtr4-like protein coordinates nuclear RNA processing for heterochromatin assembly and for telomere maintenance. *Cell* **155**, 1061–1074 (2013).



28. Shah, S., Wittmann, S., Kilchert, C. & Vasilieva, L. lncRNA recruits RNAi and the exosome to dynamically regulate *pho1* expression in response to phosphate levels in fission yeast. *Genes Dev.* **28**, 213–244 (2014).
29. Pang, K. C., Frith, M. C. & Mattick, J. S. Rapid evolution of noncoding RNAs: lack of conservation does not mean lack of function. *Trends Genet.* **22**, 1–5 (2006).
30. Ulitsky, I., Shkumatava, A., Jan, C. H., Sive, H. & Bartel, D. P. Conserved function of lincRNAs in vertebrate embryonic development despite rapid sequence evolution. *Cell* **147**, 1537–1550 (2011).
31. Rhind, N. *et al.* Comparative functional genomics of the fission yeasts. *Science* **332**, 930–936 (2011).
32. Wilhelm, B. T. *et al.* Dynamic repertoire of a eukaryotic transcriptome surveyed at single-nucleotide resolution. *Nature* **453**, 1239–1243 (2008).
33. Leong, H. S. *et al.* A global non-coding RNA system modulates fission yeast protein levels in response to stress. *Nat. Commun.* **5**, 3947 (2014).
34. Almaguer, C., Mantella, D., Perez, E. & Patton-Vogt, J. P. Inositol and phosphate regulate *GIT1* transcription and glycerophosphoinositol incorporation in *Saccharomyces cerevisiae*. *Eukaryot. Cell* **2**, 729–736 (2003).
35. Carter-O'Connell, I., Peel, M. T., Wykoff, D. D. & O'Shea, E. Genome-wide characterization of the phosphate starvation response in *Schizosaccharomyces pombe*. *BMC Genomics* **13**, 697 (2012).
36. Henry, T. C. *et al.* Systematic screen of *Schizosaccharomyces pombe* deletion collection uncovers parallel evolution of the phosphate signal transduction pathway in yeasts. *Eukaryot. Cell* **10**, 198–206 (2011).
37. Yuan, G. C. *et al.* Genome-scale identification of nucleosome positions in *S. cerevisiae*. *Science* **309**, 626–630 (2005).
38. Bitton, D. A. *et al.* Programmed fluctuations in sense/antisense transcript ratios drive sexual differentiation in *S. pombe*. *Mol. Syst. Biol.* **7**, 559 (2011).
39. Reyes-Turcu, F. E., Zhang, K., Zofall, M., Chen, E. & Grewal, S. I. Defects in RNA quality control factors reveal RNAi-independent nucleation of heterochromatin. *Nat. Struct. Mol. Biol.* **18**, 1132–1138 (2011).
40. Hansen, K. R. *et al.* Global effects on gene expression in fission yeast by silencing and RNA interference machineries. *Mol. Cell Biol.* **25**, 590–601 (2005).
41. Zafar, M. A., Carabetta, V. J., Mandel, M. J. & Silhavy, T. J. Transcriptional occlusion caused by overlapping promoters. *Proc. Natl Acad. Sci. USA* **111**, 1557–1561 (2014).
42. Greger, I. H., Aranda, A. & Proudfoot, N. Balancing transcription interference and initiation on the *GAL7* promoter of *Saccharomyces cerevisiae*. *Proc. Natl Acad. Sci. USA* **97**, 8415–8420 (2000).
43. Bumgarner, S. L. *et al.* Toggle involving cis-interfering noncoding RNAs controls variegated gene expression in yeast. *Proc. Natl Acad. Sci. USA* **106**, 18321–18326 (2009).
44. Petruk, S. *et al.* Transcriptional elongations of non-coding bxd RNAs promoted by the Trithorax TAC1 complex represses *Ubx* by a transcriptional interference mechanism. *Cell* **127**, 1209–1221 (2006).
45. Martjanov, I., Ramadass, A., Barros, A. S., Chow, N. & Akoulitchev, A. Repression of the human dihydrofolate reductase gene by a non-coding interfering transcript. *Nature* **445**, 666–670 (2007).
46. Latos, P. A. *et al.* Airn transcriptional overlap, but not its lncRNA products, induces imprinted *Igf2r* silencing. *Science* **338**, 1469–1472 (2012).
47. Lee, J. T. & Bartolomei, M. S. X-activation, imprinting, and long non-coding RNAs in health and disease. *Cell* **152**, 1308–1323 (2013).
48. Eißmann, M. *et al.* Loss of the abundant nuclear non-coding RNA MALAT1 is compatible with life and development. *RNA Biol.* **9**, 1076–1087 (2012).
49. Korostowski, L., Sedlak, N. & Engel, N. The *Kcnq1ot1* long non-coding RNA affects chromatin and expression of *Kcnq1*, but does not regulate its imprinting in the developing heart. *PLOS Genet.* **8**, e1002956 (2012).
50. Nakagawa, S. *et al.* Malat1 is not an essential component of nuclear speckles in mice. *RNA* **18**, 1487–1499 (2012).
51. Schorderet, P. & Duboule, D. Structural and functional differences in the long non-coding RNA *hotair* in mouse and human. *PLOS Genet.* **7**, e1002071 (2011).
52. Zhang, B. *et al.* The lncRNA *Malat1* is dispensable for mouse development but its transcription plays a cis-regulatory role in the adult. *Cell Rep.* **2**, 111–123 (2012).
53. Nakagawa, S., Naganuma, T., Shioi, G. & Hirose, T. Paraspeckles are subpopulation-specific nuclear bodies that are not essential in mice. *J. Cell Biol.* **193**, 31–39 (2011).
54. Moreno, S., Klar, A. & Nurse, P. Molecular genetic analysis of fission yeast *Schizosaccharomyces Pombe*. *Methods Enzymol.* **194**, 795–823 (1991).
55. Guarente, L. Yeast promoters and *lacZ* fusions designed to study expression of cloned genes in yeast. *Methods Enzymol.* **101**, 181–191 (1983).
56. Nakagawachi, T. *et al.* Silencing effect of CpG island hypermethylation and histone modifications on O6-methylguanine-DNA methyltransferase (MGMT) gene expression in human cancer. *Oncogene* **22**, 8835–8844 (2003).
57. Buscaino, A. *et al.* Raf1 is a DCAF for the Rik1 DDB1-like protein and has separable roles in siRNA generation and chromatin modification. *PLOS Genet.* **8**, e1002499 (2012).

### Acknowledgements

We would like to thank Sandra Catania, Alison Pidoux, Manu Shukla and Sharon White for their technical expertise and input. We are grateful to Takeshi Urano for the H3K9me2 (5.1.1) antibody, Tomoyasu Sugiyama and Lidia Vasilieva for strains, and Steven West and Sander Granneman for critically evaluating the manuscript. R.A. is supported by the Darwin Trust of Edinburgh. The Centre for Cell Biology is supported by core funding from the Wellcome Trust (092076/Z/10/Z). P.T. is supported by European Commission Network of Excellence EpiGeneSys (HEALTH-F4-2010-257082) to R.C.A. R.C.A. is a Wellcome Trust Principal Research Fellow and this research was supported by the Wellcome Trust (095021/Z/10/Z).

### Authors contributions

R.A. and R.C.A. conceived and designed the experiments. R.A. performed the experiments. P.T. performed bioinformatics analysis. R.A., P.T. and R.C.A. analyzed the data. R.A. and R.C.A. wrote the paper.

### Additional information

**Supplementary Information** accompanies this paper at <http://www.nature.com/naturecommunications>

**Competing financial interests:** The authors declare no competing financial interests.

**Reprints and permission** information is available online at <http://npg.nature.com/reprintsandpermissions/>

**How to cite this article:** Ard, R. *et al.* Long non-coding RNA-mediated transcriptional interference of a permease gene confers drug tolerance in fission yeast. *Nat. Commun.* 5:5576 doi: 10.1038/ncomms6576 (2014).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>