

November 25-27, 2021 THAILAND
Virtual Conference

“All accepted abstracts will be published in the JGO”

**Abstract submission:
Open on June 1, 2021 - August 15, 2021
Abstract acceptance notification by:
September 15, 2021**

**Early Registration
Open : June 21, 2021
Close : September 30, 2021**



Contact us: asgo2021@gmail.com

Online Registration: www.asgo2021.org



Editorial



How to perform sentinel node detection in high-risk endometrial cancer: one step forward

Vincent Balaya ¹, Fabrice Lecuru ^{2,3}, Enora Laas ^{2,3}

¹Service de Gynécologie-Obstétrique, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland

²Service de Chirurgie Sénologique, Gynécologique et Plastie, Labelled Ovarian Cancer Center by ESGO, Institut Curie, Paris, France

³Université de Paris, Paris, France

► See the article “Parsaortic sentinel lymph node detection in intermediate and high-risk endometrial cancer by transvaginal ultrasound-guided myometrial injection of radiotracer (TUMIR)” in volume 32, e52.

OPEN ACCESS

Received: May 10, 2021

Accepted: May 10, 2021

Correspondence to

Fabrice Lecuru

Address: Service de Chirurgie Sénologique,
Gynécologique et Plastie, Institut Curie, 26 rue
d’Ulm, 75005, Paris, France.
E-mail: fabrice.lecuru@curie.fr

Copyright © 2021. Asian Society of
Gynecologic Oncology, Korean Society of
Gynecologic Oncology, and Japan Society of
Gynecologic Oncology

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>)
which permits unrestricted non-commercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.

ORCID iDs

Vincent Balaya
<https://orcid.org/0000-0002-7561-239X>

Fabrice Lecuru
<https://orcid.org/0000-0002-4071-3932>

Enora Laas
<https://orcid.org/0000-0003-3875-4011>

Conflict of Interest

No potential conflict of interest relevant to this
article was reported.

Endometrial cancer is the most common gynecologic cancer in developed countries. Its incidence is rising in several parts of the world, in parallel to the incidence of obesity. This concerns mainly low-risk lesions, most of them being cured by a total hysterectomy with bilateral salpingo-oophorectomy whereas high-risk tumors require more aggressive treatments [1].

The nodal staging of endometrial cancer is an old debate. No staging, limited pelvic dissection, pelvic and paraaortic dissection have been proposed according to clinical stage and time. The therapeutic value of nodal dissection has also been suggested. Unfortunately, trials addressing what we call now low and intermediate-risk endometrial cancers showed no benefit of a systematic lymphadenectomy [2].

Sentinel node biopsy (SLN) has been introduced more recently. This technique appears seductive since it could provide the nodal status with a limited surgical aggressiveness. The limitation of systematic extended dissections in routine was precisely the imbalance between the potential benefit in terms of staging and prognosis vs. the feasibility and the perioperative risk due to age and comorbidity. The interest to SLN is reinforced since it is usable by minimally invasive surgery, and more recently by the spread of indocyanine green with high accuracy [3].

Concerning endometrial cancer, 2 sites of injection are possible. The most used and evaluated is the cervix, with a technique similar to that of cervical cancer [3]. This approach is easy to learn, to practice, provides a good sensitivity and negative predictive value especially for low-risk or intermediate-risk tumors. However, cervical injection does not explore the lymphatic drainage of the tumor, and the rate of paraaortic SLN (and positive paraaortic nodes) is much lower than after classical dissections [4].

In this issue, Angeles et al. [5] report on a large experience of SLN with transvaginal myometrial injection. They included 102 patients with intermediate and high-risk endometrial carcinoma. All patients underwent SLN (isotope + blue dye) and pelvic + paraaortic dissection. The detection rate was 79%. The sensitivity and negative predictive value were 87.5% and 97%; and 83% and 96.9% for paraaortic metastases. This technique

Author Contributions

Conceptualization: B.V., L.E., L.F. Supervision:
 L.F. Validation: B.V., L.E., L.F. Writing - original
 draft:: B.V., L.E., L.F. Writing - review & editing::
 B.V., L.E., L.F.

appears as effective as extended dissection to assess the nodal status of these patients, especially in the paraaortic area. However, the rate of isolated paraaortic SLN and moreover of positive isolated paraaortic SLN were low (respectively 6/81 and 3/20 patients). As in previous reports, most paraaortic involvement cases were associated to a pelvic spread. The technique presented in this paper is comparable to those previously described such as hysteroscopic injection, subserosal injection or dual injections. Moreover, it appears surprisingly well tolerated by the patients. We could argue that this technique makes realistic the search of rare isolated paraaortic metastases.

If the role of SLN in low risk and low-intermediate risk is now well established, its use in high intermediate and high risk is still controversial. Nonetheless, the recent ESGO/ESTRO/ESP guidelines for endometrial management state for SLN in this setting for lymph node staging in stage I and II, due to the high sensitivity to detect pelvic lymph node metastases and a high negative predictive value [1,6]. To be fully reliable in this context, the use of surgical SLN algorithm and experienced surgeons are mandatory.

The paucity of isolated paraaortic metastases is a question per se given that this event is rare (less than 2%). This may be explained by anatomic uterine lymphatic drainage pathway [7]. Usually, endometrial cancer spreads along the pelvic pathways. The lower and middle part of the uterus drain laterally to the parametria and then to the obturator lymph nodes, where are mostly found metastatic lymph nodes. The upper corpus and fundus drain to the interiliac lymph nodes, then to the common iliac lymph nodes and finally to the paraaortic lymph nodes. In very few cases, the fundic lymphatic drainage may reach directly to the paraaortic lymph nodes through the gonadal vessels and therefore shunt pelvic lymphatic drainage. Although the imaging (magnetic resonance imaging, positron emission tomography, or computed tomography scan) is not perfect, they permit to detect preoperatively paraaortic positive nodes with sufficient accuracy [8]. In addition, the ratio of the number of patients submitted to SLN vs. number of patients with improved prognosis due to the diagnosis and treatment of isolated aortic metastasis is questionable.

Finally, the real pending challenge is now to define the place of SLN at the light of the molecular classification [9]. The real impact of positive SLN for the adjuvant treatment is challenged by the molecular profile. Patients with POLE mutation have a negligible risk of nodal metastasis, an overall excellent prognosis and could avoid any nodal sampling. Patients with p53 mutations require mainly chemotherapy, whatever the result of nodal sampling. One could argue that positive paraaortic SLN could modify the radiation fields. We just emphasize that extended field radiation therapy has never proved any benefit in terms of overall survival. It could be hypothesized that SLN might provide an added prognostic value with diagnosis of micrometastasis and improve prognostic assessment of p53 mutated, or mismatch-repair deficient or mismatch-repair proficient patients. However, acceptable quality data are required in this subgroup before drawing such conclusion.

One of major concern is maybe first to standardize surgical techniques of SLN and to respect detection algorithm, as it could impact diagnostic accuracy and oncologic outcomes.

This paper makes things progress concerning the best approach to inject the tracer in endometrial cancer. It changes the vision of paraaortic nodal staging and keeps nodal surgery in the armamentarium of high-risk endometrial cancer. Future works have to correlate SLN and the molecular classification.

REFERENCES

1. Concin N, Matias-Guiu X, Vergote I, Cibula D, Mirza MR, Marnitz S, et al. ESGO/ESTRO/ESP guidelines for the management of patients with endometrial carcinoma. *Int J Gynecol Cancer* 2021;31:12-39.
[PUBMED](#) | [CROSSREF](#)
2. ASTEC study group Kitchener H, Swart AMC, Qian Q, Amos C, Parmar MKB. Efficacy of systematic pelvic lymphadenectomy in endometrial cancer (MRC ASTEC trial): a randomised study. *Lancet* 2009;373:125-36.
[PUBMED](#) | [CROSSREF](#)
3. Frumovitz M, Plante M, Lee PS, Sandadi S, Lilja JF, Escobar PF, et al. Near-infrared fluorescence for detection of sentinel lymph nodes in women with cervical and uterine cancers (FILM): a randomised, phase 3, multicentre, non-inferiority trial. *Lancet Oncol* 2018;19:1394-403.
[PUBMED](#) | [CROSSREF](#)
4. Mariani A, Dowdy SC, Cliby WA, Gostout BS, Jones MB, Wilson TO, et al. Prospective assessment of lymphatic dissemination in endometrial cancer: a paradigm shift in surgical staging. *Gynecol Oncol* 2008;109:11-8.
[PUBMED](#) | [CROSSREF](#)
5. Angeles MA, Migliorelli F, Vidal-Sicart S, Saco A, Ordi J, Ros C, et al. Paraaortic sentinel lymph node detection in intermediate and high-risk endometrial cancer by transvaginal ultrasound-guided myometrial injection of radiotracer (TUMIR). *J Gynecol Oncol* 2021;32:e52.
[PUBMED](#) | [CROSSREF](#)
6. Persson J, Salehi S, Bollino M, Lönnerfors C, Falconer H, Geppert B. Pelvic sentinel lymph node detection in high-risk endometrial cancer (SHREC-trial)-the final step towards a paradigm shift in surgical staging. *Eur J Cancer* 2019;116:77-85.
[PUBMED](#) | [CROSSREF](#)
7. Mariani A, Webb MJ, Keeney GL, Podratz KC. Routes of lymphatic spread: a study of 112 consecutive patients with endometrial cancer. *Gynecol Oncol* 2001;81:100-4.
[PUBMED](#) | [CROSSREF](#)
8. Gong Y, Guo Z, Tang X, Li C, Wang Q. Performance of different imaging techniques for detection of para-aortic lymph node metastasis from gynecological malignancies: a systematic review and meta-analysis. *Gynecol Obstet Invest* 2020;85:53-71.
[PUBMED](#) | [CROSSREF](#)
9. Cancer Genome Atlas Research Network Kandoth C, Schultz N, Cherniack AD, Akbani R, Liu Y, et al. Integrated genomic characterization of endometrial carcinoma. *Nature* 2013;497:67-73.
[PUBMED](#) | [CROSSREF](#)