


CORRECTION

Correction: Nano-sized Al₂O₃ reduces acute toxic effects of thiacloprid on the non-biting midge *Chironomus riparius*

Carla S. Lorenz, Anna-J. Wicht, Leyla Guluzada, Leilei Luo, Leonie Jäger, Barbara Crone, Uwe Karst, Rita Triebkorn, Yucang Liang, Reiner Anwander, Stefan B. Haderlein, Carolin Huhn, Heinz-R. Köhler

In [Fig 1](#), the y-axis is labeled incorrectly. Please see the correct [Fig 1](#) here.



OPEN ACCESS

Citation: Lorenz CS, Wicht A-J, Guluzada L, Luo L, Jäger L, Crone B, et al. (2017) Correction: Nano-sized Al₂O₃ reduces acute toxic effects of thiacloprid on the non-biting midge *Chironomus riparius*. PLoS ONE 12(6): e0179786. <https://doi.org/10.1371/journal.pone.0179786>

Published: June 12, 2017

Copyright: © 2017 Lorenz et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 1. Internal thiacloprid concentrations in *C. riparius* larvae [µg/kg] vs nominal concentration in water [µg/L]. Larvae were exposed for 96 h before they were transferred to filtered and dechlorinated tap water for 24 h to empty their guts (n = 1–3). R² of the respective regression curves were 0.91 for Thiacloprid, 0.97 for animals exposed to the mixture including 300 mg/L nano-Al₂O₃ and 0.82 for animals exposed to a mixture with 1000 mg/L nano-Al₂O₃. Nominal values are shown in this graph, whereas measured concentrations can be obtained from Table 1.

<https://doi.org/10.1371/journal.pone.0179786.g001>

Reference

1. Lorenz CS, Wicht A-J, Guluzada L, Luo L, Jäger L, Crone B, et al. (2017) Nano-sized Al₂O₃ reduces acute toxic effects of thiacloprid on the non-biting midge *Chironomus riparius*. PLoS ONE 12(5): e0176356. doi:10.1371/journal.pone.0176356 PMID: 28464012