

OPEN

# Soil aggregates stability and storage of soil organic carbon respond to cropping systems on Black Soils of Northeast China

Meng Zhou<sup>1,2,3</sup>, Chunzhu Liu<sup>1</sup>, Jie Wang<sup>3</sup>, Qingfeng Meng<sup>1</sup>, Ye Yuan<sup>4</sup>, Xianfa Ma<sup>1\*</sup>, Xiaobing Liu<sup>1,2\*</sup>, Yingxue Zhu<sup>1</sup>, Guangwei Ding<sup>5</sup>, Jizhou Zhang<sup>1,6</sup>, Xiannan Zeng<sup>7</sup> & Weiling Du<sup>3</sup>

Monoculture and improper management may reduce soil fertility and deteriorate soil structure in Black soils (Mollisols) of Northeast China. The experiment was carried out from 2015 to 2016 in Black Soils comprising five cropping systems: continuous corn (CC), soybean-corn rotation (SC), corn-soybean rotation (CS), fallow-corn (FC), and fallow-soybean (FS). Our results showed that CS and FS treatments significantly increased mean weight diameter (MWD) and fractal dimension (D) in mechanical stability aggregates (MSAs), and increased MWD and geometric mean diameter (GMD) in water-stable aggregates (WSAs) compared with CC treatment. These two treatments were also significantly increased water-stable aggregates stability rate (WSAR), but decreased percentage of aggregates destruction (PAD) than CC treatment. Meanwhile, CS and FS treatments exhibited a higher carbon accumulation than CC treatment in bulk soils. Soil organic carbon (SOC) concentration in  $WSA_{0.106-0.25}$ ,  $WSA_{2-5\text{ mm}}$  and  $WSA_{0.5-1\text{ mm}}$  had a dominant effect on aggregate stability. Simultaneously, SOC in  $WSA_{>5\text{ mm}}$  affected SOC concentration in bulk soils. As a whole, the CS and FS treatments can increase the percentage of macro-aggregates, enhance aggregate stability, as well as increase SOC concentration in bulk soils and all soil aggregate sizes.

Soil organic carbon (SOC) plays a key role in forming and stabilizing soil structure, enhancing soil physical properties, and nutrient recycling<sup>1-3</sup>. Soil aggregate, the basic unit of soil structure, mediates many physical and chemical processes in soils<sup>4-8</sup>, such as soil compaction, soil nutrient recycling, soil erosion, root penetration, and crop yield<sup>9</sup>. Aggregate stability is frequently used as an indicator of soil structure<sup>10-12</sup> because better soil structure and higher aggregate stability are vital to improve soil fertility, soil sustainability, and productivity<sup>13,14</sup>.

SOC influenced aggregate stability and soil structure<sup>15,16</sup>. The stability of organic carbon in different size aggregates is different. Organic carbon in the micro-aggregates is less susceptible to change than it is in the macro-aggregates<sup>17</sup>. The soil organic matters of different cropping systems differed based on the quantity and quality of the crop residue coverage and the environment, affecting the organic carbon contents of the soil and the aggregate stability<sup>18</sup>. The cropping systems mainly create conditions for the decomposition and transformation of soil organic matter by changing the distribution of soil organic carbon and the active habitat of microorganisms, thereby causing changes in soil aggregates<sup>19</sup>.

Soil mean weight diameter (MWD), geometric mean diameter (GMD), fractal dimension (D), percentage of aggregates destruction (PAD) and water-stable aggregates stability rate (WSAR) are all indicators of soil aggregate stability. The larger the MWD and GMD values are, the higher the average particle size agglomeration of soil aggregates are, and the stronger the stability of soil structure is<sup>20</sup>. Castrignano and Stelluti<sup>21</sup> found that the larger

<sup>1</sup>School of Resources and Environment, Northeast Agricultural University, Harbin, 150030, China. <sup>2</sup>Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Harbin, 150081, China. <sup>3</sup>Department of Biological Center, Harbin Academy of Agricultural Sciences, Harbin, 150028, China. <sup>4</sup>Mudanjiang Tobacco Science Research Institute, Heilongjiang Branch of China Tobacco Corporation, Harbin, 150076, China. <sup>5</sup>Chemistry Department, Northern State University, Aberdeen, SD, 57401, USA. <sup>6</sup>Institute of Natural Resources and Ecology, Heilongjiang Academy of Sciences, Harbin, 150040, China. <sup>7</sup>Institute of Crop Cultivation and Tillage, Heilongjiang Academy of Agricultural Sciences, Harbin, 150086, China. \*email: [mx7856@163.com](mailto:mx7856@163.com); [liuxb@iga.ac.cn](mailto:liuxb@iga.ac.cn)

the D value of soil aggregates are, the easier dispersed the aggregates were. The amount of soil micro-aggregates ( $WSA_{0.053-0.25\text{ mm}}$ ) increases with the increase of D value<sup>22</sup>. The lower the PAD value and higher WSAR value are, the more stable the soil structure is, and the better the soil erosion resistance is, and thus the higher the soil fertility is<sup>23,24</sup>.

Legume crops with non-legume crop rotation extensively applied around the world are important technologies for conserving soil resources and achieving sustainable development of agriculture<sup>25–30</sup>. The two-year rotation system of corn-soybean increased soil fertility, adjusted the balance of nutrients in the soil, and controlled soil degradation in the cold region<sup>31</sup>. Compared with continuous cropping, leguminous-graminaceous crop rotations were able to increase the farmland biodiversity with time growing<sup>32</sup>. Rotation in conjunction with manure amendment management decelerated the degradation of water stable macro-aggregates with re-aggregated structure<sup>33</sup>.

To conserve soil resources, land fallow practices have been implemented in many parts of the world. The Conservation Reserve Program (CRP) was proposed by the United States in 1986<sup>34</sup>, and this measures can control soil erosion<sup>35</sup>, improve water quality<sup>36</sup>, protect biodiversity<sup>37</sup>, create the habitat for wild animals and plants, and ameliorate agricultural non-point source pollution<sup>38</sup>. Thus, the experience is worth referencing with respect to land reclamation and land protection<sup>39–41</sup>.

Black Soils in Northeast China, also referred to as Mollisols or Chernozems, are one of the most precious soil resources for sustainable agricultural development in the country<sup>33</sup>, which play an irreplaceable role in safeguarding China's regional eco-environmental security and national food security<sup>42,43</sup>. The original Black Soils demonstrate a high fertility and excellent physicochemical characteristics with approximately  $50\text{--}80\text{ g kg}^{-1}$  SOC<sup>44,45</sup>. However, the wealth of agricultural Black Soils has undergone a decline in natural fertility and soil structure deterioration due to long-term continuous cropping and monoculture as well as excessive reclamation<sup>46</sup>.

The main cropping systems in Northeast China are continuous cropping and crop rotation with different crop sequences. Previous studies indicated that long-term tillage affected the formation and stabilization of soil aggregate<sup>10,12,14,47–49</sup>, which holds SOC stocks reckoned as an important strategy in maintaining soil structure and enhancing the condition of soil particles<sup>9,50</sup>. However, limited information is available on the impact of rotation and intermittent fallow on the distribution and aggregate stability in Black Soils of Northeast China. We rationally hypothesized that cropping systems with rotation and intermittent fallow will possibly affect the processes of aggregate formation. The objectives of this study were (1) to determine the distribution and stability of mechanical stability aggregates (MSAs) water-stable aggregates (WSAs) under different cropping systems in Black Soils; (2) to evaluate cropping impact on the SOC concentrations and stocks in bulk soils and WSAs; and (3) to understand how the cropping systems influence the correlations between the parameters of soil structure stability and WSAs, as well as SOC concentrations.

## Material and Methods

**Experimental site.** The study was conducted from 2015 to 2016 at the research farm of Northeast Agricultural University, located at the Xiangyang, Harbin, China (longitude  $126^{\circ}54'8.68''\text{E}$ , latitude  $45^{\circ}46'14.27''\text{N}$ , and altitude 184 meters). Before 2015, our experimental site was planted corn all the time. The site is in a moderate temperate zone with a semi-humid continental monsoon climate. The mean annual temperature is  $3.5^{\circ}\text{C}$  with a frost-free period of 142 days, and the mean annual precipitation is 553.9 mm. The soil type in the area is typical Black Soils with light loam texture. Soil physic-chemical properties were as following: soil pH of 6.8; soil organic matter (SOM) of  $35\text{ g kg}^{-1}$ , total nitrogen (TN) of  $1.71\text{ g kg}^{-1}$ , available nitrogen (AN), phosphorus (AP), and potassium (AK) of  $76\text{ mg kg}^{-1}$ ,  $12\text{ mg kg}^{-1}$ , and  $158\text{ mg kg}^{-1}$ , respectively, and bulk density ( $\rho_b$ ) of  $1.24\text{ g cm}^{-3}$ .

**Experimental design and management.** The experimental treatments for two consecutive years were as follows:

Continuous corn (CC): Corn was planted with hole sowing in the spring of 2015 and intertillaged with 15–20 cm soil depth in the mid-June of 2015, and then chopped corn stalk was incorporated into the soil with 25–30 cm plowing depth at harvest in the autumn of 2015. The tillage methods of 2016 was the same as 2015.

Soybean-corn rotation (SC): Soybean was planted with drill sowing in the spring of 2015 and intertillaged with 10–15 cm soil depth in the mid-June of 2015, and then chopped soybean stalk was incorporated into the soil with 25 cm plowing depth at harvest in the autumn of 2015. Next, corn was planted on the original ridge with hole sowing in the spring of 2016 and intertillaged with 15–20 cm soil depth in the mid-June of 2016, and then chopped corn stalk was incorporated into the soil with 25–30 cm plowing depth at harvest in the autumn of 2016.

Corn-soybean rotation (CS): Corn was planted with hole sowing in the spring of 2015 and intertillaged with 25 cm soil depth in the mid-June of 2015, and then chopped corn stalk was incorporated into the soil with 30–35 cm plowing depth at harvest in the autumn of 2015. Next, soybean was planted on the original ridge with drill sowing in the spring of 2016 and intertillaged with 15–20 cm soil depth in the mid-June of 2016, and then chopped soybean stalk was incorporated into the soil with 25–30 cm plowing depth at harvest in the autumn of 2016.

Fallow-corn (FC): The experimental site was abandoned without any tillage measures and restored by natural vegetation in 2015. Next, corn was planted with hole tillage in the spring of 2016 and intertillaged with 15–20 cm soil depth in the mid-June of 2016, and then chopped corn stalk was incorporated into the soil with 25–30 cm plowing depth at harvest in the autumn of 2016.

Fallow-soybean (FS): The experimental site was abandoned without any tillage measures and restored by natural vegetation in 2015. Next, soybean was planted with drill tillage in the spring of 2016 and intertillaged with 15–20 cm soil depth in the mid-June of 2016, and then chopped soybean stalk was incorporated into the soil with 25–30 cm plowing depth at harvest in the autumn of 2016.

The experimental design was a randomized complete block with three replications consisting of 15 plots in total. Each plot was 10.0 meters long and 3.9 meters wide with row spacing of 0.65 m. The soybean (*Glycine max*) variety was Jiannong No. 18 (Soybean Institute of Jilin Academy of Agricultural Sciences of China) and the corn (*Zea mays*) variety was Xianyu 335 (Pioneer Corporation Ltd., United States). For fertilization, the compound fertilizer (N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O:15-23-10) was applied at a rate of 225 kg ha<sup>-1</sup> year<sup>-1</sup> as the base fertilizer of soybean; and the compound fertilizer (N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O: 23-12-13) was applied at a rate of 375 kg ha<sup>-1</sup> year<sup>-1</sup> as the base fertilizer and urea was applied at a rate of 150 kg ha<sup>-1</sup> year<sup>-1</sup> on jointing stage of corn according to the amount of fertilizer application after soil testing in the local area.

**Soil sampling.** All soil samples (0–20 cm) were collected from each plot in the harvest of 2016. Undisturbed soil samples with dimensions of 20 cm (height) × 30 cm (length) × 30 cm (width) were obtained using a spade from five points by “S” type in each plot for the determination of WSAs. Composite bulk soil samples were collected from the same plots for the measurement of SOC and TN concentration in bulk soil.

**Laboratory method.** The distribution and stability of soil aggregates were measured according to the dry sieving and wet sieving method created by Elliott<sup>51</sup>. The specific dry sieving method was as follows. 200-gram samples of soil were passed through a series of six sieves (5, 2, 1, 0.5, 0.25, and 0.106 mm) to isolate seven aggregate size fractions and shaken them gently. Meanwhile, stones, roots and other impurities were eliminated in the soil samples of each size. Then the whole series of sieves were moved up and down for 7 minutes at 30 cycle min<sup>-1</sup> manually. Finally, the mass of dry sieve aggregates with different particle sizes was measured and weighed as  $W_{di}$ .

At the same time, the specific wet sieving method was as follows. 50-gram samples of soil were air-dried for 24 hours and evenly distributed over the nested sieve surfaces through a series of six sieves (5, 2, 1, 0.5, 0.25, and 0.106 mm) to isolate seven aggregate size fractions. The nest was set at the highest point when the oscillation cylinders were filled with distilled water. Soil samples were completely covered with water. To slake the air-dried soil, 1 L of distilled water was rapidly added to each cylinder until the soil sample and top screen were covered with water. The soils were submerged in water for 10 minutes before the start of the wet-sieving action. The apparatus specifications of oscillation time (10 minutes), stroke length (4 cm vertical) and frequency (30 cycle min<sup>-1</sup>) were held constant. Material remaining on each sieve was collected, dried at 60–80 degrees Celsius, and weighed as  $M_1$ . The water-stable aggregate distribution was based on the percentage of total mass in each aggregate fraction.

The soil particles remaining on the sieve were dried and weighed as  $M_2$ , (Eq. 1). The mass of each graded aggregate  $M_0$  was calculated from Eq. (1).

$$M_0 = M_1 - M_2 \quad (1)$$

The aggregate fraction in each sieve represented the water-stable aggregates (WSAs) according to class size.  $WSA_{>2\text{ mm}}$ ,  $WSA_{0.25-2\text{ mm}}$ , and  $WSA_{0.053-0.25\text{ mm}}$  were considered as the large macro-aggregates, small macro-aggregate, and micro-aggregates<sup>4,52,53</sup>, respectively.

$W_i$  is the proportion of each aggregate class in relation to the weight of the soil samples (Eq. 2). Wet-sieving is calculated from Eq. (3).

$$W_i = \frac{W_{di}}{200} \times 100\% \quad (2)$$

$$W_i = \frac{M_i}{50} \times 100\% \quad (3)$$

According to the data of each aggregate class obtained by the experiment,  $R_{0.25}$  means aggregates larger than 0.25 mm in diameter, the mean weight diameter (MWD)<sup>54</sup>, and geometric mean diameter (GMD)<sup>55</sup> are computed as Eqs. (4), (5), and (6), respectively.

$$R_{0.25} = \frac{M_{r>0.25}}{M_T} = 1 - \frac{M_{r<0.25}}{M_T} \quad (4)$$

$$MWD = \sum_{i=1}^n (\bar{X}_i W_i) \quad (5)$$

$$GMD = \exp\left(\frac{\sum_{i=1}^n W_i \lg \bar{X}_i}{\sum_{i=1}^n W_i}\right) \quad (6)$$

$X_i$  is the mean diameter of the class (mm).  $M(r < X_i)$  means the mass of aggregates, which are smaller than  $X_i$ .  $M_T$  means total mass of the aggregates.

The fractal dimension<sup>56</sup> is computed as Eq. 7:

$$(3-D) \lg\left(\frac{\bar{d}_i}{d_{\max}}\right) = \lg\left[\frac{W_{(\delta \leq \bar{d}_i)}}{W_0}\right] \quad (7)$$

Treatments	Aggregate size (percentage)						
	MSA <sub>&gt;5 mm</sub>	MSA <sub>2-5 mm</sub>	MSA <sub>1-2 mm</sub>	MSA <sub>0.5-1 mm</sub>	MSA <sub>0.25-0.5 mm</sub>	MSA <sub>0.106-0.25 mm</sub>	MSA <sub>&lt;0.106 mm</sub>
CC	45.02 ± 0.52 b	18.46 ± 0.32 d	11.70 ± 0.11 bc	15.01 ± 0.21 b	5.84 ± 0.05 b	2.07 ± 0.04 b	1.90 ± 0.04 a
SC	35.42 ± 0.48 c	21.91 ± 0.27 bc	13.58 ± 0.15 ab	17.44 ± 0.19 a	6.49 ± 0.07 a	2.92 ± 0.03 a	2.26 ± 0.07 a
CS	43.58 ± 0.67 b	20.31 ± 0.22 c	12.38 ± 0.13 bc	14.20 ± 0.08 b	5.31 ± 0.08 c	2.02 ± 0.02 b	2.20 ± 0.06 a
FC	34.68 ± 0.41 c	24.82 ± 0.16 a	14.63 ± 0.18 a	15.98 ± 0.18 ab	5.58 ± 0.06 bc	2.26 ± 0.03 b	2.06 ± 0.05 a
FS	50.22 ± 1.03 a	23.16 ± 0.18 ab	11.02 ± 0.09 c	11.81 ± 0.15 c	1.86 ± 0.03 d	0.76 ± 0.01 c	1.16 ± 0.02 b

**Table 1.** Distribution of soil mechanical stability aggregate size fractions (percentage) under different cropping systems. **Note:** CC, SC, CS, FC, and FS represent continuous corn, soybean-corn rotation, corn-soybean rotation, fallow-corn, and fallow-soybean treatments. MSA indicates mechanical stability aggregate. MSA<sub>>5 mm</sub>, MSA<sub>2-5 mm</sub>, MSA<sub>1-2 mm</sub>, MSA<sub>0.5-1 mm</sub>, MSA<sub>0.25-0.5 mm</sub>, MSA<sub>0.106-0.25 mm</sub> and MSA<sub><0.106 mm</sub> represent > 5 mm, 2–5 mm, 1–2 mm, 0.5–1 mm, 0.25–0.5 mm, 0.106–0.25 mm and <0.106 mm aggregate fractions in mechanical stability aggregate. Values are given as mean ± standard error (n = 9). Mean value ± SE in the same column followed by the different lowercase letters indicate significant differences among different treatments for each size of soil aggregates (L.S.D. test,  $P < 0.05$ ).

Take  $\lg\left[\frac{W(\delta \leq \bar{d}_i)}{W_0}\right]$  and  $\lg\left(\frac{\bar{d}_i}{d_{\max}}\right)$  as the horizontal and vertical axis, respectively. It draws the conclusion that 3-D is the slope of straight line between  $\lg\left(\frac{\bar{d}_i}{d_{\max}}\right)$  and  $\lg\left[\frac{W(\delta \leq \bar{d}_i)}{W_0}\right]$ . Soil fractal dimension is marked as D.

The formula for calculating the percentage of aggregates destruction and water-stable aggregates stability rate were calculated as seen Eqs. (8) and (9).

$$PAD = \frac{M_d - M_w}{M_d} \times 100\% \quad (8)$$

$$WSAR = \frac{WSA}{A} \times 100\% \quad (9)$$

where PAD is the percentage of aggregates destruction (%),  $M_d$  and  $M_w$  represent the aggregate mass fractions of dry sieve and wet sieve with >0.25 mm particle sizes, respectively. WSAR is the water-stable aggregates stability rate (%), WSA and A mean the water-stable aggregate weight (g) and mechanical stability aggregate weight (g), respectively.

Composite bulk soil samples for each treatments with three replications were also air dried at room temperature (22 degrees Celsius), passed through a 0.25 mm diameter sieve, and stored at room temperature prior to the analysis of SOC concentrations. SOC in the bulk soils and WSAs were determined by 0.8 mol L<sup>-1</sup> 1/6 K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> oxidation and FeSO<sub>4</sub> titration<sup>57</sup>. Soil available phosphorus and potassium (AP and AK) were measured by 0.5 mol L<sup>-1</sup> NaHCO<sub>3</sub> and molybdenum antimony anti-colorimetric method, and 1 mol L<sup>-1</sup> CH<sub>3</sub>COONH<sub>4</sub> extraction method, respectively<sup>57</sup>. The SOC stock calculated using the following Eq. (10)<sup>58</sup>.

$$M'_i = C_i \times SOC_i \times BD \times H \times 10^{-1} \quad (10)$$

where,  $M'_i$  means SOC stock of i-level aggregates (t hm<sup>-2</sup>),  $C_i$  and  $SOC_i$  mean relative mass fraction and SOC concentration of i-level aggregates, respectively. BD means soil bulk density in the 0–20 cm (g cm<sup>-3</sup>), H is the thickness of soil layer and we take it as 20 cm in our research.

**Statistical analysis.** All statistical analysis was carried out using SPSS (Statistical Package for Social Science) 20.0 for Windows and all graphs were drawn using Origin 7.5. Significant differences among treatments for MSAs, WSAs, SOC concentrations and stocks, as well as MWD, GMD, D, PAD, and WSAD were determined with one-way analysis of variance in the different treatments with the same aggregate sizes followed by the least significant difference (LSD) test at  $P < 0.05$  (n = 9). The correlations among the measured soil attributes were determined using stepwise regression analysis<sup>59</sup>.

## Results

**Size distribution of mechanical stability aggregates.** MSAs were mainly concentrated in MSA<sub>>5 mm</sub>, ranging from 35.4% in SC treatment to 50.2% in FC treatment (Table 1). The lowest aggregate content was found in the MSA<sub><0.106 mm</sub>, accounting for about 2%. The highest proportions in MSA<sub>>5 mm</sub>, MSA<sub>2-5 mm</sub>, and MSA<sub>1-2 mm</sub> were obtained in FS (50.2%), FC (24.8%), and FC (14.6%) treatments, respectively. Meanwhile, we were surprised to find that SC treatment documented the highest proportion in the MSA<sub>0.5-1 mm</sub> (17.4%), MSA<sub>0.25-0.5 mm</sub> (6.5%), MSA<sub>0.106-0.25 mm</sub> (2.9%), and MSA<sub><0.106 mm</sub> (2.3%). On the other hand, the lowest proportions in the MSA<sub>>5 mm</sub> and MSA<sub>2-5 mm</sub> were identified in FC treatment (34.7%) and CC treatment (18.5%), respectively. While, the FS treatment had the lowest proportions in the MSA<sub>1-2 mm</sub> (11%), MSA<sub>0.5-1 mm</sub> (11.8%), MSA<sub>0.25-0.5 mm</sub> (1.9%), MSA<sub>0.106-0.25 mm</sub> (0.8%), and MSA<sub><0.106 mm</sub> (1.2%).

Compared with CC treatment, SC treatment significantly ( $P < 0.05$ ) increased the proportion in MSA<sub>2-5 mm</sub>, MSA<sub>0.5-1 mm</sub>, MSA<sub>0.25-0.5 mm</sub>, and MSA<sub>0.106-0.25 mm</sub> by 18.7%, 16.2%, 11.1%, and 41.1%, respectively. CS treatment only significantly ( $P < 0.05$ ) increased the proportion in MSA<sub>2-5 mm</sub> by 10%. However, FC treatment both

Treatments	Aggregate size (percentage)						
	WSA <sub>&gt;5 mm</sub>	WSA <sub>2-5 mm</sub>	WSA <sub>1-2 mm</sub>	WSA <sub>0.5-1 mm</sub>	WSA <sub>0.25-0.5 mm</sub>	WSA <sub>0.106-0.25 mm</sub>	WSA <sub>&lt;0.106 mm</sub>
CC	2.04 ± 0.05 c	3.90 ± 0.04 d	2.90 ± 0.02 e	8.94 ± 0.37 d	20.59 ± 1.15 a	30.46 ± 1.04 a	31.17 ± 1.13 a
SC	4.31 ± 0.06 a	6.27 ± 0.07 b	3.85 ± 0.03 d	14.88 ± 0.62 c	19.99 ± 0.83 a	22.05 ± 0.94 c	28.66 ± 1.10 ab
CS	2.07 ± 0.03 c	6.45 ± 0.06 b	8.83 ± 0.06 b	20.70 ± 0.48 a	13.85 ± 0.45 b	26.02 ± 0.66 b	22.09 ± 0.57 c
FC	0.27 ± 0.01 d	5.80 ± 0.04 c	9.27 ± 0.07 a	13.82 ± 0.62 c	18.55 ± 0.64 a	23.60 ± 0.54 c	28.69 ± 1.09 ab
FS	3.39 ± 0.05 b	7.57 ± 0.05 a	7.01 ± 0.08 c	16.77 ± 0.54 b	20.49 ± 0.59 a	18.72 ± 0.88 d	26.05 ± 0.61 b

**Table 2.** Distribution of soil water-stable aggregate size fractions (percentage) under different cropping systems. **Note:** CC, SC, CS, FC, and FS represent continuous corn, soybean-corn rotation, corn-soybean rotation, fallow-corn, and fallow-soybean treatments. WSA indicates mechanical stability aggregate. WSA<sub>>5 mm</sub>, WSA<sub>2-5 mm</sub>, WSA<sub>1-2 mm</sub>, WSA<sub>0.5-1 mm</sub>, WSA<sub>0.25-0.5 mm</sub>, WSA<sub>0.106-0.25 mm</sub> and WSA<sub><0.106 mm</sub> represent > 5 mm, 2–5 mm, 1–2 mm, 0.5–1 mm, 0.25–0.5 mm, 0.106–0.25 mm and < 0.106 mm aggregate fractions in water-stable aggregate. Values are given as mean ± standard error (n = 9). Mean value ± SE in the same column followed by the different lowercase letters indicate significant differences among different treatments for each size of soil aggregates (L.S.D. test,  $P < 0.05$ ).

Treatments	MWD		GMD		D		PAD (%)	WSAR (%)
	MSAs	WSAs	MSAs	WSAs	MSAs	WSAs		
CC	3.79 ± 0.03 c	0.55 ± 0.02 c	1.00 ± 0.02 a	0.21 ± 0.01 c	2.20 ± 0.01 a	2.86 ± 0.02 a	60.05 ± 0.82 a	40.03 ± 0.54 e
SC	4.30 ± 0.02 b	0.84 ± 0.03 a	1.00 ± 0.01 a	0.28 ± 0.01 b	2.17 ± 0.02 b	2.83 ± 0.01 a	48.02 ± 0.97 c	51.56 ± 0.46 c
CS	4.84 ± 0.02 a	0.78 ± 0.02 b	1.01 ± 0.02 a	0.33 ± 0.02 a	2.03 ± 0.01 c	2.79 ± 0.02 a	45.82 ± 0.75 d	54.47 ± 0.62 b
FC	3.83 ± 0.04 c	0.59 ± 0.01 c	1.00 ± 0.01 a	0.27 ± 0.03 b	2.19 ± 0.03 a	2.83 ± 0.03 a	50.14 ± 0.66 b	49.24 ± 0.58 d
FS	4.34 ± 0.05 b	0.87 ± 0.04 a	1.00 ± 0.01 a	0.32 ± 0.02 a	2.17 ± 0.02 b	2.80 ± 0.02 a	43.69 ± 0.81 e	57.92 ± 0.41 a

**Table 3.** MWD, GMD, D, PAD and WSAR under different cropping systems in MSAs and WSAs. **Note:** CC, SC, CS, FC, and FS represent continuous corn, soybean-corn rotation, corn-soybean rotation, fallow-corn, and fallow-soybean treatments. Soil properties of MWD, GMD, D, PAD, and WSAR indicate mean weight diameter, geometric mean diameter, fractal dimension, percentage of aggregates destruction, and water-stable aggregates stability rate, respectively. MSAs and WSAs indicate mechanical stability aggregates and water-stable aggregates. Values are given as mean ± standard error (n = 9). Mean value ± SE in the same column followed by the different lowercase letters indicate significant differences among different treatments (L.S.D. test,  $P < 0.05$ ).

significantly ( $P < 0.05$ ) increased the proportion in MSA<sub>2-5 mm</sub> and MSA<sub>1-2 mm</sub> by 34.5% and 25%. Simultaneously, FS treatment significantly ( $P < 0.05$ ) increased the proportion in MSA<sub>2-5 mm</sub> and MSA<sub>>5 mm</sub> by 11.6% and 25.5%, respectively.


**Size distribution of water-stable aggregates.** After the two-year experiment, we found that among the five treatments, the highest proportions in the WSA<sub>>5 mm</sub>, WSA<sub>2-5 mm</sub>, WSA<sub>1-2 mm</sub> and WSA<sub>0.5-1 mm</sub> were obtained in SC (4.3%), FS (7.6%), FC (9.3%), and CS (20.7%) treatments, respectively (Table 2). However, the lowest proportion in the WSA<sub>>5 mm</sub> was the FC treatment (0.3%), and The lowest proportions in WSA<sub>2-5 mm</sub>, WSA<sub>1-2 mm</sub> and WSA<sub>0.5-1 mm</sub> were all found in CC treatment with 3.9%, 2.9%, and 8.9%, respectively. We were surprised to note that the CC treatment documented the highest proportion in the WSA<sub>0.25-0.5 mm</sub> (20.6%), WSA<sub>0.106-0.25 mm</sub> (30.5%), and WSA<sub><0.106 mm</sub> (31.2%). On the other hand, the lowest proportions in the WSA<sub>0.25-0.5 mm</sub>, WSA<sub>0.106-0.25 mm</sub>, and WSA<sub><0.106 mm</sub> were identified in CS (13.9%), FS (18.7%), and CS (22.1%) treatments, respectively. The FS treatment showed the highest proportion in the WSA<sub>>0.25 mm</sub> (55.2%), followed by CS (51.9%), SC (49.3%), and FC (47.7%) treatments (Table 2). The CC treatment only had 38.4% in WSA<sub>>5 mm</sub>.

Compared with the CC treatment, the SC and FS treatments significantly ( $P < 0.05$ ) increased the proportion in WSA<sub>>5 mm</sub> by 111.3% and 66.2%, respectively. The SC, CS, FC, and FS treatments all significantly ( $P < 0.05$ ) increased the proportion of WSA<sub>2-5 mm</sub> (60.8%, 65.4%, 48.7% and 94.1%), WSA<sub>1-2 mm</sub> (32.8%, 204.5%, 219.7% and 141.7%), and WSA<sub>0.5-1 mm</sub> (66.4%, 131.5%, 54.6% and 87.6%), respectively. However, all the treatments significantly ( $P < 0.05$ ) decreased the proportion of WSA<sub>0.106-0.25 mm</sub>. The proportion of WSA<sub><0.106 mm</sub> was significantly ( $P < 0.05$ ) decreased in the CS and FS treatments by 29.1% and 16.4%, respectively.

**MWD, GMD, D, PAD, and WSAR within soil aggregates.** From the point of MSAs, the highest MWD and lowest D were both obtained in the CS treatment with 4.84 and 2.03, respectively (Table 3). Meantime, the lowest MWD and highest D were both found in the CC treatment, respectively (Table 3). In comparison to CC treatments, SC, CS, and FS treatments all significantly ( $P < 0.05$ ) increased MWD with 13.5%, 27.7%, and 14.5%; and decreased D with 1.4%, 7.7%, and 1.4%, respectively (Table 3). However, no significant ( $P < 0.05$ ) difference was found for GMD values among the five treatments.

From the point of WSAs, the highest MWD and GMD were observed in the FS and CS treatments, respectively, while the lowest MWD and GMD were both characterized in the CC treatment (Table 3). Compared with the CC treatment, the SC, CS, and FS treatments significantly ( $P < 0.05$ ) increased MWD and GMD by 52.7%,





**Figure 1.** Soil organic carbon (SOC) concentration in the soil aggregate sizes under different treatments. Values are given as mean  $\pm$  standard error ( $n = 9$ ). Different lowercase letters indicate significant differences among different treatments for each size of soil aggregates (L.S.D. test,  $P < 0.05$ ). CC, SC, CS, FC, and FS represent continuous corn, soybean-corn rotation, corn-soybean rotation, fallow-corn, and fallow-soybean treatments, respectively.

41.8%, and 58.2%; and by 33.3%, 57.1%, and 52.4%, respectively. The maximum and minimum D were obtained in the CC and CS treatments (Table 3). However, no significant ( $P < 0.05$ ) difference was observed for D values among the five treatments.


The PAD declined and WSAR increased in the order of  $CC < FC < SC < CS < FS$  treatment (Table 3). In comparison with CC treatment, the FC, SC, CS, and FS treatments significantly ( $P < 0.05$ ) decreased PAD and increased WSAR by 16.5%, 20%, 23.7%, and 27.2%, and by 23%, 28.8%, 36.1%, and 44.7%, respectively.

**SOC concentrations and stocks within water-stable aggregate sizes and bulk soils.** In general, compared with the CC treatment, SC treatment increased SOC concentration in each water-stable aggregate size and bulk soils, while, the FC treatment decreased SOC concentration in the WSAs (Fig. 1) and bulk soils (Fig. 2a). Specifically, the SC treatment had the highest SOC concentration in the  $WSA_{1-2\text{ mm}}$  with  $23.38\text{ g kg}^{-1}$  (Fig. 1). The highest SOC concentrations in the  $WSA_{>5\text{ mm}}$ ,  $WSA_{0.5-1\text{ mm}}$ , and  $WSA_{<0.106\text{ mm}}$  were obtained by the CS treatment with  $24.02\text{ g kg}^{-1}$ ,  $23.61\text{ g kg}^{-1}$ , and  $15.60\text{ g kg}^{-1}$ , respectively (Fig. 1). Meanwhile, the highest SOC concentrations in the  $WSA_{2-5\text{ mm}}$ ,  $WSA_{0.25-0.5\text{ mm}}$ , and  $WSA_{0.106-0.25\text{ mm}}$ , WSAs were obtained in the FS treatment with  $22.13\text{ g kg}^{-1}$ ,  $21.33\text{ g kg}^{-1}$ , and  $20.58\text{ g kg}^{-1}$ , respectively (Fig. 1).

The SOC concentration in the  $WSA_{2-5\text{ mm}}$ ,  $WSA_{0.5-1\text{ mm}}$ ,  $WSA_{0.25-0.5\text{ mm}}$ ,  $WSA_{0.106-0.25\text{ mm}}$ , and  $WSA_{<0.106\text{ mm}}$  were all significantly ( $P < 0.05$ ) increased by 15.2%, 26.2%, 20.7%, 41.6%, and 28.7% from SC treatment; by 11%, 35.6%, 24.5%, 34.2%, and 33.8% from CS treatment; and by 20.2%, 25.8%, 29.7%, 43.5%, and 27.4% from FS treatment in comparison with the CC treatment, respectively (Fig. 1). Simultaneously, compared with CC treatment, CS and FS treatments both significantly ( $P < 0.05$ ) increased SOC concentration in the  $WSA_{>5\text{ mm}}$  by 22.4% and 19.4%, as well as SC and CS treatments both significantly ( $P < 0.05$ ) increased SOC concentration in the  $WSA_{1-2\text{ mm}}$  by 21.4% and 14.1%, respectively (Fig. 1). In addition, the CS and FS treatments both significantly ( $P < 0.05$ ) increased SOC concentration by 17.6% and 14.1% compared with the CC treatment in bulk soils (Fig. 2a).

Across all treatments, the SOC stock in the seven aggregates' sizes showed a similar tendency in the SOC concentration, although the bulk density differed a little among the treatments (Fig. 2b). Bulk density for the five treatments was in the range of  $1.12\text{--}1.18\text{ g cm}^{-3}$  (Fig. 2b). CS treatment had the highest SOC stock in the  $WSA_{>5\text{ mm}}$ ,  $WSA_{0.5-1\text{ mm}}$ , and  $WSA_{<0.106\text{ mm}}$  with  $8.89\text{ t hm}^{-2}$ ,  $8.59\text{ t hm}^{-2}$ , and  $3.75\text{ t hm}^{-2}$ , respectively (Fig. 3). While the FS treatment had the highest SOC stock in the  $WSA_{2-5\text{ mm}}$  ( $7.64\text{ t hm}^{-2}$ ),  $WSA_{0.25-0.5\text{ mm}}$  ( $7.10\text{ t hm}^{-2}$ ), respectively (Fig. 3). Furthermore, the SC treatment demonstrated the biggest SOC stock in the  $WSA_{1-2\text{ mm}}$  ( $8.80\text{ t hm}^{-2}$ ) and  $WSA_{0.106-0.25\text{ mm}}$  ( $6.64\text{ t hm}^{-2}$ ), respectively (Fig. 3). Except for  $WSA_{0.106-0.25\text{ mm}}$  and  $WSA_{<0.106\text{ mm}}$ , the FC treatment documented the lowest SOC stock in all five other aggregate sizes.

Furthermore, the SOC stock in the  $WSA_{2-5\text{ mm}}$ ,  $WSA_{1-2\text{ mm}}$ ,  $WSA_{0.5-1\text{ mm}}$ ,  $WSA_{0.25-0.5\text{ mm}}$ ,  $WSA_{0.106-0.25\text{ mm}}$ , and  $WSA_{<0.106\text{ mm}}$  from the SC treatment, the stock in the  $WSA_{>5\text{ mm}}$ ,  $WSA_{0.5-1\text{ mm}}$ ,  $WSA_{0.25-0.5\text{ mm}}$ ,  $WSA_{0.106-0.25\text{ mm}}$ , and  $WSA_{<0.106\text{ mm}}$  from the CS treatment were all significantly ( $P < 0.05$ ) increased by 8%, 19.6%, 29.5%, 18.3%, 63.1%, and 34.7%; and by 16.7%, 43.1%, 20.6%, 40.2%, and 39.4% compared with CC treatment, respectively (Fig. 3). Similarly, the SOC stock in the  $WSA_{0.106-0.25\text{ mm}}$  and  $WSA_{<0.106\text{ mm}}$  from the FC treatment, and the stock in the  $WSA_{>5\text{ mm}}$ ,  $WSA_{2-5\text{ mm}}$ ,  $WSA_{0.5-1\text{ mm}}$ ,  $WSA_{0.25-0.5\text{ mm}}$ ,  $WSA_{0.106-0.25\text{ mm}}$ , and  $WSA_{<0.106\text{ mm}}$  from the FS treatment were also significantly ( $P < 0.05$ ) increased by 10.7%, and 23.8%; and by 12.3%, 13.8%, 24.7%, 32.4%, 62.3%, and 27.9% in comparison with the CC treatment, respectively (Fig. 3).



**Figure 2.** SOC concentration (a) and bulk density (b) in bulk soils under different treatments. SOC indicates soil organic carbon. CC, SC, CS, FC, and FS represent continuous corn, soybean-corn rotation, corn-soybean rotation, fallow-corn, and fallow-soybean treatments, respectively (n = 9).


On the whole, the SOC concentration and stocks in WSAs both accounted for almost half of the total in small macro-aggregates ( $WSA_{0.25-2\text{ mm}}$ ) under the five cropping systems (Fig. 4a,b). For instance, SOC concentrations and stocks were 45.6% and 48.1% in the SC treatments; 45.5% and 47.6% in the CS treatments; and 44.1% and 44.7% in the FS treatment (Fig. 4a,b).

**Correlations among measured soil attributes and parameters.** Linear regression models between measured soil attributes and parameters, obtained by stepwise regression analysis, were listed in Table 4. The data indicated that the MWD in WSAs was significantly and positively correlated to SOC concentration in  $WSA_{0.106-0.25\text{ mm}}$ , as shown in Eq. (10) ( $R^2 = 0.942$ ;  $P = 0.006$ ). Similarly, significant and positive correlation was obtained between SOC concentrations in bulk soils and SOC in the  $WSA_{>5\text{ mm}}$ , as shown in Eq. (11) ( $R^2 = 0.988$ ;  $P = 0.001$ ). Simultaneously, significant correlations were obtained between the GMD and D in WSAs and WSAs of each particle size. In specification, significant and positive correlations were exhibited between GMD in WSAs and  $WSA_{2-5\text{ mm}}$ , and  $WSA_{0.5-1\text{ mm}}$ , as shown in Eq. (12) ( $R^2 = 0.936$ ;  $P = 0.007$ ). However, a significant and negative correlation was demonstrated between D in WSAs and  $WSA_{0.5-1\text{ mm}}$ , as shown in Eq. (13) ( $R^2 = 0.940$ ;  $P = 0.006$ ).

## Discussion

In the perspective of size distribution, the proportion of macro-aggregates with MSAs under different cropping systems in our experiment were significantly higher than that of WSAs, which may be due to the fact that the soil is more and more compacted with the increasing of agricultural mechanization operations<sup>60-62</sup>.

Water stable aggregates were the indicator of soil anti-disintegration<sup>63</sup>, and the aggregate sizes elucidated the effects of management on soil structural stability. Relevant studies demonstrated that soil aggregates were



**Figure 3.** Soil organic carbon (SOC) stocks in the soil aggregate sizes under different treatments. Values are given as mean  $\pm$  standard error ( $n = 9$ ). Different lowercase letters indicate significant differences among different treatments for each size of soil aggregates (L.S.D. test,  $P < 0.05$ ). CC, SC, CS, FC, and FS represent continuous corn, soybean-corn rotation, corn-soybean rotation, fallow-corn, and fallow-soybean treatments, respectively.

divided into macro-aggregates ( $WSA_{>0.25 \text{ mm}}$ ) and micro-aggregates ( $WSA_{<0.25 \text{ mm}}$ )<sup>53,64</sup>. Six *et al.*<sup>65</sup> believed that macro-aggregates were the best structures in the soil, and the higher the content, the better agglomeration and stability of soil aggregates.

In the current study, the proportion of  $WSA_{>0.25 \text{ mm}}$  was in the order of  $FS > CS > SC > FC > CC$  treatments, which suggested that fallow and rotation were more conducive to the formation of  $WSA_{>0.25 \text{ mm}}$  than continuous corn. Higher microbial biomass in the rhizosphere with soybean or rotation might be responsible for the formation of macro-aggregates<sup>66</sup>. Alternatively, Nakamoto and Suzuki<sup>67</sup> held a different opinion. They indicated that the root growth of soybean illustrated a stronger destructive effect on macro-aggregates in the soil. Planting soybean could reduce the number of macro-aggregates compared with corn even though the root density of soybean was relatively small. Small macro-aggregates ( $WSA_{0.25-2 \text{ mm}}$ ) were found to be the leading size fractions in the SOC concentrations and stocks in our study, which was consistent with other findings<sup>68,69</sup>.

In the present study, fallow-corn and soybean-corn rotations were significantly greater than continuous corn in MWD and GMD, while D was smaller in the CS and FS treatments than the CC treatment. Our findings indicated that fallow-corn and soybean-corn rotations had made the soil aggregate structure more stable. Soil aggregates stability is expressed by MWD of the size range, which is proportional to the amount of larger WSAs<sup>70</sup>. Devine *et al.*<sup>71</sup> in Horseshoe Bend of USA showed that no-tillage increased MSAs. Furthermore, MWD in WSAs compared with conventional tillage, MWD increased with the increasing of soil depths.


Many studies believed that MWD was related to soil organic matter content. Das *et al.*<sup>70</sup> found that the increased amounts of macro-aggregates ( $WSA_{>2 \text{ mm}}$ ) for applying green manure, cereal residues, and farmyard manure on wheat and rice were associated with MWD. The correlation of strong linearity between MWD and SOC suggested that SOC content in the  $WSA_{0.106-0.25 \text{ mm}}$  played a major role in soil aggregates stability.

In this study, the four treatments of soybean-corn, corn-soybean, fallow-corn, fallow-soybean all had lower PAD and higher WSAR than that of continuous corn treatment, which indicated that the effect of rotation and intermittent fallow practices on soil aggregates stability was more favorable than planting crops. Six *et al.*<sup>64</sup> had shown that tillage accelerated the renewal rate of macro-aggregates, which was not conducive to the formation of micro-aggregates in macro-aggregates. The reduction of tillage disturbance increased the stability of soil aggregates<sup>19,72,73</sup>, which may be because the frequent tillage of farmland destroys the soil particle structure, increases the soil aeration, and deteriorates the protection of soil particles, resulting in loose soil structure and increase damage to the soil structure<sup>74</sup>.

Our experiment found that the stability of soil aggregates treated by corn-soybean and soybean-corn in both legumes and gramineae rotation system was higher than continuous corn treatment, which may be because that the rotation of legumes and gramineae plants could rapidly increase the storage of soil organic carbon and promote the formation of macro-aggregates and stabilization of micro-aggregates<sup>75</sup>. We conclude that the rotation system can increase the energy required for microbial life activities in the soil, produce cementing substances that form soil aggregates, weaken the destruction of soil aggregates, and reduce the damage to soil structure.

In our research, MWD and D in MSAs demonstrated that CS treatment had the best soil aggregate stability. However, MWD in WSAs as well as PAD and WSAR showed that FS treatment had the highest soil aggregate stability. GMD in WSAs showed the highest stability was corn-soybean treatment, but our study found that corn-soybean and fallow-soybean treatments were not significantly different. These five indicators indicate that the effects of soil





**Figure 4.** Proportion of SOC (a) and SOC stock (b) in the soil aggregate sizes under different treatments. SOC indicates soil organic carbon. CC, SC, CS, FC, and FS represent continuous corn, soybean-corn rotation, corn-soybean rotation, fallow-corn, and fallow-soybean treatments, respectively (n = 9).

Indicators in Y	Indicators in X	Regression model (Equation number)	R <sup>2</sup>	F	P
Aggregate stability (MWD) in water-stable aggregates	SOC in water-stable aggregate particle sizes	$Y_{MWD} = 0.069 \text{ SOC}_{\text{WSA } 0.106-0.25 \text{ mm}} + 0.025$ (10)	0.942	49.123	0.006
SOC in bulk soils	SOC in water-stable aggregate particle sizes	$Y_{\text{SOC}} = 1.814 \text{ SOC}_{\text{WSA} > 5 \text{ mm}} + 2.036$ (11)	0.988	255.224	0.001
Aggregate stability (GMD) in water-stable aggregates	water-stable aggregate particle sizes	$Y_{\text{GMD}} = 0.014 \text{ WSA}_{2-5 \text{ mm}} + 0.007 \text{ WSA}_{0.5-1 \text{ mm}} + 0.088$ (12)	0.936	43.656	0.007
Aggregate stability (D) in water-stable aggregates	water-stable aggregate particle sizes	$Y_{\text{D}} = -0.006 \text{ WSA}_{0.5-1 \text{ mm}} + 2.916$ (13)	0.940	46.777	0.006

**Table 4.** Relationship among soil aggregate stability in WSAs and SOC in WSAs, soil aggregate stability in WSAs and WSAs, as well as SOC concentration in bulk soils and WSAs of all water-stable aggregate size fractions under five treatments. Note: Eq. (10)  $Y_{MWD}$  MWD value in WSAs;  $\text{SOC}_{\text{WSA } 0.106-0.25 \text{ mm}}$  SOC in water-stable aggregate 0.106–0.25 mm size fraction. Eq. (11)  $Y_{\text{SOC}}$  SOC in bulk soils;  $\text{SOC}_{\text{WSA} > 5 \text{ mm}}$  SOC in water-stable aggregate > 5 mm size fraction. Eq. (12)  $Y_{\text{GMD}}$  GMD value in WSAs;  $\text{WSA}_{2-5 \text{ mm}}$  water-stable aggregate 2–5 mm size fraction;  $\text{WSA}_{0.5-1 \text{ mm}}$  water-stable aggregate 0.5–1 mm size fraction. Eq. (13)  $Y_{\text{D}}$  D value in WSAs;  $\text{WSA}_{0.5-1 \text{ mm}}$  water-stable aggregate 0.5–1 mm size fraction. MWD, GMD, D, SOC and WSAs represent mean weight diameter, geometric mean diameter, fractal dimension, soil organic carbon and water-stable aggregates, n = 45.

aggregate stability are basically similar, and they can be mutually verified with each other. Therefore, we concluded that CS and FS treatment had higher soil aggregate stability combining these five indicators.

The SOC contents in soil aggregates of each particle size is a microscopic characterization between soil organic matter balance and mineralization rate, which has dual significance in soil fertility and soil carbon sink<sup>76</sup>. Our research found that SOC concentration exhibited the “M” type among the seven particle sizes in that the WSA<sub>1–2 mm</sub> and WSA<sub>0.106–0.25 mm</sub>, had higher SOC concentration in the five different cropping systems. Furthermore, through the regression models, we obtained that WSA<sub>2–5 mm</sub> and WSA<sub>0.5–1 mm</sub> dominated the primary particle sizes in soil structure stability. Therefore, we speculated that the small macro-aggregates (WSA<sub>0.25–2 mm</sub>) had larger SOC concentration accumulation and higher soil aggregates stability. These results corresponded with Puge’s viewpoint that macro-aggregates were a source of organic carbon enrichment<sup>77,78</sup>. Tisdall and Oades<sup>79</sup> proposed that the formation and stability of micro-aggregates and macro-aggregates were interrelated processes. A wealth of research had manifested that macro-aggregates composed of organic binding agents<sup>80–84</sup>. Consequently, the protective mechanism of macro-aggregates on SOC was better than micro-aggregates<sup>85,86</sup>.

However, some researchers had the distinctive standpoints, for instance, Christensen<sup>87</sup>, De Jonge *et al.*<sup>88</sup>, and Li *et al.*<sup>89</sup> research demonstrated that organic carbon was mainly distributed in micro-aggregates (WSA<sub><0.25 mm</sub>) and that organic carbon contents increased with the decrease of aggregate particle size’s decrease. Li *et al.*<sup>69</sup> found that organic carbon is distributed in a “V” shape in aggregates as the result of the organic carbon contents of aggregates  $d > 2 \text{ mm}$  and  $d < 0.25 \text{ mm}$  are high.

In our study, CS and FS treatments did show more C accumulation compared with CC treatment in bulk soils, which might be beneficial to the formation of soil structure and the enhancement of soil structure stability<sup>90</sup>. Alternatively, the formation of aggregates affects the decomposition of SOC<sup>4</sup>. Huang *et al.*<sup>91</sup> believed that the factors affecting soil aggregates all influenced soil carbon, while the quantity and quality of soil carbon were closely related to aggregates. Meng *et al.*<sup>92</sup> showed that the main reason for the decline of aggregate stability and the decrease of water-stable aggregates were the reduction of soil organic matter.

It was interesting to illustrate in this study that the SOC concentration in fallow-soybean treatment was higher than continuous corn, and fallow-corn treatments in the WSA<sub>>5 mm</sub>, WSA<sub>2–5 mm</sub>, WSA<sub>0.25–0.5 mm</sub>, and WSA<sub>0.106–0.25 mm</sub>. This might be due to the accelerated mineralization of SOC in the soil with corn stalk, which acts as a cementing material to facilitate the formation of macro-aggregates<sup>93</sup>.

## Conclusions

Small macro-aggregates (WSA<sub>0.25–2 mm</sub>) dominated the concentration of SOC in the five cropping systems. SOC concentration in the WSA<sub>0.106–0.25 mm</sub>, WSA<sub>2–5 mm</sub>, and WSA<sub>0.5–1 mm</sub> had a dominant effect on aggregate stability as well as SOC in WSA<sub>>5 mm</sub> affected SOC concentration in bulk soils. Corn-soybean and fallow-soybean can increase macro-aggregates, enhance aggregate stability, and increase SOC concentrations in bulk soils and all aggregate sizes. These results are likely related to decrease the soil aeration with no tillage in farmland, and rapidly increasing the storage of SOC with the rotation of legumes and gramineae plants, and then reinforcing soil structure and promoting stabilization of soil aggregates.

Received: 27 June 2019; Accepted: 18 December 2019;

Published online: 14 January 2020

## References

1. Beare, M. H., Cabrera, M. L., Hendrix, P. F. & Coleman, D. C. Aggregate protected and unprotected organic matter pools in conventional and no-tillage soils. *Soil Sci. Soc. Am. J.* **58**, 787–795 (1994).
2. Martens, D. A. Plant residue biochemistry regulates soil carbon cycling and carbon sequestration. *Soil Biol. Biochem.* **32**, 361–369 (2000).
3. Lal, R. & Shukla, M. K. *Principles of soil physics*. (Marcel Dekker 2013).
4. Six, J., Paustian, K., Elliott, E. T. & Combrink, C. Soil structure and organic matter: I. Distribution of aggregate-size classes and aggregate-associated carbon. *Soil Sci. Soc. Am. J.* **64**, 681–689 (2000).
5. Albalasmeh, A. A., Berli, M., Shafer, D. S. & Ghezzehei, T. A. Degradation of moist soil aggregates by rapid temperature rise under low intensity fire. *Plant Soil* **362**, 335–344 (2013).
6. Gupta, V. V. S. R. & Germida, J. J. Soil aggregation: Influence on microbial biomass and implications for biological processes. *Res. J. Soil. Biol.* **80**, 1–7 (2014).
7. Trivedi, P. *et al.* Soil aggregate size mediates the impacts of cropping regimes on soil carbon and microbial communities. *Soil Biol. Biochem.* **91**, 169–181 (2015).
8. Cates, A. M., Ruark, M. D., Hedtcke, J. L. & Posner, J. L. Long-term tillage, rotation and perennialization effects on particulate and aggregate soil organic matter. *Soil Tillage Res.* **155**, 371–380 (2016).
9. Bronick, C. J. & Lal, R. Soil structure and management: a review. *Geoderma* **124**, 3–22 (2005).
10. Angers, D. A. Changes in soil aggregation and organic carbon under corn and alfalfa. *Soil Sci. Soc. Am. J.* **56**, 1244–1249 (1992).
11. Xie, H. T. *et al.* Long-term manure amendments reduced soil aggregate stability via redistribution of the glomalin-related soil protein in macroaggregates. *Sci. Rep.* **5**, 14687 (2015).
12. Xie, J. Y., Yang, Y., Zhang, S. L., Sun, B. H. & Yang, X. Y. Soil aggregation and aggregate associated organic carbon and total nitrogen under long-term contrasting soil management regimes in loess soil. *J. Integr. Agric.* **14**, 2405–2416 (2015).
13. Zhang, J. G. *et al.* Effects of straw incorporation on soil nutrients, enzymes, and aggregate stability in tobacco fields of China. *Sustainability* **8**, 710 (2016).
14. Zhang, S. L., Wang, R. J., Yang, X. Y., Sun, B. H. & Li, Q. H. Soil aggregation and aggregating agents as affected by long term contrasting management of an Anthrosol. *Sci. Rep.* **6**, 39107 (2016).
15. Onweremadu, E. U., Onyia, V. N. & Anikwe, M. A. N. Carbon and nitrogen distribution in water-stable aggregates under two tillage techniques in Fluvisols of Owerri area, southeastern Nigeria. *Soil Tillage Res.* **97**, 195–206 (2007).
16. Durigan, M. R. *et al.* Soil organic matter responses to anthropogenic forest disturbance and land use change in the Eastern Brazilian Amazon. *Sustainability* **9**, 379 (2017).
17. Cambardella, C. A. & Elliott, E. T. Carbon and nitrogen distribution in aggregates from cultivated and native grassland soils. *Soil Sci. Soc. Am. J.* **57**, 1071–1076 (1993).
18. Novara, A. *et al.* Litter contribution to soil organic carbon in the processes of agriculture abandon. *Solid Earth* **6**, 425–432 (2015).

19. Qi, Y. C., Wang, Y. Q., Liu, J., Yu, X. S. & Zhou, C. J. Comparative study on composition of soil aggregates with different land use patterns and several kinds of soil aggregate stability index. *Trans CSAE* **27**, 340–347 (in Chinese with English abstract) (2011).
20. Nimmo, J. R. & Perkins, K. S. Method of soil analysis. *Aggregates stability and size distribution. Part 4-physical methods*. 317–328. (E-Publishing Inc. Soil Sci. Soc. Am. J. 2002).
21. Castrignano, A. & Stelluti, M. Fractal geometry and geostatistics for describing the field variability of soil aggregation. *J. Agric. Eng. Res.* **73**, 13–18 (1999).
22. Nweke, I. A. & Nnabude, P. C. Aggregate stability of four soils as evaluated by different indices. *J. Exp. Biol. Agric. Sci.* **3**, 246–252 (2015).
23. Ahmadi, A., Neyshabouri, M. R., Rouhipour, H. & Asadi, H. Fractal dimension of soil aggregates as an index of soil erodibility. *J. Hydrol.* **400**, 305–311 (2011).
24. Li, J., Han, W. C., Chen, C., Ye, S. L. & Li, J. B. Effects of land use types on soil aggregate characteristics in hilly-gully region of Loess Plateau. *J. Soil Water Conserv.* **31**, 248–253+259 (in Chinese with English abstract) (2017).
25. Olsen, R. J., Hensler, R. F., Attoe, O. J., Witzel, S. A. & Peterson, L. A. Fertilizer nitrogen and crop rotation in relation to movement of nitrate nitrogen through soil profiles. *Soil Sci. Soc. Am. J.* **34**, 448–452 (1970).
26. Vance, C. P. Symbiotic nitrogen fixation and phosphorus acquisition. Plant nutrition in a world of declining renewable resources. *Plant Physiol.* **127**, 390–397 (2001).
27. Brisson, N. *et al.* Why are wheat yields stagnating in Europe? A comprehensive data analysis for France. *Field Crops Res.* **119**, 0–212 (2010).
28. Jensen, E. S. *et al.* Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries. a review. *Agron. Sustainable Dev.* **32**, 329–364 (2012).
29. Drury, C. F. *et al.* Impacts of 49–51 years of fertilization and crop rotation on growing season nitrous oxide emissions, nitrogen uptake and corn yields. *Can. J. Soil Sci.* **94**, 421–433 (2014).
30. Gan, Y. *et al.* Diversifying crop rotations with pulses enhances system productivity. *Sci. Rep.* **5**, 14625 (2015).
31. Samarajeewa, K. B. D. P., Horiuchi, T. & Oba, S. Effect of Chinese milk vetch (*Astragalus sinicus* L.) as a cover crop on weed control, growth and yield of wheat under different tillage systems. *Plant Prod. Sci.* **8**, 79–85 (2005).
32. Orr, C. H., James, A., Leifert, C., Cooper, J. M. & Cummings, S. P. Diversity and activity of free-living nitrogen-fixing bacteria and total bacteria in organic and conventionally managed soils. *Appl. Environ. Microbiol.* **77**, 911–919 (2011).
33. Zou, W. X., Si, B. C., Han, X. Z. & Jiang, H. The effect of long-term fertilization on soil water storage and water deficit in the Black Soil Zone in northeast China. *Can. J. Soil Sci.* **92**, 439–448 (2012).
34. Wu, Q. & Xie, H. L. A review and implication of land fallow systems research. *J. Resour. Ecol.* **8**, 223–231 (2017).
35. Landgraf, D., Böhm, C. & Makeschin, F. Dynamic of different C and N fractions in a Cambisol under five year succession fallow in Saxony (Germany). *J. Plant Nutr. Soil Sci.* **166**, 319–325 (2003).
36. Ribaud, M. O. Targeting the conservation reserve program to maximize water quality benefits. *Land Econ.* **65**, 320–332 (2010).
37. Szentandrási, S., Polasky, S., Berrens, R. & Leonard, J. Conserving biological diversity and the conservation reserve program. *Growth Change* **26**, 383–404 (2010).
38. Lant, C. L. Potential of the conservation reserve program to control agricultural Surface water pollution. *Environ. Manage.* **15**, 507–518 (1991).
39. Lal, R. Mechanized tillage systems effects on soil erosion from an alfisol in Watersheds cropped to maize. *Soil Tillage Res.* **4**, 349–360 (1984).
40. Yang, X. M. *et al.* Conservation tillage systems in North America and their significance for China. *Chin. J. Appl. Ecol.* **15**, 335–340 (2004).
41. Hazarika, S. *et al.* Effect of tillage system and straw management on organic matter dynamics. *Agron. Sustainable Dev.* **29**, 525–533 (2009).
42. Liu, X. B. *et al.* Overview of Mollisols in the world: distribution, land use and management. *Can. J. Soil Sci.* **92**, 383–402 (2012).
43. Liu, J. J. *et al.* Soil carbon content drives the biogeographical distribution of fungal communities in the black soil zone of northeast China. *Soil Biol. Biochem.* **83**, 29–39 (2015).
44. Wang, G. H. *et al.* Effects of fertilization on bacterial community structure and function in a black soil of Dehui region estimated by Biolog and PCR-DGGE methods. *Acta Ecologica Sinica.* **28**, 220–226 (2008).
45. Liu, X. B. *et al.* Soil degradation: a problem threatening the sustainable development of agriculture in Northeast China. *Plant Soil Environ.* **56**, 87–97 (2010).
46. Wang, Z. Q., Liu, B. Y., Wang, X. Y., Gao, X. F. & Liu, G. Erosion effect on the productivity of black soil in Northeast China. *Sci. Chi.* **52**, 1005–1021 (2009).
47. Noellemeyer, E., Frank, F., Alvarez, C., Morazzo, G. & Quiroga, A. Carbon contents an aggregation related to soil physical and biological properties under a land-use sequence in the semiarid region of central Argentina. *Soil Tillage Res.* **99**, 179–190 (2008).
48. Sodhi, G. P. S., Beri, V. & Benbi, D. K. Soil aggregation and distribution of carbon and nitrogen in different fractions under long-term application of compost in rice-wheat system. *Soil Tillage Res.* **103**, 412–418 (2009).
49. Fernandez, R., Quiroga, A., Zorati, C. & Noellemeyer, E. Carbon contents and respiration rates of aggregate size fractions under no-till and conventional tillage. *Soil Tillage Res.* **109**, 103–109 (2010).
50. Mikha, M. M. & Rice, C. W. Tillage and manure effects on soil and aggregate-Associated carbon and nitrogen. *Soil Sci. Soc. Am. J.* **68**, 809–816 (2004).
51. Elliott, E. T. Aggregate structure and carbon, nitrogen, and phosphorus in native and cultivated soils. *Soil Sci. Soc. Am. J.* **50**, 627–633 (1986).
52. Sainju, U. M., Caesar-TonThat, T. & Jabro, J. D. Carbon and nitrogen fractions in dryland soil aggregates affected by long-term tillage and cropping sequence. *Soil Sci. Soc. Am. J.* **73**, 1488–1495 (2009).
53. Xiao, S. S., Zhang, W., Ye, Y. Y., Zhao, J. & Wang, K. L. Soil aggregate mediates the impacts of land uses on organic carbon, total nitrogen, and microbial activity in a Karst ecosystem. *Sci. Rep.* **7**, 41402 (2017).
54. Bavel, C. H. M. V. Mean weight-diameter of soil aggregates as a statistical index of aggregation. *Soil Sci. Soc. Am. J.* **14**, 20–23 (1950).
55. Kemper, W. D. & Rosenau, K. Methods of soil analysis, Klute edition. *Size distribution of aggregates, in American. Part 1-physical and mineralogical methods*. 425–442 (Am. Soc. Agron. 1986).
56. Yang, P. L., Luo, Y. P. & Shi, Y. C. Characterized soil fractal characteristics by weight distribution of particle size. *Chin. Sci. Bull.* **38**, 1896–1899 (in Chinese with English abstract) (1993).
57. Bao, S. D. *Analysis of soil agrochemistry*. (China Agricultural Press 2008).
58. Li, J. *et al.* Effects of 15-year conservation tillage on soil and aggregate organic carbon sequestration in the Loess hilly region of China. *Sci. Agric. Sin.* **48**, 4690–4697 (in Chinese with English abstract) (2015).
59. Tan, R. B. & Mei, X. R. *SPSS statistical analysis of practical course*. (Science 2007).
60. Brown, H. J., Cruse, R. M., Erbach, D. C. & Melvin, S. W. Tractive device effects on soil physical properties. *Soil Tillage Res.* **22**, 41–53 (1992).
61. Schaffer, B., Attinger, W. & Schulin, R. Compaction of restored soil by heavy agricultural machinery-soil physical and mechanical aspects. *Soil Tillage Res.* **93**, 28–43 (2007).
62. Alkukku, L. *et al.* Prevention strategies for field traffic-induced subsoil compaction: a review. Part 1. Machine/soil interactions. *Soil Tillage Res.* **73**, 145–160 (2003).

63. Biswas, A. K., Mohanty, M., Hati, K. M. & Misra, A. K. Distillery effluents effect on soil organic carbon and aggregate stability of a Vertisol in India. *Soil Tillage Res.* **104**, 241–246 (2009).
64. Six, J., Elliott, E. T. & Paustian, K. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biol. Biochem.* **32**, 2099–2103 (2000).
65. Six, J., Elliott, E. T. & Paustian, K. Soil structure and soil organic mater: II. A normalized stability index and the effect of mineralogy. *Soil Sci. Soc. Am. J.* **64**, 1042–1049 (2000).
66. Neu, T. R. & Kuhllicke, U. Fluorescence Lectin bar-coding of glycoconjugates in the extracellular matrix of biofilm and bioaggregate forming microorganisms. *Microorganisms* **5**, 5 (2017).
67. Nakamoto, T. & Suzuki, K. Influence of soybean and maize roots on the seasonal Change in soil aggregate size and stability. *Plant Prod. Sci.* **4**, 317–319 (2001).
68. Wick, A. E., Stahl, P. D. & Ingram, L. J. Aggregate-associated carbon and nitrogen in reclaimed sandy loam soils. *Soil Sci. Soc. Am. J.* **73**, 1852–1860 (2009).
69. Kumari, M. *et al.* Soil aggregation and associated organic carbon fractions as affected by tillage in a rice-wheat rotation in north India. *Soil Sci. Soc. Am. J.* **75**, 560–567 (2011).
70. Das, B. *et al.* Effect of organic inputs on strength and stability of soil aggregates under rice-wheat rotation. *Int. Agrophys.* **28**, 163–168 (2014).
71. Devine, S., Markewitz, D., Hendrix, P. & Coleman, D. Soil aggregates and associated organic matter under conventional tillage, no-tillage, and forest succession after three decades. *Plos One* **9**, e84988 (2014).
72. Pinheiro, E. F. M., Pereira, M. G. & Anjos, L. H. C. Aggregate distribution and soil organic matter under different tillage systems for vegetable crops in a Red Latosol from Brazil. *Soil Tillage Res.* **77**, 19–84 (2004).
73. Zhang, X. F., Zhu, A. N. & Zhang, J. B. The long-term effect research of various tillage managements on the soil aggregates and organic carbon in Fluvo-Aquic soil. *Sci. Agric. Sin.* **48**, 4639–4648 (in Chinese with English abstract) (2015).
74. Su, F. L., Zhao, G. H., Wang, T. L., Li, H. F. & Li, Y. M. Characteristics of surface soil macro aggregates under different land use patterns. *Pratacultural Sci.* **34**, 924–931 (in Chinese with English abstract) (2017).
75. Li, L. Q., Zhang, X. H. & Pan, G. X. Changes in organic carbon storage in aggregates of the surface horizon in a degraded Paleudlt upon vegetation recovery. *Chin. J. Soil Sci.* **31**, 193–195 (in Chinese with English abstract) (2000).
76. Wu, J. G., Zhang, X. Q. & Xu, D. Y. Impact of land use change on soil carbon storage. *Chin. J. Appl. Ecol.* **4**, 593–599 (2004).
77. Puget, P., Angers, D. A. & Chenu, C. Nature of carbohydrates associated with water stable aggregates of two cultivated soils. *Soil Biol. Biochem.* **31**, 55–63 (1998).
78. Puget, P., Chenu, C. & Balesdent, J. Dynamics of soil organic matter associated with particle-size fractions of water-stable aggregates. *Eur. J. Soil Sci.* **51**, 595–605 (2000).
79. Tisdall, J. M. & Oades, J. M. Landmark papers: no. 1. Organic matter and water-stable aggregates in soils. *Eur. J. Soil Sci.* **63**, 8–21 (2012).
80. An, S. S., Mentler, A., Mayer, H. & Blum, W. E. H. Soil aggregation, aggregate stability, organic carbon and nitrogen in different soil aggregate fractions under forest and shrub vegetation on the Loess Plateau, China. *Catena* **81**, 226–233 (2010).
81. Peng, X., Yan, X., Zhou, H., Zhang, Y. Z. & Sun, H. Assessing the Contributions of sesquioxides and soil organic matter to aggregation in an Ultisol under long-term fertilization. *Soil Tillage Res.* **146**, 89–98 (2015).
82. Yu, M. *et al.* Impact of land-use changes on soil hydraulic properties of Calcaric Regosols on the Loess Plateau, NW China. *J. Plant Nutr. Soil Sci.* **178**, 486–498 (2015).
83. Somasundaram, J., Reeves, S., Wang, W. J., Heenan, M. & Dalal, R. Impact of 47 year of no tillage and stubble retention on soil aggregation and carbon distribution in a vertisol. *Land Degrad. Dev.* **28**, 1589–1602 (2016).
84. Kalhoro, S. A. *et al.* Effects of different land-use systems on soil aggregates: a case study of the loess plateau (Northern China). *Sustainability* **9**, 1349 (2017).
85. Chivenge, P., Vanlauwe, B., Gentile, R. & Six, J. Comparison of organic versus mineral resource effects on short-term aggregate carbon and nitrogen dynamics in a sandy soil versus a fine textured soil. *Agric. Ecosyst. Environ.* **140**, 361–371 (2011).
86. Hurisso, T. T. *et al.* Rapid changes in microbial biomass and aggregate size distribution in response to changes in organic matter management in grass pasture. *Geoderma* **193**, 68–75 (2013).
87. Christensen, B. T. Straw incorporation and soil organic matter in macro aggregates and particle size separates. *Eur. J. Soil Sci.* **37**, 125–135 (1986).
88. De Jonge, L. W., Jacobsen, O. H. & Moldrup, P. Soil water repellency effects of Water content, temperature and particle size. *Soil Sci. Soc. Am. J.* **63**, 437–442 (1999).
89. Li, H. X., Yuan, Y. H., Huang, Q. R., Hu, F. & Pan, G. X. Effects of fertilization on soil organic carbon distribution in various aggregates of red paddy soil. *Acta Pedologica Sin.* **3**, 422–428 (in Chinese with English abstract) (2006).
90. Wang, X. J. *et al.* Effects of organic manure application on dry land soil organic matter and water stable aggregates. *Chin. J. Appl. Ecol.* **23**, 159–165 (in Chinese with English abstract) (2012).
91. Huang, X. Q., Xin, Z. B., Zhao, Y. J. & Ma, F. Y. Soil aggregate composition and distribution characteristics of soil organic carbon of typical plantations in mountainous area of Beijing. *J. Soil Water Conserv.* **30**, 239–246 (in Chinese with English abstract) (2016).
92. Meng, Q. F. *et al.* Distribution of carbon and nitrogen in water-stable aggregates and soil stability under long-term manure application in solonchic soils of the Songnen plain, northeast China. *J. Soils Sediments* **14**, 1041–1049 (2014).
93. Gürsoy, S., Sessiz, A. & Malhi, S. S. Short-term effects of tillage and residue management following cotton on grain yield and quality of wheat. *Field Crops Res.* **119**, 260–268 (2010).

## Acknowledgements

We appreciate two anonymous reviewers for their insightful comments on this paper. This research was supported by the National Key Research and Development Program of China (No. 2018YFD0200407) and the Research Program of Heilongjiang Branch of China Tobacco Corporation (No. 20182300002700081).

## Author contributions

X.F.M. conceived and designed the experiments; M.Z. and C.Z.L. performed the experiments; M.Z. and Y.Y. analysed the data; M.Z., X.F.M. and X.B.L. wrote the paper; X.B.L., Q.F.M., Y.X.Z. and G.W.D. revised the manuscript; J.Z.Z., X.N.Z., J.W. and W.L.D. collected and processed samples.

## Competing interests

The authors declare no competing interests.

## Additional information

**Correspondence** and requests for materials should be addressed to X.M. or X.L.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020