

RESEARCH ARTICLE

Open Access



Bioinformatics analysis of differentially expressed genes in rotator cuff tear patients using microarray data

Yi-Ming Ren[†], Yuan-Hui Duan[†], Yun-Bo Sun[†], Tao Yang and Meng-Qiang Tian^{*}

Abstract

Background: Rotator cuff tear (RCT) is a common shoulder disorder in the elderly. Muscle atrophy, denervation and fatty infiltration exert secondary injuries on torn rotator cuff muscles. It has been reported that satellite cells (SCs) play roles in pathogenic process and regenerative capacity of human RCT via regulating of target genes. This study aims to complement the differentially expressed genes (DEGs) of SCs that regulated between the torn supraspinatus (SSP) samples and intact subscapularis (SSC) samples, identify their functions and molecular pathways.

Methods: The gene expression profile GSE93661 was downloaded and bioinformatics analysis was made.

Results: Five hundred fifty one DEGs totally were identified. Among them, 272 DEGs were overexpressed, and the remaining 279 DEGs were underexpressed. Gene ontology (GO) and pathway enrichment analysis of target genes were performed. We furthermore identified some relevant core genes using gene–gene interaction network analysis such as GNG13, GCG, NOTCH1, BCL2, NMUR2, PMCH, FFAR1, AVPR2, GNA14, and KALRN, that may contribute to the understanding of the molecular mechanisms of secondary injuries in RCT. We also discovered that GNG13/calcium signaling pathway is highly correlated with the denervation atrophy pathological process of RCT.

Conclusion: These genes and pathways provide a new perspective for revealing the underlying pathological mechanisms and therapy strategy of RCT.

Keywords: Rotator cuff muscle, Satellite cells, Differentially expressed genes, Bioinformatics analysis, Calcium signaling, Denervation

Introduction

The rotator cuff muscle complex of the shoulder is comprised of four distinct muscles (supraspinatus, infraspinatus, teres minor, and subscapularis), which controls essential shoulder movements [1, 2]. The rotator cuff tear (RCT) is a common cause of impact pain, nocturnal pain and shoulder joint dysfunction, which seriously affect the life and working ability of patients, and reduce the quality of life of patients [3, 4]. Most tears require reparative surgery; however, recurrence of tears following surgery is common, with failure rates ranging from 30 to 94% [5]. Rotator cuff tendon tears are accompanied by secondary changes in the rotator cuff muscles, including muscle

atrophy, denervation, and fatty infiltration, which may explain the progressive loss of function after an acute injury and also the high rate of surgical failure. However, the underlying mechanism is not well understood.

Satellite cells (SCs) are mitotically quiescent muscle stem cells located between the basal lamina and the muscle membrane, which are known to play a key role in the adaptive response of muscle to exercise, and in the maintenance of the regenerative capacity of muscle. Hepatocyte growth factor (HGF) and nitric oxide (NO) could regulate transit of a SC from the quiescent G0 state into the G1 (activated) stage of the cell cycle [6]. Recently, Deanna et al. discovered possible supraspinatus denervation in RCT and suggested NO-donor treatment combined with stretching has potential to promote growth in atrophic supraspinatus muscle after RCT and improve functional outcome [7, 8]. Lundgreen et al.

* Correspondence: tmjoint@126.com

[†]Yi-Ming Ren, Yuan-Hui Duan and Yun-Bo Sun contributed equally to this work. Department of Joint and Sport Medicine, Tianjin Union Medical Center, Jieyuan Road 190, Hongqiao District, Tianjin 300121, People's Republic of China



showed patients with full-thickness tears had a reduced density of SCs, fewer proliferating cells, and atrophy of myofibers [9]. With muscle atrophy, fatty infiltration into skeletal muscles is thought to cause muscle degeneration by impairing the myogenic function of SCs [10].

Here, we downloaded the gene expression profile GSE93661 from the Gene Expression Omnibus database (GEO) and made bioinformatics analysis to investigate differentially expressed genes (DEGs) of SCs that regulated between torn supraspinatus (SSP) samples and intact subscapularis (SSC) samples from RCT patients. By doing this, we hope that the key target genes and pathways involved in the pathological process of RCT could be identified and existing molecular mechanisms could be revealed.

Materials and methods

Gene expression microarray data

The gene expression profile GSE93661 was downloaded from the Gene Expression Omnibus (GEO, www.ncbi.nlm.nih.gov/geo/). GSE93661 was based on Agilent-026652 Whole Human Genome Microarray 4x44K v2 platform. GSE93661 dataset contained four samples, including two torn SSP samples, and two intact SSC samples.

DEGs in torn SSP and intact SSC samples

The raw data files used for the analysis included TXT files. The analysis was carried out using GEO2R, which can perform comparisons on original submitter-supplied processed data tables using the GEO query and limma R packages from Bioconductor project. The P value < 0.05 and log fold change (FC) > 2.0 or log FC < -2.0 were used as the cut-off criteria. The DEGs with statistical significance between the torn SSP samples and intact SSC samples were selected and identified.

GO and KEGG analysis of DEGs

Target genes list were submitted to the DAVID 6.8 (<https://david.ncifcrf.gov/tools.jsp>) and ClueGO version 2.33 (based on Cytoscape software version 3.4.0 (www.cytoscape.org)) to identify overrepresented GO categories and pathway categories. Gene ontology (GO) analysis was used to predict the potential functions of the DEGs in biological process (BP), molecular function (MF), and cellular component (CC). The Kyoto Encyclopedia of Genes and Genomes (KEGG) is a knowledge base for systematic analysis of gene functions, linking genomic information with higher-level systemic functions. Finally, the overrepresented pathway categories were considered statistically significant using KEGG pathway enrichment analysis.

Gene interaction network construction

A large number of DEGs we obtained may be RCT-associated genes, and it is suggested that these

DEGs in torn SSP samples may participate in the progression of RCT. Firstly, DEGs list was submitted to the Search Tool for the Retrieval of Interacting Genes (STRING) database (<http://www.string-db.org/>), and an interaction network chart with a combined score > 0.4 was saved and exported. Subsequently, the interaction regulatory network of RCT-associated genes was visualized using Cytoscape software version 3.4.0. The distribution of core genes in the interaction network was made by NetworkAnalyzer in Cytoscape. Then, the plugin Molecular Complex Detection (MCODE) was applied to screen the modules of the gene interaction network in Cytoscape. Venn diagram was drawn using Venny 2.1 (<http://bioinfogp.cnb.csic.es/tools/venny/>).

Result

Identification of DEGs

The gene expression profile GSE93661 was downloaded from the GEO, and the GEO2R method was used to identify DEGs in torn SSP samples compared with intact SSC samples. P value < 0.05 , log FC > 2.0 , or log FC < -2.0 were used as the cut-off criteria. After analyzing, differentially expression gene profiles were obtained. Totally, 551 DEGs were identified including 272 upregulated DEGs and 279 downregulated DEGs screened in torn SSP samples compared with intact SSC samples. Parts of DEGs were listed in Table 1.

GO term enrichment analysis of DEGs

Functional annotation of the 551 DEGs was clarified using the DAVID 6.8 online tool. GO analysis indicated that these DEGs were significantly enriched in muscle contraction, aging, regulation of ion transmembrane transport, mesenchymal cell development, and other biological processes (Fig. 1). For MF, the DEGs were enriched in ion channel activity, calcium ion binding, structural molecule activity, and others. In addition, GO CC analysis also showed that the DEGs were significantly enriched in keratin filament, integral component of plasma membrane, axon, cornified envelope, cortical cytoskeleton, and others.

KEGG pathway analysis of DEGs

The result of KEGG pathway analysis revealed that target genes were enriched in butanoate metabolism, ABC transporters, notch signaling pathway, arachidonic acid metabolism, hedgehog signaling pathway, cell adhesion molecules (CAMs), prolactin signaling pathway, neuroactive ligand-receptor interaction, dopaminergic synapse, GABAergic synapse, calcium signaling pathway, cGMP-PKG signaling pathway, drug metabolism, B cell receptor signaling pathway, NF-kappa B signaling pathway, estrogen signaling pathway, cAMP signaling pathway, and others. These key pathways were showed in Fig. 2.

Table 1 The top 10 Regulated DEGs in SCs of torn rotator cuff muscle with *P* value < 0.05

ID	<i>P</i> value	logFC	Gene symbol
Upregulated			
9522	0.0024644	5.71	FAM196B
39,990	0.0031637	5.05	SCN2A
8187	0.0000663	4.91	NNT
41,585	0.0035824	4.74	TDRD3
8204	0.0040603	4.46	LOC100131129
28,406	0.0002489	4.42	C8orf42
8118	0.0041306	4.42	KIAA1751
12,220	0.0043186	4.32	LRRC2
7774	0.0044309	4.27	CYP2E1
37,687	0.0007211	4.24	CUBN
Downregulated			
32,273	0.0024714	-4.62	PTPRC
45,097	0.0037977	-4.61	TCF7L2
12,505	0.0016694	-4.49	tcag7.1023
35,725	0.0001639	-4.11	FAM101A
45,093	0.0083048	-4.08	C8orf67
8427	0.0005044	-3.84	PTCH2
38,947	0.0004951	-3.6	PLLP
31,098	0.0001216	-3.55	TRPV5
31,980	0.0064577	-3.54	SPNS3
29,856	0.0004283	-3.52	ACP2

SCs satellite cells, DEGs differentially expressed genes, FC fold change

Besides, these core pathways and their associated genes found were summarized in Table 2. The first-ranking butanoate metabolism signaling pathway had the 10.71% associated genes, which included ACSM2B, ACSM4, and ACSM6. The second-placed ABC transporters signaling pathway had the 8.89% associated genes, which included ABCA3, ABCD1, ABCG2, and ABCG4. The third-placed notch signaling pathway had the 8.33% associated genes, which included JAG2, MAML3, MFNG, and NOTCH1.

Interaction network of DEGs and core genes in the interaction network

Based on the information in the STRING database, the gene interaction network contained 386 nodes and 440 edges. The nodes indicated the DEGs, and the edges indicated the interactions between the DEGs. NetworkAnalyzer in Cytoscape software was used to analyze these genes, and core genes were ranked according to the predicted scores. The top 10 high-degree hub nodes included GNG13, GCG, NOTCH1, BCL2, NMUR2, PMCH, FFAR1, AVPR2, GNA14, and KALRN. Among these genes, GNG13 showed the highest node degree which was 32. The core genes and their corresponding degree were shown in Table 3. The distribution of core genes in the interaction network was revealed in Fig. 3. The correlation between the data points and corresponding points on the line is approximately 0.993. The *R*² value is 0.902, giving a relatively high confidence that the underlying model is indeed linear. Then, we used MCODE to screen the modules of the gene interaction network, and eight modules were showed in Fig. 4.

The score of top 1 module including GCG, GNG13, NMUR2, and KALRN was 14, which had 14 nodes and 91 edges. The score of top 2 module including BCL2 and CD22 was 6, which had 6 nodes and 15 edges. The





Fig. 1 Gene ontology (GO)-enrichment analysis of biological processes (a) molecular functions (b) and cellular components (c). The labels in *Y* axis mean enrichment score (-log₁₀ *P* value), and labels in *X* axis mean GO terms



score of top 3 module including CBLB, RNF6, TRIM9, and FBXO44 was 6, which had 6 nodes and 15 edges. Lastly, the interaction network of the top 10 high-degree hub nodes (core genes) was made by STRING database in Fig. 5. GNG13, GCG, NOTCH1, BCL2, NMUR2, PMCH, FFAR1, AVPR2, GNA14, and KALRN, which regulate 7, 7, 2, 2, 6, 6, 6, 3, 6, and 7 targets, respectively, showed the good connectivity.

Interestingly, Chaudhury et al. [9, 11] reported that gene expression profiles of different-sized human rotator cuff tendon tears versus normal rotator cuff tendons. In order to seek the possibly common target genes, we pooled together the top 10 high-degree core genes as mentioned earlier and the 77 significantly DEGs of Chaudhury's research using Venn diagram [9, 11]. GNG13 was discovered as the only common target gene in Fig. 6.

Discussion

RCT is common and painful. Even after surgery, joint stability and function may not recover [11]. SCs play a major role in muscle regeneration. However, human SCs in muscles with atrophy, denervation, and fatty infiltration are unclear due to the difficulty in isolating from small samples, and the mechanism has not been elucidated [12–14]. In the present study, the gene expression profile of GSE93661 was downloaded and a bioinformatics analysis was performed. The results showed that there were 551 DEGs in SCs of torn rotator cuff tendons and normal rotator cuff tendons. Furthermore, GO, KEGG pathway, and gene–gene interaction network analysis were performed to obtain the biomarkers or the major genes related to pathogenicity mechanism of RCT.

In order to disclose the underlying molecular mechanisms between SCs and RCT, we characterized the possible GO functional terms and signaling pathways of DEGs. Considering the results of GO function analysis, we linked the DEGs with aging, regulation of ion transmembrane transports, ion channel activity, and calcium ion binding, which are very important for the development process of RCT. When muscle is injured, exercised, overused, or mechanically stretched, SCs are activated to enter the cell cycle, divide, differentiate, and fuse with the adjacent muscle fiber. In this way, SCs are responsible for regeneration and work-induced hypertrophy of muscle fibers. Ryuichi's results suggested that the activation mechanism is a cascade of molecular events including an influx of calcium ions and their binding to calmodulin, nitric oxide synthase (NOS) activation, NO radical production by eNOS, matrix metalloproteinase activation, HGF release from the matrix, and presentation of HGF to the signaling receptor c-met. Understanding the mechanisms of SC activation is essential when planning procedures that could enhance muscle growth and repair [15].


As previous articles reported, our KEGG pathway analysis showed that notch signaling pathway, hedgehog signaling pathway, dopaminergic synapse, GABAergic synapse, calcium signaling pathway, NF-kappa B signaling pathway, and estrogen signaling pathway were among the most relevant pathways for SCs in RCT. Pasut et al. found that in normal muscle, high levels of notch signaling is required to maintain the uncommitted state of SCs. Notch signaling plays a role in SC fate as activation of Notch1 strongly promotes the lineage

Table 2 Core pathways and their associated genes found

GOID	GO term	Term P value	% associated genes	Associated genes found
GO:000650	Butanoate metabolism	0.02	10.71	[ACSM2B, ACSM4, ACSM6]
GO:0002010	ABC transporters	0.01	8.89	[ABCA3, ABCD1, ABCG2, ABCG4]
GO:0004330	Notch signaling pathway	0.02	8.33	[JAG2, MAML3, MFNG, NOTCH1]
GO:0000590	Arachidonic acid metabolism	0.04	6.45	[CYP2B6, CYP2E1, PTGS1, TBXAS1]
GO:0004340	Hedgehog signaling pathway	0.07	6.38	[BCL2, IHH, PTCH2]
GO:0004514	Cell adhesion molecules (CAMs)	0.00	6.21	[CD22, CD226, CD86, CLDN15, ITGAM, NCAM2, NLGN1, PTPRC, VCAM1]
GO:0004917	Prolactin signaling pathway	0.05	5.71	[CISH, MAPK10, PRL, TH]
GO:0005217	Basal cell carcinoma	0.10	5.45	[PTCH2, TCF7L2, WNT10A]
GO:0004658	Th1 and Th2 cell differentiation	0.04	5.43	[IL12RB1, JAG2, MAML3, MAPK10, NOTCH1]
GO:0004080	Neuroactive ligand-receptor interaction	0.00	5.40	[AVPR2, CHRN3, CHRNE, CYSLTR1, EDNRB, GABBR1, GIPR, GLP2R, HRH3, LTB4R, NMUR2, NPY2R, NPY4R, OPRL1, PRL]
GO:0004728	Dopaminergic synapse	0.02	5.38	[CACNA1C, CALML6, GNG13, KCNJ6, MAPK10, SCN1A, TH]
GO:0004610	Complement and coagulation cascades	0.07	5.06	[F7, ITGAM, MASP2, VWF]
GO:0005210	Colorectal cancer	0.12	5.00	[BCL2, MAPK10, TCF7L2]
GO:0004742	Taste transduction	0.09	4.82	[CACNA1C, GABBR1, GNG13, SCN2A]
GO:0004911	Insulin secretion	0.09	4.71	[CACNA1C, FFAR1, GCG, KCNU1]
GO:0005214	Glioma	0.14	4.69	[CALML6, PDGFB, PLCG2]
GO:0000830	Retinol metabolism	0.14	4.62	[ADH6, CYP2B6, DHRS4L1]
GO:0004727	GABAergic synapse	0.10	4.55	[CACNA1C, GABBR1, GNG13, KCNJ6]
GO:0000010	Glycolysis/gluconeogenesis	0.15	4.48	[ACSS1, ADH6, LDHC]
GO:0005031	Amphetamine addiction	0.16	4.41	[CACNA1C, CALML6, TH]
GO:0004020	Calcium signaling pathway	0.03	4.40	[ATP2A1, ATP2B3, CACNA1C, CALML6, CYSLTR1, EDNRB, GNA14, PLCG2]
GO:0004022	cGMP-PKG signaling pathway	0.05	4.29	[ATP2A1, ATP2B3, CACNA1C, CALML6, EDNRB, KCNU1, NPPB]
GO:0000982	Drug metabolism	0.17	4.29	[ADH6, CYP2B6, CYP2E1]
GO:0004662	B cell receptor signaling pathway	0.17	4.23	[CD22, PLCG2, RASGRP3]
GO:0005418	Fluid shear stress and atherosclerosis	0.07	4.23	[BCL2, CALML6, MAPK10, PDGFB, PIAS4, VCAM1]
GO:0004064	NF-kappa B signaling pathway	0.12	4.21	[BCL2, PIAS4, PLCG2, VCAM1]
GO:0004713	Circadian entrainment	0.13	4.17	[CACNA1C, CALML6, GNG13, KCNJ6]
GO:0004915	Estrogen signaling pathway	0.14	4.08	[CALML6, GABBR1, HSPA6, KCNJ6]
GO:0000980	Metabolism of xenobiotics by cytochrome P450	0.19	4.05	[ADH6, CYP2B6, CYP2E1]
GO:0004024	cAMP signaling pathway	0.06	4.04	[ATP2B3, CACNA1C, CALML6, FXYD1, GABBR1, GHRL, GIPR, MAPK10]
GO:0004933	AGE-RAGE signaling pathway in diabetic complications	0.14	4.04	[BCL2, MAPK10, PLCG2, VCAM1]


Table 3 The core genes and their corresponding degree


Gene	Degree	Gene	Degree	Gene	Degree	Gene	Degree
GNG13	32	PMCH	18	PTPRC	14	MPO	13
GCG	22	FFAR1	15	CYSLTR1	14	OPN4	13
NOTCH1	21	AVPR2	15	EDNRB	14	GNRHR2	13
BCL2	21	GNA14	15	UTS2D	13	LTB4R	13
NMUR2	18	KALRN	15	UTS2	13	PRL	13



switch from myogenic towards brown adipogenic fate [16]. Khayrullin et al. reported that upregulation of Notch signaling suppresses myogenesis and maintains muscle SC quiescence and miRNAs targeting Notch are likely to play important roles in alcohol-related myopathy in zebrafish model [17]. SC self-renewal is an


essential process to maintaining the robustness of skeletal muscle regenerative capacity. Ogura’s study demonstrates that TNF-like weak inducer of apoptosis cytokine suppresses SC self-renewal through activating NF-kappa B and repressing Notch signaling [18]. Kamizaki’s findings indicate that Ror1 has a critical role in regulating





SC proliferation via NF-kappa B activation during skeletal muscle regeneration of injured muscle [19]. In addition, estrogen regulates myosin heavy chain expression in SCs related to muscle function mainly through an estrogen receptor α -mediated pathway [20]. In Voronova's study, the formation of skeletal muscle during embryogenesis and adult muscle regeneration is regulated by myocyte enhancer factors and myogenic regulatory factors (such as MyoD). Hedgehog signaling could regulate MyoD expression during embryogenesis and adult muscle regeneration in SCs [21].

The gene interaction network analysis revealed top 10 high-degree hub nodes of DEGs including GNG13, GCG, NOTCH1, BCL2, NMUR2, PMCH, FFAR1, AVPR2, GNA14, and KALRN. Most of them were not reported in SCs and RCT research. Only NOTCH1, who is a receptor that mediates intercellular signaling through a pathway conserved across the metazoan, had been studied [22].



Rando et al. found that activation of Notch1 signaling stimulates the proliferation of SCs and leads to the expansion of proliferating myoblasts. And, inhibition of Notch1 signaling abolishes SC activation and impairs muscle regeneration [23]. Also, recent studies found Notch1 is active in quiescent muscle SCs, and Notch1 signaling is critical for maintaining the quiescence of muscle SCs [24, 25]. As a supplement, Fujimaki et al. indicated that Notch1 and Notch2 coordinately maintain the SC pool in the quiescent state by preventing activation and regulate SC-fate decision in the activated state, governing adult muscle regeneration [26]. To sum up, upregulation of Notch1 may result in SC proliferation and self-renewal in the activated state, controlling muscle regeneration and improving muscle atrophy in RCT.

Furthermore, we analyzed the top 10 high-degree core genes and the 77 significantly DEGs of different-sized human rotator cuff tendon tears versus normal rotator cuff tendons in Chaudhury's research using Venn diagram [27]. GNG13 was discovered as the only common core gene. Heterotrimeric G proteins, which consist of alpha, beta, and gamma subunits, function as signal transducers for the 7-transmembrane-helix G protein-coupled receptors. GNG13 is a gamma subunit that is expressed in taste, retinal, and neuronal tissues, and plays a key role in taste transduction [28]. Through KEGG pathway results, we discovered that GNG13 acts as an important node in dopaminergic synapse signaling, and targets PLC and AC5 in calcium signaling pathway. Many previous studies had emphasized the importance of calcium signaling pathway and skeletal muscle development, homeostasis, and regeneration. Calcium-ion is an important component of the signaling promoting muscle formation, muscle homeostasis, and regeneration. In particular, calcium-ion changes may direct muscle SCs to maintain their quiescent state, proliferate, or differentiate into functional muscle [29]. What is more, we are still learning how calcium signaling pathway and neuromuscular connections are restored on regenerating muscle. The establishment of connections between the motor nerve terminal and a post-synaptic region of membrane on regenerating fibers is essential to re-innervation and functional contractility, which is important for clarifying denervation atrophy pathological process of injured rotator cuff muscle. Recent researches have implicated SC signaling in the process of muscle re-innervation. SC have potential to influence axon growth and the reappearance of neuromuscular connections by their secretion of semaphorin 3A (Sema3A). Sema3A is a neural chemorepellent that is thought to coordinate the reconnection of motor axons with a differentiating fiber in a regenerating muscle. These direct proofs encourage a possible implication of SCs in the spatiotemporal regulation of extracellular Sema3A concentrations, which potentially ensures coordinating a delay in neurite sprouting

and re-attachment of motoneuron terminals onto damaged muscle fibers early in muscle regeneration in synchrony with recovery of muscle-fiber integrity [30–32]. In addition, calcium signaling pathway is also linked with the activation of neuromuscular connections, and calcium-ion influx through voltage-gated calcium channels regulates the neuron's responsiveness to Sema2A-dependent chemorepulsion exerted by the muscle [33]. Our research highlights the crucial role of nerve–muscle interaction in restoring innervation after RCT, and hypothesizes that SC-mediated GNG13 could affect neuromuscular connections and cause denervation atrophy via calcium signaling after rotator cuff muscle injured.

Conclusions

In summary, 551 DEGs were identified including 272 up-regulated DEGs and 279 downregulated DEGs screened in SCs of torn SSP samples compared with intact SSC samples. GO and KEGG pathway analysis provided a series of related key genes and pathways to contribute to the understanding of the molecular mechanisms between SCs and RCT, thus yielding clues to speculate the GNG13/calcium signaling pathway is highly correlated with the denervation atrophy pathological process of RCT. Furthermore, further experimental validation should be made in future studies.

Abbreviations

BP: Biological process; CAMs: Cell adhesion molecules; CC: Cellular component; FC: Fold change; GEO: Gene Expression Omnibus database; GO: Gene ontology; HGF: Hepatocyte growth factor; KEGG: Kyoto Encyclopedia of Genes and Genomes; MCODE: Molecular Complex Detection; MF: Molecular function; NO: Nitric oxide; NOS: Nitric oxide synthase; RCT: Rotator cuff tear; SCs: Satellite cells; Sema3A: Semaphorin 3A; SSC: Subscapularis; SSP: Supraspinatus; STRING: Search Tool for the Retrieval of Interacting Genes

Acknowledgements

None.

Funding

This research is supported by Foundation of Tianjin Union Medical Center (2017YJ018).

Availability of data and materials

The data was freely downloaded from the public GEO database.

Authors' contributions

YMR, YHD, and YBS conceived the design of the study. TY and YMR performed and collected the data, and contributed to the design of the study. TY and YMR analyzed the data. YMR and MQT prepared and revised the manuscript. All authors read and approved the final content of the manuscript.

Ethics approval and consent to participate

Not applicable. This paper does not involve research on humans.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 19 May 2018 Accepted: 29 October 2018

Published online: 13 November 2018

References

1. Yamaguchi K, Ditsios K, Middleton WD, et al. The demographic and morphological features of rotator cuff disease. *J Bone Joint Surg (Am Vol)*. 2006;88(8):1699–704.
2. Liu X, Ravishankar B, Ning A, et al. Knocking-out matrix metalloproteinase-13 exacerbates rotator cuff muscle fatty infiltration. *Muscles Ligaments Tendons J*. 2017;7(2):202–7.
3. Schmidt CC, Jarrett CD, Brown BT. Management of rotator cuff tears. *J Hand Surg*. 2015;40(2):399–408.
4. Liu X, Ning AY, Chang NC, et al. Investigating the cellular origin of rotator cuff muscle fatty infiltration and fibrosis after injury. *Muscles Ligaments Tendons J*. 2016;6(1):6.
5. Isaac C, Gharaibeh B, Witt M, et al. Biologic approaches to enhance rotator cuff healing after injury. *J Shoulder Elb Surg*. 2012;21(2):181–90.
6. Dhawan J, Rando TA. Stem cells in postnatal myogenesis: molecular mechanisms of satellite cell quiescence, activation and replenishment[J]. *Trends in Cell Biology*. 2005;15(12):666–673.
7. Deanna G, Leiter JRS, Macdonald PB, et al. Altered satellite cell responsiveness and denervation implicated in progression of rotator-cuff injury. *PLoS One*. 2016;11(9):e0162494.
8. Gigliotti D, Leiter JR, Macek B, et al. Atrophy, inducible satellite cell activation and possible denervation of supraspinatus muscle in injured human rotator-cuff muscle. *Am J Physiol Cell Physiol*. 2015;309(6):C383.
9. Lundgreen K, Lian OB, Engebretsen L, et al. Lower muscle regenerative potential in full-thickness supraspinatus tears compared to partial-thickness tears. *Acta Orthop*. 2013;84(6):565.
10. Pisani DF, Clement N, Loubat A, et al. Hierarchization of myogenic and adipogenic progenitors within human skeletal muscle. *Stem Cells*. 2010; 28(12):2182–94.
11. Cofield RH, Parvizi J, Hoffmeyer PJ, et al. Surgical repair of chronic rotator cuff tears. A prospective long-term study. *J Bone Joint Surg Am*. 2001;83-A(1):71–7.
12. Zammit PS, Partridge TA, Yablonkareveni Z. The skeletal muscle satellite cell: the stem cell that came in from the cold. *J Histochem Cytochem*. 2006; 54(11):1177–91.
13. Oliva F, Piccirilli E, Bossa M, et al. I.S.Mu.L.T - Rotator Cuff Tears Guidelines. *Muscles Ligaments Tendons J*. 2016;5(4):227.
14. Hall KE, Sarkissian EJ, Sharpe O, et al. Identification of differentially expressed micro-RNA in rotator cuff tendinopathy. *Muscles Ligaments Tendons J*. 2018;8:8–14.
15. Ryuichi T, Ronalde A. Mechano-biology of resident myogenic stem cells: molecular mechanism of stretch-induced activation of satellite cells. *Anim Sci J*. 2008;79(3):279–90.
16. Pasut A, Chang NC, Rodriguez UG, et al. Notch signaling rescues loss of satellite cells lacking Pax7 and promotes brown adipogenic differentiation. *Cell Rep*. 2016;16(2):333–43.
17. Khayrullin A, Smith L, Mistry D, et al. Chronic alcohol exposure induces muscle atrophy (myopathy) in zebrafish and alters the expression of microRNAs targeting the Notch pathway in skeletal muscle. *Biochem Biophys Res Commun*. 2016;479(3):590–5.
18. Ogura Y, Mishra V, Hindi SM, et al. Proinflammatory cytokine tumor necrosis factor (TNF)-like weak inducer of apoptosis (TWEAK) suppresses satellite cell self-renewal through inversely modulating notch and NF-kappa B signaling pathways. *J Biol Chem*. 2013;288(49):35159–69.
19. Kamizaki K, Doi R, Hayashi M, et al. The Ror1 receptor tyrosine kinase plays a critical role in regulating satellite cell proliferation during regeneration of injured muscle. *J Biol Chem*. 2017;292(38):15939.
20. Guo T, Liu W, Konermann A, et al. Estradiol modulates the expression pattern of myosin heavy chain subtypes via an ERα-mediated pathway in muscle-derived tissues and satellite cells. *Cell Physiol Biochem*. 2014;33(3):681–91.
21. Voronova A, Coyne E, Madhoun AA, et al. Hedgehog signaling regulates MyoD expression and activity. *J Biol Chem*. 2013;288(6):4389–404.
22. Conboy IM, Conboy MJ, Smythe GM, et al. Notch-mediated restoration of regenerative potential to aged muscle. *Science*. 2003;302(5650):1575–7.

23. Luo D, Renault VM, Rando TA. The regulation of notch signaling in muscle stem cell activation and postnatal myogenesis. *Semin Cell Dev Biol.* 2005;16(5): 612–22.
24. Mourikis P, Sambasivan R, Castel D, et al. A critical requirement for notch signaling in maintenance of the quiescent skeletal muscle stem cell state. *Stem Cells.* 2012;30(2):243–52.
25. Bjornson CRR, Cheung TH, Liu L, et al. Notch signaling is necessary to maintain quiescence in adult muscle stem cells. *Stem Cells.* 2012;30(2):232–42.
26. Fujimaki S, Seko D, Kitajima Y, et al. Notch1 and Notch2 Coordinately Regulate Stem Cell Function in the Quiescent and Activated States of Muscle Satellite Cells[J]. *Stem Cells.* 2018;36(2):278–285.
27. Chaudhury S, Xia Z, Thakkar D, et al. Gene expression profiles of changes underlying different-sized human rotator cuff tendon tears. *J Shoulder Elbow Surg.* 2016;25(10):1561–70.
28. Blake BL, Wing MR, Zhou JY, et al. G beta association and effector interaction selectivities of the divergent G gamma subunit G gamma(13). *J Biol Chem.* 2001;276(52):49267–74.
29. Tu MK, Levin JB, Hamilton AM, et al. Calcium signaling in skeletal muscle development, maintenance and regeneration. *Cell Calcium.* 2016;59(2–3):91–7.
30. Anderson JE, Do MQ, Daneshvar N, et al. The role of semaphorin3A in myogenic regeneration and the formation of functional neuromuscular junctions on new fibres[J]. *Biological Reviews.* 2016;92(3):1389–1405.
31. Sato Y, Do MK, Suzuki T, et al. Satellite cells produce neural chemorepellent semaphorin 3A upon muscle injury. *Anim Sci J Nihon chikusan Gakkaiho.* 2013;84(2):185–9.
32. Tatsumi R, Sankoda Y, Anderson JE, et al. Possible implication of satellite cells in regenerative motoneuritogenesis: HGF upregulates neural chemorepellent Sema3A during myogenic differentiation. *Am J Physiol Cell Physiol.* 2009;297(2):C238.
33. Vonhoff F, Keshishian H. In vivo calcium signaling during synaptic refinement at the *Drosophila* neuromuscular junction. *J Neurosci.* 2017; 37(22):5511–26.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

