

# Reply: Roll out of bedaquiline without drug susceptibility testing leads to emergence of bedaquiline-resistance in high burden countries with tuberculosis - Danger of losing a novel drug

Dear Sir,

In response to the correspondence by Udwardia Zarir and Jennifer<sup>[1]</sup> “India should speed up access to bedaquiline based all-oral regimens, not procrastinate further,” I would like to submit that my key message in the Editorial was grossly misinterpreted by authors. The article “WHO consolidated guidelines on tuberculosis 2020 moving towards fully oral regimens: Should country act in hurry?”<sup>[2]</sup> have been written with concern to save bedaquiline for long-term use in the management of drug-resistant tuberculosis. The article does not question the efficacy of bedaquiline in the management of drug-resistant tuberculosis with its judicious use.

The same concern has been expressed internationally too, on rolling out WHO consolidated guidelines, Kranzer *et al.*<sup>[3]</sup> have written “New World Health Organization Treatment Recommendations for Multidrug-Resistant Tuberculosis: Are We Well Enough Prepared?” and Andres *et al.*<sup>[4]</sup> have commented, “Bedaquiline Resistant-Tuberculosis: Dark Clouds on the Horizon” both the articles published in prestigious American Journal of Respiratory and Critical Care Medicine.

The same enthusiasm was there when rifampicin was discovered for treatment for tuberculosis. Drug-resistant tuberculosis (DR-TB) was not a major problem till year 1997–2000 when RNTCP was rolled out with weaker regimens, as a result of which India has become home to 24% of world’s Multi DR-TB (MDR-TB)<sup>[5]</sup> problem, second highest in the world within 20 years of its implementation (WHO Global TB Report 2019).

The Editorial<sup>[2]</sup> answers almost all the questions raised by Udwardia and Furin, however for the sake of clarity I would like to respond further. Udwardia and Furin write about touting of Bangladesh regimen, there is no question of touting it, there are enough published data on the efficacy of this regimen in programmatic management of DR-TB in many countries, over 80 countries are implementing short MDR-regimen maintaining a cure rate of over 80%<sup>[6-10]</sup> and the same has been proven in well-controlled clinical study STREAM (a trial of Shorter Regimen for Rifampicin-Resistant Tuberculosis).<sup>[11]</sup> This regimen has been suggested to buy time awaiting results of large number of clinical trials in pipeline including TB-PRACTECAL,<sup>[12]</sup>

SIMPLICI-TB,<sup>[13]</sup> ZeNIX,<sup>[14]</sup> NEXT-TB,<sup>[15]</sup> STREAM II,<sup>[16]</sup> END-TB,<sup>[17]</sup> MDR-END,<sup>[18]</sup> and BEAT TB<sup>[19]</sup> the results of which will be available by end of 2022. By this time country can establish a strong infrastructure of DST laboratories for DST to bedaquiline, clofazimine, linezolid, and delamanid as WHO validated accurate reliable and reproducible methods have been established for these drugs (WHO consolidated guidelines on Drug-Resistant TB, 2020).<sup>[20]</sup>

Currently, bedaquiline DST is available at only 3 national reference laboratories, rolling out of WHO Consolidated Guidelines 2020 in such situation without the establishment of DST methods for drugs used in the regimen (s) may result in the development of bedaquiline-resistant TB which will be a disaster for the nation.

Primary bedaquiline-resistance exists in 5%–6% of drug-resistant TB cases<sup>[21-23]</sup> and insufficient protection of bedaquiline-clofazimine (BDQ/CFZ) by the remaining companion drugs because of resistance or drug toxicity especially to fluoroquinolone and linezolid, amplification of resistance to BDQ/CFZ is likely to occur.<sup>[4]</sup>

Many countries that have rolled out bedaquiline containing regimens largely in the absence of programmatic drug-susceptibility testing (DST) are facing problems of high frequency of acquired resistance to bedaquiline (15%–45%).<sup>[24-26]</sup> Andres *et al.*, in 2020<sup>[4]</sup> have proposed monthly phenotype and genotype testing to determine the emergence of resistance to drugs used and in an early view, Chesov *et al.*<sup>[27]</sup> in European Respiratory Journal (2021), have expressed that in high burden countries of tuberculosis bedaquiline resistance can emerge in more than 15% of mycobacterial TB complex strains obtained from follow-up isolates of MDR-TB patients and globally emerging resistant strains threaten the effectivity of novel MDR-TB treatment regimens. It is rightly written by Köser *et al.* that “THOSE WHO CANNOT REMEMBER THE PAST ARE CONDEMNED TO REPEAT IT.”<sup>[28]</sup>

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

**Nirmal Kumar Jain**

Department of Respiratory Medicine, JNU Medical College,  
Jaipur, Rajasthan, India  
E-mail: jainmkdr1971@gmail.com

Submitted: 13-Jan-2022 Accepted: 09-Feb-2022

Published: 01-Jul-2022

## REFERENCES

- Udwadia Zarir F, Jennifer F. India should speed up access to bedaquiline-based all-oral regimens, not procrastinate further. *Lung India* 2021;38:590-1.
- Jain NK. WHO consolidated guidelines on tuberculosis 2020 moving toward fully oral regimen: Should country act in hurry? *Lung India* 2021;38:303-6.
- Kranzer K, Kalsdorf B, Heyckendorf J, Andres S, Merker M, Hofmann-Thiel S, et al. New world health organization treatment recommendations for multidrug-resistant tuberculosis: Are we well enough prepared? *Am J Respir Crit Care Med* 2019;200:514-5.
- Andres S, Merker M, Heyckendorf J, Kalsdorf B, Rumetshofer R, Indra A, et al. Bedaquiline-resistant tuberculosis: Dark clouds on the horizon. *Am J Respir Crit Care Med* 2020;201:1564-8.
- World Health Organisation, Global Tuberculosis Report; 2019. Available from: <https://www.who.int/publications/i/item/9789241565714>. [Last accessed on 2021 Dec 25].
- Aung KJ, Van Deun A, Declercq E, Sarker MR, Das PK, Hossain MA, et al. Successful '9-month Bangladesh regimen' for multidrug-resistant tuberculosis among over 500 consecutive patients. *Int J Tuberc Lung Dis* 2014;18:1180-7.
- Trébucq A, Decroo T, Van Deun A, Piubello A, Chiang CY, Koura KG, et al. Short-Course Regimen for Multidrug-Resistant Tuberculosis: A Decade of Evidence. *J Clin Med* 2019;9:55.
- Piubello A, Harouna SH, Souleymane MB, Boukary I, Morou S, Daouda M, et al. High cure rate with standardised short-course multidrug-resistant tuberculosis treatment in Niger: No relapses. *Int J Tuberc Lung Dis* 2014;18:1188-94.
- Trébucq A, Schwoebel V, Kashongwe Z, Bakayoko A, Kuaban C, Noeske J, et al. Treatment outcome with a short multidrug-resistant tuberculosis regimen in nine African countries. *Int J Tuberc Lung Dis* 2018;22:17-25.
- Kuaban C, Noeske J, Rieder HL, Ait-Khaled N, Abena Foe JL, Trébucq A. High effectiveness of a 12-month regimen for MDR-TB patients in Cameroon. *Int J Tuberc Lung Dis* 2015;19:517-24.
- Nunn AJ, Phillips PP, Meredith SK, Chiang CY, Conradie F, Dalai D, et al. A trial of a shorter regimen for rifampin-resistant tuberculosis. *N Engl J Med* 2019;380:1201-13.
- TB-PRACTICAL. Programmatic Clinical Trial for More Effective, Concise and Less Toxic MDR-TB Treatment Regimen (s). *ClinicalTrials.gov* Identifier: NCT02589782. Available from: <https://clinicaltrials.gov/ct2/show/NCT04062201> [Last accessed on 2021 Dec 25].
- SIMPLI-TB (TB Alliance). Trial to Evaluate the Efficacy, Safety and tolerability of BPaMZ in Drug-Sensitive (DS-TB) adult patient and Drug-Resistant (DR-TB) Adult Patients. *ClinicalTrials.gov* Identifier: NCT03338621. Available from: <https://clinicaltrials.gov/ct2/show/NCT03338621>. [Last accessed on 2021 Dec 25].
- ZeNix (TB Alliance). Safety and Efficacy of Various Doses and Treatment Duration of Linezolid Plus Bedaquiline and Pretomanid in Participants with Pulmonary TB, XDR-TB, Pre-XDR-TB or Non-Responsive/Intolerant MDR-TB. *ClinicalTrials.gov* Identifier: NCT03086486. Available from: <https://clinicaltrials.gov/ct2/show/NCT03086486>. [Last accessed on 2021 Dec 25].
- NEXT-TB. An Open Label RCT to Evaluate a New Treatment Regimen for Patients with Multidrug-Resistant Tuberculosis. *ClinicalTrials.gov* Identifier: NCT02454205. Available from: <https://clinicaltrials.gov/ct2/show/NCT02454205>. [Last accessed on 2021 Dec 25].
- STREAM-II (Working Group for NEW TB Drugs). The Evaluation of a Standard Treatment Regimen of Anti-Tuberculosis Drugs for Patients with Multidrug-Resistant Tuberculosis. *ClinicalTrials.gov* Identifier: NCT02409290. Available from: <https://clinicaltrials.gov/ct2/show/NCT02409290>. [Last accessed on 2021 Dec 25].
- END-TB. Evaluating New Approved Drugs for Multidrug-Resistant TB. *ClinicalTrials.gov* Identifier: NCT02754765. Available from: <https://clinicaltrials.gov/ct2/show/NCT02754765>. [Last accessed on 2021 Dec 25].
- MDR-END. Treatment Shortening of MDR-TB using Existing and New Drugs. *ClinicalTrials.gov* Identifier: NCT02619994. Available from: <https://clinicaltrials.gov/ct2/show/NCT02619994>. [Last accessed on 2021 Dec 25].
- BEAT-TB. Building Evidence for Advancing New Treatment for Rifampin Resistant (RR-TB) Comparing a Short Course of Treatment (Containing Bedaquiline, Delamanid and Linezolid) with the Current South African Standard of Care. *ClinicalTrials.gov* Identifier: NCT04062201. Available from: <https://clinicaltrials.gov/ct2/show/NCT04062201>. [Last accessed on 2021 Dec 25].
- WHO Consolidated Guidelines on Tuberculosis, Module 2: Screening- Systematic Screening for Tuberculosis Disease. Available from: <https://www.who.int/publications/i/item/9789240022676>. [Last accessed on 2021 Dec 25].
- Xu J, Wang B, Hu M, Huo F, Guo S, Jing W, et al. Primary clofazimine and bedaquiline resistance among isolates from patients with multidrug-resistant tuberculosis. *Antimicrob Agents Chemother* 2017;61:e00239-17.
- Villellas C, Coeck N, Meehan CJ, Lounis N, de Jong B, Rigouts L, et al. Unexpected high prevalence of resistance-associated Rv0678 variants in MDR-TB patients without documented prior use of clofazimine or bedaquiline. *J Antimicrob Chemother* 2017;72:684-90.
- Nimmo C, Millard J, Brien K, Moodley S, van Dorp L, Lutchmarain K, et al. Bedaquiline resistance in drug-resistant tuberculosis HIV co-infected patients. *Eur Respir J* 2020;55:1902383.
- Derendinger B, Dippenaar A, De Vos M, Albert's R, Sirgel F, Dolby T, et al. High frequency of Bedaquiline resistance in programmatically treated drug-resistant TB patients with sustained culture-positivity in Cap Town, South Africa. *Int J Mycobacteriol* 2021;9:9.
- Mokrousov I, Akhmedova G, Molchanov V, Fundovnaya E, Kozlova E, Ostankova Y, et al. Frequent acquisition of bedaquiline resistance by epidemic extensively drug-resistant *Mycobacterium tuberculosis* strains in Russia during long-term treatment. *Clin Microbiol Infect* 2021;27:478-80.
- Ghodousi A, Rizvi AH, Baloch AQ, Ghafoor A, Khanzada FM, Qadir M, et al. Acquisition of cross-resistance to bedaquiline and clofazimine following treatment for tuberculosis in Pakistan. *Antimicrob Agents Chemother* 2019;63:e00915-19.
- Chesov E, Chesov D, Maurer FP, Andres S, Utpatel C, Barilar I, et al. Emergence of bedaquiline-resistance in a high-burden country of tuberculosis. *Eur Respir J* 2021;59:2100621.
- Köser CU, Maurer FP, Kranzer K. 'Those who cannot remember the past are condemned to repeat it': Drug-susceptibility testing for bedaquiline and delamanid. *Int J Infect Dis* 2019;80S: S32-5.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online	
<b>Quick Response Code:</b> 	<b>Website:</b> <a href="http://www.lungindia.com">www.lungindia.com</a>
	<b>DOI:</b> <a href="https://doi.org/10.4103/lungindia.lungindia_31_22">10.4103/lungindia.lungindia_31_22</a>

**How to cite this article:** Jain NK. Reply: Roll out of bedaquiline without drug susceptibility testing leads to emergence of bedaquiline-resistance in high burden countries with tuberculosis - Danger of losing a novel drug. *Lung India* 2022;39:389-90.

© 2022 Indian Chest Society | Published by Wolters Kluwer - Medknow