



POSTER PRESENTATION

Open Access

A transgenic in vitro cell model for the analysis of proinflammatory effects of naturally occurring genetic variants of caspase-1

F Schulze*, E Hengst, S Winkler, A Rösen-Wolff

From 8th International Congress of Familial Mediterranean Fever and Systemic Autoinflammatory Diseases Dresden, Germany. 30 September - 3 October 2015

Introduction

Caspase-1 (Interleukin-1 Converting Enzyme, ICE) is a proinflammatory enzyme mediating cleavage and secretion of the proinflammatory cytokines IL-1 β and IL-18. Caspase-1 plays pivotal roles in the innate immune system and in inflammatory diseases like periodic fever syndromes, arthritis, or type-II-diabetes. In previous studies published by our group, genetic variants of procaspase-1 had been detected in patients suffering from autoinflammatory symptoms. Analyses of the procaspase-1 variants in HEK293T cells revealed reduced enzymatic activity of caspase-1 but enhanced ability to activate NF- κ B signaling. The latter was mediated by enhanced CARD/CARD interactions of procaspase-1 with RIP2.

Objectives

The primary objective of this study was to analyze the effects of procaspase-1 variants in a monocyte/macrophage cell model. Furthermore, protein expression and enzymatic activity of procaspase-1 with or without a C-terminal FLAG-tag was analyzed.

Materials and methods

Genetically engineered THP-1 monocytes were used for the *in vitro* study. First, THP-1 cells were transduced with lentiviral vectors expressing shRNA against procaspase-1 mRNA. Subsequently, procaspase-1 wildtype (wt) or variants with or without a C-terminal FLAG-tag were reconstituted using a second lentiviral transduction. THP-1 cells were differentiated into macrophages and stimulated with different inflammasome activators. Activation and release of caspase-1 and IL-1 β was assessed

using immunoblotting (caspase-1) or immunoblotting and cytometric bead arrays (IL-1 β). The activation of NF- κ B was estimated by measuring the NF- κ B regulated cytokines IL-6 and IL-8. Cell death following inflammasome activation was analyzed by measuring LDH in the cell culture supernatant.

Results

The protein expression level of reconstituted FLAG-tagged procaspase-1 variants was significantly reduced compared to expression of endogenous procaspase-1 or reconstituted procaspase-1 variants without FLAG-tag. In line with this data, the release of IL-1 β after inflammasome stimulation was reduced in cells expressing FLAG-tagged procaspase-1 compared to cells expressing procaspase-1 without FLAG-tag. Interestingly, the mRNA expression of the FLAG-tagged procaspase-1 variants was not reduced compared to reconstituted procaspase-1 without FLAG-tag. Furthermore, cells expressing the procaspase-1 variants released reduced amounts of IL-1 β and showed a reduced frequency of cell death following inflammasome stimulation. No differences in IL-6 or IL-8 secretion were detected when cells expressing wt or enzymatically inactive procaspase-1 were compared.

Conclusion

This study shows that even short protein-tags can influence protein expression significantly. Therefore, phenotypic rescue in knockdown studies can be complicated when using tagged proteins. Using the genetically engineered THP-1 cells we were able to show a reduced IL-1 β release and reduced frequency of pyroptosis following inflammasome stimulation without detecting any differential regulation of the proinflammatory cytokines IL-6 or IL-8.

University Hospital Carl Gustav Carus, Department of Pediatrics, Dresden, Germany

This study was supported by the German Research Foundation (DFG, KFO 249) and by a MeDDrive project (University of Technology, Medical Faculty) to SW.

Published: 28 September 2015

doi:10.1186/1546-0096-13-S1-P18

Cite this article as: Schulze *et al.*: A transgenic in vitro cell model for the analysis of proinflammatory effects of naturally occurring genetic variants of caspase-1. *Pediatric Rheumatology* 2015 **13**(Suppl 1):P18.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

