

# Population-based cohort study of the impact on postoperative mortality of anastomotic leakage after anterior resection for rectal cancer

P. Boström<sup>1</sup> , M. M. Haapamäki<sup>1</sup>, J. Rutegård<sup>1</sup>, P. Matthiessen<sup>2</sup> and M. Rutegård<sup>1</sup> 

<sup>1</sup>Surgery Unit, Department of Surgical and Perioperative Sciences, Umeå University, Umeå, and <sup>2</sup>Department of Surgery, Faculty of Medicine and Health, School of Health and Medical Sciences, Örebro University, Örebro, Sweden

Correspondence to: Dr P. Boström, Department of Surgical and Perioperative Sciences, Surgery, Umeå University, Umeå University Hospital, SE-901 85 Umeå, Sweden (e-mail: petrus.bostrom@surgery.umu.se)

**Background:** Anastomotic leakage following anterior resection for rectal cancer may result in death. The aim of this study was to yield an updated, population-based estimate of postoperative mortality and evaluate possible interacting factors.

**Methods:** This was a retrospective national cohort study of patients who underwent anterior resection between 2007 and 2016. Data were retrieved from a prospectively developed database. Anastomotic leakage constituted exposure, whereas outcome was defined as death within 90 days of surgery. Logistic regression analyses, using directed acyclic graphs to evaluate possible confounders, were performed, including interaction analyses.

**Results:** Of 6948 patients, 693 (10.0 per cent) experienced anastomotic leakage and 294 (4.2 per cent) underwent reintervention due to leakage. The mortality rate was 1.5 per cent in patients without leakage and 3.9 per cent in those with leakage. In multivariable analysis, leakage was associated with increased mortality only when a reintervention was performed (odds ratio (OR) 5.57, 95 per cent c.i. 3.29 to 9.44). Leaks not necessitating reintervention did not result in increased mortality (OR 0.70, 0.25 to 1.96). There was evidence of interaction between leakage and age on a multiplicative scale ( $P = 0.007$ ), leading to a substantial mortality increase in elderly patients with leakage.

**Conclusion:** Anastomotic leakage, in particular severe leakage, led to a significant increase in 90-day mortality, with a more pronounced risk of death in the elderly.

## Funding information

Cancer Research Foundation in Northern Sweden  
Landstinget Västernorrland

Paper accepted 22 August 2018

Published online 15 October 2018 in Wiley Online Library (www.bjsopen.com). DOI: 10.1002/bjs5.50106

## Introduction

Every anastomosis has a risk of anastomotic leakage. For reasons only partly understood, this risk is greater for extraperitoneal colorectal anastomoses than for other enteric anastomoses<sup>1</sup>. Anastomotic leakage can lead to faecal peritonitis, sepsis and multiple organ failure, and is thus associated with substantial morbidity and mortality. Modern population-based prospective studies<sup>2,3</sup> on anterior resection for rectal cancer have estimated the risk of anastomotic leakage, depending on whether or not a stoma was created, to be 7.8–9 and 11.6–12 per cent respectively.

The mortality rate for patients with leakage was 5.7–7 per cent in these studies.

Following the introduction of total mesorectal excision for rectal cancer, the incidence of anastomotic leakage increased initially<sup>4,5</sup>. More recently, however, both the incidence and severity of anastomotic leak have decreased owing to a greater understanding of the risk factors for anastomotic leakage, the impact of protective measures, especially the construction of a diverting stoma, which mitigates the effects of leakage<sup>2,5–9</sup>, and improved postoperative management. Modern intensive care can support vital functions for longer than the traditional 30-day period during which postoperative complications

(including death) were considered to occur. For this reason, more recent studies<sup>10,11</sup> on postoperative complications have used an extended interval of 90 days so that the risk of adverse outcomes after surgery is not underestimated.

The aim of this study was to make an updated, population-based estimate of the incidence of anastomotic leakage and associated postoperative mortality. Possible interacting factors were also evaluated, including the use of a diverting stoma.

## Methods

This was a retrospective study of patients with rectal cancer included in the Swedish Cancer Registry. Since 2007, the Regional Oncological Centre in each healthcare region in Sweden has provided data for the national Swedish Colorectal Cancer Registry. All patients with colorectal cancer in Sweden have been reported to this registry, which has frequently been checked against the National Cancer Registry to ensure completeness. Data on patient characteristics, surgery, postoperative treatment, pathological assessment and follow-up for 5 years are recorded, with new cases stemming from clinicians and pathologists. The registry has been validated on a number of occasions, showing a level of completeness of 97 per cent regarding rectal cancer surgery<sup>12</sup>. The registry uses several categories for surgical complications, including wound infection, wound dehiscence, intra-abdominal infection, postoperative bleeding, anastomotic leakage, stoma complication, urinary catheter at discharge, or not specified.

Patients were included in the present study if their operation was registered as anterior resection for rectal cancer and performed in 2007–2016. Rectal cancer was defined as histologically proven adenocarcinoma, with its lower border within 15 cm of the anal verge, measured by rigid sigmoidoscopy. No exclusion criteria were used.

## Exposure and outcome

Primary exposure was defined as anastomotic leakage within 30 days of surgery or intervention for anastomotic leak, as recorded in the registry. Anastomotic leak was defined as leakage of the colorectal anastomosis, pelvic abscess or rectovaginal fistula<sup>13</sup>. As secondary exposure, leakage categorized as leaks not necessitating reintervention, as well as those where reintervention was used, was examined, to outline groups of patients with different severity of anastomotic leakage. Reoperation and radiologically performed drainage were defined as reinterventions by the registry, without defining which reintervention was used.

Postoperative mortality was defined as a registered death in the Swedish total population registry, within 90 days of surgery. This dichotomous outcome variable was used for logistic regression analyses.

## Hypotheses

The main hypothesis was that anastomotic leakage is associated with an increased 90-day mortality rate. A dose–response relationship was expected between the severity of anastomotic leakage and mortality, and that leakage would interact with patient-related and operative risk factors. In addition, the aim was to calculate an updated, population-based estimate of the relative risk of death within 90 days after anastomotic leakage, in a recent time period.

## Statistical analyses

Univariable associations were evaluated with the  $\chi^2$  test and Fisher's exact test, as appropriate. Unconditional logistic regression was performed to yield single estimates, measured as odds ratios (ORs) with 95 per cent c.i. of the independent risk of postoperative death at 90 days.

To adjust for confounding the concept of directed acyclic graphs<sup>14–16</sup> was used, and biologically mechanistic reasoning as to how different possible confounders influence exposure and outcome. This model (*Fig. S1*, supporting information) is considered an unbiased measure of the total effect of anastomotic leakage and reintervention for leakage on postoperative mortality. It could be calculated by adjusting for age (continuous), hospital volume (divided by tertiles), ASA grade (I, II, III–IV), presence of a diverting stoma (yes or no) and intraoperative blood loss (using the median: 300 ml or less *versus* more than 300 ml). Mixed models were used to accommodate for clustering within hospitals. The hypothetical dose–response relationship was evaluated by grading leakage severity according to need for reintervention. Multiplicative interaction analyses were conducted of overall anastomotic leakage with every co-variable, using a backward-elimination procedure. To visualize the interaction results, predicted probabilities of death were calculated and plotted in graphical form.

The regression models were analysed using the Hosmer–Lemeshow test for goodness-of-fit, and tested for collinearity by calculating the variance inflation factor and evaluating *P* values. A complete-case analysis was used, thereby excluding observations with missing data. All tests for significance were two-sided, with the level of significance set at 5 per cent. All analyses were conducted in STATA® 13.1 (StataCorp, College Station, Texas, USA).

**Table 1** Clinical variables stratified by mortality within 90 days after surgery, for patients undergoing anterior resection for rectal cancer in Sweden 2007–2016

	Postoperative mortality within 90 days		P <sup>¶</sup>
	No (n = 6827)	Yes (n = 121)	
Anastomotic leakage			< 0.001
No	6161 (90.2)	94 (77.7)	
Yes	666 (9.8)	27 (22.3)	
Reintervention for leakage			< 0.001
No	6555 (96.0)	99 (81.8)	
Yes	272 (4.0)	22 (18.2)	
Age (years)			< 0.001
< 65	2668 (39.1)	11 (9.1)	
65–75	2851 (41.8)	44 (36.4)	
> 75	1308 (19.2)	66 (54.5)	
ASA fitness grade			< 0.001
I	1659 (24.3)	6 (5.0)	
II	3914 (57.3)	66 (54.5)	
III–IV	1138 (16.7)	48 (39.7)	
Tumour stage (pTNM)			0.651
I	1822 (26.7)	30 (24.8)	
II	1886 (27.6)	37 (30.6)	
III	2335 (34.2)	41 (33.9)	
IV	468 (6.9)	5 (4.1)	
Radiotherapy			0.004
Short course*	1983 (29.0)	38 (31.4)	
Long course†	488 (7.2)	1 (0.8)	
Other	21 (0.3)	1 (0.8)	
None	4335 (63.5)	81 (66.9)	
Tumour level (cm)‡			0.336
≤ 6	509 (7.5)	9 (7.4)	
7–12	4512 (66.1)	73 (60.3)	
13–15	1806 (26.5)	39 (32.2)	
Type of anastomosis			0.173
End-to-end	1906 (27.9)	40 (33.1)	
Side-to-end/J pouch	4496 (65.9)	71 (58.7)	
Surgical technique			0.039
Open surgery	5545 (81.2)	99 (81.8)	
Laparoscopy, not converted	1003 (14.7)	12 (9.9)	
Laparoscopy, converted	238 (3.5)	9 (7.4)	
Diverting stoma			0.015
Yes	5249 (76.9)	82 (67.8)	
No	1578 (23.1)	39 (32.2)	
Blood loss (ml)			0.013#
≤ 300	3413 (50.0)	52 (43.0)	
> 300	3414 (50.0)	69 (57.0)	
Surgical complication§			< 0.001
No	5402 (79.1)	74 (61.2)	
Yes	1425 (20.9)	47 (38.8)	
Neurological complication			0.038
No	6811 (99.8)	119 (98.3)	
Yes	16 (0.2)	2 (1.7)	

**Table 1** Continued

	Postoperative mortality within 90 days		P <sup>¶</sup>
	No (n = 6827)	Yes (n = 121)	
Non-surgical infection			< 0.001
No	6383 (93.5)	92 (76.0)	
Yes	444 (6.5)	29 (24.0)	
Cardiovascular complication			< 0.001
No	6653 (97.5)	87 (71.9)	
Yes	174 (2.5)	34 (28.1)	

Values in parentheses are percentages; percentages may not sum to 100 owing to missing values for some variables. \*5 × 5 Gy; †1.8–2.0 Gy to a total of 46–50.4 Gy. ‡Measured from anal verge. §Surgical complication noted in the registry as any of the following: wound infection, wound dehiscence, intra-abdominal infection, postoperative bleeding, anastomotic leakage, stoma complication, urinary catheter at discharge or not specified. ¶ $\chi^2$  or Fisher's exact test, except #Mann–Whitney *U* test (blood loss analysed as a continuous variable).

## Results

Some 6948 patients were identified using the registry. The majority were men (4103, 59.1 per cent) with only mild systemic disease (ASA grade II). The median age was 67 years, and half the population was aged 60–74 years. Tumour stages I–III were roughly equally common, whereas a minority of tumours were stage IV cancers. Most tumours were located at 7–12 cm, and only 518 (7.5 per cent) were 6 cm or less from the anal verge. A total of 121 patients (1.7 per cent) died within 90 days of surgery (median 25 (i.q.r. 6–46) days).

A diverting stoma was constructed in 5331 patients (76.7 per cent). Some 1262 patients (18.2 per cent) were operated on laparoscopically. Of these operations, 247 were converted to open surgery (conversion rate 19.6 per cent). A J pouch or side-to-end anastomosis was favoured over end-to-end anastomosis. *Table 1* outlines the frequencies of patient characteristics, operative data and postoperative complications, stratified by mortality.


## Anastomotic leakage and postoperative mortality

In total, 693 of 6948 patients (10.0 per cent) developed anastomotic leakage. The mortality rate in these patients was 3.9 per cent, compared with 1.5 per cent in patients without leakage. In univariable logistic regression analysis, anastomotic leak was associated with increased mortality (OR 2.66, 95 per cent c.i. 1.72 to 4.11). This association was similar in multivariable analysis (OR 2.64, 1.65 to 4.22) (*Table 2*). In the 294 patients (4.2 per cent) who underwent reintervention for leakage, the mortality rate was 7.5 per cent.

**Table 2** Frequency and logistic regression analysis of postoperative mortality within 90 days after anastomotic leak and reintervention for leakage

	No. of leaks	No. of deaths/leak*	Univariable analysis Odds ratio†	Multivariable analysis‡ Odds ratio†
All leaks	693	27 (3.9)	2.66 (1.72, 4.11)	2.64 (1.65, 4.22)
Leak without reintervention	399	5 (1.3)	0.83 (0.34, 2.06)	0.70 (0.25, 1.96)
Leak with reintervention	294	22 (7.5)	5.30 (3.28, 8.57)	5.57 (3.29, 9.44)

Values in parentheses are \*percentages and †95 per cent confidence intervals. ‡Adjusted for age, ASA grade, diverting stoma, hospital volume, intraoperative blood loss and clustering within hospitals.

**Fig. 1** Predicted 90-day postoperative mortality after anterior resection for rectal cancer according to age, for patients with and without anastomotic leakage

In the secondary analysis of leakage severity, only anastomotic leakage with reintervention was associated with increased mortality (OR 5.57, 95 per cent c.i. 3.29 to 9.44). For leakage without intervention, no statistically significant association was observed (OR 0.70, 0.25 to 1.96) (Table 2).

### Interaction analyses

On a multiplicative scale, stepwise interaction analyses were performed to eliminate non-significant variables (hospital volume, intraoperative bleeding, ASA grade and diverting stoma). The only interaction variable that remained was age (OR 1.09, 95 per cent c.i. 1.02 to 1.17). The variable age had a high variance inflation factor (41.5), but a low *P* value (*P*=0.007), and was therefore retained in the analyses. To visualize these interaction results, predicted probabilities were calculated (Fig. 1). With increasing age, mortality increased substantially, most prominently after anastomotic leakage.

### Discussion

This population-based national cohort study suggests that anastomotic leakage results in increased mortality in the

prolonged postoperative period. The overall mortality rate was 3.9 per cent in patients with anastomotic leakage. No increased mortality was observed in patients with anastomotic leakage that did not require reintervention. Substantial interaction was detected between leakage and age, indicating that this complication, if present, is more detrimental in older patients.

The major strength of the study is its large, population-based, cohort design. Estimates of this study are therefore robust, reflecting the whole population of interest, namely patients undergoing anterior resection for rectal cancer today. Small numbers, however, may still have affected some secondary analyses.

The ambiguity in the term anastomotic leakage is problematic for all research aimed at quantifying the rate of leakage<sup>17</sup>. The registry used allows the surgeon to define leakage arbitrarily, thereby threatening the validity of the data on which this study was based. It also provides clinical heterogeneity, perhaps explaining why an association was found only for patients requiring reintervention. In addition, the registry does not allow differentiation between different reinterventions. It would be interesting to demarcate further the group at greatest risk of death after

leakage, by separating relaparotomy from simple drainage. Another drawback is the delay in diagnosing leakage; at present, approximately one-third of all leaks are diagnosed after 30 postoperative days<sup>18</sup>. In addition, as the registry probably underestimates the true incidence even within that time period<sup>19,20</sup>, the estimates presented here are most likely conservative measures.

In the present study, the mortality rates observed following anastomotic leakage are similar to those of other recently published studies on population-based data<sup>17,21</sup>. The association between anastomotic leak and increased postoperative mortality is reasonable pathophysiologically, and has been shown previously for colorectal surgery<sup>2,22</sup>. For rectal cancer surgery specifically, Snijders and colleagues<sup>23</sup> tried to quantify this effect in a meta-analysis of 22 studies; they concluded that one-third of all postoperative mortality is caused directly by anastomotic leakage. In contrast, the association between age and complication rates, morbidity and mortality is less obvious and difficult to study, partly as randomization of the exposure variable is impossible. Age has, however, been correlated with increased postoperative mortality in studies of colorectal surgery<sup>24,25</sup>.

The driving force behind this increased mortality is not clear, and studies of the correlation between age and complication rates must be well designed to control for the increased co-morbidity and frailty of the elderly population<sup>26</sup>. However, Marusch and co-workers<sup>27</sup> conducted a multicentre study of 309 hospitals, including 19 080 patients who had surgery for colorectal carcinoma, and demonstrated increased complication rates among octogenarians, when controlling for confounders. Aquina *et al.*<sup>28</sup> evaluated 24 426 patients with stage I–III colonic cancer, while controlling for confounders, and similarly found a dose–response relationship between increasing complication rates and increasing age.

Regarding the association between advanced age and the development of anastomotic leak, the picture is less clear. Den Dulk and colleagues<sup>2</sup> summarized data from five European trials of colorectal surgery between 1987 and 2003, and found an association between age and mortality after leakage, but not between age and leakage itself. For anterior resection specifically, an observational single-centre study by Lin *et al.*<sup>29</sup>, of 999 patients in 1993–2003, showed age above the median to be a risk factor for anastomotic leakage in multivariable analysis (OR 2.2, 95 per cent c.i. 1.21 to 3.88). In contrast, Parthasarathy and co-workers<sup>30</sup> used prospectively collected data on 17 518 patients in 2013 and found a higher risk of anastomotic leakage among younger patients undergoing colorectal resection. A meta-analysis by

Pommergaard and colleagues<sup>31</sup> on resection of colorectal cancer found 21 studies that evaluated the effect of age, of which only two showed an association; seven of these studies were used to calculate a pooled OR of 0.99 (0.89 to 1.10).

Although advanced age is not an established risk factor for anastomotic leak, the consequences of leakage seem to be aggravated in the elderly. It is therefore important to consider the use of possible protective measures, including a temporary or definitive stoma. Better individual risk-profiling is warranted, but the findings of the present study may help to counsel patients and improve shared decision-making in the preoperative setting.

### Acknowledgements

This study was supported by the Cancer Research Foundation in Northern Sweden and Landstinget Västernorrland.

*Disclosure:* The authors declare no conflict of interest.

### References

- 1 Platell C, Barwood N, Dorfmann G, Makin G. The incidence of anastomotic leaks in patients undergoing colorectal surgery. *Colorectal Dis* 2007; **9**: 71–79.
- 2 den Dulk M, Marijnen CA, Collette L, Putter H, Pahlman L, Folkesson J *et al.* Multicentre analysis of oncological and survival outcomes following anastomotic leakage after rectal cancer surgery. *Br J Surg* 2009; **96**: 1066–1075.
- 3 Bakker IS, Snijders HS, Wouters MW, Havenga K, Tollenaar RA, Wiggers T *et al.* High complication rate after low anterior resection for mid and high rectal cancer; results of a population-based study. *Eur J Surg Oncol* 2014; **40**: 692–698.
- 4 Carlsen E, Schlichting E, Guldvog I, Johnson E, Heald RJ. Effect of the introduction of total mesorectal excision for the treatment of rectal cancer. *Br J Surg* 1998; **85**: 526–529.
- 5 Peeters KC, Tollenaar RA, Marijnen CA, Klein Kranenbarg E, Steup WH, Wiggers T *et al.*; Dutch Colorectal Cancer Group. Risk factors for anastomotic failure after total mesorectal excision of rectal cancer. *Br J Surg* 2005; **92**: 211–216.
- 6 Matthiessen P, Hallböök O, Andersson M, Rutegård J, Sjö Dahl R. Risk factors for anastomotic leakage after anterior resection of the rectum. *Colorectal Dis* 2004; **6**: 462–469.
- 7 Matthiessen P, Hallböök O, Rutegård J, Simert G, Sjö Dahl R. Defunctioning stoma reduces symptomatic anastomotic leakage after low anterior resection of the rectum for cancer: a randomized multicenter trial. *Ann Surg* 2007; **246**: 207–214.
- 8 Bertelsen CA, Andreasen AH, Jørgensen T, Harling H; Danish Colorectal Cancer Group. Anastomotic leakage after anterior resection for rectal cancer: risk factors. *Colorectal Dis* 2010; **12**: 37–43.

- 9 Marusch F, Koch A, Schmidt U, Geibetaler S, Dralle H, Saeger HD *et al.* Value of a protective stoma in low anterior resections for rectal cancer. *Dis Colon Rectum* 2002; **45**: 1164–1171.
- 10 van Gestel YR, Lemmens VE, de Hingh IH, Steevens J, Rutten HJ, Nieuwenhuijzen GA *et al.* Influence of comorbidity and age on 1-, 2-, and 3-month postoperative mortality rates in gastrointestinal cancer patients. *Ann Surg Oncol* 2013; **20**: 371–380.
- 11 Damhuis RA, Wijnhoven BP, Plaisier PW, Kirkels WJ, Kranse R, van Lanschot JJ. Comparison of 30-day, 90-day and in-hospital postoperative mortality for eight different cancer types. *Br J Surg* 2012; **99**: 1149–1154.
- 12 Pählman L, Bohe M, Cedermark B, Dahlberg M, Lindmark G, Sjö Dahl R *et al.* The Swedish Rectal Cancer Registry. *Br J Surg* 2007; **94**: 1285–1292.
- 13 Rahbari NN, Weitz J, Hohenberger W, Heald RJ, Moran B, Ulrich A *et al.* Definition and grading of anastomotic leakage following anterior resection of the rectum: a proposal by the International Study Group of Rectal Cancer. *Surgery* 2010; **147**: 339–351.
- 14 Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology* 1999; **10**: 37–48.
- 15 Shrier I, Platt RW. Reducing bias through directed acyclic graphs. *BMC Med Res Methodol* 2008; **8**: 70.
- 16 Morgan SL. *Handbook of Causal Analysis for Social Research*. Springer: New York, 2013.
- 17 Frouws MA, Snijders HS, Malm SH, Liefers GJ, Van de Velde CJH, Neijenhuis PA *et al.* Clinical relevance of a grading system for anastomotic leakage after low anterior resection: analysis from a National Cohort Database. *Dis Colon Rectum* 2017; **60**: 706–713.
- 18 Morks AN, Ploeg RJ, Sijbrand Hofker H, Wiggers T, Havenga K. Late anastomotic leakage in colorectal surgery: a significant problem. *Colorectal Dis* 2013; **15**: e271–e275.
- 19 Gunnarsson U, Seligsohn E, Jestin P, Pählman L. Registration and validity of surgical complications in colorectal cancer surgery. *Br J Surg* 2003; **90**: 454–459.
- 20 Rutegård M, Kverneng Hultberg D, Angenete E, Lydrup ML. Substantial underreporting of anastomotic leakage after anterior resection for rectal cancer in the Swedish Colorectal Cancer Registry. *Acta Oncol* 2017; **56**: 1741–1745.
- 21 Matsubara N, Miyata H, Gotoh M, Tomita N, Baba H, Kimura W *et al.* Mortality after common rectal surgery in Japan: a study on low anterior resection from a newly established nationwide large-scale clinical database. *Dis Colon Rectum* 2014; **57**: 1075–1081.
- 22 Krarup PM, Nordholm-Carstensen A, Jorgensen LN, Harling H. Association of comorbidity with anastomotic leak, 30-day mortality, and length of stay in elective surgery for colonic cancer: a nationwide cohort study. *Dis Colon Rectum* 2015; **58**: 668–676.
- 23 Snijders HS, Wouters MW, van Leersum NJ, Kolfschoten NE, Henneman D, de Vries AC *et al.* Meta-analysis of the risk for anastomotic leakage, the postoperative mortality caused by leakage in relation to the overall postoperative mortality. *Eur J Surg Oncol* 2012; **38**: 1013–1019.
- 24 Faiz O, Haji A, Bottle A, Clark SK, Darzi AW, Aylin P. Elective colonic surgery for cancer in the elderly: an investigation into postoperative mortality in English NHS hospitals between 1996 and 2007. *Colorectal Dis* 2011; **13**: 779–785.
- 25 Morris EJ, Taylor EF, Thomas JD, Quirke P, Finan PJ, Coleman MP *et al.* Thirty-day postoperative mortality after colorectal cancer surgery in England. *Gut* 2011; **60**: 806–813.
- 26 Saxton A, Velanovich V. Preoperative frailty and quality of life as predictors of postoperative complications. *Ann Surg* 2011; **253**: 1223–1229.
- 27 Marusch F, Koch A, Schmidt U, Steinert R, Ueberrueck T, Bittner R *et al.*; Working Group Colon/Rectum Cancer. The impact of the risk factor ‘age’ on the early postoperative results of surgery for colorectal carcinoma and its significance for perioperative management. *World J Surg* 2005; **29**: 1013–1021.
- 28 Aquina CT, Mohile SG, Tejani MA, Becerra AZ, Xu Z, Hensley BJ *et al.* The impact of age on complications, survival, and cause of death following colon cancer surgery. *Br J Cancer* 2017; **116**: 389–397.
- 29 Lin JK, Yueh TC, Chang SC, Lin CC, Lan YT, Wang HS *et al.* The influence of fecal diversion and anastomotic leakage on survival after resection of rectal cancer. *J Gastrointest Surg* 2011; **15**: 2251–2261.
- 30 Parthasarathy M, Greensmith M, Bowers D, Groot-Wassink T. Risk factors for anastomotic leakage after colorectal resection: a retrospective analysis of 17 518 patients. *Colorectal Dis* 2017; **19**: 288–298.
- 31 Pommergaard HC, Gessler B, Burcharth J, Angenete E, Haglind E, Rosenberg J. Preoperative risk factors for anastomotic leakage after resection for colorectal cancer: a systematic review and meta-analysis. *Colorectal Dis* 2014; **16**: 662–671.

### Supporting information

Additional supporting information can be found online in the Supporting Information section at the end of the article.