

Biology and biotechnology of *Trichoderma*

André Schuster · Monika Schmoll

Received: 22 February 2010 / Revised: 16 April 2010 / Accepted: 17 April 2010 / Published online: 12 May 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract Fungi of the genus *Trichoderma* are soilborne, green-spored ascomycetes that can be found all over the world. They have been studied with respect to various characteristics and applications and are known as successful colonizers of their habitats, efficiently fighting their competitors. Once established, they launch their potent degradative machinery for decomposition of the often heterogeneous substrate at hand. Therefore, distribution and phylogeny, defense mechanisms, beneficial as well as deleterious interaction with hosts, enzyme production and secretion, sexual development, and response to environmental conditions such as nutrients and light have been studied in great detail with many species of this genus, thus rendering *Trichoderma* one of the best studied fungi with the genome of three species currently available. Efficient biocontrol strains of the genus are being developed as promising biological fungicides, and their weaponry for this function also includes secondary metabolites with potential applications as novel antibiotics. The cellulases produced by *Trichoderma reesei*, the biotechnological workhorse of the genus, are important industrial products, especially with respect to production of second generation biofuels from cellulosic waste. Genetic engineering not only led to significant improvements in industrial processes but also to intriguing insights into the biology of these fungi and is now complemented by the availability of a sexual cycle in

T. reesei/Hypocrea jecorina, which significantly facilitates both industrial and basic research. This review aims to give a broad overview on the qualities and versatility of the best studied *Trichoderma* species and to highlight intriguing findings as well as promising applications.

Keywords *Trichoderma* · *Hypocrea* · Cellulase · Biocontrol · Heterologous protein expression · Emerging human pathogen · Green mold disease · Biodiversity · Application · Biofuels

Biodiversity and phylogeny of *Trichoderma*

The first description of a fungus named *Trichoderma* dates back to 1794 (Persoon 1794), and in 1865, a link to the sexual state of a *Hypocrea* species was suggested (Tulasne and Tulasne 1865). However, the different species assigned to the genus *Trichoderma/Hypocrea* were difficult to distinguish morphologically. It was even proposed to reduce taxonomy to only a single species, *Trichoderma viride*. Hence, it took until 1969 that development of a concept for identification was initiated (Rifai 1969; Samuels 2006). Thereafter, numerous new species of *Trichoderma/Hypocrea* were discovered, and by 2006, the genus already comprised more than 100 phylogenetically defined species (Druzhinina et al. 2006a). In some cases, especially in earlier reports, misidentifications of certain species occurred, for example with the name *Trichoderma harzianum* which has been used for many different species (Kullnig et al. 2001). However, it is difficult to safely correct these mistakes without analyzing the strains originally used, and therefore, we describe the data using the names as originally reported. In recent years, safe identification of new species was significantly facilitated by

A. Schuster · M. Schmoll (✉)
Research Area Gene Technology and Applied Biochemistry,
Institute of Chemical Engineering,
Vienna University of Technology,
Getreidemarkt 9/166-5,
1060 Vienna, Austria
e-mail: monika.schmoll@tuwien.ac.at

A. Schuster
e-mail: andre.schuster@tuwien.ac.at

development of an oligonucleotide barcode (TrichoKEY) and a customized similarity search tool (TrichoBLAST), both available online at www.isth.info (Druzhinina et al. 2005; Kopchinskiy et al. 2005). A further useful tool for characterization of newly isolated *Trichoderma* species (but also recombinant strains) are phenotype microarrays, which allow for investigation of carbon utilization patterns for 96 carbon sources (Bochner et al. 2001; Kubicek et al. 2003; Druzhinina et al. 2006b). The continued efforts to elucidate diversity and geographical occurrence of *Trichoderma/Hypocrea* resulted in detailed documentations of the genus in Europe and worldwide (Samuels et al. 2002a; Chaverri and Samuels 2003; Jaklitsch 2009; <http://nt.ars-grin.gov/taxadescriptions/keys/TrichodermaIndex.cfm>)

The Index Fungorum database (<http://www.indexfungorum.org/Names/Names.asp>) currently even lists 471 different names for *Hypocrea* species and 165 records for *Trichoderma*. However, many of these names have been introduced long before molecular methods for species identification were available and thus are likely to have become obsolete in the meantime. At present, the International Subcommittee on *Trichoderma/Hypocrea* lists 104 species (<http://www.isth.info/biodiversity/index.php>), which have been characterized at the molecular level. Seventy-five species of *Hypocrea* have been identified in temperate Europe (Jaklitsch 2009). Nevertheless, a considerable number of putative *Hypocrea* strains and even more *Trichoderma* strains, for which sequences have been deposited in GenBank, are still without safe identification (Druzhinina et al. 2006a) and remain to be studied further. Species of the genus produce a broad array of pigments from bright greenish-yellow to reddish in color, although some are also colorless. Similarly, conidial pigmentation varies from colorless to various green shades and sometimes also gray or brown. Other than pigmentation, species identification within the genus is difficult because of the narrow range of variation of the simplified morphology in *Trichoderma* (Gams and Bissett 1998).

Characteristics of *Trichoderma* spp.

Trichoderma spp. are ubiquitous colonizers of cellulosic materials and can thus often be found wherever decaying plant material is available (Kubicek et al. 2008; Jaklitsch 2009) as well as in the rhizosphere of plants, where they can induce systemic resistance against pathogens (Harman 2000). Some characteristic features of different *Trichoderma* spp. are shown in Fig. 1. The search for potent biomass degrading enzymes and organisms also led to isolation of these fungi from unexpected sources, such as cockroaches (Yoder et al. 2008), marine mussels and shellfish (Sallenave et al. 1999; Sallenave-Namont et al. 2000), or termite guts

(Sreerama and Veerabhadrapa 1993). *Trichoderma* spp. are characterized by rapid growth, mostly bright green conidia and a repetitively branched conidiophore structure (Gams and Bissett 1998).

Despite the early suggested link between *Trichoderma* and *Hypocrea* (Tulasne and Tulasne 1865), this anamorph–teleomorph relationship was only confirmed more than 100 years later for *Trichoderma reesei* and *Hypocrea jecorina* (Kuhls et al. 1996). Nevertheless, *T. reesei* was then termed a clonal, asexual derivative of *H. jecorina* because all attempts to cross the available strains of this species had failed. It took more than a decade until a sexual cycle was reported in any *Trichoderma* species (Seidl et al. 2009a), and a detailed study on molecular evolution of this species led to the discovery of a described sympatric agamospecies *Trichoderma parareesei* (Druzhinina et al. 2010). Especially because of the industrial importance of *T. reesei*, the availability of a sexual cycle was a groundbreaking discovery and now paves the way for elucidation of sexual development also in other members of the genus.

Trichoderma spp. are highly successful colonizers of their habitats, which is reflected both by their efficient utilization of the substrate at hand as well as their secretion capacity for antibiotic metabolites and enzymes. They are able to deal with such different environments as the rich and diversified habitat of a tropical rain forest as well as with the dark and sterile setting of a biotechnological fermentor or shake flask. Under all these conditions, they respond to their environment by regulation of growth, conidiation, enzyme production, and hence adjust their lifestyle to current conditions, which can be exploited for the benefit of mankind. One of these environmental factors is the presence or absence of light. *Trichoderma* has a long tradition of research toward the effect of light on its physiology and development, which already started in 1957 and largely paralleled that of *Phycomyces blakesleeana* (Schmoll et al. 2010). Besides effects on growth, reproduction, and secondary metabolite biosynthesis, which are common light responses in fungi, also a surprising influence of light on cellulase gene expression has been found (Schmoll et al. 2005). This link between light response and metabolic processes was further substantiated by a study on carbon source utilization using phenotype microarrays in light and darkness (Friedl et al. 2008). Studies on the molecular basis of these light effects revealed that interconnections between the signaling pathways of light response, heterotrimeric G-proteins, the cAMP-pathway, sulfur metabolism, and oxidative stress are operative in *Trichoderma* (Schmoll et al. 2010; Tisch and Schmoll 2010).

In recent years, research with *Trichoderma* has been facilitated significantly by sequencing of the genomes of three strains representing the most important applications of this genus: The genome sequence of *T. reesei*, the industrial workhorse (Martinez et al. 2008; <http://genome.jgi-psf.org/>




Fig. 1 Characteristic features of *Trichoderma* spp. **a** *T. reesei* and **b** *T. atroviride* growing on plates, **c** *T. reesei* or *H. jecorina* growing in daylight and showing light responsive conidiation, **d** fruiting body formation of *T. reesei* upon crossing with a nature isolate of *H.*

jecorina, (**e, f**) *T. longibrachiatum* germinating and growing on human cells, **g, i** *T. reesei* (left) during confrontation with *Pythium ultimum* (right), **h** *T. atroviride* (left) during confrontation with *R. solani* (right)

Trire2/Trire2.home.html), surprisingly revealed that, despite its importance in industrial cellulase production, its genome comprises the fewest amount of genes encoding cellulolytic and hemicellulolytic enzymes. Analysis and annotation of the genomes of *Trichoderma atroviride* and *Trichoderma virens*, two important biocontrol species (<http://genome.jgi-psf.org/Triat1/Triat1.home.html>; <http://genome.jgi-psf.org/Trive1/Trive1.home.html>), is still in progress. Interestingly, the genomes of *T. atroviride* and *T. virens* are significantly larger than that of *T. reesei*, and they comprise roughly 2000 genes more than does *T. reesei*. It will be interesting to learn the significance of this considerable difference in genome sizes in the physiology of these fungi. These

milestones in research with *Trichoderma* enabled detailed studies, which provided intriguing insights into their lifestyle, physiology, and the underlying mechanisms at the molecular level (Brunner et al. 2008; Martinez et al. 2008; Schmoll 2008; Le Crom et al. 2009; Seidl et al. 2009b).

Tools for genetic manipulation of *Trichoderma*

Due to the industrial application of *T. reesei*, the genetic toolkit for this fungus is the most extensive of the genus, although also research with other species is not limited by technical obstacles and most tools can also be used for all

species with slight modifications. Transformation of many species is possible, and different approaches such as protoplasting (Gruber et al. 1990), *Agrobacterium*-mediated transformation (Zeilinger 2004), or biolistic transformation (Lorito et al. 1993) were developed. The range of selectable marker cassettes, which includes hygromycin (Mach et al. 1994) and benomyl resistance (Peterbauer et al. 1992; Schuster et al. 2007), the *Aspergillus nidulans amdS* gene, which enables growth on acetamide as sole nitrogen source (Penttila et al. 1987) as well as the auxotrophic markers, *pyr4* (Gruber et al. 1990), *arg2* (Baek and Kenerley 1998), and *hvk1* (Guangtao et al. 2010) allows for construction of multiple mutants, which is now facilitated by the availability of a *T. reesei* strain with perturbed nonhomologous end-joining pathway (Guangtao et al. 2009). Sequential deletions despite a limited number of selection markers became possible by the use of a blaster cassette comprising direct repeats for homologous recombination and excision of the marker gene (Hartl and Seiboth 2005). Besides knockout strategies for functional analysis of genes, also expression of antisense constructs for knockdown (Rocha-Ramirez et al. 2002; Moreno-Mateos et al. 2007; Schmoll et al. 2009) was reported for *Trichoderma*, and RNAi has been shown to function in *T. reesei* (Brody and Suchindra 2009). Last but not least, the recent discovery of a sexual cycle in *T. reesei* (Seidl et al. 2009a) further boosts the versatility of this fungus for research and industry.

Defense mechanisms and their exploitation

Successful colonization of a given habitat by any organism is crucially dependent on its potential to defend its ecological niche and to thrive and prosper despite competition for nutrients, space, and light. Many fungi and especially those of the genus *Trichoderma* are masters of this game (Herrera-Estrella and Chet 2004; Harman 2006; Vinale et al. 2008). Their defense mechanisms comprise both enzymatic and chemical weapons, which make *Trichoderma* spp. efficient mycoparasites, antagonists, and biocontrol agents—characteristics that can be exploited by using *Trichoderma* spp. or the metabolites secreted by these fungi as biological fungicides to fight plant diseases caused by pathogenic fungi (Spiegel and Chet 1998; Vinale et al. 2006; Navazio et al. 2007; Vinale et al. 2009). Thereby *Trichoderma* spp. play an important role in the three-way interaction with the plant and the pathogen (Lu et al. 2004; Woo et al. 2006).

Trichoderma's strategies for combat

After publication of *Trichoderma lignorum* (later found to be *T. atroviride*) acting as a parasite on other fungi in 1932 (Weindling 1932), research on antagonistic properties of

Trichoderma spp. progressed rapidly. Nowadays, the most important species in this field are *T. atroviride* (in earlier reports sometimes misidentified as *T. harzianum*), *T. harzianum*, *T. virens*, and *Trichoderma asperellum* (Benitez et al. 2004), while *T. reesei*, the biotechnological workhorse, can rather be seen as a model organism used because of the established molecular biological methods and available recombinant strains (Seidl et al. 2006). *Trichoderma* spp. are able to control ascomycetes, basidiomycetes, and oomycetes (Monte 2001; Benitez et al. 2004), and recently, also their effect on nematodes was reported (Dababat et al. 2006; Kyalo et al. 2007; Goswami et al. 2008)

In their defensive actions, *Trichoderma* spp. apply lytic enzymes (Kubicek et al. 2001; Viterbo et al. 2002), proteolytic enzymes (Kredics et al. 2005; Suarez et al. 2007; Chen et al. 2009), ABC transporter membrane pumps (Ruocco et al. 2009), diffusible or volatile metabolites (Calistru et al. 1997; Eziashi et al. 2006), and other secondary metabolites (Reino et al. 2008) as active measures against their hosts or they succeed by their impairing growth conditions of pathogens (Benitez et al. 2004). Interestingly, the success of these actions is not independent of the surrounding temperature (Mukherjee and Raghu 1997), which can be crucial for the use as a biocontrol agent in different climates. Large-scale studies on gene expression during biocontrol at least in part reflect these findings (Grinyer et al. 2005; Marra et al. 2006; Samolski et al. 2009) and reveal additional components with potential effectivity such as a superoxide dismutase (Grinyer et al. 2005) and amino acid oxidase (Tseng et al. 2008) to be secreted under these conditions. Moreover, the response of *Trichoderma* to its host has also been shown to involve stress response, response to nitrogen shortage, cross pathway control, lipid metabolism, and signaling processes (Seidl et al. 2009b).

Regulatory mechanisms triggering the defense of *Trichoderma*

Signal transduction pathways triggering the genes involved in biocontrol and mycoparasitism have been studied in considerable depth and include heterotrimeric G-protein signaling, mitogen-activated protein kinase (MAPK) cascades, and the cAMP pathway (Zeilinger and Omann 2007). Especially the MAP-kinase TVK1, characterized in *T. virens* (Mendoza-Mendoza et al. 2003; Mukherjee et al. 2003; Mendoza-Mendoza et al. 2007) as well as its orthologs in *T. asperellum* (TmkA; Viterbo et al. 2005) and *T. atroviride* (TMK1; Reithner et al. 2007), is important in regulation of signaling mechanisms targeting output pathways relevant for efficient biocontrol. Transcript levels of the respective genes increased upon interaction with

plant roots in *T. virens* and *T. asperellum* (Viterbo et al. 2005). Deletion of *T. atroviride* *tmk1* causes higher antifungal activity and improved protection against *Rhizoctonia solani* but reduced mycoparasitic activity (Reithner et al. 2007). In agreement with this study, lack of *T. virens* TVK1 considerably increases biocontrol effectivity of this fungus (Mendoza-Mendoza et al. 2003). Hence, although deletion of the respective genes causes reduced mycoparasitic efficiency, the biocontrol abilities of the mutant strains are enhanced.

As for the action of the pathway of heterotrimeric G-protein signaling, two genes have been studied so far with respect to biocontrol related mechanisms in *Trichoderma* spp.: the class I (adenylate cyclase inhibiting) G-alpha subunits TGA1 of *T. atroviride* and TgaA of *T. virens* as well as the class III (adenylate cyclase activating) G-alpha subunits TGA3 of *T. atroviride* and GNA3 of *T. reesei*. TGA1 plays an important role in regulation of coiling around host hyphae and regulates production of antifungal metabolites. Lack of TGA1 results in enhanced growth inhibition of host fungi (Rocha-Ramirez et al. 2002; Reithner et al. 2005). For TgaA, a host specific involvement as shown in case of the action of MAP-kinases has been reported (Mukherjee et al. 2004). TGA3 on the other hand is crucial for biocontrol since deletion of the corresponding gene resulted in avirulent strains (Zeilinger et al. 2005). Since constitutive activation of GNA3 in *T. reesei* is suggested to positively influence mycoparasitism, a similar mechanism, may be at work in this fungus (Silva et al. 2009). These results are in agreement with analysis of cAMP signaling components, which indicate a positive role of cAMP in biocontrol (Mukherjee et al. 2007). Recently, also an important role in biocontrol of *T. virens* has been reported for the homolog of the VELVET protein, so far mainly known as light-dependent regulator protein (Mukherjee and Kenerley 2010).

Attempts were made to identify characteristics among all these genes and enzymes regulated upon interaction of *Trichoderma* with a pathogen, which could be used to distinguish efficient from nonefficient biocontrol strains isolated from nature (Nagy et al. 2007; Scherm et al. 2009). However, only further, extensive studies will reveal the reliability of standardized marker gene assays for evaluation of potential biocontrol strains.

Trichoderma as a protector of plant health

The beneficial action of *Trichoderma* spp. is not limited to fighting pathogens; they have also been shown to be opportunistic plant symbionts, enhancing systemic resistance of plants (Yedidia et al. 1999; Shores et al. 2010), a response which is improved by ceratoplatenin family

proteins (Djonovic et al. 2006; Seidl et al. 2009b). Perception of the signals transmitted by *Trichoderma* in the plant requires the function of a MAPK (Shores et al. 2006), and also in the fungus itself, a MAPK signaling is crucial for full induction of systemic response in the plant (Viterbo et al. 2005). By colonizing plant roots, which is significantly enhanced by swollenin (Brotman et al. 2008) or invading them, they are also carried through soil and occupy new niches. This interaction with plants as well as their rhizosphere competence leads to enhanced root proliferation, better growth, and protection of the plants against toxic chemicals, against which *Trichoderma* spp. themselves show a remarkable resistance. Hence, these fungi are promising agents that can be applied for remediation of polluted soil and water by treatment of appropriate plants with spores (Harman et al. 2004).

Secondary metabolites

In order to survive and compete in their ecological niche, fungi apply not only enzymatic weapons but also have a potent arsenal for chemical warfare at their disposal (Vinale et al. 2008). Thereby, not only potential antibiotics (for example the peptaibols) but also mycotoxins and more than 100 metabolites with antibiotic activity including polyketides, pyrones, terpenes, metabolites derived from amino acids, and polypeptides (Sivasithamparam and Ghisalberti 1998) were detected in *Trichoderma* spp. and have been suggested to be used for chemotaxonomy of these species. However, the evolution of peptaibol formation seems to be too complex to allow for prediction of peptaibol production profiles from phylogenetic relationships (Daniel and Filho 2007; Kubicek et al. 2007; Neuhof et al. 2007; Degenkolb et al. 2008). One of the first characterized secondary metabolites of *Trichoderma* spp. was the peptide antibiotic paracelsin (Bruckner and Graf 1983; Bruckner et al. 1984). A wide variety of peptaibols was identified in *Trichoderma* thereafter (Degenkolb et al. 2003, 2007, 2008; Stoppacher et al. 2008). Interestingly, the four trichothecene mycotoxin-producing species (*Trichoderma brevicompactum*, *Trichoderma arundinaceum*, *Trichoderma turrialbense*, and *Trichoderma protrudens*) are not closely related to those species used in biocontrol, which not only means that the application of biocontrol in agriculture does not pose a risk in this respect but also indicates that these mycotoxins do not play a major role in the defense mechanisms of these fungi (Nielsen et al. 2005; Degenkolb et al. 2008). As many other fungi, also *Trichoderma* spp. have been shown to produce a broad array of volatile organic compounds, which recently have received closer attention (Stoppacher et al. 2010). With respect to regulation of peptaibol biosynthesis in *Trichoderma* spp., several

factors are known to be relevant. Environmental cues such as light, pH, nutrients, starvation, or mechanical injury impact this process. Efficient production of peptaibols predominantly occurs in solid cultivation and correlates with conidiation (Kubicek et al. 2007; Tisch and Schmoll 2010). Signaling molecules involved range from the blue light photoreceptors BLR1 and BLR2 to the G-alpha subunits GNA3 (TGA3) and GNA1 as well as protein kinase A (Reithner et al. 2005; Komon-Zelazowska et al. 2007a). Thereby, GNA3 is essential for peptaibol formation, but on the other hand, stimulation of peptaibol formation in the absence of BLR1 and BLR2 is still possible (Komon-Zelazowska et al. 2007a).

***Trichoderma* spp. as industrial workhorses**

Shortly after the discovery of *T. viride* QM6a by the US army during World War II (Reese 1976), the outstanding efficiency of its cellulases led to extensive research toward industrial applications of these enzymes. Later on, this species was renamed *T. reesei* in honor of Elwyn T. Reese (Simmons 1977) and became the most important cellulase producer worldwide. Until now, this species is the most important one of the genus for industrial purposes.

Cellulases and plant cell wall-degrading enzymes

Rising energy costs and the imminent climate change led to an increased attention to biofuel production (Somerville 2007; Rubin 2008). As a potent cellulase producer, research with *T. reesei* is nowadays particularly focused on improvement of efficiency of the enzyme cocktail produced in order to decrease overall costs of production of bioethanol from cellulosic waste material (Kumar et al. 2008), although applications in the pulp and paper industry (Buchert et al. 1998) and textile industry (Galante et al. 1998a) are also important. After the early mutation programs (El-Gogary et al. 1998) and strain improvement, the protein secretion capacity of industrial strains now reaches 100 g/l, with up to 60% of the major cellulase Cel7a (CBHI) and 20% of Cel6a (CBHII). High levels of cellulase and hemicellulase gene expression can be achieved upon cultivation on cellulose, xylan, or a mixture of plant polymers (Mach and Zeilinger 2003) as well as on lactose (Seiboth et al. 2007), all of which are agricultural or industrial byproducts. The natural inducer of at least a subset of these enzymes is believed (yet not definitely proven) to be sophorose, a transglycosylation product of cellobiose (Sternberg and Mandels 1979; Vaheri et al. 1979). Targeted strategies to further enhance the efficiency of the enzymes secreted include elucidation of regulatory mechanisms both at the promoter level (Mach and Zeilinger

2003; Schmoll and Kubicek 2003) as well as with respect to signal transduction (Schmoll et al. 2010). However, auxiliary components acting on the substrate could also enhance efficiency of its degradation (Saloheimo et al. 2002; Schmoll and Kubicek 2005).

Metabolic engineering in recent years provided intriguing insights into these processes (Kubicek et al. 2009), and exploration of the genome sequence of *T. reesei* revealed that this industrial workhorse possesses the smallest amount of genes within Sordariomycetes encoding the enzymes which made it so popular—plant cell wall-degrading enzymes (Foreman et al. 2003; Martinez et al. 2008). Availability of the genome sequence also spurred genome wide analysis of early mutant strains and identification of putatively beneficial mutations, which caused their high efficiency (Le Crom et al. 2009). Interestingly, it seems that even early mutants such as RutC-30 bear considerable alterations of their genome (Seidl et al. 2008). These novel tools also facilitated characterization of the enzyme cocktails secreted by these strains (Herpoel-Gimbert et al. 2008). In addition to these efforts enzyme engineering approaches (Bansal et al. 2009), improvement of the secretion machinery (Conesa et al. 2001; Kruszewska et al. 2008) as well as screening of the enormous variety of plant cell wall-degrading enzymes from nature isolates (Kubicek et al. 1996) or other organisms secreting cellulases (Dashtban et al. 2009) and directed evolution (Nakazawa et al. 2009) complement the optimization of the regulatory mechanism of available production strains. Hence, with the aid of *Trichoderma*, economically reasonable production of second generation biofuels from waste products is on the way.

Heterologous protein production

Filamentous fungi are versatile cell factories and frequently used for heterologous protein expression (Adrio and Demain 2003), especially if they have generally regarded as safe status (Nevalainen et al. 2005), as has *T. reesei* (Nevalainen et al. 1994). The industrial use of *T. reesei* as a producer of heterologous proteins started more than 20 years ago with the production of calf chymosin (Harkki et al. 1989; Uusitalo et al. 1991). Shortly thereafter, even expression of immunologically active antibody fragments (Nyyssonen et al. 1993) in *T. reesei* was achieved and numerous enzymes and performance proteins followed. Nowadays, *T. reesei* is one of the most commonly used filamentous fungi for heterologous protein production (Penttila 1998; Nevalainen et al. 2005).

Based on the efficient expression as well as the considerable knowledge on regulation of cellulase genes, their promoters are routinely used for heterologous protein production (Penttila 1998; Schmoll and Kubicek 2003). Consequently, improvements in cellulase transcription are

also beneficial for these applications. In many cases, the signal peptide of Cel7a (CBHI) is used to facilitate efficient secretion of the product into the culture medium. Nevertheless, also alternative promoters were also shown to be useful for certain applications (Keränen and Penttilä 1995). In general, the high efficiency and the inducibility of the cellulase promoters have proven beneficial in many applications. Using the cellulase promoters, also relatively cheap carbon sources such as cellulose or lactose can be used for production. Nevertheless, it must be considered that the large amount of enzymes secreted into the culture medium can be an issue in specific purification of the heterologous protein, and the complex substrates used could induce extracellular proteases, which are deleterious for the yield of the process (Keränen and Penttilä 1995). For further improvement, promoter modifications, for example with the *chb1*-promotor (Liu et al. 2008), can increase yields of the protein to be expressed.

Food industry

With their long history of safe industrial scale enzyme production, *Trichoderma* spp. have also been extensively applied for production of food additives and related products (Nevalainen et al. 1994; Blumenthal 2004). Currently, various *Trichoderma* enzymes are applied to improve the brewing process (β -glucanases), as macerating enzymes in fruit juice production (pectinases, cellulases, hemicellulases), as feed additive in livestock farming (xylanases) and for pet food. Cellulases are mainly applied in baking, malting, and grain alcohol production (Galante et al. 1998b). However, not only enzymes but also metabolites of *Trichoderma* spp. are used as additives. One of the first products isolated from *T. viride* was a chemical with characteristic coconut-like aroma, a 6-pentyl- α -pyrone with antibiotic properties, the production of which was constantly improved to reach concentrations of more than 7 g/L in extractive fermentation cultures in *T. atroviride* nowadays (Collins and Halim 1972; Oda et al. 2009). An interesting idea is the application of cell wall-degrading enzymes, for example of *T. harzianum*, as food preservatives because of their antifungal effect (Fuglsang et al. 1995), but so far this suggestion has not found broad application. With a similar aim, *T. harzianum* mutanase can be used in toothpaste to prevent accumulation of mutan in dental plaque (Wiater et al. 2005).

Black sheep in the genus *Trichoderma*

In addition to the highly beneficial and frequently used species, the genus *Trichoderma* also comprises opportunist

human pathogens, which show efficient growth at body temperature and mycoparasitic species, which are a significant threat to mushroom farms.

Human pathogenic species

Besides such long-known and well-studied pathogenic fungi as *Candida*, *Aspergillus*, or *Cryptococcus*, also the genus *Trichoderma* comprises opportunistic human pathogens, which pose a serious and often lethal threat—especially to HIV-infected persons and other immunocompromised patients. Belonging to the emerging fungal pathogens, these fungi are often not recognized or diagnosed in a stadium when efficient treatment is problematic (Walsh et al. 2004). *Trichoderma* species have been reported to cause respiratory problems due to volatile organic compounds they produce (Larsen et al. 1998), but more importantly, they can infect immunocompromised patients (*Trichoderma citrinoviride*, *T. harzianum*, and *Trichoderma longibrachiatum* and *Hypocrea orientalis*) after transplantations or suffering from leukemia or HIV (Kredics et al. 2003). The typically poor prognosis of such infections is (besides delayed diagnosis) predominantly due to the low susceptibility of these fungi to commonly used antifungal agents (Chouaki et al. 2002; Kratzer et al. 2006), which often necessitates combined treatment with different drugs (Kratzer et al. 2006; Alanio et al. 2008). Nevertheless, few data on investigation of virulence factors of these fungi are available (Kredics et al. 2004). Among the clinical isolates, *T. longibrachiatum* and *H. orientalis* are the most common ones. Interestingly, no specific phylogenetic characteristics of the clinical isolates as compared to environmental isolates could be found, and no correlation between virulence or pathogenicity and genomic structure was detected (Antal et al. 2005, 2006; Druzhinina et al. 2008). However, most intriguingly, *T. longibrachiatum* not only causes disease; at the same time, it seems to be a source for potential antifungal drugs efficient against *Candida* and *Aspergillus* species (Vicente et al. 2001).

Green mold disease

Cultivation of the edible mushrooms *Agaricus bisporus* (champignon) and *Pleurotus ostreatus* (oyster mushrooms) on mushroom farms all over the world is of considerable economic importance. In the 1980s, a mixture of strains at first identified as *T. harzianum* was found to cause deleterious infections in these farms (Seaby 1987) with losses between 30% and 100% (Seaby 1998). Actually, these strains represented two new species, *Trichoderma aggressivum* fsp. *europaeae* and *T. aggressivum* fsp. *aggressivum* (Samuels et al. 2002b). Since then, this “green mold disease” has spread all over the world (Komon-Zelazowska

et al. 2007b) and was shown to be mainly caused by *Trichoderma pleurotum*, *Trichoderma pleuroticola* in *P. ostreatus* (Park et al. 2006), and *T. aggressivum* in *A. bisporus*. Nevertheless, also *T. harzianum*, *T. longibrachiatum*, *Trichoderma ghanense*, *T. asperellum*, and *T. atroviride* have been detected in *Agaricus* compost and *Pleurotus* substrates (Hatvani et al. 2007), but aggressive colonization of the substrate has not been proven for these species. This threat to commercial mushroom production has recently also led to the development of methods for rapid and specific detection of these fungi in cultivation substrates (Kredics et al. 2009). A similar objective led to the development of a key for identification of *Trichoderma* species commonly associated with commercially grown mushrooms (Muthumeenakshi et al. 1998; Samuels et al. 2002b).

Future prospects

More than ever before sustainable economy and protection of our environment are dominant topics in our everyday life and one alarming report about contaminated landscapes or catastrophes caused by climate change follows another. Today, 87% of energy used in the world comes from nonrenewable sources like natural gas, oil, and coal (Merino and Cherry 2007). Although biofuel production is now being pushed in order to decrease the requirement for fossil fuels, the raw materials therefore originate from commodities and land also needed for food. In this respect, production of the so-called second generation biofuels from agricultural waste products by the aid of cellulases and hemicellulases produced for example by *T. reesei* and fermentation of the resulting oligosaccharides by yeast provides an alternative strategy. However, for an economically competitive process an increase in efficiency of more than 40-fold would be necessary, which is a formidable challenge for research with *Trichoderma*.

Sustainability is also the major driving force for investigation of biocontrol with *Trichoderma*. As opportunistic plant symbionts and effective mycoparasites, numerous species of this genus have the potential to become commercial biofungicides. The challenge in this field of research will be the development of reliable screening techniques, which allow for prediction of the biocontrol efficiency of a given isolate by determination of the key factors for this process. Nevertheless, also the ecological effects of widespread application of a single (or few) fungal species in agriculture remain to be investigated in order to ensure a truly beneficial effect for the environment.

Besides these major applications of *Trichoderma* spp., also the fields of green and white biotechnology become increasingly important for environmentally safe production

of enzymes and antibiotics. These industrial applications will also benefit from studies on molecular physiology and regulatory processes, which continuously reveal novel and valuable metabolites and enzymes as well as components to be modified or adjusted for cost effective high yield production.

Last but not least, the extensive studies on diverse physiological traits available and still progressing for *Trichoderma* make these fungi versatile model organisms for research on both industrial fermentations as well as natural phenomena.

Acknowledgments This work was supported by a grant from the Austrian Science Fund FWF (P20004) to MS. We want to thank C. P. Kubicek for critically reading the manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Adrio JL, Demain AL (2003) Fungal biotechnology. *Int Microbiol* 6:191–199
- Alanio A, Brethon B, Feuilhade de Chauvin M, de Kerviler E, Leblanc T, Lacroix C, Baruchel A, Menotti J (2008) Invasive pulmonary infection due to *Trichoderma longibrachiatum* mimicking invasive Aspergillosis in a neutropenic patient successfully treated with voriconazole combined with caspofungin. *Clin Infect Dis* 46:e116–e118
- Antal Z, Kredics L, Pakarinen J, Doczi I, Andersson M, Salkinoja-Salonen M, Manczinger L, Szekeres A, Hatvani L, Vagvolgyi C, Nagy E (2005) Comparative study of potential virulence factors in human pathogenic and saprophytic *Trichoderma longibrachiatum* strains. *Acta Microbiol Immunol Hung* 52:341–350
- Antal Z, Varga J, Kredics L, Szekeres A, Hatvani L, Manczinger L, Vagvolgyi C, Nagy E (2006) Intraspecific mitochondrial DNA polymorphism within the emerging filamentous fungal pathogen *Trichoderma longibrachiatum*. *J Med Microbiol* 55:31
- Baek JM, Kenerley CM (1998) The *arg2* gene of *Trichoderma virens*: cloning and development of a homologous transformation system. *Fungal Genet Biol* 23:34–44
- Bansal P, Hall M, Realff MJ, Lee JH, Bommarius AS (2009) Modeling cellulase kinetics on lignocellulosic substrates. *Bio-technol Adv* 27:833–848
- Benitez T, Rincon AM, Limon MC, Codon AC (2004) Biocontrol mechanisms of *Trichoderma* strains. *Int Microbiol* 7:249–260
- Blumenthal CZ (2004) Production of toxic metabolites in *Aspergillus niger*, *Aspergillus oryzae*, and *Trichoderma reesei*: justification of mycotoxin testing in food grade enzyme preparations derived from the three fungi. *Regul Toxicol Pharmacol* 39:214–228
- Bochner BR, Gadzinski P, Panomitros E (2001) Phenotype microarrays for high-throughput phenotypic testing and assay of gene function. *Genome Res* 11:1246–1255
- Brody H, Suchindra M (2009) RNAi-mediated gene silencing of highly expressed genes in the industrial fungi *Trichoderma reesei* and *Aspergillus niger*. *Ind Biotechnol* 5:53–60
- Brotman Y, Briff E, Viterbo A, Chet I (2008) Role of swollenin, an expansin-like protein from *Trichoderma*, in plant root colonization. *Plant Physiol* 147:779–789

- Bruckner H, Graf H (1983) Paracelsin, a peptide antibiotic containing alpha-aminoisobutyric acid, isolated from *Trichoderma reesei* Simmons. Part A *Experientia* 39:528–530
- Bruckner H, Graf H, Bokel M (1984) Paracelsin; characterization by NMR spectroscopy and circular dichroism, and hemolytic properties of a peptaibol antibiotic from the cellulolytically active mold *Trichoderma reesei*. Part B *Experientia* 40:1189–1197
- Brunner K, Omann M, Pucher ME, Delic M, Lehner SM, Domnanich P, Kratochwill K, Druzhinina I, Denk D, Zeilinger S (2008) *Trichoderma* G protein-coupled receptors: functional characterisation of a cAMP receptor-like protein from *Trichoderma atroviride*. *Curr Genet* 54:283–299
- Buchert J, Oksanen T, Pere J, Siika-Aho M, Suurnäkki A, Viikari L (1998) Applications of *Trichoderma reesei* enzymes in the pulp and paper industry. In: Harman GE, Kubicek CP (eds) *Trichoderma and Gliocladium*. Vol. 2. Taylor and Francis, London, pp 343–363
- Calistru C, McLean M, Berjak P (1997) In vitro studies on the potential for biological control of *Aspergillus flavus* and *Fusarium moniliforme* by *Trichoderma* species. A study of the production of extracellular metabolites by *Trichoderma* species. *Mycopathologia* 137:115–124
- Chaverri P, Samuels GJ (2003) *Hypocrea/Trichoderma* (Ascomycota, Hypocreales, Hypocreaceae): species with green ascospores. *Stud Mycol* 48:1–116
- Chen LL, Liu LJ, Shi M, Song XY, Zheng CY, Chen XL, Zhang YZ (2009) Characterization and gene cloning of a novel serine protease with nematocidal activity from *Trichoderma pseudokoningii* SMF2. *FEMS Microbiol Lett* 299:135–142
- Chouaki T, Lavarde V, Lachaud L, Raccurt CP, Hennequin C (2002) Invasive infections due to *Trichoderma* species: report of 2 cases, findings of in vitro susceptibility testing, and review of the literature. *Clin Infect Dis* 35:1360–1367
- Collins RP, Halim AF (1972) Characterization of the major aroma constituent of the fungus *Trichoderma viride*. *J Agric Food Chem* 20:437–438
- Conesa A, Punt PJ, van Lwijk N, van den Hondel CA (2001) The secretion pathway in filamentous fungi: a biotechnological view. *Fungal Genet Biol* 33:155–171
- Dababat AA, Sikora RA, Hauschild R (2006) Use of *Trichoderma harzianum* and *Trichoderma viride* for the biological control of *Meloidogyne incognita* on tomato. *Commun Agric Appl Biol Sci* 71:953–961
- Daniel JF, Filho ER (2007) Peptaibols of *Trichoderma*. *Nat Prod Rep* 24:1128–1141
- Dashtban M, Schraft H, Qin W (2009) Fungal bioconversion of lignocellulosic residues; opportunities & perspectives. *Int J Biol Sci* 5:578–595
- Degenkolb T, Berg A, Gams W, Schlegel B, Grafe U (2003) The occurrence of peptaibols and structurally related peptaibiotics in fungi and their mass spectrometric identification via diagnostic fragment ions. *J Pept Sci* 9:666–678
- Degenkolb T, Kirschbaum J, Bruckner H (2007) New sequences, constituents, and producers of peptaibiotics: an updated review. *Chem Biodivers* 4:1052–1067
- Degenkolb T, von Dohren H, Nielsen KF, Samuels GJ, Bruckner H (2008) Recent advances and future prospects in peptaibiotics, hydrophobin, and mycotoxin research, and their importance for chemotaxonomy of *Trichoderma* and *Hypocrea*. *Chem Biodivers* 5:671–680
- Djonovic S, Pozo MJ, Dangott LJ, Howell CR, Kenerley CM (2006) Sm1, a proteinaceous elicitor secreted by the biocontrol fungus *Trichoderma virens* induces plant defense responses and systemic resistance. *Mol Plant Microbe Interact* 19:838–853
- Druzhinina IS, Kopchinskiy AG, Komon M, Bissett J, Szakacs G, Kubicek CP (2005) An oligonucleotide barcode for species identification in *Trichoderma* and *Hypocrea*. *Fungal Genet Biol* 42:813–828
- Druzhinina IS, Kopchinskiy AG, Kubicek CP (2006a) The first 100 *Trichoderma* species characterized by molecular data. *Mycoscience* 47:55–64
- Druzhinina IS, Schmoll M, Seiboth B, Kubicek CP (2006b) Global carbon utilization profiles of wild-type, mutant, and transformant strains of *Hypocrea jecorina*. *Appl Environ Microbiol* 72:2126–2133
- Druzhinina IS, Komon-Zelazowska M, Kredics L, Hatvani L, Antal Z, Belayneh T, Kubicek CP (2008) Alternative reproductive strategies of *Hypocrea orientalis* and genetically close but clonal *Trichoderma longibrachiatum*, both capable of causing invasive mycoses of humans. *Microbiology* 154:3447–3459
- Druzhinina IS, Komon-Zelazowska M, Atanasova L, Seidl V, Kubicek CP (2010) Evolution and ecophysiology of the industrial producer *Hypocrea jecorina* (anamorph *Trichoderma reesei*) and a new sympatric agamospecies related to it. *PLoS ONE* 5:e9191
- El-Gogary S, Leite A, Crivellaro O, El-Dorry H, Eveleigh DE (1998) *Trichoderma reesei* cellulase—from mutants to induction. In: Kubicek CP, Eveleigh DE, Esterbauer H, Steiner W, Kubicek-Pranz EM (eds) *Trichoderma reesei* cellulases. Royal Society of Chemistry, Cambridge, pp 200–211
- Eziashi EI, Uma NU, Adekunle AA, Airede CE (2006) Effect of metabolites produced by *Trichoderma* species against *Ceratomyces paradoxa* in culture medium. *Afr J Biotechnol* 5:703–706
- Foreman PK, Brown D, Dankmeyer L, Dean R, Diener S, Dunn-Coleman NS, Goedegebuur F, Houfek TD, England GJ, Kelley AS, Meerman HJ, Mitchell T, Mitchinson C, Olivares HA, Teunissen PJ, Yao J, Ward M (2003) Transcriptional regulation of biomass-degrading enzymes in the filamentous fungus *Trichoderma reesei*. *J Biol Chem* 278:31988–31997
- Friedl MA, Schmoll M, Kubicek CP, Druzhinina IS (2008) Photostimulation of *Hypocrea atroviridis* growth occurs due to a crosstalk of carbon metabolism, blue light receptors and response to oxidative stress. *Microbiology* 154:1229–1241
- Fuglsang CC, Johansen C, Christgau S, Adler-Nissen J (1995) Antimicrobial enzymes: applications and future potential in the food industry. *Trends Food Sci Technol* 6:390–396
- Galante YM, De Conti A, Monteverdi R (1998a) Application of *Trichoderma* enzymes in the textile industry. In: Harman GE, Kubicek CP (eds) *Trichoderma and Gliocladium*. Taylor and Francis, London, pp 311–326
- Galante YM, De Conti A, Monteverdi R (1998b) Application of *Trichoderma* enzymes in the food and feed industries. In: Harman GE, Kubicek CP (eds) *Trichoderma and Gliocladium*. Taylor and Francis, London, pp 327–342
- Gams W, Bissett J (1998) Morphology and identification of *Trichoderma*. In: Harman GE, Kubicek CP (eds) *Trichoderma and Gliocladium*. Taylor and Francis, London, pp 3–34
- Goswami J, Pandey RK, Tewari JP, Goswami BK (2008) Management of root knot nematode on tomato through application of fungal antagonists, *Acremonium strictum* and *Trichoderma harzianum*. *J Environ Sci Health B* 43:237–240
- Grinyer J, Hunt S, McKay M, Herbert BR, Nevalainen H (2005) Proteomic response of the biological control fungus *Trichoderma atroviride* to growth on the cell walls of *Rhizoctonia solani*. *Curr Genet* 47:381–388
- Gruber F, Visser J, Kubicek CP, de Graaff LH (1990) The development of a heterologous transformation system for the cellulolytic fungus *Trichoderma reesei* based on a *pyrG*-negative mutant strain. *Curr Genet* 18:71–76
- Guangtao Z, Hartl L, Schuster A, Polak S, Schmoll M, Wang T, Seidl V, Seiboth B (2009) Gene targeting in a nonhomologous end joining deficient *Hypocrea jecorina*. *J Biotechnol* 139:146–151

- Guangtao Z, Seiboth B, Wen C, Yaohua Z, Xian L, Wang T (2010) A novel carbon source-dependent genetic transformation system for the versatile cell factory *Hypocrea jecorina* (anamorph *Trichoderma reesei*). FEMS Microbiol Lett. doi:10.1111/j.1574-6968.2009.01851.x
- Harkki A, Uusitalo J, Bailey M, Penttilä M, Knowles JKC (1989) A novel fungal expression system: secretion of active calf chymosin from the filamentous fungus *Trichoderma reesei*. Nat Biotechnol 7:596–603
- Harman GE (2000) Myths and dogmas of biocontrol. Changes in perceptions derived from research on *Trichoderma harzianum* T22. Plant Dis 84:377–393
- Harman GE (2006) Overview of mechanisms and uses of *Trichoderma* spp. Phytopathology 96:190–194
- Harman GE, Lorito M, Lynch JM (2004) Uses of *Trichoderma* spp. to alleviate or remediate soil and water pollution. Adv Appl Microbiol 56:313–330
- Hartl L, Seiboth B (2005) Sequential gene deletions in *Hypocrea jecorina* using a single blaster cassette. Curr Genet 48:204–211
- Hatvani L, Antal Z, Manczinger L, Szekeres A, Druzhinina IS, Kubicek CP, Nagy A, Nagy E, Vagvolgyi C, Kredics L (2007) Green mold diseases of *Agaricus* and *Pleurotus* spp. are caused by related but phylogenetically different *Trichoderma* species. Phytopathology 97:532–537
- Herpoel-Gimbert I, Margeot A, Dolla A, Jan G, Molle D, Lignon S, Mathis H, Sigoillot JC, Monot F, Asther M (2008) Comparative secretome analyses of two *Trichoderma reesei* RUT-C30 and CL847 hypersecretory strains. Biotechnol Biofuels 1:18
- Herrera-Estrella A, Chet I (2004) The biological control agent *Trichoderma*—from fundamentals to applications. In: Arora DK (ed) Fungal biotechnology in agricultural, food and environmental applications. Marcel Dekker, New York, pp 147–156
- Jaklitsch WM (2009) European species of *Hypocrea*. Part I. The green-spored species. Stud Mycol 63:1–91
- Keränen S, Penttilä M (1995) Production of recombinant proteins in the filamentous fungus *Trichoderma reesei*. Curr Opin Biotechnol 6:534–537
- Komon-Zelazowska M, Neuhofer T, Dieckmann R, von Dohren H, Herrera-Estrella A, Kubicek CP, Druzhinina IS (2007a) Formation of atroviridin by *Hypocrea atroviridis* is conidiation associated and positively regulated by blue light and the G protein GNA3. Eukaryot Cell 6:2332–2342
- Komon-Zelazowska M, Bissett J, Zafari D, Hatvani L, Manczinger L, Woo S, Lorito M, Kredics L, Kubicek CP, Druzhinina IS (2007b) Genetically closely related but phenotypically divergent *Trichoderma* species cause green mold disease in oyster mushroom farms worldwide. Appl Environ Microbiol 73:7415–7426
- Kopchinskiy A, Komon M, Kubicek CP, Druzhinina IS (2005) TrichoBLAST: a multilocus database for *Trichoderma* and *Hypocrea* identifications. Mycol Res 109:658–660
- Kratzer C, Tobudic S, Schmoll M, Graninger W, Georgopoulos A (2006) In vitro activity and synergism of amphotericin B, azoles and cationic antimicrobials against the emerging pathogen *Trichoderma* spp. J Antimicrob Chemother 58:1058–1061
- Kredics L, Antal Z, Doczi I, Manczinger L, Kevei F, Nagy E (2003) Clinical importance of the genus *Trichoderma*. A review. Acta Microbiol Immunol Hung 50:105–117
- Kredics L, Antal Z, Szekeres A, Manczinger L, Doczi I, Kevei F, Nagy E (2004) Production of extracellular proteases by human pathogenic *Trichoderma longibrachiatum* strains. Acta Microbiol Immunol Hung 51:283–295
- Kredics L, Antal Z, Szekeres A, Hatvani L, Manczinger L, Vagvolgyi C, Nagy E (2005) Extracellular proteases of *Trichoderma* species. A review. Acta Microbiol Immunol Hung 52:169–184
- Kredics L, Kocsube S, Nagy L, Komon-Zelazowska M, Manczinger L, Sajben E, Nagy A, Vagvolgyi C, Kubicek CP, Druzhinina IS, Hatvani L (2009) Molecular identification of *Trichoderma* species associated with *Pleurotus ostreatus* and natural substrates of the oyster mushroom. FEMS Microbiol Lett 300:58–67
- Kruszewska JS, Perlinska-Lenart U, Gorka-Niec W, Orłowski J, Zembek P, Palamarczyk G (2008) Alterations in protein secretion caused by metabolic engineering of glycosylation pathways in fungi. Acta Biochim Pol 55:447–456
- Kubicek CP, Bolzlbauer UM, Kovacs W, Mach RL, Kuhls K, Lieckfeldt E, Borner T, Samuels GJ (1996) Cellulase formation by species of *Trichoderma* sect. Longibrachiatum and of *Hypocrea* spp. with anamorphs referable to *Trichoderma* sect. Longibrachiatum. Fungal Genet Biol 20:105–114
- Kubicek CP, Mach RL, Peterbauer CK, Lorito M (2001) *Trichoderma*: from genes to biocontrol. J Plant Pathology 83:11–23
- Kubicek CP, Bissett J, Druzhinina I, Kullnig-Gradinger C, Szakacs G (2003) Genetic and metabolic diversity of *Trichoderma*: a case study on South-East Asian isolates. Fungal Genet Biol 38:310–319
- Kubicek CP, Komon-Zelazowska M, Sandor E, Druzhinina IS (2007) Facts and challenges in the understanding of the biosynthesis of peptaibols by *Trichoderma*. Chem Biodivers 4:1068–1082
- Kubicek CP, Komon-Zelazowska M, Druzhinina IS (2008) Fungal genus *Hypocrea/Trichoderma*: from barcodes to biodiversity. J Zhejiang Univ Sci B 9:753–763
- Kubicek CP, Mikus M, Schuster A, Schmoll M, Seiboth B (2009) Metabolic engineering strategies for the improvement of cellulase production by *Hypocrea jecorina*. Biotechnol Biofuels 2:19
- Kuhls K, Lieckfeldt E, Samuels GJ, Kovacs W, Meyer W, Petrin O, Gams W, Borner T, Kubicek CP (1996) Molecular evidence that the asexual industrial fungus *Trichoderma reesei* is a clonal derivative of the ascomycete *Hypocrea jecorina*. Proc Natl Acad Sci U S A 93:7755–7760
- Kullnig CM, Krupica T, Woo SL, Mach RL, Rey M, Benítez T, Lorito M, Kubicek CP (2001) Confusion abounds over identities of *Trichoderma* biocontrol isolates. Mycol Res 105:769–772
- Kumar R, Singh S, Singh OV (2008) Bioconversion of lignocellulosic biomass: biochemical and molecular perspectives. J Ind Microbiol Biotechnol 35:377–391
- Kyalo G, Affokpon A, Coosemans J, Coyne DL (2007) Biological control effects of *Pochonia chlamydsosporia* and *Trichoderma* isolates from Benin (West-Africa) on root-knot nematodes. Commun Agric Appl Biol Sci 72:219–223
- Larsen FO, Clementsen P, Hansen M, Maltbaek N, Ostenfeldt-Larsen T, Nielsen KF, Gravesen S, Skov PS, Nom S (1998) Volatile organic compounds from the indoor mould *Trichoderma viride* cause histamine release from human bronchoalveolar cells. Inflamm Res 47(Suppl 1):S5–S6
- Le Crom S, Schackwitz W, Pennacchio L, Magnuson JK, Culley DE, Collett JR, Martin J, Druzhinina IS, Mathis H, Monot F, Seiboth B, Cherry B, Rey M, Berka R, Kubicek CP, Baker SE, Margeot A (2009) Tracking the roots of cellulase hyperproduction by the fungus *Trichoderma reesei* using massively parallel DNA sequencing. Proc Natl Acad Sci U S A 106:16151–16156
- Liu T, Wang T, Li X, Liu X (2008) Improved heterologous gene expression in *Trichoderma reesei* by cellobiohydrolase I gene (*cbh1*) promoter optimization. Acta Biochim Biophys Sin (Shanghai) 40:158–165
- Lorito M, Hayes CK, Di Pietro A, Harman GE (1993) Biolistic transformation of *Trichoderma harzianum* and *Gliocladium virens* using plasmid and genomic DNA. Curr Genet 24:349–356
- Lu Z, Tombolini R, Woo S, Zeilinger S, Lorito M, Jansson JK (2004) In vivo study of *Trichoderma*–pathogen–plant interactions, using constitutive and inducible green fluorescent protein reporter systems. Appl Environ Microbiol 70:3073–3081
- Mach RL, Zeilinger S (2003) Regulation of gene expression in industrial fungi: *Trichoderma*. Appl Microbiol Biotechnol 60:515–522

- Mach RL, Schindler M, Kubicek CP (1994) Transformation of *Trichoderma reesei* based on hygromycin B resistance using homologous expression signals. *Curr Genet* 25:567–570
- Marra R, Ambrosino P, Carbone V, Vinale F, Woo SL, Ruocco M, Ciliento R, Lanzuise S, Ferraioli S, Soriente I, Gigante S, Turra D, Fogliano V, Scala F, Lorito M (2006) Study of the three-way interaction between *Trichoderma atroviride*, plant and fungal pathogens by using a proteomic approach. *Curr Genet* 50:307–321
- Martinez D, Berka RM, Henrissat B, Saloheimo M, Arvas M, Baker SE, Chapman J, Chertkov O, Coutinho PM, Cullen D, Danchin EG, Grigoriev IV, Harris P, Jackson M, Kubicek CP, Han CS, Ho I, Larrondo LF, de Leon AL, Magnuson JK, Merino S, Misra M, Nelson B, Putnam N, Robbertse B, Salamov AA, Schmoll M, Terry A, Thayer N, Westerholm-Parvinen A, Schoch CL, Yao J, Barabote R, Nelson MA, Detter C, Bruce D, Kuske CR, Xie G, Richardson P, Rokhsar DS, Lucas SM, Rubin EM, Dunn-Coleman N, Ward M, Brettin TS (2008) Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma reesei* (syn. *Hypocrea jecorina*). *Nat Biotechnol* 26:553–560
- Mendoza-Mendoza A, Pozo MJ, Grzegorski D, Martinez P, Garcia JM, Olmedo-Monfil V, Cortes C, Kenerley C, Herrera-Estrella A (2003) Enhanced biocontrol activity of *Trichoderma* through inactivation of a mitogen-activated protein kinase. *Proc Natl Acad Sci U S A* 100:15965–15970
- Mendoza-Mendoza A, Rosales-Saavedra T, Cortes C, Castellanos-Juarez V, Martinez P, Herrera-Estrella A (2007) The MAP kinase TVK1 regulates conidiation, hydrophobicity and the expression of genes encoding cell wall proteins in the fungus *Trichoderma virens*. *Microbiology* 153:2137–2147
- Merino ST, Cherry J (2007) Progress and challenges in enzyme development for biomass utilization. *Adv Biochem Eng Biotechnol* 108:95–120
- Monte E (2001) Understanding *Trichoderma*: between biotechnology and microbial ecology. *Int Microbiol* 4:1–4
- Moreno-Mateos MA, Delgado-Jarana J, Codon AC, Benitez T (2007) pH and Pac1 control development and antifungal activity in *Trichoderma harzianum*. *Fungal Genet Biol* 44:1355–1367
- Mukherjee PK, Kenerley CM (2010) Regulation of morphogenesis and biocontrol properties in *Trichoderma virens* by a VELVET protein, Vell1. *Appl Environ Microbiol* 76:2345–2352
- Mukherjee PK, Raghu K (1997) Effect of temperature on antagonistic and biocontrol potential of *Trichoderma* sp. on *Sclerotium rolfsii*. *Mycopathologia* 139:151–155
- Mukherjee PK, Latha J, Hadar R, Horwitz BA (2003) TmkA, a mitogen-activated protein kinase of *Trichoderma virens*, is involved in biocontrol properties and repression of conidiation in the dark. *Eukaryot Cell* 2:446–455
- Mukherjee PK, Latha J, Hadar R, Horwitz BA (2004) Role of two G-protein alpha subunits, TgaA and TgaB, in the antagonism of plant pathogens by *Trichoderma virens*. *Appl Environ Microbiol* 70:542–549
- Mukherjee M, Mukherjee PK, Kale SP (2007) cAMP signalling is involved in growth, germination, mycoparasitism and secondary metabolism in *Trichoderma virens*. *Microbiology* 153:1734–1742
- Muthumeenakshi S, Brown AE, Mills PR (1998) Genetic comparison of the aggressive weed mould strains of *Trichoderma harzianum* from mushroom compost in North America and the British Isles. *Mycol Res* 102:385–390
- Nagy V, Seidl V, Szakacs G, Komon-Zelazowska M, Kubicek CP, Druzhinina IS (2007) Application of DNA bar codes for screening of industrially important fungi: the haplotype of *Trichoderma harzianum* sensu stricto indicates superior chitinase formation. *Appl Environ Microbiol* 73:7048–7058
- Nakazawa H, Okada K, Onodera T, Ogasawara W, Okada H, Morikawa Y (2009) Directed evolution of endoglucanase III (Cel12A) from *Trichoderma reesei*. *Appl Microbiol Biotechnol* 83:649–657
- Navazio L, Baldan B, Moscatiello R, Zuppini A, Woo SL, Mariani P, Lorito M (2007) Calcium-mediated perception and defense responses activated in plant cells by metabolite mixtures secreted by the biocontrol fungus *Trichoderma atroviride*. *BMC Plant Biol* 7:41
- Neuhof T, Dieckmann R, Druzhinina IS, Kubicek CP, von Dohren H (2007) Intact-cell MALDI-TOF mass spectrometry analysis of peptaibol formation by the genus *Trichoderma/Hypocrea*: can molecular phylogeny of species predict peptaibol structures? *Microbiology* 153:3417–3437
- Nevalainen H, Suominen P, Taimisto K (1994) On the safety of *Trichoderma reesei*. *J Biotechnol* 37:193–200
- Nevalainen KM, Te'o VS, Bergquist PL (2005) Heterologous protein expression in filamentous fungi. *Trends Biotechnol* 23:468–474
- Nielsen KF, Grafenhan T, Zafari D, Thrane U (2005) Trichothecene production by *Trichoderma brevicompactum*. *J Agric Food Chem* 53:8190–8196
- Nyssonen E, Penttila M, Harkki A, Saloheimo A, Knowles JK, Keranen S (1993) Efficient production of antibody fragments by the filamentous fungus *Trichoderma reesei*. *Biotechnology (N Y)* 11:591–595
- Oda S, Isshiki K, Ohashi S (2009) Production of 6-pentyl-[alpha]-pyrone with *Trichoderma atroviride* and its mutant in a novel extractive liquid-surface immobilization (Ext-LSI) system. *Process Biochem* 44:625–630
- Park MS, Bae KS, Yu SH (2006) Two new species of *Trichoderma* associated with green mold of oyster mushroom cultivation in Korea. *Mycobiology* 34:11–113
- Penttila M (1998) Heterologous protein production in *Trichoderma*. In: Harman GE, Kubicek CP (eds) *Trichoderma and Gliocladium*. Taylor and Francis, London, pp 365–382
- Penttila M, Nevalainen H, Ratto M, Salminen E, Knowles J (1987) A versatile transformation system for the cellulolytic filamentous fungus *Trichoderma reesei*. *Gene* 61:155–164
- Persoon CH (1794) *Disposita methodica fungorum*. Römer's Neues Mag Bot 1:81–128
- Peterbauer CK, Heidenreich E, Baker RT, Kubicek CP (1992) Effect of benomyl and benomyl resistance on cellulase formation by *Trichoderma reesei* and *Trichoderma harzianum*. *Can J Microbiol* 38:1292–1297
- Reese ET (1976) History of the cellulase program at the U. S. Army Natick Development Center. *Biotechnol Bioeng Symp* 6:9–20
- Reino JL, Guerrero RF, Hernandez-Galan R, Collado IG (2008) Secondary metabolites from species of the biocontrol agent *Trichoderma*. *Phytochemistry* 7:89–123
- Reithner B, Brunner K, Schuhmacher R, Peissl I, Seidl V, Krska R, Zeilinger S (2005) The G protein alpha subunit Tgal of *Trichoderma atroviride* is involved in chitinase formation and differential production of antifungal metabolites. *Fungal Genet Biol* 42:749–760
- Reithner B, Schuhmacher R, Stoppacher N, Pucher M, Brunner K, Zeilinger S (2007) Signaling via the *Trichoderma atroviride* mitogen-activated protein kinase Tmk I differentially affects mycoparasitism and plant protection. *Fungal Genet Biol* 44:1123–1133
- Rifai MA (1969) A revision of the genus *Trichoderma*. *Mycol Pap* 116:1–56
- Rocha-Ramirez V, Omero C, Chet I, Horwitz BA, Herrera-Estrella A (2002) *Trichoderma atroviride* G-protein alpha-subunit gene *tgai* is involved in mycoparasitic coiling and conidiation. *Eukaryot Cell* 1:594–605
- Rubin EM (2008) Genomics of cellulosic biofuels. *Nature* 454:841–845

- Ruocco M, Lanzuise S, Vinale F, Marra R, Turra D, Woo SL, Lorito M (2009) Identification of a new biocontrol gene in *Trichoderma atroviride*: the role of an ABC transporter membrane pump in the interaction with different plant-pathogenic fungi. *Mol Plant Microbe Interact* 22:291–301
- Sallenave C, Pouchus YF, Bardouil M, Lassus P, Roquebert MF, Verbist JF (1999) Bioaccumulation of mycotoxins by shellfish: contamination of mussels by metabolites of a *Trichoderma koningii* strain isolated in the marine environment. *Toxicon* 37:77–83
- Sallenave-Namont C, Pouchus YF, Robiou du Pont T, Lassus P, Verbist JF (2000) Toxicogenic saprophytic fungi in marine shellfish farming areas. *Mycopathologia* 149:21–25
- Saloheimo M, Paloheimo M, Hakola S, Pere J, Swanson B, Nyssonen E, Bhatia A, Ward M, Penttila M (2002) Swollenin, a *Trichoderma reesei* protein with sequence similarity to the plant expansins, exhibits disruption activity on cellulosic materials. *Eur J Biochem* 269:4202–4211
- Samolski I, de Luis A, Vizcaino JA, Monte E, Suarez MB (2009) Gene expression analysis of the biocontrol fungus *Trichoderma harzianum* in the presence of tomato plants, chitin, or glucose using a high-density oligonucleotide microarray. *BMC Microbiol* 9:217
- Samuels GJ (2006) *Trichoderma*: systematics, the sexual state, and ecology. *Phytopathology* 96:195–206
- Samuels GJ, Dodd SL, Gams W, Castlebury LA, Petrini O (2002a) *Trichoderma* species associated with the green mold epidemic of commercially grown *Agaricus bisporus*. *Mycologia* 94:146
- Samuels GJ, Chaverri P, Farr DF, McCray EB (2002a) *Trichoderma* Online. Systematic Mycology and Microbiology Laboratory, ARS, USDA; <http://nt.ars-grin.gov/taxadescriptions/keys/TrichodermaIndex.cfm>. Accessed 16 Apr 2010
- Scherm B, Schmoll M, Balmas V, Kubicek CP, Migheli Q (2009) Identification of potential marker genes for *Trichoderma harzianum* strains with high antagonistic potential against *Rhizoctonia solani* by a rapid subtraction hybridization approach. *Curr Genet* 55:81–91
- Schmoll M (2008) The information highways of a biotechnological workhorse—signal transduction in *Hypocrea jecorina*. *BMC Genomics* 9:430
- Schmoll M, Kubicek CP (2003) Regulation of *Trichoderma* cellulase formation: lessons in molecular biology from an industrial fungus. A review. *Acta Microbiol Immunol Hung* 50:125–145
- Schmoll M, Kubicek CP (2005) *ooC1*, a unique gene expressed only during growth of *Hypocrea jecorina* (anamorph: *Trichoderma reesei*) on cellulose. *Curr Genet* 48:126–133
- Schmoll M, Franchi L, Kubicek CP (2005) Envoy, a PAS/LOV domain protein of *Hypocrea jecorina* (anamorph *Trichoderma reesei*), modulates cellulase gene transcription in response to light. *Eukaryot Cell* 4:1998–2007
- Schmoll M, Schuster A, Silva Rdo N, Kubicek CP (2009) The G-alpha protein GNA3 of *Hypocrea jecorina* (anamorph *Trichoderma reesei*) regulates cellulase gene expression in the presence of light. *Eukaryot Cell* 8:410–420
- Schmoll M, Esquivel-Naranjo UE, Herrera-Estrella A (2010) *Trichoderma* in the light of day—physiology and development. *Fungal Genet Biol* (in press)
- Schuster A, Kubicek CP, Friedl MA, Druzhinina IS, Schmoll M (2007) Impact of light on *Hypocrea jecorina* and the multiple cellular roles of ENVOY in this process. *BMC Genomics* 8:449
- Seaby D (1987) Infection of mushroom compost by *Trichoderma* species. *Mushroom J* 179:355–361
- Seaby D (1998) *Trichoderma* as a weed mould or pathogen in mushroom cultivation. In: Harman GE, Kubicek CP (eds) *Trichoderma & Gliocladium*. Taylor & Francis, London, pp 267–288
- Seiboth B, Gamauf C, Pail M, Hartl L, Kubicek CP (2007) The D-xylose reductase of *Hypocrea jecorina* is the major aldose reductase in pentose and D-galactose catabolism and necessary for beta-galactosidase and cellulase induction by lactose. *Mol Microbiol* 66:890–900
- Seidl V, Schmoll M, Scherm B, Balmas V, Seiboth B, Migheli Q, Kubicek CP (2006) Antagonism of *Pythium* blight of zucchini by *Hypocrea jecorina* does not require cellulase gene expression but is improved by carbon catabolite derepression. *FEMS Microbiol Lett* 257:145–151
- Seidl V, Gamauf C, Druzhinina IS, Seiboth B, Hartl L, Kubicek CP (2008) The *Hypocrea jecorina* (*Trichoderma reesei*) hypercellulolytic mutant RUT C30 lacks a 85 kb (29 gene-encoding) region of the wild-type genome. *BMC Genomics* 9:327
- Seidl V, Seibel C, Kubicek CP, Schmoll M (2009a) Sexual development in the industrial workhorse *Trichoderma reesei*. *Proc Natl Acad Sci U S A* 106:13909–13914
- Seidl V, Song L, Lindquist E, Gruber S, Koptchinsky A, Zeilinger S, Schmoll M, Martinez P, Sun J, Grigoriev I, Herrera-Estrella A, Baker SE, Kubicek CP (2009b) Transcriptomic response of the mycoparasitic fungus *Trichoderma atroviride* to the presence of a fungal prey. *BMC Genomics* 10:567
- Shoresh M, Gal-On A, Leibman D, Chet I (2006) Characterization of a mitogen-activated protein kinase gene from cucumber required for *Trichoderma*-conferred plant resistance. *Plant Physiol* 142:1169–1179
- Shoresh M, Harman GE, Mastouri F (2010) Induced systemic resistance and plant responses to fungal biocontrol agents. *Annu Rev Phytopathol* (in press)
- Silva R, Steindorff AS, Ulhoa CJ, Felix RC (2009) Involvement of G-alpha protein GNA3 in production of cell wall-degrading enzymes by *Trichoderma reesei* (*Hypocrea jecorina*) during mycoparasitism against *Pythium ultimum*. *Biotechnol Lett* 31:531–536
- Simmons EG (1977) Classification of some cellulase-producing *Trichoderma* species. In: Bigelow HE, Simmons EG (eds) 2nd International Mycological Congress. University of South Florida, Tampa, p 618
- Sivasithamparam K, Ghisalberti EL (1998) Secondary metabolism in *Trichoderma* and *Gliocladium*. In: Harman GE, Kubicek CP (eds) *Trichoderma and Gliocladium*. Taylor and Francis, London, pp 139–192
- Somerville C (2007) Biofuels. *Curr Biol* 17:R115–R119
- Spiegel Y, Chet I (1998) Evaluation of *Trichoderma* spp. as a biocontrol agent against soilborne fungi and plant-parasitic nematodes in Israel. *Integr Pest Manage Rev* 3:169–175
- Sreerama L, Veerabhadrapa PS (1993) Isolation and properties of carboxylesterases of the termite gut-associated fungus, *Xylaria nigripes*. K., and their identity from the host termite, *Odentotermes horni*. W., mid-gut carboxylesterases. *Int J Biochem* 25:1637–1651
- Sternberg D, Mandels GR (1979) Induction of cellulolytic enzymes in *Trichoderma reesei* by sophorose. *J Bacteriol* 139:761
- Stoppacher N, Zeilinger S, Omann M, Lassahn PG, Roitinger A, Krska R, Schuhmacher R (2008) Characterisation of the peptaibome of the biocontrol fungus *Trichoderma atroviride* by liquid chromatography/tandem mass spectrometry. *Rapid Commun Mass Spectrom* 22:1889–1898
- Stoppacher N, Kluger B, Zeilinger S, Krska R, Schuhmacher R (2010) Identification and profiling of volatile metabolites of the biocontrol fungus *Trichoderma atroviride* by HS-SPME-GC-MS. *J Microbiol Methods* 81(2):187–193
- Suarez MB, Vizcaino JA, Llobell A, Monte E (2007) Characterization of genes encoding novel peptidases in the biocontrol fungus *Trichoderma harzianum* CECT 2413 using the TrichoEST functional genomics approach. *Curr Genet* 51:331–342
- Tisch D, Schmoll M (2010) Light regulation of metabolic pathways in fungi. *Appl Microbiol Biotechnol* 85:1259–1277

- Tseng SC, Liu SY, Yang HH, Lo CT, Peng KC (2008) Proteomic study of biocontrol mechanisms of *Trichoderma harzianum* ETS 323 in response to *Rhizoctonia solani*. *J Agric Food Chem* 56:6914–6922
- Tulasne LR, Tulasne C (1865) *Selecta fungorum carpologia*. Jussu, Paris
- Uusitalo JM, Nevalainen KM, Harkki AM, Knowles JK, Penttila ME (1991) Enzyme production by recombinant *Trichoderma reesei* strains. *J Biotechnol* 17:35–49
- Vaheri M, Leisola M, Kauppinen V (1979) Transglycosylation products of cellulase system of *Trichoderma reesei*. *Biotechnol Lett* 1:41–46
- Vicente MF, Cabello A, Platas G, Basilio A, Diez MT, Dreikorn S, Giacobbe RA, Onishi JC, Meinz M, Kurtz MB, Rosenbach M, Thompson J, Abruzzo G, Flattery A, Kong L, Tsipouras A, Wilson KE, Pelaez F (2001) Antimicrobial activity of ergokonin A from *Trichoderma longibrachiatum*. *J Appl Microbiol* 91:806–813
- Vinale F, Marra R, Scala F, Ghisalberti EL, Lorito M, Sivasithamparam K (2006) Major secondary metabolites produced by two commercial *Trichoderma* strains active against different phytopathogens. *Lett Appl Microbiol* 43:143–148
- Vinale F, Sivasithamparam K, Ghisalberti EL, Marra R, Woo SL, Lorito M (2008) *Trichoderma*–plant–pathogen interactions. *Soil Biol Biochem* 40:1–10
- Vinale F, Ghisalberti EL, Sivasithamparam K, Marra R, Ritieni A, Ferracane R, Woo S, Lorito M (2009) Factors affecting the production of *Trichoderma harzianum* secondary metabolites during the interaction with different plant pathogens. *Lett Appl Microbiol* 48:705–711
- Viterbo A, Ramot O, Chemin L, Chet I (2002) Significance of lytic enzymes from *Trichoderma* spp. in the biocontrol of fungal plant pathogens. *Antonie Leeuwenhoek* 81:549–556
- Viterbo A, Harel M, Horwitz BA, Chet I, Mukherjee PK (2005) *Trichoderma* mitogen-activated protein kinase signaling is involved in induction of plant systemic resistance. *Appl Environ Microbiol* 71:6241–6246
- Walsh TJ, Groll A, Hiemenz J, Fleming R, Roilides E, Anaissie E (2004) Infections due to emerging and uncommon medically important fungal pathogens. *Clin Microbiol Infect* 10(Suppl 1):48–66
- Weindling R (1932) *Trichoderma lignorum* as a parasite of other soil fungi. *Phytopathology* 22:837–845
- Wiater A, Szczodrak J, Pleszczyńska M (2005) Optimization of conditions for the efficient production of mutan in streptococcal cultures and post-culture liquids. *Acta Biol Hung* 56:137–150
- Woo SL, Scala F, Ruocco M, Lorito M (2006) The molecular biology of the interactions between *Trichoderma* spp., phytopathogenic fungi, and plants. *Phytopathology* 96:181–185
- Yedidia II, Benhamou N, Chet II (1999) Induction of defense responses in cucumber plants (*Cucumis sativus* L.) by the biocontrol agent *Trichoderma harzianum*. *Appl Environ Microbiol* 65:1061–1070
- Yoder JA, Glenn BD, Benoit JB, Zettler LW (2008) The giant Madagascar hissing-cockroach (*Gromphadorhina portentosa*) as a source of antagonistic moulds: concerns arising from its use in a public setting. *Mycoses* 51:95–98
- Zeilinger S (2004) Gene disruption in *Trichoderma atroviride* via *Agrobacterium*-mediated transformation. *Curr Genet* 45:54–60
- Zeilinger S, Omann M (2007) *Trichoderma* biocontrol: signal transduction pathways involved in host sensing and mycoparasitism. *Gene Regul Syst Biol* 1:227–234
- Zeilinger S, Reithner B, Scala V, Peissl I, Lorito M, Mach RL (2005) Signal transduction by Tga3, a novel G protein alpha subunit of *Trichoderma atroviride*. *Appl Environ Microbiol* 71:1591–1597