

POSTER PRESENTATION

Open Access

Design and development of novel non-peptide agonists at NPR-C

Filipa Mota Quinteiro^{1*}, Paul Gane¹, Anne-Sophie Rebstock¹, Roberta Worthington¹, Michela Simone¹, Snezana Djordjevic², Adrian Hobbs³, David Selwood¹

From 5th International Conference on cGMP: Generators, Effectors and Therapeutic Implications Halle, Germany. 24-26 June 2011

Background

Endothelium-derived C-type natriuretic peptide (CNP) possesses cytoprotective and anti-atherogenic functions that regulate vascular tone and smooth-muscle relaxation and might be key in protecting against ischaemia-reperfusion injury [1]. The finding that many of the vasoprotective effects of CNP are mediated by the natriuretic peptide receptor type-C (NPR-C) suggests that this receptor might represent a novel therapeutic target for the treatment of cardiovascular diseases. Thus, we have designed and developed small molecule drug-like mimetics of CNP agonists at NPRC.

Methods and results

We employ a multi-disciplinary approach that comprises molecular modelling, chemical synthesis and biological and functional assays. The crystal structure of NPR-C was used as the starting point for the design of peptidic and subsequently non-peptidic ligands to the receptor [2]. We have determined which fragments of CNP are crucial for binding to NPR-C and modified the NPR-C antagonist AP-811 using pharmacophore searches to replace the peptide component, which led to the design of a library of compounds that were subsequently synthesised, tested and optimised [3].

Conclusion

Novel and selective non-peptide NPR-C agonists have been identified that relax rat isolated mesenteric arteries in vitro. We foresee that such molecules will facilitate the development of potential therapeutic agents for cardiovascular diseases.

* Correspondence: filipa.quinteiro.10@wibr.ucl.ac.uk

¹Wolfson Institute for Biomedical Research, University College London, London, WC1E 6BT, UK

Full list of author information is available at the end of the article

Author details

¹Wolfson Institute for Biomedical Research, University College London, London, WC1E 6BT, UK. ²Institute for Structural and Molecular Biology, University College London, London WC1E 6BT, UK. ³Neuroscience, Physiology & Pharmacology, University College London, London, WC1E 6BT, UK.

Published: 1 August 2011

References

1. Ahluwalia A, Hobbs AJ: Endothelium-derived C-type natriuretic peptide: more than just a hyperpolarizing factor. *Trends Pharmacol Sci* 2005, **26**:162-167.
2. He XL, Chow DC, Martick MM, Garcia KC: Allosteric activation of a spring-loaded natriuretic peptide receptor dimer by hormone. *Science* 2001, **293**:1657-1662.
3. Veale CA, Alford VC, Aharony D, Banville DL, Bialecki RA, Brown FJ, Damewood JR, Dantzman CL, Edwards PD, Jacobs RT, Mauger RC, Murphy MM, Palmer WE, Pine KK, Rumsey WL, Garcia-Davenport LE, Shaw A, Steelman GB, Surian JM, Vacek EP: The discovery of non-basic atrial natriuretic peptide clearance receptor antagonists. *Bioorg Med Chem Lett* 2000, **10**:1949-1952, Part 1.

doi:10.1186/1471-2210-11-S1-P54

Cite this article as: Quinteiro et al.: Design and development of novel non-peptide agonists at NPR-C. *BMC Pharmacology* 2011 **11**(Suppl 1):P54.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

BioMed Central