

Research Article

The Forgotten Hemodynamic (PCO₂ Gap) in Severe Sepsis

Zouheir Ibrahim Bitar , Ossama Sajeh Maadarani , AlAsmar Mohammed El-Shably, Ragab Desouky Elshabasy, and Tamer Mohamed Zaalouk

Critical Care Unit, Ahmadi Hospital, Kuwait Oil Company, P.O. Box 46468, Postal Code 64015, Ahmadi, Kuwait

Correspondence should be addressed to Zouheir Ibrahim Bitar; zbitar2@hotmail.com

Received 23 September 2019; Accepted 25 November 2019; Published 7 January 2020

Academic Editor: Samuel A. Tisherman

Copyright © 2020 Zouheir Ibrahim Bitar et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Central venous-arterial carbon dioxide difference (PCO₂ gap) can be a marker of cardiac output adequacy in global metabolic conditions that are less affected by the impairment of oxygen extraction capacity. We investigated the relation between the PCO₂ gap, serum lactate, and cardiac index (CI) and prognostic value on admission in relation to fluid administration in the early phases of resuscitation in sepsis. We also investigated the chest ultrasound pattern A or B. **Method.** We performed a prospective observational study and recruited 28 patients with severe sepsis and septic shock in a mixed ICU. We determined central venous PO₂, PCO₂, PCO₂ gap, lactate, and CI at 0 and 6 hours after critical care unit (CCU) admission. The population was divided into two groups based on the PCO₂ gap (cutoff value 0.8 kPa). **Results.** The CI was significantly lower in the high PCO₂ gap group ($P = 0.001$). The high PCO₂ gap group, on admission, required more administered fluid and vasopressors ($P = 0.01$ and $P = 0.009$, respectively). There was also a significant difference between the two groups for low mean pressure ($P = 0.01$), central venous O₂ ($P = 0.01$), and lactate level ($P = 0.003$). The mean arterial pressure was lower in the high PCO₂ gap group, and the lactate level was higher, indicating global hypoperfusion. The hospital mortality rate for all patients was 24.5% (7/28). The in-hospital mortality rate was 20% (2/12) for the low gap group and 30% (5/16) for the high gap group; the odds ratio was 1.6 (95% CI 0.5–5.5; $P = 0.53$). Patients with a persistent or rising PCO₂ gap larger than 0.8 kPa at $T = 6$ and 12 hours had a higher mortality change ($n = 6$; in-hospital mortality was 21.4%) than patients with a PCO₂ gap of less than 0.8 kPa at $T = 6$ ($n = 1$; in-hospital mortality was 3%); this odds ratio was 5.3 (95% CI 0.9–30.7; $P = 0.08$). The PCO₂ gap had no relation with the chest ultrasound pattern. **Conclusion.** The PCO₂ gap is an important hemodynamic variable in the management of sepsis-induced circulatory failure. The PCO₂ gap can be a marker of the adequacy of the cardiac output status in severe sepsis. A high PCO₂ gap value (>0.8 kPa) can identify situations in which increasing CO can be attempted with fluid resuscitation in severe sepsis. The PCO₂ gap carries an important prognostic value in severe sepsis.

1. Introduction

Severe sepsis remains associated with high mortality, and the early recognition of the signs of tissue hypoperfusion is crucial in its management [1]. The effectiveness of oxygen-derived parameters as resuscitation goals has been questioned, and the latest data have failed to demonstrate clinical advantage [2]. The venous oxygen saturation (SvO₂) is <70% in the majority of patients with severe sepsis on admission to the intensive care unit (ICU) [3]. The venous-to-arterial carbon dioxide difference (Pv-aCO₂) can indicate the adequacy of microvascular blood flow in the early phases of

resuscitation in sepsis [4]. Hence, other resuscitation goals, such as PCO₂ gap, have been suggested, due to their ability to predict adverse clinical outcomes and simplicity in patients achieving normal oxygen-derived parameters during the early phases of resuscitation in septic shock.

The optimization of cardiac output during the early resuscitation of patients with sepsis is an important goal in the management of severe sepsis. The Pv-aCO₂ usually does not go beyond 0.8 kPa (6 mmHg) under physiological conditions [5] and indicates the adequacy of venous blood flow and cardiac output (CO) [6]. At the circulatory level, an inverse relationship between the PCO₂ gap and cardiac

index (CI) has been described in critically ill patients with severe sepsis [7]. Combining ScvO₂ values as a surrogate for global tissue hypoxia and the PCO₂ gap as a substitute for CI may be practical during resuscitation of critically ill patients. However, in patients with sepsis, microcirculatory level distributive changes may be independent of CI [8]. This means that, on a regional level, in agreement with the persistent tissue hypoxia despite normal ScvO₂ levels, the accumulation of carbon dioxide (CO₂) may occur in sepsis despite adequate CI.

We studied the relationship between the Pv-aCO₂ and CI and tackled the question of whether the Pv-aCO₂ can add a value to the outcome prediction.

1.1. Study Design. This work was a prospective, blinded, observational study of an enrolled sample of patients with sepsis who required ICU admission.

1.2. Population and Setting. The study was performed in the Ahmadi Hospital critical care unit (CCU). The hospital is a subsidiary of the Kuwait Oil Company (KOC). The hospital serves KOC members and their families, including their parents, with an annual ED census of 60,000 visits. Ultrasound examinations are currently used in daily practice in the CCU. The protocol was reviewed and approved by an Ahmadi Hospital Ethical Committee.

Patients were included if they were >18 years old and had sepsis according to international guidelines [9] between September 2018 and March 30, 2019. Additionally, only patients for whom there was a clinical indication for additional hemodynamic monitoring using transthoracic echocardiographic cardiac output [10] and ultrasound of the chest [9] were included. Informed consent was obtained from the patient or legal representative. Central lines and arterial lines were inserted into every patient admitted to the CCU. Cardiac output and index were measured by a certified intensivist in critical care ultrasound who was blinded to the blood results. Hemodynamic data were collected at the time of admission to the CCU and then every 6 hours (*T*₀, *T*₁, and *T*₂).

Cardiac output was measured by the assessment of stroke volume (SV) by PW Doppler. The site for measurement was the left ventricular outflow tract (LVOT) [10], where SV is the multiplication of the cross-sectional area (CSA) of the LVOT by the velocity-time integral (VTI) at that same location ($SV = CSA \times VTI$). VTI is the sum of the instantaneous velocities measured at a specific location by PW Doppler and is equal to the area under the curve of the Doppler velocity profile. It can be thought of as the distance the average blood cell travels during systole [10]. To calculate CO, the following steps are required [10]:

- (1) Measure the diameter (*D*) of the LVOT. Obtained from the image of the surface parasternal long-axis view, or the transesophageal route is used when the parasternal view is not clear. Start the measurement from the junction of the atrioventricular leaflet with the septal endocardium to the junction of the other

leaflet with the mitral valve posteriorly, using inner edge to inner edge, in early systole [7]. Use the largest of 3 to 5 repeated measurements to avoid underestimation of the true diameter.

- (2) Calculate the CSA by using the following formula: $CSA = D^2 \times 0.785$.
- (3) Measure the velocity at the LVOT using an apical 3-chamber or 5-chamber view. Put the PW sample volume 5 mm above the aortic valve and record a Doppler waveform with a well-defined envelope.
- (4) Take the average of 3 VTI measurements by tracing 3 consecutive waveforms.
- (5) Calculate the SV using the following formula: $SV = CSA \times VTI$, and then multiply the SV with heart rate to calculate the CO.

Radiometer ABL90 Flex analyzer (Copenhagen, Denmark) is used to analyze blood samples in the ICU, to ensure rapid assessment and avoid poor quality measurements. The severity score Simplified Acute Physiologic Score (SAPS) was calculated.

1.3. Statistics. The admitted patients were divided into two groups: patients with a normal PCO₂ gap (<0.8 kPa) and patients with a high PCO₂ gap (>0.8 kPa) at *T*₀. The two-tailed statistical tests were performed by the Statistical Package for the Social Sciences (IBM SPSS 19). Student's *t*-test was used to assess the differences between both groups in the case of a normal distribution. The chi-square test or Fisher's exact test was used for categorical data. For each time point (*T*₀–*T*₄), the difference between arterial CO₂ partial pressure (paCO₂) and central venous CO₂ partial pressure (pvCO₂) (i.e., the PCO₂ gap) was calculated. Additionally, for each time point, the average CI, mean arterial pressure (MAP), ScvO₂ lactate, and infusion rate of norepinephrine were calculated. The agreement between the CI and the PCO₂ gap was assessed by the mean bias and 95% limits of agreement (mean bias ± 1.969 standard deviation). The odds ratio for mortality between patients with a high PCO₂ gap and patients with a low PCO₂ gap at both *T*₀ and *T*₄ was calculated. Data are displayed as the mean ± SD. Statistical significance was assumed at *P* < 0.05.

2. Results

We enrolled 31 patients. Of these, three patients were excluded due to lack of consent. Baseline characteristics and outcome of the total population and both groups are shown in Table 1.

When the CI was plotted against the PCO₂ gap for all paired measurements (Figure 1), there was an inverse logarithmic relationship with increasing central PCO₂ gap as CI decreased ($R^2 = 0.07$; $P = 0.001$). CI was significantly lower in the population with a high PCO₂ gap ($P = 0.001$).

The high CO₂ gap group, on admission, required more bolus fluid administration and vasopressors ($P = 0.01$ and $P = 0.009$, respectively) (Table 1). There was also a significant difference between the two groups for low mean

TABLE 1: Baseline characteristics.

	Total population (<i>n</i> = 28)	Low gap (<i>n</i> = 12)	High gap (<i>n</i> = 16)	<i>P</i> value
Age (years)	71 ± 14	69 ± 17	73.62 ± 13	0.7
Gender (m/f)		7/5	10/6	
SOFA	10.39 ± 3.9	8.5 ± 3.4	11.75 ± 3.8	0.03*
Diagnosis				
Abdominal	8	2	6	
Respiratory	11	6	5	
UTI	6	3	3	
Endo	3	1	2	
Lab and therapy				
Lactate (mmol/L)	2.7 ± 2.3	1.3 ± 0.9	3.76 ± 2.6	0.003*
CI (L/min/m ²)	2.6 ± 0.41	3.2 ± 0.37	2.1 ± 0.46	0.001*
Ultrasound chest A/B profiles		8/4	8/8	0.8
Mechanical ventilation	8	5	3	0.38
Renal replacement therapy	5	2	3	0.6
Vasopressor	18	5	13	0.009*
Fluid boluses	28	10	18	0.01*
Mean BP	60 ± 10	65 ± 9	55 ± 11	0.01*
SvO ₂ (%)	71.9 ± 10.7	73.2 ± 9.1	70.3 ± 6.6	0.01*
Hematocrit (%)	31 ± 1	30 ± 5	31 ± 6	0.49

*Statistically significant difference. SOFA score: sequential organ failure assessment score; Endo: endocarditis.




FIGURE 1: Correlation between central venous PCO₂ difference and cardiac index in the total population for all paired measurements at *T* = 0. DPCO₂T₀: deltaPCO₂ on admission; CIT₀: cardiac index on admission.

pressure (*P* = 0.01), central venous O₂ (*P* = 0.01), and lactate level (*P* = 0.003). The mean arterial pressure was lower in the high CO₂ gap group, and the lactate level was higher, indicating global hypoperfusion.

The hospital mortality rate for all patients was 24.5% (7/28). The in-hospital mortality rate was 20% (2/12) for the low gap group and 30% (5/16) for the high gap group; the odds ratio was 1.6 (95% CI 0.5–5.5; *P* = 0.53). Patients with a persistent or rising PCO₂ gap larger than 0.8 kPa at *T* = 6 and 12 had a higher mortality change (*n* = 6; in-hospital mortality was 21.4%) compared to patients with a PCO₂ gap less than 0.8 kPa at *T* = 6 (*n* = 1; in-hospital mortality was 3%); this odds ratio was 5.3 (95% CI 0.9–30.7; *P* = 0.08).

The PCO₂ gap had no relation with the chest ultrasound pattern.

3. Discussion

The venous and arterial PCO₂ are dependent on circulatory flow and pulmonary gas exchange, respectively [11]. Thus, a decreased flow will increase the difference between the arterial and venous PCO₂ [12]. The cardiac output is measured from the ratio between alveolar oxygen uptake and the oxygen content difference between arterial-venous blood (cardiac output = VO₂/CaO₂CvO₂) or from the ratio of carbon dioxide production and arteriovenous carbon dioxide content difference (cardiac output = VCO₂/CaCO₂CvCO₂), when applying the Fick principle to cardiac output. By substituting CO₂ pressure for content, the inverse relationship between cardiac output and the PCO₂ gap can be seen [13]. An increase in the venous-arterial PCO₂ difference occurs in states of decreased flow, regardless of the reason for the circulatory failure, and this has been demonstrated in clinical trials to be inversely related to cardiac output [14].

In critically ill patients of a mixed population, Cuschieri et al. [7] found a significant inverse relation between the PCO₂ gap and CI, taking into consideration that one-third of their population were patients with cardiogenic shock. Our findings are in line with physiological theory, which describes an inverse curvilinear relation between cardiac output and PCO₂ gap, according to a modified Fick equation for a range of CO₂ production isopleths [15]. Various studies have described such an inverse relationship between the delta PCO₂ gap and CI in septic circulatory failure [12, 16]. Due to the heterogeneity of microcirculatory blood flow, inadequate washout of CO₂ in microcirculatory weak units, despite normal or even elevated cardiac output, has been observed during sepsis [8]. Vallee et al. [17] tested this hypothesis in patients with septic shock who were supposedly adequately resuscitated at the systemic level, with a ScvO₂ C70% [18]. An increase in the PCO₂ gap may be a

marker of microcirculatory dysfunction in septic shock, especially when SvO₂ is normal [4].

The prognostic value of a high PCO₂ gap is a matter of debate. There is no mortality difference between the two groups on admission. However, higher mortality is noted in the group with a persistently high PCO₂ gap. A persistently elevated PCO₂ gap could indicate poor prognosis and nonresponse to current therapy or could indicate severe microcirculatory dysfunction. However, the study was not powered to test whether closing the PCO₂ gap will improve the prognosis of the high CO₂ gap group.

The chest ultrasound is used for the hemodynamic assessment of shock patients of different etiologies with the need for fluid assessment [19]. This finding depends on the ability of ultrasound to detect interstitial syndrome, which is considered as a direct marker of clinical volume [18]. Our results did not show a significant relation between the chest ultrasound pattern and delta PCO₂, taking into consideration that our population is only those with septic circulatory failure. Most likely, delta PCO₂ reflects microcirculatory dysfunction, and the chest ultrasound pattern reflects intracardiac pressure.

The small number of patients and the fact that this work is a single-center study are both limitations of the study, but we tried to be blinded to avoid bias. Additionally, the population was only those with severe sepsis or septic shock, and the results cannot be applied to other less critically ill patients or those with shock due to other etiologies.

4. Conclusion

The PCO₂ gap is an important hemodynamic variable in the management of sepsis-induced circulatory failure. The PCO₂ gap can be a marker of the adequacy of cardiac output in severe sepsis. A high PCO₂ gap value (>0.8 kPa) can identify situations in which increasing CO can be attempted with fluid resuscitation in severe sepsis. The PCO₂ gap carries an important prognostic value in severe sepsis.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Disclosure

The research was performed as part of the employment of the authors in Kuwait Oil Company.

Conflicts of Interest

The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Authors' Contributions

Zouheir Ibrahim Bitar wrote the article, Ossama Sajeh Maadarani and AlAsmar Mohammed El-Shably were involved in the discussion, and Ragab Desouky Elshabasy and

Tamer Mohamed Zaalouk collected the data and revised the manuscript.

Acknowledgments

The authors are grateful for the suggestions and comments from the manager of Ahmadi Hospital and chief clinical officer and to Dr. Emad Alawd and Dr. Suad Alajmi that have substantially contributed to the final shape of the paper.

References

- [1] J.-L. Vincent and D. De Backer, "Circulatory shock," *The New England Journal of Medicine*, vol. 369, no. 18, pp. 1726–1734, 2013.
- [2] D. M. Yealy, D. M. Yealy, J. A. Kellum et al., "A randomized trial of protocol-based care for early septic shock," *The New England Journal of Medicine*, vol. 370, no. 18, pp. 1683–1693, 2014.
- [3] P. A. van Beest, J. J. Hofstra, M. J. Schultz, E. C. Boerma, P. E. Spronk, and M. A. Kuiper, "The incidence of low venous oxygen saturation on admission to the intensive care unit: a multi-center observational study in the Netherlands," *Critical Care*, vol. 12, no. 2, p. R33, 2008.
- [4] G. A. Ospina-Tascón, M. Umaña, W. F. Bermúdez et al., "Can venous-to-arterial carbon dioxide differences reflect microcirculatory alterations in patients with septic shock?," *Intensive Care Medicine*, vol. 42, no. 2, pp. 211–221, 2016.
- [5] A. Guyton and J. Hall, "Transport of oxygen and carbon dioxide in the blood and body fluids," in *Textbook of Medical Physiology*, A. Guyton and J. Hall, Eds., pp. 463–472, WB Saunders Co., Philadelphia, PA, USA, 2000.
- [6] B. A. Johnson and M. H. Weil, "Redefining ischemia due to circulatory failure as dual defects of oxygen deficits and of carbon dioxide excesses," *Critical Care Medicine*, vol. 19, no. 11, pp. 1432–1438, 1991.
- [7] J. Cuschieri, E. P. Rivers, M. W. Donnino et al., "Central venous-arterial carbon dioxide difference as an indicator of cardiac index," *Intensive Care Medicine*, vol. 31, no. 6, pp. 818–822, 2005.
- [8] J. Creteur, D. De Backer, Y. Sakr, M. Koch, and J.-L. Vincent, "Sublingual capnometry tracks microcirculatory changes in septic patients," *Intensive Care Medicine*, vol. 32, no. 4, pp. 516–523, 2006.
- [9] M. Singer, C. S. Deutschman, C. W. Seymour et al., "The third international consensus definitions for sepsis and septic shock (sepsis-3)," *JAMA*, vol. 315, no. 8, pp. 801–810, 2016.
- [10] M. A. Quiñones, C. M. Otto, M. Stoddard, A. Waggoner, and W. A. Zoghbi, "Recommendations for quantification of Doppler echocardiography: a report from the Doppler quantification task force of the nomenclature and standards committee of the American society of echocardiography," *Journal of the American Society of Echocardiography*, vol. 15, no. 2, pp. 167–184, 2002.
- [11] M. H. Weil, E. C. Rackow, R. Trevino, W. Grundler, J. L. Falk, and M. I. Griffel, "Difference in acid-base state between venous and arterial blood during cardiopulmonary resuscitation," *The New England Journal of Medicine*, vol. 315, no. 3, pp. 153–156, 1986.
- [12] J. Bakker, J.-L. Vincent, P. Gris, M. Leon, M. Coffernils, and R. J. Kahn, "Veno-arterial carbon dioxide gradient in human septic shock," *Chest*, vol. 101, no. 2, pp. 509–515, 1992.
- [13] M. Dres, X. Monnet, and J.-L. Teboul, "Hemodynamic management of cardiovascular failure by using PCO(2)

- venous-arterial difference,” *Journal of Clinical Monitoring and Computing*, vol. 26, no. 5, pp. 367–374, 2012.
- [14] J. P. Ducey, J. M. Lamiell, and G. E. Gueller, “Arterial-venous carbon dioxide tension difference during severe hemorrhage and resuscitation,” *Critical Care Medicine*, vol. 20, no. 4, pp. 518–522, 1992.
- [15] B. Lamia, X. Monnet, and J. L. Teboul, “Meaning of arterio-venous PCO₂ difference in circulatory shock,” *Minerva Anestesiologica*, vol. 72, no. 6, pp. 597–604, 2006.
- [16] P. A. Van Beest, M. C. Lont, N. D. Holman, B. Loef, M. A. Kuiper, and E. C. Boerma, “Central venous-arterial pCO₂ difference as a tool in resuscitation of septic patients,” *Intensive Care Medicine*, vol. 39, no. 6, pp. 1034–1039, 2013.
- [17] F. Vallée, B. Vallet, O. Mathe et al., “Central venous-to-arterial carbon dioxide difference: an additional target for goal-directed therapy in septic shock?,” *Intensive Care Medicine*, vol. 34, no. 12, pp. 2218–2225, 2008.
- [18] X. Monnet and J. L. Teboul, “My patient has received fluid. How to assess its efficacy and side effects?,” *Annals Intensive Care*, vol. 8, no. 1, p. 54, 2018.
- [19] D. Lichtenstein, “FALLS-protocol: lung ultrasound in hemodynamic assessment of shock,” *Heart, Lung and Vessels*, vol. 5, no. 3, pp. 142–147, 2013.