

Biodiversity of the Great Barrier Reef—how adequately is it protected?

Zoe T. Richards^{1,2} and Jon C. Day³

¹ Trace and Environmental DNA Laboratory, School of Molecular and Life Sciences, Curtin University of Technology, Perth, WA, Australia

² Aquatic Zoology Department, Western Australian Museum, Welshpool, WA, Australia

³ ARC Centre of Excellence for Coral Reef Studies, James Cook University of North Queensland, Townsville, QLD, Australia

ABSTRACT

Background: The Great Barrier Reef (GBR) is the world's most iconic coral reef ecosystem, recognised internationally as a World Heritage Area of outstanding significance. Safeguarding the biodiversity of this universally important reef is a core legislative objective; however, ongoing cumulative impacts including widespread coral bleaching and other detrimental impacts have heightened conservation concerns for the future of the GBR.

Methods: Here we review the literature to report on processes threatening species on the GBR, the status of marine biodiversity, and evaluate the extent of species-level monitoring and reporting. We assess how many species are listed as threatened at a global scale and explore whether these same species are protected under national threatened species legislation. We conclude this review by providing future directions for protecting potentially endangered elements of biodiversity within the GBR.

Results: Most of the threats identified to be harming the diversity of marine life on the GBR over the last two–three decades remain to be effectively addressed and many are worsening. The inherent resilience of this globally significant coral reef ecosystem has been seriously compromised and various elements of the biological diversity for which it is renowned may be at risk of silent extinction. We show at least 136 of the 12,000+ animal species known to occur on the GBR (approximately 20% of the 700 species assessed by the IUCN) occur in elevated categories of threat (*Critically Endangered*, *Endangered* or *Vulnerable*) at a global scale. Despite the wider background level of threat for these 136 species, only 23 of them are listed as threatened under regional or national legislation.

Discussion: To adequately protect the biodiversity values of the GBR, it may be necessary to conduct further targeted species-level monitoring and reporting to complement ecosystem management approaches. Conducting a vigorous value of information analysis would provide the opportunity to evaluate what new and targeted information is necessary to support dynamic management and to safeguard both species and the ecosystem as a whole. Such an analysis would help decision-makers determine if further comprehensive biodiversity surveys are needed, especially for those species recognised to be facing elevated background levels of threat. If further monitoring is undertaken, it will be important to ensure it aligns with and informs the GBRMPA Outlook five-year reporting schedule. The potential also exists to incorporate new environmental DNA technologies into routine

Submitted 27 October 2017

Accepted 21 April 2018

Published 8 May 2018

Corresponding author

Zoe T. Richards,
zoe.richards@curtin.edu.au

Academic editor

James Reimer

Additional Information and
Declarations can be found on
page 15

DOI 10.7717/peerj.4747

© Copyright

2018 Richards and Day

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

monitoring to deliver high-resolution species data and identify indicator species that are cursors of specific disturbances. Unless more targeted action is taken to safeguard biodiversity, we may fail to pass onto future generations many of the values that comprise what is universally regarded as the world's most iconic coral reef ecosystem.

Subjects Biodiversity, Marine Biology, Climate Change Biology, Natural Resource Management, Environmental Impacts

Keywords World heritage area, Values, Threatened species, IUCN, Species, Climate change, Value of information analysis, Silent extinctions, Monitoring and reporting, Marine conservation

INTRODUCTION

The Great Barrier Reef (GBR) is a diverse ecosystem extending for more than 2,300 km along Australia's northeast coast. It is recognised internationally as being of outstanding universal value (*United Nations Educational, Scientific and Cultural Organisation, 1981; Lucas et al., 1997; GBRMPA, 2014a*). Its diversity includes but is not restricted to over 410 species of hard coral, over 1,620 species of fish, 2,000 species of sponge, 14 species of sea snake, six of the world's seven species of marine turtle, at least 300 mollusc species, 630 species of echinoderm, and 500 species of marine alga. No other World Heritage Area on the planet contains such diversity (*Day, 2016*) and the GBR's exceptional biodiversity values were specifically mentioned in two of the four criteria for natural heritage (ix and x) when it was World Heritage listed in 1981 (*United Nations Educational, Scientific and Cultural Organisation, 1981*).

The long-term protection and conservation of biodiversity values is at the core of the primary legislative objective for the GBR (s. 2A of the Act, *Great Barrier Reef Marine Park Act, 1975*) of all recent GBR planning and management documents such as the 2014 Outlook Report (*GBRMPA, 2014a*), the 2014 Strategic Assessment (*GBRMPA, 2014b*) and the Reef 2050 Sustainability Plan (Reef 2050 Plan) (*Commonwealth of Australia, 2015*) reflect this. However, a recent report on the feasibility of the Reef 2050 Plan suggests that maintaining the GBR's Outstanding Universal Value and biodiversity as we know it may no longer be realistic, and it is recommended that the key managerial focus should be preserving the ecological function of GBR ecosystems (*Roth et al., 2017*).

This recommendation is based on the ongoing habitat deterioration of the GBR, even in sectors where there is minimal human influence (*Hughes et al., 2017*). It is also based on the finding that windows of opportunity for recovery after disturbances are narrowing (*Hughes et al., 2018*)—a consequence of the lack of progress that has been made towards global emission and local water quality targets. While such a policy change may be inevitable, it should not preclude biodiversity conservation. In addition to the moral and aesthetic reasons for protecting biodiversity, the conservation of diversity is important for protecting the functioning and resilience of the ecosystem as a whole. This is because inherent variation in species responses to, and recovery after disturbances provides ecological insurance in the face of change (*Nyström et al., 2008*). Furthermore, diversity buffers ecosystems against environmental change and increases the stability

and functioning of ecosystem processes (Griffin *et al.*, 2009; Loreau & de Mazancourt, 2013). Conversely, species loss can accelerate change in ecosystem processes (Stachowicz, Bruno & Duffy, 2007; Perrings *et al.*, 2011; Hooper *et al.*, 2012) and the extinction of a species may have unforeseen impacts (Dulvy, Sadovy & Reynolds, 2003).

There is little doubt that quantitative scenarios for the future of biodiversity in the 21st century are bleak (Pereira *et al.*, 2010), however there is still a chance to intervene through better policies. Given biodiversity is so intimately related to ecosystem functioning, in this review, we focus on the biodiversity values of the GBR. We acknowledge that in complex and socio-economically valuable ecosystems where there are multiple stakeholders and users, conservation decisions should never be made with biodiversity data alone. However, the purpose of this review is to introduce the biodiversity values of the GBR, summarise the threats directly or indirectly impacting these values, evaluate the status of marine life and examine the influence of scale on threatened species management approaches. Our objective is not to provide an exhaustive commentary on the biodiversity conservation literature (see Yoccoz, Nichols & Boulinier, 2001; Stem *et al.*, 2005; Ferraro & Pattanayak, 2006 for general reviews of monitoring and evaluation in conservation), or the challenges faced by environmental managers and policymakers (see Anthony, 2016), but rather to highlight opportunities for optimizing the evaluation and protection of biodiversity on the GBR.

SURVEY METHODOLOGY

The 2014 GBR Outlook Report (GBRMPA, 2014a) was chosen as the initial primary source of key references of the key threats/pressures impacting upon the GBR. The 2014 Outlook Report is a comprehensive compilation of information about the GBR underpinned by many references. Searches of the Outlook Report were undertaken against the following keywords: climate change, water quality, coastal development, shipping, unsustainable fishing, diseases, pests, and marine debris. Eighty-two articles that referred to the consequences, impacts or implications of the threats/pressures for species on the GBR were retained and included in Table S1. Table S1 was further augmented by conducting the same keyword searches of the James Cook University library, with only those articles directly relevant to species on the GBR retained. Articles that did not refer to consequences, impacts or implications of the threats/pressures for species were excluded. Precedence was given to articles published within the last 15 years; however, five articles published in the 1990s were included in Table S1 as they provided critical information not available elsewhere. In total, 125 titles were included in our review of the key threats/pressures impacting species within the GBR and these are listed in Table S1. We also conducted Google Scholar searches focusing on the ecological, environmental management, and conservation planning literature for articles relating to the challenges presented by monitoring and managing diversity in coral reef ecosystems, coral reef surrogates and proxy metrics. We examined the 2014 GBR Outlook Report (GBRMPA, 2014a) to quantify how many species were known from the GBR and the threatened status of these fauna based on national (EPBC Act: <http://www.environment.gov.au/marine/marine-species/marine-species-list>) and international




Figure 1 Cumulative impacts on the Great Barrier Reef. Cumulative impacts on the Great Barrier Reef that are directly or indirectly affecting species. For more information on key threats/pressures and the implications for species see [Table S1](#). Water quality image—Photographer: C. Honchin, Copyright Commonwealth of Australia (GBRMPA); Marine debris image—Photographer: S. Whiting, Copyright Commonwealth of Australia (GBRMPA); Shipping image—Photographer: P. Howorth, Copyright Commonwealth of Australia (GBRMPA); Coastal Development image courtesy Queensland Government Department of Heritage and Protection under Creative Commons Attribution 3.0 Australia (CC BY) license (<http://www.ehp.qld.gov.au/legal/copyright.html>). All other images by Zoe Richards.

Full-size DOI: [10.7717/peerj.4747/fig-1](https://doi.org/10.7717/peerj.4747/fig-1)

threatened species lists (IUCN red-list <http://www.iucnredlist.org/>). This comparison was undertaken for approximately 700 species (including hard corals, sea cucumbers, giant clams, grouper, wrasse, parrotfish, sharks, rays, sea snakes, turtles, whales, dolphins, and dugong) whose conservation status has been assessed at a global level. Lastly we used a Google Scholar search to locate titles that relate to optimal coral reef monitoring and alternative coral reef monitoring technologies.

RESULTS AND DISCUSSION

Current status of the diversity of marine life on the GBR

Despite significant management actions (*Fernandes et al., 2005*; *Commonwealth of Australia, 2015*), the diversity of marine life on the GBR is threatened by seven key pressures (climate change, declining water quality, coastal development, shipping impacts, fishing impacts, diseases and pest species, and marine debris) ([Fig. 1](#)). These direct or indirect pressures have led to at least 43 impacts which may adversely impact species on the GBR ([Fig. 1](#), see also [Table S1](#)). The latest extensive impact, the back-to-back thermal stress events in 2015–2016, led to 91% of surveyed GBR reefs experiencing coral bleaching

(Hughes et al., 2017). This large-scale coral bleaching event heightens concerns about whether the biodiversity values of the GBR are intact because it occurred at a time when parts of the GBR system were recovering from three category five cyclones (2010–2015, Perry et al., 2014), and a legacy of cumulative pressures (e.g. the commercial harvesting of dugongs, green turtles, pearl shell, trochus, and sea cucumbers; catchment management issues, including land clearing, changes to natural water flows and water pollution (GBRMPA, 2013a; Anthony, 2016)).

Prior to the 2016 bleaching event, both habitats and species in the southern two-thirds of the GBR, and particularly in the inshore areas, had reportedly declined (Brodie & Waterhouse, 2012; GBRMPA, 2009, 2014a; Osborne et al., 2011). A temporal analysis of the level of coral cover indicated that from 1985 to 2012 reef-wide coral cover declined by an average of 0.53% per year from 28.0% to 13.8% (De'ath et al., 2012). Between 1988 and 2003 coral calcification rates are also reported to have declined from 1.96 g cm⁻² year⁻¹ (±0.05) to 1.59 cm⁻² year⁻¹ (±0.04), equivalent to a decline of 14–21% over this period (Cooper et al., 2008; De'ath Lough & Fabricius, 2009). Degradation was not, however, restricted to the corals; sea grass health, particularly in central GBR was poor (Fairweather, McKenna & Rasheed, 2011; McKenzie, Collier & Waycott, 2015), dugong numbers had declined abruptly (Marsh et al., 2001; Sobotzick et al., 2012), hawksbill turtles were in decline (Bell, Schwarzkopf & Manicom, 2012) and some sharks, rays, and large fish populations were in decline, especially in coastal and inshore areas (Chin et al., 2010; Ceccarelli et al., 2014). Overall, even before the 2016 bleaching event, the inherent resilience of this globally significant reef ecosystem had been seriously compromised and there was a concern that numerous elements of the biological diversity for which it is renowned may have been at risk (World Heritage Committee, 2014, 2015; Day, 2015).

In complex and dynamic ecosystems like coral reefs, biodiversity in the true sense of the word—encompassing genetic, species, community, and ecosystem diversity is enigmatic (Gaston, 1998). On the GBR at least 12,000 species of marine vertebrates and invertebrates have been reported (Table 1; Pitcher et al., 2007; Hutchings, Kingsford & Hoegh-Guldberg, 2008). However, this estimate of species diversity is highly conservative. For many groups, only preliminary estimates are available and these are often based on survey data that were collected 20–40 years ago. Furthermore, condition and trend information is only available for only a limited number of species (Fabricius & De'ath, 2001; GBRMPA, 2014a); hence even for many large charismatic species of high conservation significance, the status of populations is uncertain (Hamann & Chin, 2015). Cryptic lineages (Schmidt-Roach et al., 2013); new species (e.g. Hooper & Van Soest, 2006; Miller & Downie, 2009; Sutcliffe, Hooper & Pitcher, 2010; Hunter & Cribb, 2012; Schmidt-Roach, Miller & Andreakis, 2013; Capa & Murray, 2015) and even entire habitats such as sponge gardens, mesophotic reefs, and deep water corals, are still being discovered and mapped (e.g. Bridge & Guinotte, 2012; Bridge et al., 2012; Beaman, 2012; Harris et al., 2013; López-Cabrera et al., 2016; McNeil et al., 2016).

Table 1 Threatened species on the Great Barrier Reef.

	Number of species recorded on the GBR*	Listed threatened species	
		Australia's environment protection and biodiversity conservation act 1999	Global red list index (critically endangered, endangered or vulnerable)
Sponges [#]	2,500	0	Not assessed
Jellyfish [#]	100	0	Not assessed
Soft corals and sea pens [#]	150	0	Not assessed
Ascidians/tunicates [#]	720	0	Not assessed
Bryozoans [#]	950	0	Not assessed
Anemones [#]	40	0	Not assessed
Hard corals [#]	450	0	88
Echinoderms [#]	630	0	10
Crustaceans [#]	1,300	0	Not assessed
Molluscs [#]	3,000	0	2
Insects and arachnids [#]	25	0	Not assessed
Worms [#]	500	0	Not assessed
Bony fishes	1,625	1	5
Sharks and rays [∞]	136	9	21
Breeding sea snakes	14	0	0
Marine turtles	6	6	5
Whales and dolphins	30	6	4
Dugong	1	1	1
Total	12,177	23	136

Notes:
The number of animal species known to occur on the GBR contrasted with the number listed on national and global threatened species lists. Adapted from the 2014 Outlook Report (*GBRMPA, 2014a*) and the 2016 IUCN Red List.

* Excludes crocodile, nesting seabirds, shorebirds, plankton and marine flora.

[#] Best available estimate.

[∞] A. Chin, 2016, unpublished data.

Current approaches to informing coral reef management for biodiversity conservation

The task of protecting marine biodiversity is immense and exacerbated by the logistic challenges of conducting species-level surveys and the high level of taxonomic expertise needed to identify species. Hence, including a large number of species in routine monitoring can be regarded as both impractical and in some cases ineffective (*Bottrill et al., 2008*). Moreover, in contemporary conservation science there is a growing view that monitoring can be a waste of resources rather than a prerequisite for optimal management (*Legg & Nagy, 2006*). This perspective has been fuelled by the decades of monitoring studies that have reported population declines with no apparent link to management objectives (*Nichols & Williams, 2006*), or without any responsive action being taken. Furthermore, the amount of money and capacity required to protect all biodiversity is considered astronomical and far beyond the current investment in conservation action (*James, Gaston & Balmford, 2001*). Hence, by necessity, ecosystem management approaches adopt a triage approach that involves prioritizing the investment

of scarce resources in a reduced set of factors that are more manageable (*Bottrill et al., 2008; Wilson et al., 2011*).

The principal way managers are informed about the status of biodiversity is through surrogate information (defined in *Hunter et al., 2016*). In some cases, surrogate information can relate to subsets of data about indicator species (*Gardner et al., 2008*), cross-taxon surrogates (*Rodrigues & Brooks, 2007*), broad habitat-based proxy metrics (*Dalleau et al., 2010*) or abiotic surrogates (*Beier et al., 2015*). Some have argued that proxies reduce the time and cost required for data collection (*Humphries, Williams & Vane-Wright, 1995; Favareau et al., 2006*) suggesting that developing indicator, surrogate or proxy metrics that adequately represent diversity trends is an important and pragmatic conservation objective (*Baillie et al., 2008*). Numerous other studies, however, have questioned the ability for proxy metrics to effectively represent diversity (*Araújo et al., 2001; Rodrigues & Brooks, 2007; Andleman & Fagan, 2000*), highlighting that all proxy metrics have limitations (*Pressey, 2004*) especially if their performance is not evaluated with empirical data (*Vellend, Lilley & Starzomski, 2008*). One study examining the efficacy of biological surrogacy in seabed assemblages on the GBR indicated that no one taxonomic group was a particularly good surrogate for another and recommended that examining multiple taxonomic groups together was the preferred approach (*Sutcliffe et al., 2012*).

On the GBR, the overwhelming majority of diversity surrogate information is collected (and reported) at a habitat level. Relatively little data is available at the species-level. For example, all but two of the references cited in the 2014 Strategic Assessment (*GBRMPA, 2014b*) and the 2014 Outlook Report (*GBRMPA, 2014a*) to substantiate the condition and trend of coral species diversity on the GBR relate to the percentage of live coral cover. Likewise, a recent paper documenting the impact of the 2016 coral bleaching event (*Hughes et al., 2017*) reports species-level responses for only two of the 410 coral species known to occur on the GBR despite claiming to document the resistant corals and susceptible species. Moreover, 21 taxa were recorded at the level of genera, two genera were further broken down into growth forms, three taxa were reported at the family level and soft corals were reported at the level of order. Despite their important role as ecosystem engineers, for hard corals, there has not been a comprehensive species-level assessment of coral diversity on the GBR since 2001 (*DeVantier et al., 2006*) and species-level information is not routinely collected or reported on, hence the only species-level data available to underpin management decisions about the status of coral biodiversity on the GBR is now over 15 years old.

The situation is worse for other neglected marine taxa—no current information on status or trends exists for the overwhelming majority of marine taxa including highly targeted and vulnerable taxa such as sea cucumbers and giant clams (*Purcell et al., 2014*). Thus, given multiple disturbances (e.g. including bleaching and mortality events, cyclones, freshwater flood plumes, and a new outbreak of crown of thorn (COT) sea stars (*Acanthaster planci*) have impacted the GBR over the last decade, the questions remain—how can we be sure we are adequately protecting biodiversity if we have only a partial understanding of what is there? Do we have adequate

data to evaluate how biodiversity has responded to both management efforts and disturbance events?

Coral cover is a poor surrogate for biodiversity

On coral reefs, habitat proxies are commonly used to quantify the condition of coral reefs, with percent live hard coral cover being the most widely used metric in monitoring studies (*Bruno & Selig, 2007; Eakin et al., 2010; De'ath et al., 2012*). It is stated in the Reef 2050 Plan that the extent, condition and trend of habitat provide the best indicators of biodiversity (*Commonwealth of Australia, 2015*). However, despite its broad use, hard coral cover is not a robust indicator of coral diversity. A study undertaken at Lizard Island on the GBR showed that coral cover was not an effective proxy for coral diversity because coral cover is not related to species richness as a positive linear function (*Richards, 2013*). To further explore the generality of this finding, the study was repeated at four additional locations (Ashmore Reef, Kosrae-Micronesia, Majuro-Marshall Is., and Christmas Island; *Richards & Hobbs, 2014; Ryan, Richards & Hobbs, 2014*). In those studies, percent live coral cover consistently performed poorly as an indicator of coral species richness (*Fig. 2*).

Numerous studies have maintained that the level of coral cover can be used to make inferences about fish biomass (*Chabanet et al., 1997*); the prevalence of coral diseases (*Bruno et al., 2007*); reef accretion potential (*Perry et al., 2012*) and arguably reef aesthetics (*Pocock, 2002*) or the economic value of reefs relating to tourism (*Stoeckl et al., 2011*). However, focusing solely on collecting data on habitat condition using general indicators such as coral cover is counter-productive to species conservation because when used in isolation, habitat proxies are not informative about ecological condition (*Hughes et al., 2010*), functionality (*Alvarez-Filip et al., 2013*) or diversity (*Richards, 2013*). Moreover, the finding of high coral cover can be deceptive because cover can be driven by the dominance of a small number of species whilst a high level of diversity can be sustained even when coral cover is low or moderate. Thus, it is important to note that a habitat with high coral cover does not guarantee functional diversity (defined by *Cadotte, 2011* as the trait variation in an assemblage), or community reassembly after disturbance (*Hughes et al., 2010*). Communities with high coral cover may be dominated by one or a small number of species and a single disturbance event (i.e. selective bleaching, disease outbreak or COTs infestation) could wipe out these dominant habitat formers with cascading ecosystem effects. With greater diversity comes the increased likelihood that some species would survive disturbance events and that critical functions such as reef building will be maintained. Thus, monitoring coral cover alone can fail to alert decision-makers to declines in resilience and it may in-fact mask losses of resilience.

The risk of silent extinctions in the absence of an effective understanding of biodiversity

Without appropriate empirical baseline data (e.g. current population size and trend data), it is impossible to accurately detect or predict population growth, depletions, range shifts, to identify species with superior tolerance or to understand how species




Figure 2 The relationship between mean coral species richness and mean percent live coral cover.

(A). The results of a survey conducted at Lizard Island based on 28 sites showed that there was a polynomial relationship between coral species richness and percent live coral cover (adj $r^2 = 0.351$, SE 10.789, $df = 25$, $p = 0.001$) and significance of the quadratic term in multiple regression suggested a non-linear relationship where species richness peaked at intermediate levels of coral cover (Adapted from Richards, 2013). (B). When repeated at Christmas Island, no significant ($p > 0.05$) linear or polynomial relationships were found between percent live coral cover and species richness at any scale (site, depth or transect) (Adapted from Ryan, Richards & Hobbs, 2014). (C). When the relationship was analysed at three additional locations Pacific and Indian Ocean locations, the overall r^2 values were considered too low (i.e. below 0.7) to provide meaningful estimates of coral species richness regardless of the scale of analysis (black bars: site means; white bars: transect level) (Adapted from Richards & Hobbs, 2014).

Full-size DOI: 10.7717/peerj.4747/fig-2

respond to management effort. Furthermore, without baseline data it is likely that extinctions may be occurring as unrecorded *silent extinctions* (Myers & Ottensmeyer, 2005). Silent extinctions may initially take place as ecological extinctions (when a species is reduced to such low abundance that, although still present, it no longer plays its typical ecological role); but they may also occur as undetected local extinctions

(the disappearance of a species from part of its range) (*Estes, Duggins & Rathbun, 1989*). Both of these types of extinctions are precursors to wider extinction events.

Even though there have been few, if any, known extinctions on the GBR, there is reason to believe that the threat of extinction is growing in tandem with the increasing frequency and velocity of disturbance events. The GBR Strategic Assessment (*GBRMPA, 2014b*) states there are significant concerns about a small number of species, including the spear tooth shark which may now be extinct in the GBR Region. While some potentially at risk species on the GBR are protected by other legislative mechanisms (e.g. all species of sawfish which are protected under Queensland legislation, *GBRMPA, 2013b*)—others are not. For example, two species of sea cucumbers (*Holothuria whitmaei* and *H. fuscogilva*) that are listed as *Endangered* and *Vulnerable* on the IUCN red-list index (RLI) (*IUCN, 2015*) have been heavily fished, and the annual catch of another Vulnerable species (*Stichopus herrmanni*) rose at an average annual pace of 200% from 2007 to 2011 (*Eriksson & Byrne, 2013*) however none of these species are protected under Australian legislation.

Other habitat-forming species such as hard corals are listed as no-take species under the GBRMPA Act 1975 which affords them protection from extractive threats. However, no stony coral species are specifically listed as threatened species under the *EPBC Act*, and their actual population status is not known nor is the effectiveness of current legislation as a regulatory control. Furthermore, with no post-impact data recorded at a species-level, it is impossible to report the impact that disturbance events (such as the recent widespread coral bleaching event) have had, nor is it possible to substantiate species recovery.

The significant and escalating situation on the GBR was placed in the international spotlight by UNESCO's World Heritage Committee (WHC) in 2011, and the Committee has since continued to raise concerns about the status of the GBR World Heritage Area (e.g. *World Heritage Committee, 2013, 2014, 2015*). In 2014, the WHC requested that the Australian government provide concrete and consistent management measures to ensure the overall long-term conservation of the property, including addressing cumulative impacts and increase reef resilience. In response, the Reef 2050 Plan was drafted (*Commonwealth of Australia, 2015*). As highlighted elsewhere (*Australian Academy of Science, 2015; Hughes, Day & Brodie, 2015; Roth et al., 2017*), the Reef 2050 Plan falls short on responding to these requests especially in the context of addressing climate change, shipping impacts, setting realistic, and measurable targets, cumulative impacts and a commitment to the level of funding required. The Reef 2050 Plan does, however, provide some useful actions and targets to monitor population trends for some mega fauna such as dugongs and turtles. The provision of strategies to evaluate and report on the population trends of other non-charismatic taxa like benthic invertebrates is notably absent. For example, the condition and trend for the 'other invertebrates' section in the GBR Strategic Assessment is listed as 'very good' and 'stable' but there is no confidence in the data and very limited evidence (*GBRMPA, 2014b*). Also in the 'sea snakes' section of the same assessment, it states abundance estimates are only available for a few species or for small areas, and there is little information about population trends.

Scale-based mismatches in threatened species assessments

For most of the 12,000+ animal species that are documented from the GBR (Table 1), regional status information is notably absent (exceptions are nesting seabirds and shorebirds). However, for approximately 700 species including hard corals, sea cucumbers, giant clams, grouper, wrasse, parrotfish, sharks, rays, sea snakes, turtles, whales, dolphins, and dugong (Carpenter *et al.*, 2008; Elfes *et al.*, 2013; Sadovy de Mitcheson *et al.*, 2013; Conand *et al.*, 2014; Dulvy *et al.*, 2014), conservation status has been assessed at a global level using the Red List of Threatened Species (RLI) (IUCN, 2015). The global assessments indicate that at least 136 species on the GBR (approximately 20% of species assessed) occur in elevated categories of threat (*Critically Endangered*, *Endangered* or *Vulnerable*) (Table 1) including at least 89 species of hard coral, 10 species of sea cucumbers, 21 species of sharks and rays, two species of giant clam, and five species of bony fish (Table S2).

Despite the wider background level of threat for these 136 species, only 23 of them (17%) are listed as threatened under the *Environment Protection and Biodiversity Conservation Act, 1999* (Australian Government, 2015) or defined as protected species under the *Great Barrier Reef Marine Park Regulations, 1983*. The regional status of the other 113 assessed species that have been recognised as highly threatened on a global scale by their respective taxonomic experts is not known (e.g. the Narrow Sawfish (*A. cuspidata*), Purple Eagle Ray (*Myliobatus hamlyni*), Humphead Wrasse (*Cheilinus undulatus*, see Fig. 3) and five species of sea cucumbers (Holothridae)). Nor is the status of an additional 35 species that are listed on the RLI as Data Deficient (Table S2, only three of which are protected under EPBC), or the other 11,000+ animal species known or likely to occur on the GBR which have not been assessed at any scale (global, national or regional, see Table 1). These species are not protected under the *Environment Protection and Biodiversity Conservation Act, 1999* because they are not endemic to Australia and the global IUCN assessments have little bearing on decisions made under the EPBC Act unless the species in question is endemic to Australia.

Future directions

In order to effectively manage the GBR a multi-pronged approach is needed (McCook *et al.*, 2009). Pragmatism has led to a preference for system-wide ecosystem management with the result that species-level initiatives have been de-emphasised. However, obtaining a comprehensive and up-to-date understanding of the status of GBR marine life through targeted taxonomic and ecological research on key knowledge gaps and long-term species monitoring remains a fundamental way to make informed management decisions (GBRMPA, 2009, 2013a, 2014a) and creates opportunities to engage with the public and bolster support for biodiversity conservation. The key argument against species-level surveys is that they are prohibitively expensive. But is this true? By way of example, a large multi-institutional, multi-taxon marine biodiversity project (*Marine Life of Kimberley* project) was recently conducted in NW Australia. Just like the GBR, the Kimberley marine bioregion comprises diverse coastal and offshore habitats and a large number of islands with fringing reef habitats. However, it covers a larger and





Figure 3 Examples of species that are under threat at a global scale but their regional status is not known. (A) *Tridacna gigas*, a vulnerable clam species (see <http://www.iucnredlist.org/details/22137/0>). (B) *Acropora donei*, a vulnerable staghorn coral species (see <http://www.iucnredlist.org/details/133223/0>). (C) *Cheilinus undulatus*, an endangered species of wrasse (see <http://www.iucnredlist.org/details/4592/0>). Photos by Zoe Richards. Full-size  DOI: 10.7717/peerj.4747/fig-3

even more remote area (Kimberley project area is 476,000 km² (Bryce, Bryce & Radford, 2018); the GBR is 347,800 km² (GBRMMPA, 2014a)). The Kimberley Marine Life project had two phases. The first phase (2008–2011) involved the compilation and analyses of

historic data from several Australian museums and the WA Herbarium to give a state of current knowledge for the region's marine life. Phase 2 (2011–2015) involved a series of surveys to increase the resolution of the marine diversity data and fill knowledge gaps identified during Phase 1. The project targeted the region's molluscs, crustaceans, fish, hard corals, soft corals, sponges, echinoderms, worms, marine algae, and sea grasses (Bryce & Sampey, 2014; Fromont & Sampey, 2014; Hosie et al., 2015; Huisman & Sampey, 2014; Hutchings et al., 2014; Moore et al., 2015; Richards, Sampey & Marsh, 2014; Sampey & Marsh, 2015; Willan, Bryce & Slack-Smith, 2015). This \$2.7 AUD million dollar project (plus in-kind support) has provided a comprehensive reference dataset to guide in the design of the newly gazetted marine parks in the region; informed a range of research projects (WAMSI; <http://www.wamsi.org.au>); led to award-winning educational outreach opportunities (<http://museum.wa.gov.au/btw/>) and established a knowledge legacy for future generations. If an equivalent annual financial investment was made into updating our understanding of the status of marine life on the GBR, this would equate to less than 0.42% of the estimated annual worth of the reef to Australia's economy (valued at \$6.4 billion annually, *Deloitte Access Economics, 2017*).

If new biodiversity studies are undertaken on the GBR, it would be crucial for this new information to be made available to decision-makers in a timely manner to ensure enduring and constructive links are made between monitoring and subsequent management actions (Day, 2008). More specifically, it must align with the five-year GBR outlook reporting schedule (the next report is due in 2019). In lieu of new surveys, a value of information analysis (Runge, Converse & Lyons, 2011; Moore & Runge, 2012; Yokomizo, Coutts & Possingham, 2014) could be undertaken to provide scientifically credible advice on how, when, and what new information is needed to broaden and accelerate efforts to conserve the biodiversity values of the GBR. Such an analysis may, for example, find there is a need to expand the existing monitoring program for the GBR to include key diversity indices. For some taxa, biodiversity proxies have been established e.g. particular families have been identified as good indicators of wider reef fish diversity (Allen & Werner, 2002), however for most, this information is not available and more research is needed in this area.

Any expansion of the existing GBR long-term monitoring program would need to be carefully considered and negotiated amongst a body of experts, managers, stakeholders, independent advisors, and preferably, informed by a value of information analysis. As a start, some possible options for adapting and expanding the monitoring program include:

1. Targeted top and tail monitoring which involves collating species-level data on a selection of the most common (keystone) species in addition to some of the most endangered and vulnerable species and/or invasive species.
2. Rotational monitoring whereby different taxa are monitored in a four-year cycle to increase the breadth of species captured in the overall monitoring program (see Fig. 4).
3. Stratified monitoring that involves in-depth taxonomic and ecological assessments at a smaller number of sites that overlap with key long-term monitoring stations.




Figure 4 Rotational biodiversity monitoring is one example of how existing monitoring programs could be adapted. If long-term monitoring programs are adapted, all decisions relating to target species and yearly taxa subsets would require expert discussion following baseline surveys or a Value of Information Analysis. Such a program would require resourcing and co-ordination but the vision could be for it to be conducted by two to three scientists per year (rotating based on expertise) and undertaken alongside the existing long-term monitoring program. Credits: Aerial—Photographer: M. Cowlishaw, Copyright Commonwealth of Australia (GBRMPA); Humpback Whale—Photographer: M. Simmons, Copyright Commonwealth of Australia (GBRMPA); Sea grass—Photographer: J. Jones, Copyright Commonwealth of Australia (GBRMPA); Great White Shark—Photographer: K. Hoppen, Copyright Commonwealth of Australia (GBRMPA); *Halimeda*—Photographer: G. Goby, Copyright Commonwealth of Australia (GBRMPA); Dugong—Photographer: B. Cropp, Copyright Commonwealth of Australia (GBRMPA); Turtle—Photographer: E. Goodwin, Copyright Commonwealth of Australia (GBRMPA). All other images by Zoe Richards. [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90_img.jpg\) DOI: 10.7717/peerj.4747/fig-4](https://doi.org/10.7717/peerj.4747/fig-4)

While these are just a few examples of possible ways the monitoring and reporting program could evolve, they illustrate that if preventing coral biodiversity loss is a priority to coral reef management authorities, it is essential to begin the process of collecting species-level data, using that information to choose appropriate indicators or surrogates (Beier *et al.*, 2015) and adapting monitoring programs to collect additional complementary data that more effectively informs managers and the public about the status of biodiversity.

In addition, there are other reporting opportunities. In some cases, existing photographs and video records could be retrospectively analysed to identify trends for key species of interest (e.g. giant clams). Moreover, citizen science projects (such as ReefLife Surveys) hold promise for providing data on charismatic or easily identified

fauna. For some taxa, estimates of generic diversity (Richards, 2013, Richards & Hobbs, 2014) or functional diversity (Cadotte, 2011; Cernansky, 2017) may be useful and pragmatic additions to monitoring programs. An additional promising approach is to audit species composition is through the application of environmental DNA (eDNA) meta-barcoding technologies. Advances in DNA sequencing technologies and the swift drop in the cost of sequencing has led to a rapid rise in the applications of eDNA, particularly in the marine environment (e.g. Foote et al., 2012; Lejzerowicz et al., 2015; Thomsen et al., 2012; Valentini et al., 2016; Miya et al., 2015). The power and limitations surrounding eDNA as a tool for marine surveys of diversity, diet, food webs and invasive species is currently being explored by numerous molecular laboratories. So far it appears that so long as careful attention is paid to meta-barcoding workflows, assay development and the development of taxonomically sound reference datasets, eDNA represents a powerful new tool. eDNA is increasingly being used to evaluate the composition and health of marine communities (Foote et al., 2012; Murray, Coghlan & Bunce, 2015; Clarke et al., 2017) including coral reefs (Stat et al., 2017; Shinzato et al., 2018) and may be a useful complement to traditional monitoring approaches.

CONCLUSION

Most of the threats identified to be harming the diversity of marine life on the GBR over the last two–three decades remains to be effectively addressed and many are worsening. The habitat degradation that has already occurred will take decades to reverse and far greater resources than are currently being expended are required to document and protect biodiversity on the GBR (Day, 2015; Brodie, 2016). What is urgently required is decisive and effective conservation action by key leaders to ensure the legal obligations of protecting biodiversity are met. Establishing sound taxonomic datasets and ensuring biodiversity is adequately and representatively monitored are urgent prerequisites for achieving efficient conservation plans and mitigating biodiversity loss (Wilson, 2016; Troudet et al., 2017; Thomson et al., 2018). The problem of having a limited understanding of the species that inhabit our oceans and the threats they face is not restricted to the GBR or Australia (Webb & Mindel, 2015). However, for Australia, the problem could also be an opportunity for best-practice marine conservation. The changing nature of the GBR necessitates we consider what additional dynamic approaches to monitoring and reporting are needed to more fully understand the current status of biodiversity and mitigate the risk that silent extinction events will occur. Unless more action is taken to evaluate and manage biodiversity and to provide current and comprehensive assessments of extinction risk (Costello, May & Stork, 2013; Bland et al., 2015; Costello, 2015), we may fail to pass onto future generations many of the values that comprise what is universally regarded as the world's most iconic coral reef ecosystem.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the WA Museum and Woodside Energy and a Curtin Research Fellowship awarded to Zoe T. Richards. The work was undertaken as part of

Linkage Project LP160101508. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
WA Museum and Woodside Energy and a Curtin Research Fellowship.
Linkage Project: LP160101508.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Zoe T. Richards conceived and designed the experiments, analysed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Jon C. Day prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Data Availability

The following information was supplied regarding data availability:
This research did not generate any data or code.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4747#supplemental-information>.

REFERENCES

- Allen GR, Werner TB. 2002. Coral reef fish assessment in the ‘coral triangle’ of southeastern Asia. *Environmental Biology of Fishes* 65:209–214 DOI 10.1023/a:1020093012502.
- Alvarez-Filip L, Carricart-Ganivet JP, Horta-Puga G, Iglesias-Prieto R. 2013. Shifts in coral assemblage composition do not ensure persistence of reef functionality. *Scientific Reports* 3(1):3486 DOI 10.1038/srep03486.
- Andleman SL, Fagan WF. 2000. Umbrellas and flagships: efficient conservation surrogates or expensive mistakes? *Proceedings of the National Academy of Sciences of the United States of America* 97(11):5954–5959 DOI 10.1073/pnas.100126797.
- Anthony KRN. 2016. Coral reefs under climate change and ocean acidification: challenges and opportunities for management and policy. *Annual Review of Environment and Resources* 41(1):59–81 DOI 10.1146/annurev-environ-110615-085610.
- Araújo MB, Humphries CJ, Densham PJ, Lampinen R, Hagemeyer WJM, Mitchell-Jones AJ, Gasc JP. 2001. Would environmental diversity be a good surrogate for species diversity? *Ecography* 24(1):103–110 DOI 10.1034/j.1600-0587.2001.240112.x.
- Australian Academy of Science. 2015. Reef 2050 long-term sustainability plan—position statement. Available at <https://www.science.org.au/supporting-science/science-policy/position-statements/reef-2050-long-term>.
- Australian Government. 2015. EPBC act list of threatened fauna. Available at <http://www.environment.gov.au/cgi-bin/sprat/public/publicthreatenedlist.pl> (accessed 12 June 2015).

- Baillie JE, Collen B, Amin R, Akcakaya HR, Butchart SHM, Brummitt N, Meagher TR, Ram M, Hilton-Taylor C, Mace GM. 2008. Toward monitoring global biodiversity. *Conservation Letters* 1(1):18–26 DOI 10.1111/j.1755-263x.2008.00009.x.
- Beaman RJ. 2012. New light on dark depths: Australia's Coral Sea landscape. In: *12th International Coral Reef Symposium, 9–13 July 2012*. Cairns: ARC Centre for Coral Reef Studies.
- Beier P, Sutcliffe P, Hjort J, Faith DP, Pressey RL, Albuquerque F. 2015. A review of selection-based tests of abiotic surrogates for species representation. *Conservation Biology* 29(3):668–679 DOI 10.1111/cobi.12509.
- Bell IP, Schwarzkopf L, Manicom C. 2012. High survivorship of an annually decreasing aggregation of hawksbill turtles, *Eretmochelys imbricata*, found foraging in the northern Great Barrier Reef. *Aquatic Conservation: Marine and Freshwater Ecosystems* 22(5):673–682 DOI 10.1002/aqc.2245.
- Bland LM, Collen B, Orme CDL, Bielby J. 2015. Predicting the conservation status of data-deficient species. *Conservation Biology* 29(1):250–259 DOI 10.1111/cobi.12372.
- Bottrill MC, Joseph LN, Carwardine J, Bode M, Cook C, Game ET, Grantham H, Kark S, Linke S, McDonald-Madden E, Pressey RL, Walker S, Wilson KA, Possingham HP. 2008. Is conservation triage just smart decision making? *Trends in Ecology & Evolution* 23(12):649–654 DOI 10.1016/j.tree.2008.07.007.
- Bridge T, Beaman R, Done T, Webster J. 2012. Predicting the location and spatial extent of submerged coral reef habitat in the Great Barrier Reef world heritage area, Australia. *PLOS ONE* 7(10):e48203 DOI 10.1371/journal.pone.0048203.
- Bridge T, Guinotte J. 2012. Mesophotic coral reef ecosystems in the Great Barrier Reef world heritage area: their potential distribution and possible role as refugia from disturbance, Research Publication No 109, Great Barrier Reef Marine Park Authority, Townsville. Available at <http://elibrary.gbrmpa.gov.au/jspui/handle/11017/2776>.
- Brodie J. 2016. Great Barrier Reef report to UN shows the poor progress on water quality. *The Conversation*, 2 December 2016. Available at <https://theconversation.com/great-barrier-reef-report-to-un-shows-the-poor-progress-on-water-quality-69779>.
- Brodie J, Waterhouse J. 2012. A critical review of environmental management of the 'not so Great' Barrier Reef. *Estuarine, Coastal and Shelf Science* 104–105:1–22 DOI 10.1016/j.ecss.2012.03.012.
- Bruno JF, Selig ER. 2007. Regional decline of coral cover in the Indo-Pacific: timing, extent, and subregional comparisons. *PLOS ONE* 2(8):e711 DOI 10.1371/journal.pone.0000711.
- Bruno JF, Selig ER, Casey KS, Page CA, Willis BL, Harvell CD, Sweatman H, Melendy AM. 2007. Thermal stress and coral cover as drivers of coral disease outbreaks. *PLOS Biology* 5(6):e124 DOI 10.1371/journal.pbio.0050124.
- Bryce C, Bryce M, Radford B. 2018. Project methods and station geomorphology related to a multi-taxon survey (2009–2014) of the Kimberley. *Records of the Western Australian Museum, Supplement* 85(1):1–43 DOI 10.18195/issn.0313-122x.85.2018.001-043.
- Bryce M, Sampey A. 2014. Kimberley marine biota. Historical data: soft corals and sea fans (Octocorallia). *Records of the Western Australian Museum, Supplement* 84(1):101–110 DOI 10.18195/issn.0313-122x.84.2014.101-110.
- Cadotte MW. 2011. The new diversity: management gains through insights into the functional diversity of communities. *Journal of Applied Ecology* 48(5):1067–1069 DOI 10.1111/j.1365-2664.2011.02056.x.
- Capa M, Murray A. 2015. A taxonomic guide to the fanworms (Sabellidae, Annelida) of Lizard Island, Great Barrier Reef, Australia, including new species and new records. *Zootaxa* 4019(1):98–167 DOI 10.11646/zootaxa.4019.1.8.

- Carpenter KE, Abrar M, Aeby G, Aronson RB, Banks S, Bruckner A, Chiriboga A, Cortes J, Delbeek JC, DeVantier L, Edgar GJ, Edwards AJ, Fenner D, Guzman HM, Hoeksema BW, Hodgson G, Johan O, Licuanan WY, Livingstone SR, Lovell ER, Moore JA, Obura DO, Ochavillo D, Polidoro BA, Precht WF, Quibilan MC, Reboton C, Richards ZT, Rogers AD, Sanciangco J, Sheppard A, Sheppard C, Smith J, Stuart S, Turak E, Veron JEN, Wallace C, Weil E, Wood E. 2008. One-third of reef-building corals face elevated extinction risk from climate change and local impacts. *Science* 321(5888):560–563 DOI 10.1126/science.1159196.
- Ceccarelli DM, Frisch AJ, Graham NA, Ayling AM, Beger M. 2014. Habitat partitioning and vulnerability of sharks in the Great Barrier Reef Marine Park. *Reviews in Fish Biology and Fisheries* 24(1):169–197 DOI 10.1007/s11160-013-9324-8.
- Cernansky R. 2017. Biodiversity moves beyond counting species. *Nature* 546(7656):22–24 DOI 10.1038/546022a.
- Chabanet P, Ralambondrainy H, Amanieu M, Faure G, Galzin R. 1997. Relationships between coral reef substrata and fish. *Coral Reefs* 16(2):93–102 DOI 10.1007/s003380050063.
- Chin A, Kyne PM, Walker TI, McAuley R. 2010. An integrated risk assessment for climate change: analysing the vulnerability of sharks and rays on Australia's Great Barrier Reef. *Global Change Biology* 16(7):1936–1953 DOI 10.1111/j.1365-2486.2009.02128.x.
- Clarke LJ, Beard JM, Swalding KM, Deagle BE. 2017. Effect of marker choice and thermal cycling protocol on zooplankton DNA metabarcoding studies. *Ecology and Evolution* 7(3):873–883 DOI 10.1002/ece3.2667.
- Commonwealth of Australia. 2015. Reef 2050 long-term sustainability plan. Available at <http://www.environment.gov.au/marine/gbr/publications/reef-2050-long-term-sustainability-plan>.
- Conand C, Polidoro BA, Mercier A, Gamboa RU, Hamel JE, Purcell SW. 2014. The IUCN red list assessment of aspidochirotid sea cucumbers and its implications. *SPC Beche-de-mer Information Bulletin* 34:3–7.
- Cooper TF, De'ath G, Fabricius KE, Lough JM. 2008. Declining coral calcification in massive *Porites* in two nearshore regions of the northern Great Barrier Reef. *Global Change Biology* 14(3):529–538 DOI 10.1111/j.1365-2486.2007.01520.x.
- Costello MJ. 2015. Biodiversity: the known, unknown, and rates of extinction. *Current Biology* 25(9):R368–R371 DOI 10.1016/j.cub.2015.03.051.
- Costello MJ, May RM, Stork NE. 2013. Can we name earth's species before they go extinct? *Science* 339(6118):413–416 DOI 10.1126/science.1230318.
- Dalleau M, Andréfouët S, Wabnitz CC, Payri C, Wantiez L, Pichon M, Friedman KI, Vigliola L, Benzoni F. 2010. Use of habitats as surrogates of biodiversity for efficient coral reef conservation planning in Pacific Ocean islands. *Conservation Biology* 24(2):541–552 DOI 10.1111/j.1523-1739.2009.01394.x.
- Day J. 2008. The need and practice of monitoring, evaluating and adapting marine planning and management—lessons from the Great Barrier Reef. *Marine Policy* 32(5):823–831 DOI 10.1016/j.marpol.2008.03.023.
- Day JC. 2015. The Barrier Reef is not listed as in danger, but the threats remain. *The Conversation*, 30 May 2015. Available at <https://theconversation.com/the-barrier-reef-is-not-listed-as-in-danger-but-the-threats-remain-42548>.
- Day JC. 2016. The Great Barrier Reef Marine Park—the grandfather of modern MPAs. In: Fitzsimmons J, Wescott G, eds. *Chapter 5 in Big, Bold and Blue: Lessons from Australia's Marine Protected Areas*. Victoria: CSIRO Publishing, 65–97.

- De'ath G, Fabricius KE, Sweatman H, Puotinen M. 2012. The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proceedings of the National Academy of Sciences of the United States of America* **109**(44):17995–17999 DOI [10.1073/pnas.1208909109](https://doi.org/10.1073/pnas.1208909109).
- De'ath G, Lough JM, Fabricius KE. 2009. Declining coral calcification on the Great Barrier Reef. *Science* **323**(5910):116–119 DOI [10.1126/science.1165283](https://doi.org/10.1126/science.1165283).
- Deloitte Access Economics. 2017. At what price? The economic, social and icon value of the Great Barrier Reef. Report to the Great Barrier Reef Foundation. Available at <https://www2.deloitte.com/content/dam/Deloitte/au/Documents/Economics/deloitte-au-economics-great-barrier-reef-230617.pdf> (accessed 28 July 2017).
- DeVantier LM, De'ath G, Turak E, Done TJ, Fabricius KE. 2006. Species richness and community structure of reef-building corals on the nearshore Great Barrier Reef. *Coral Reefs* **25**(3):329–340 DOI [10.1007/s00338-006-0115-8](https://doi.org/10.1007/s00338-006-0115-8).
- Dulvy NK, Davidson LN, Kyne PM, Simpfendorfer CA, Harrison LR, Carlson JK, Fordham SV. 2014. Ghosts of the coast: global extinction risk and conservation of sawfishes. *Aquatic Conservation: Marine and Freshwater Ecosystems* **26**(1):134–153 DOI [10.1002/aqc.2525](https://doi.org/10.1002/aqc.2525).
- Dulvy NK, Sadovy Y, Reynolds JD. 2003. Extinction vulnerability in marine populations. *Fish and Fisheries* **4**(1):25–64 DOI [10.1046/j.1467-2979.2003.00105.x](https://doi.org/10.1046/j.1467-2979.2003.00105.x).
- Eakin CM, Nim CJ, Brainard RE, Aubrecht C, Elvidge C, Gledhill DK, Muller-Karger F, Mumby PJ, Skirving WJ, Strong AE, Wang M, Weeks S, Wentz F, Ziskin D. 2010. Monitoring coral reefs from space. *Oceanography* **23**(4):118–133 DOI [10.5670/oceanog.2010.10](https://doi.org/10.5670/oceanog.2010.10).
- Elfes CT, Livingstone SR, Lane A, Lukoschek V, Sanders KL, Courtney AJ, Gatus JL, Guinea M, Lobo AS, Milton D, Rasmussen AR. 2013. Fascinating and forgotten: the conservation status of the world's sea snakes. *Herpetological Conservation & Biology* **8**:37–52.
- Eriksson H, Byrne M. 2013. The sea cucumber fishery in Australia's Great Barrier Reef Marine Park follows global patterns of serial exploitation. *Fish and Fisheries* **16**(2):329–341 DOI [10.1111/faf.12059](https://doi.org/10.1111/faf.12059).
- Estes JA, Duggins DO, Rathbun GB. 1989. The ecology of extinctions in kelp forest communities. *Conservation Biology* **3**(3):252–264 DOI [10.1111/j.1523-1739.1989.tb00085.x](https://doi.org/10.1111/j.1523-1739.1989.tb00085.x).
- Fabricius KE, De'ath G. 2001. Biodiversity on the Great Barrier Reef: large-scale patterns and turbidity-related local loss of soft coral taxa. In: Wolanski E, ed. *Oceanographic Processes of Coral Reefs, Physical and Biological Links in the Great Barrier Reef*. Boca Raton: CRC Press, 127–144.
- Fairweather C, McKenna S, Rasheed M. 2011. *Long-Term Seagrass Monitoring in the Port of Mourilyan: November 2010*. Cairns: DEEDI Publication, Fisheries Queensland, 27.
- Favareau JM, Drew CA, Hess GR, Rubino MJ, Koch FH, Eschelbach KA. 2006. Recommendations for assessing the effectiveness of surrogate species approaches. *Biodiversity and Conservation* **15**(12):3949–3969 DOI [10.1007/s10531-005-2631-1](https://doi.org/10.1007/s10531-005-2631-1).
- Fernandes L, Day JC, Lewis A, Slegers S, Kerrigan B, Breen DA, Cameron D, Jago B, Hall J, Lowe D, Innes J, Tanzer J, Chadwick V, Thompson L, Gorman K, Simmons M, Barnett B, Sampson K, De'ath G, Mapstone B, Marsh H, Possingham H, Ball I, Ward T, Dobbs K, Aumend J, Slater D, Stapleton K. 2005. Establishing representative no-take areas in the Great Barrier Reef: large-scale implementation of theory on marine protected areas. *Conservation Biology* **19**(6):1733–1744 DOI [10.1111/j.1523-1739.2005.00302.x](https://doi.org/10.1111/j.1523-1739.2005.00302.x).
- Ferraro PJ, Pattanayak SK. 2006. Money for nothing? A call for empirical evaluation of biodiversity conservation investments. *PLOS Biology* **4**(4):e105 DOI [10.1371/journal.pbio.0040105](https://doi.org/10.1371/journal.pbio.0040105).

- Foote AD, Thomsen PF, Sveegaard S, Wahlberg M, Kielgast J, Kyhn LA, Salling AB, Galatius A, Orlando L, Gilbert MT. 2012.** Investigating the potential use of environmental DNA (eDNA) for genetic monitoring of marine mammals. *PLOS ONE* 7(8):e41781 DOI 10.1371/journal.pone.0041781.
- Fromont J, Sampey A. 2014.** Kimberley marine biota. Historical data: sponges (Porifera). *Records of the Western Australian Museum, Supplement* 84(1):69–100 DOI 10.18195/issn.0313-122x.84.2014.069-100.
- Gardner TA, Barlow J, Araujo IS, Ávila-Pires TC, Bonaldo AB, Costa JE, Esposito MC, Ferreira LV, Hawes J, Hernandez MI, Hoogmoed MS, Leite RN, Lo-Man-Hung NF, Malcolm JR, Martins MB, Mestre LAM, Miranda-Santos R, Overall WL, Parry L, Peters SL, Ribeiro-Junior MA, Da Silva MNF, Da Silva Motta C, Peres CA. 2008.** The cost-effectiveness of biodiversity surveys in tropical forests. *Ecology Letters* 11(2):139–150 DOI 10.1111/j.1461-0248.2007.01133.x.
- Gaston KJ. 1998.** Biodiversity. In: Sutherland WJ, ed. *Conservation Science and Action*. Malden: Blackwell Science Ltd, 1–19.
- GBRMPA. 2009.** Great Barrier Reef outlook report 2009. Great Barrier Reef Marine Park Authority, Townsville. Available at http://www.gbrmpa.gov.au/__data/assets/pdf_file/0018/3843/OutlookReport_Full.pdf.
- GBRMPA. 2013a.** Great Barrier Reef biodiversity conservation strategy 2013, Great Barrier Reef Marine Park Authority, Townsville. Available at http://elibrary.gbrmpa.gov.au/jspui/bitstream/11017/2787/1/GBR%20BCS%2029%20April%202013_BD2_MAR.pdf.
- GBRMPA. 2013b.** Fitzroy Basin Assessment, Fitzroy Basin Association Natural Resource Management region: Assessment of ecosystem services within the Fitzroy Basin focusing on understanding and improving the health and resilience of the Great Barrier Reef, Great Barrier Reef Marine Park Authority, Townsville. Available at <http://elibrary.gbrmpa.gov.au/jspui/handle/11017/2896>.
- GBRMPA. 2014a.** Great Barrier Reef outlook report 2014. Great Barrier Reef Marine Park Authority, Townsville. Available at <http://elibrary.gbrmpa.gov.au/jspui/handle/11017/2855>.
- GBRMPA. 2014b.** Great Barrier Reef region strategic assessment: strategic assessment report 2014. Great Barrier Reef Marine Park Authority, Townsville. Available at <http://www.gbrmpa.gov.au/managing-the-reef/strategic-assessment>.
- Great Barrier Reef Marine Park Act. 1975.** Available at <http://www.gbrmpa.gov.au/about-us/legislation-regulations-and-policies/legislation>.
- Griffin JN, O’Gorman EJ, Emmerson MC, Jenkins SR, Klein A-M, Loreau M, Symstad A. 2009.** Biodiversity and the stability of ecosystem functioning. In: Naeem S, Bunker D, Hector A, Loreau M, Perrings C, eds. *Biodiversity, Ecosystem Functioning, and Human Wellbeing: An Ecological and Economic Perspective*. Oxford: Oxford University Press, 78–93.
- Hamann M, Chin A. 2015.** We’ve only monitored a fraction of the Great Barrier Reef’s species. *The Conversation*, 28 May 2015. Available at <http://theconversation.com/weve-only-monitored-a-fraction-of-the-barrier-reefs-species-39382>.
- Harris PT, Bridge TCL, Beaman RJ, Webster JM, Nichol SL, Brooke BP. 2013.** Submerged banks in the Great Barrier Reef, Australia, greatly increase available coral reef habitat. *ICES Journal of Marine Science* 70(2):284–293 DOI 10.1093/icesjms/fss165.
- Hooper DU, Adair EC, Cardinale BJ, Byrnes JE, Hungate BA, Matulich KL, Gonzalez A, Duffy JE, Gamfeldt L, O’Connor MI. 2012.** A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* 486(7401):105–108 DOI 10.1038/nature11118.

- Hooper JN, Van Soest RW. 2006.** A new species of Amphimedon (Porifera, Demospongiae, Haplosclerida, Niphatidae) from the Capricorn-Bunker group of Islands, Great Barrier Reef, Australia: target species for the 'sponge genome project'. *Zootaxa* **1314**:31–39.
- Hosie AM, Sampey A, Davie PJ, Jones DS. 2015.** Kimberley marine biota. Historical data: crustaceans. *Records of the Western Australian Museum, Supplement* **84(1)**:247–285
DOI [10.18195/issn.0313-122x.84.2015.247-285](https://doi.org/10.18195/issn.0313-122x.84.2015.247-285).
- Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H, Hobbs J-PA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Pandolfi JM, Pratchett M, Schoepf V, Torda G, Wilson SK. 2018.** Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science* **359(6371)**:80–83 DOI [10.1126/science.aan8048](https://doi.org/10.1126/science.aan8048).
- Hughes TP, Day JC, Brodie J. 2015.** Securing the future of the Great Barrier Reef. *Nature Climate Change* **5(6)**:508–511 DOI [10.1038/nclimate2604](https://doi.org/10.1038/nclimate2604).
- Hughes TP, Graham NA, Jackson JBC, Mumby PJ, Steneck RS. 2010.** Rising to the challenge of sustaining coral reef resilience. *Trends in Ecology & Evolution* **25(11)**:633–642
DOI [10.1016/j.tree.2010.07.011](https://doi.org/10.1016/j.tree.2010.07.011).
- Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, Babcock RC, Beger M, Bellwood DR, Berkelmans R, Bridge TC, Butler IR, Byrne M, Cantin NE, Comeau S, Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF, Gilmour JP, Harrison HB, Heron SF, Hoey AS, Hobbs J-PA, Hoogenboom MO, Kennedy EV, Kuo C-Y, Lough JM, Lowe RJ, Liu G, McCulloch MT, Malcolm HA, McWilliam MJ, Pandolfi JM, Pears RJ, Pratchett MS, Schoepf V, Simpson T, Skirving WJ, Sommer B, Torda G, Wachenfeld DR, Willis BL, Wilson SK. 2017.** Global warming and recurrent mass bleaching of corals. *Nature* **543(7645)**:373–377 DOI [10.1038/nature21707](https://doi.org/10.1038/nature21707).
- Huisman JM, Sampey A. 2014.** Kimberley marine biota. Historical data: marine plants. *Records of the Western Australian Museum, Supplement* **84(1)**:45–67
DOI [10.18195/issn.0313-122x.84.2014.045-067](https://doi.org/10.18195/issn.0313-122x.84.2014.045-067).
- Humphries CJ, Williams PH, Vane-Wright RI. 1995.** Measuring biodiversity value for conservation. *Annual Review of Ecology and Systematics* **26(1)**:93–111
DOI [10.1146/annurev.ecolsys.26.1.93](https://doi.org/10.1146/annurev.ecolsys.26.1.93).
- Hunter JA, Cribb TH. 2012.** A cryptic complex of species related to *Transversotrema licinum* Manter, 1970 from fishes of the Indo-West Pacific, including descriptions of ten new species of *Transversotrema* Witenberg, 1944 (Digenea: Transversotrematidae). *Zootaxa* **3176**:1–44.
- Hunter M, Westgate M, Barton P, Calhoun A, Pierson J, Tulloch A, Beger M, Branquinho C, Caro T, Gross J, Heino J, Lane P, Longo C, Martin K, McDowell WH, Mellin C, Salo H, Lindenmayer D. 2016.** Two roles for ecological surrogacy: indicator surrogates and management surrogates. *Ecological Indicators* **63**:121–125 DOI [10.1016/j.ecolind.2015.11.049](https://doi.org/10.1016/j.ecolind.2015.11.049).
- Hutchings P, Glasby C, Capa M, Sampey A. 2014.** Kimberley marine biota. Historical data: polychaetes (Annelida). *Records of the Western Australian Museum, Supplement* **84(1)**:133–159
DOI [10.18195/issn.0313-122x.84.2014.133-159](https://doi.org/10.18195/issn.0313-122x.84.2014.133-159).
- Hutchings P, Kingsford M, Hoegh-Guldberg O. 2008.** *The Great Barrier Reef: Biology, Environment and Management*. Collingwood: CSIRO Publishing.
- James A, Gaston KJ, Balmford A. 2001.** Can we afford to conserve biodiversity? *BioScience* **51(1)**:43–52 DOI [10.1641/0006-3568\(2001\)051\[0043:CWATCB\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0043:CWATCB]2.0.CO;2).
- IUCN. 2015.** *The IUCN red list of threatened species*. Version 2015-4. Available at <http://www.iucnredlist.org>.

- Legg CJ, Nagy L. 2006.** Why most conservation monitoring is, but need not be, a waste of time. *Journal of Environmental Management* **78**(2):194–199 DOI [10.1016/j.jenvman.2005.04.016](https://doi.org/10.1016/j.jenvman.2005.04.016).
- Lejzerowicz F, Esling P, Pillet LL, Wilding TA, Black KD, Pawlowski J. 2015.** High-throughput sequencing and morphology perform equally well for benthic monitoring of marine ecosystems. *Scientific Reports* **5**(1):13932 DOI [10.1038/srep13932](https://doi.org/10.1038/srep13932).
- López-Cabrera FJ, Puga-Bernabéu A, Webster JM, Beaman RJ. 2016.** Morphometric analysis of the submarine landslides in the central Great Barrier Reef margin, north-eastern Australia. *GEOGACETA* **60**:43–46.
- Loreau M, de Mazancourt C. 2013.** Biodiversity and ecosystem stability: a synthesis of underlying mechanisms. *Ecology Letters* **16**:106–115 DOI [10.1111/ele.12073](https://doi.org/10.1111/ele.12073).
- Lucas PHC, Webb T, Valentine PS, Marsh H. 1997.** The outstanding universal value of the Great Barrier Reef world heritage area. *Great Barrier Reef Marine Park Authority*. Available at <http://elibrary.gbrmpa.gov.au/jspui/handle/11017/301>.
- Marsh H, De'ath G, Gribble NA, Lane B. 2001.** Shark control records hindcast serious decline in dugong numbers off the urban coast of Queensland. *Great Barrier Reef Marine Park Authority, Townsville*. Available at <http://elibrary.gbrmpa.gov.au/jspui/handle/11017/350>.
- McCook LJ, Almany GR, Berumen ML, Day JC, Green AL, Jones GP, Leis JM, Planes S, Russ GR, Sale PF, Thorrold SR. 2009.** Management under uncertainty: guide-lines for incorporating connectivity into the protection of coral reefs. *Coral Reefs* **28**(2):353–366 DOI [10.1007/s00338-008-0463-7](https://doi.org/10.1007/s00338-008-0463-7).
- McKenzie LJ, Collier C, Waycott M. 2015.** Reef Rescue Marine Monitoring Program: Inshore seagrass. Annual report for the sampling period 1st July 2012–31st May 2013, TropWATER, James Cook University, Cairns. Available at <http://elibrary.gbrmpa.gov.au/jspui/handle/11017/2970>.
- McNeil MA, Webster JM, Beaman RJ, Graham TL. 2016.** New constraints on the spatial distribution and morphology of the *Halimeda* bioherms of the Great Barrier Reef, Australia. *Coral Reefs* **35**(4):1343–1355 DOI [10.1007/s00338-016-1492-2](https://doi.org/10.1007/s00338-016-1492-2).
- Miller TL, Downie AJ. 2009.** Morphological disparity despite genetic similarity; new species of *Lobosorchis* Miller & Cribb, 2005 (Digenea: Cryptogonimidae) from the Great Barrier Reef and the Maldives. *Zootaxa* **1992**:37–52.
- Miya M, Sato Y, Fukunaga T, Sado T, Poulsen JY, Sato K, Minamoto T, Yamamoto S, Yamanaka H, Araki H, Kondoh M, Iwasaki W. 2015.** Mifish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: detection of more than 230 subtropical marine species. *Royal Society Open Science* **2**(7):150088 DOI [10.1098/rsos.150088](https://doi.org/10.1098/rsos.150088).
- Moore G, Morrison S, Hutchins JB, Allen GR, Sampey A. 2015.** Kimberley marine biota. Historical data: fishes. *Records of the Western Australian Museum, Supplement* **84**(1):161–206 DOI [10.18195/issn.0313-122x.84.2014.161-206](https://doi.org/10.18195/issn.0313-122x.84.2014.161-206).
- Moore JL, Runge MC. 2012.** Combining structured decision making and value-of-information analyses to identify robust management strategies. *Conservation Biology* **26**(5):810–820 DOI [10.1111/j.1523-1739.2012.01907.x](https://doi.org/10.1111/j.1523-1739.2012.01907.x).
- Murray DC, Coghlan ML, Bunce M. 2015.** From benchtop to desktop: important considerations when designing amplicon sequencing workflows. *PLOS ONE* **10**(4):e0124671 DOI [10.1371/journal.pone.0124671](https://doi.org/10.1371/journal.pone.0124671).
- Myers R, Ottensmeyer A. 2005.** Extinction risk in marine species. In: Norse EA, Crowder LB, eds. *Marine Conservation Biology: The Science of Maintaining the Sea's Biodiversity*. Washington, D.C.: Island Press, 470.

- Nichols JD, Williams BK. 2006. Monitoring for conservation. *Trends in Ecology & Evolution* 21(12):668–673 DOI 10.1016/j.tree.2006.08.007.
- Nyström M, Graham NAJ, Lokrantz J, Norström AV. 2008. Capturing the cornerstones of coral reef resilience: linking theory to practice. *Coral Reefs* 27(4):795–809 DOI 10.1007/s00338-008-0426-z.
- Osborne K, Dolman AM, Burgess SC, Johns KA. 2011. Disturbance and the dynamics of coral cover on the Great Barrier Reef (1995–2009). *PLOS ONE* 6(3):e17516 DOI 10.1371/journal.pone.0017516.
- Pereira HM, Leadley PW, Proença V, Alkemade R, Scharlemann JPW, Fernandez-Manjarrés JF, Araújo MB, Balvanera P, Biggs R, Cheung WWL, Chini L, Cooper HD, Gilman EL, Guénette S, Hurtt GC, Huntington HP, Mace GM, Oberdorff T, Revenga C, Rodrigues P, Scholes RJ, Sumaila UR, Walpole M. 2010. Scenarios for global biodiversity in the 21st century. *Science* 330(6010):1496–1501 DOI 10.1126/science.1196624.
- Perrings C, Naeem S, Ahrestani FS, Bunker DE, Burkill P, Canziani G, Elmqvist T, Fuhrman JA, Jaksic FM, Kawabata ZI, Kinzig A, Mace GM, Mooney H, Prieur-Richard A-H, Tschirhart J, Weisser W. 2011. Ecosystem services, targets, and indicators for the conservation and sustainable use of biodiversity. *Frontiers in Ecology and the Environment* 9(9):512–520 DOI 10.1890/100212.
- Perry CT, Edinger EN, Kench PS, Murphy GN, Smithers SG, Steneck RS, Mumby PJ. 2012. Estimating rates of biologically driven coral reef framework production and erosion: a new census-based carbonate budget methodology and applications to the reefs of Bonaire. *Coral Reefs* 31(3):853–868 DOI 10.1007/s00338-012-0901-4.
- Perry CT, Smithers SG, Kench PS, Pears B. 2014. Impacts of Cyclone Yasi on nearshore, terrigenous sediment-dominated reefs of the central Great Barrier Reef, Australia. *Geomorphology* 222:92–105 DOI 10.1016/j.geomorph.2014.03.012.
- Pitcher CR, Doherty PJ, Arnold P, Hooper JNA, Gribble N. 2007. Seabed biodiversity on the continental shelf of the Great Barrier Reef World Heritage Area. *AIMS/CSIRO/QM/QDPI Final Report to CRC Reef Research*, 320. Available at <http://epubs.aims.gov.au/handle/11068/7785;jsessionid=1451EAA5BF1D7937A0F97D3796BA12AA>.
- Pocock C. 2002. Sense matters: aesthetic values of the Great Barrier Reef. *International Journal of Heritage Studies* 8(4):365–381 DOI 10.1080/1352725022000037191g.
- Pressey RL. 2004. Conservation planning and biodiversity: assembling the best data for the job. *Conservation Biology* 18(6):1677–1681 DOI 10.1111/j.1523-1739.2004.00434.x.
- Purcell SW, Polidoro BA, Hamel JF, Gamboa RU, Mercier A. 2014. The cost of being valuable: predictors of extinction risk in marine invertebrates exploited as luxury seafood. *Proceedings of the Royal Society B: Biological Sciences* 281(1781):20133296 DOI 10.1098/rspb.2013.3296.
- Richards ZT. 2013. A comparison of proxy performance in coral biodiversity monitoring. *Coral Reefs* 32(1):287–292 DOI 10.1007/s00338-012-0963-3.
- Richards ZT, Hobbs JPA. 2014. Predicting coral species richness: the effect of input variables, diversity and scale. *PLOS ONE* 9(1):e83965 DOI 10.1371/journal.pone.0083965.
- Richards Z, Sampey A, Marsh L. 2014. Kimberley marine biota. Historical data: scleractinian corals. *Records of the Western Australian Museum, Supplement* 84(1):111–132 DOI 10.18195/issn.0313-122x.84.2014.111-132.
- Rodrigues ASL, Brooks TM. 2007. Shortcuts for biodiversity conservation planning: the effectiveness of surrogates. *Annual Review of Ecology, Evolution, and Systematics* 38(1):713–737 DOI 10.1146/annurev.ecolsys.38.091206.095737.

- Roth CH, Addison J, Anthony K, Dale A, Eberhard R, Hobday A, Horner NJ, Jarvis D, Kroon K, Stone-Jovicich S, Walshe T. 2017. Reef 2050 Plan Review Options. Final Report Submitted to the Department of the Environment and Energy. Canberra: CSIRO.
- Ryan N, Richards Z, Hobbs JP. 2014. Optimal monitoring of coral biodiversity at Christmas Island. *Raffles Bulletin of Zoology* 30:399–405.
- Runge MC, Converse SJ, Lyons JE. 2011. Which uncertainty? Using expert elicitation and expected value of information to design an adaptive program. *Biological Conservation* 144(4):1214–1223 DOI 10.1016/j.biocon.2010.12.020.
- Sadovy de Mitcheson Y, Craig MT, Bertocini AA, Carpenter KE, Cheung WW, Choat JH, Cornish AS, Fennessy ST, Ferreira BP, Heemstra PC, Liu M, Myers RF, Pollard DA, Rhodes KL, Rocha LA, Russell BC, Samoily MA, Sanciangco J. 2013. Fishing groupers towards extinction: a global assessment of threats and extinction risks in a billion dollar fishery. *Fish and Fisheries* 14(2):119–136 DOI 10.1111/j.1467-2979.2011.00455.x.
- Sampey A, Marsh LM. 2015. Kimberley marine biota. Historical data: echinoderms. *Records of the Western Australian Museum, Supplement* 84(1):207–246 DOI 10.18195/issn.0313-122x.84.2015.207-246.
- Schmidt-Roach S, Lundgren P, Miller KJ, Gerlach G, Noreen AM, Andreakis N. 2013. Assessing hidden species diversity in the coral *Pocillopora damicornis* from Eastern Australia. *Coral Reefs* 32(1):161–172 DOI 10.1007/s00338-012-0959-z.
- Schmidt-Roach S, Miller KJ, Andreakis N. 2013. *Pocillopora aliciae*: a new species of scleractinian coral (Scleractinia, Pocilloporidae) from subtropical Eastern Australia. *Zootaxa* 3626(4):576–582 DOI 10.11646/zootaxa.3626.4.11.
- Shinzato C, Zayasu Y, Kanda M, Kawamitsu M, Satoh N, Yamashita H, Suzuki G. 2018. Using seawater to document coral-zooxanthella diversity: a new approach to coral reef monitoring using environmental DNA. *Frontiers in Marine Science* 5:28 DOI 10.3389/fmars.2018.00028.
- Sobtzick S, Hagihara R, Grech A, Marsh H. 2012. Aerial survey of the urban coast of Queensland to evaluate the response of the dugong population to the widespread effects of the extreme weather events of the summer of 2010–11, Final report to the Australian Marine Mammal Centre and the National Environmental Research Program. Townsville: James Cook University. Available at <http://www.nerptropical.edu.au/publication/project-12-technical-report-aerial-survey-urban-coast-queensland-evaluate-response>.
- Stachowicz JJ, Bruno JF, Duffy JE. 2007. Understanding the effects of marine biodiversity on communities and ecosystems. *Annual Review of Ecology, Evolution, and Systematics* 38(1):739–766 DOI 10.1146/annurev.ecolsys.38.091206.095659.
- Stat M, Huggett MJ, Bernasconi R, DiBattista J, Berry TE, Newman SJ, Harvey ES, Bunce M. 2017. Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in a tropical marine environment. *Scientific Reports* 7(1):12240 DOI 10.1038/s41598-017-12501-5.
- Stem C, Margoluis R, Salafsky N, Brown M. 2005. Monitoring and evaluation in conservation: a review of trends and approaches. *Conservation Biology* 19(2):295–309 DOI 10.1111/j.1523-1739.2005.00594.x.
- Stoeckl N, Hicks CC, Mills M, Fabricius K, Esparon M, Kroon F, Kaur K, Costanza R. 2011. The economic value of ecosystem services in the Great Barrier Reef: our state of knowledge. *Annals of the New York Academy of Sciences* 1219(1):113–133 DOI 10.1111/j.1749-6632.2010.05892.x.
- Sutcliffe PR, Hooper JN, Pitcher CR. 2010. The most common sponges on the Great Barrier Reef seabed, Australia, include species new to science (Phylum Porifera). *Zootaxa* 2616:1–30.

- Sutcliffe PR, Pitcher CR, Caley MJ, Possingham HP. 2012. Biological surrogacy in tropical seabed assemblages fails. *Ecological Applications* 22(6):1762–1771 DOI 10.1890/11-0990.1.
- Thomsen PF, Kielgast J, Iversen LL, Møller PR, Rasmussen M, Willerslev E. 2012. Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLOS ONE* 7(8):e41732 DOI 10.1371/journal.pone.0041732.
- Thomson SA, Pyle RL, Ah Yong ST, Alonso-Zarazaga M, Ammirati J, Araya JF, Ascher JS, Audisio TL, Azevedo-Santos VM, Bailly N, Baker WJ, Balke M, Barclay MVL, Barrett RL, Benine RC, Bickerstaf JRM, Bouchard P, Bour R, Bourgoïn T, Boyko CB, Breure ASH, Brothers DJ, Byng JW, Campbell D, Ceriaco LMP, Cernák I, Cerretti P, Chang CH, Cho S, Copus JM, Costello MJ, Cseh A, Csuzdi C, Culham A, D'Elia G, d'Udekem d'Acoz C, Daneliya ME, Dekker R, Dickinson EC, Dickinson TA, van Dijk PP, Dijkstra KDB, Dima B, Dmitriev DA, Duistermaat L, Dumbacher JP, Eiserhardt WL, Ekrem T, Evenhuis NL, Faille A, Fernández-Triana JL, Fiesler E, Fishbein M, Fordham BG, Freitas AVL, Friol NR, Fritz U, Frøslev T, Funk VA, Gaimari SD, Garbino GST, Garraffoni ARS, Geml J, Gill AC, Gray A, Grazziotin FG, Greenslade P, Gutiérrez EE, Harvey MS, Hazevoet CJ, He K, He X, Helfer S, Helgen KM, van Heteren AH, Garcia FH, Holstein N, Horváth MK, Hovenkamp PH, Hwang WS, Hyvönen J, Islam MB, Iverson JB, Ivie MA, Jaafar Z, Jackson MD, Jayat JP, Johnson NF, Kaiser H, Klitgård BB, Knapp DG, Kojima J, Kõljalg U, Kontschán J, Krell FT, Krisai-Greilhuber I, Kullander S, Latella L, Lattke JE, Lencioni V, Lewis GP, Lhano MG, Lujan NK, Luksenburg JA, Mariaux J, Marinho-Filho J, Marshall CJ, Mate JF, McDonough MM, Michel E, Miranda VFO, Mitroiu MD, Molinari J, Monks S, Moore AJ, Moratelli R, Murányi D, Nakano T, Nikolaeva S, Noyes J, Ohl N, Oleas NH, Orrell T, Pál Gergely B, Pape T, Papp V, Parenti KR, Patterson D, Pavlinov IT, Pine RH, Poczai P, Prado J, Prathapan D, Rabeler RK, Randall JE, Rheindt FE, Rhodin AGJ, Rodríguez SM, Rogers DC, Roque FdO, Rowe KC, Ruedas LA, Salazar-Bravo J, Salvador RB, Sangster G, Sarmiento CE, Schigel DS, Schmidt S, Schueler FW, Segers H, Snow N, Souza-Dias PGB, Stals R, Stenroos S, Stone RD, Sturm CF, Štys P, Teta P, Thomas DC, Timm RM, Tindall BJ, Todd JA, Triebel D, Valdecasas AG, Vizzini A, Vorontsova MS, de Vos JM, Wagner P, Watling L, Weakley A, Welter-Schultes F, Whitmore D, Wilding N, Will K, Williams J, Wilson K, Winston JE, Wüster W, Yanega D, Yeates DK, Zaher H, Zhang G, Zhang Z-Q, Zhou H-Z. 2018. Taxonomy based on science is necessary for global conservation. *PLOS Biology* 16(3):e2005075 DOI 10.1371/journal.pbio.2005075.
- Troudet J, Grandcolas P, Blin A, Vignes-Lebbe R, Legendre F. 2017. Taxonomic bias in biodiversity data and societal preferences. *Scientific Reports* 7(1):9132 DOI 10.1038/s41598-017-09084-6.
- United Nations Educational, Scientific and Cultural Organisation. 1981. Outstanding universal value of the Great Barrier Reef. Available at <http://whc.unesco.org/en/list/154>.
- Valentini A, Taberlet P, Miaud C, Civade R, Herder J, Thomsen PF, Bellemain E, Besnard A, Coissac E, Boyer F, Gaboriaud C, Jean P, Poulet N, Roset N, Copp GH, Geniez P, Pont D, Argillier C, Baudoin J-M, Peroux T, Crivelli AJ, Olivier A, Acqueberge M, Le Brun M, Møller PR, Willerslev E, Dejean T. 2016. Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Molecular Ecology* 25(4):929–942 DOI 10.1111/mec.13428.
- Vellend M, Lilley PL, Starzomski BM. 2008. Using subsets of species in biodiversity surveys. *Journal of Applied Ecology* 45(1):161–169 DOI 10.1111/j.1365-2664.2007.01413.x.
- Webb TJ, Mindel BL. 2015. Global patterns of extinction risk in marine and non-marine systems. *Current Biology* 25(4):506–511 DOI 10.1016/j.cub.2014.12.023.

- Willan RC, Bryce C, Slack-Smith SM. 2015.** Kimberley marine biota. Historical data: molluscs. *Records of the Western Australian Museum, Supplement* **84(1)**:287–343 DOI [10.18195/issn.0313-122x.84.2015.287-343](https://doi.org/10.18195/issn.0313-122x.84.2015.287-343).
- Wilson EO. 2016.** *Half-Earth: Our Planet's Fight for Life*. New York: WW Norton & Company.
- Wilson HB, Joseph LN, Moore AL, Possingham HP. 2011.** When should we save the most endangered species? *Ecology Letters* **14(9)**:886–890 DOI [10.1111/j.1461-0248.2011.01652.x](https://doi.org/10.1111/j.1461-0248.2011.01652.x).
- World Heritage Committee. 2013.** *Committee Decision 37 COM 7B.10*, Paragraph 8, UNESCO. Available at <http://whc.unesco.org/en/soc/1874>.
- World Heritage Committee. 2014.** *Committee Decision 38 COM 7B.63*, Paragraph 10, UNESCO. Available at <http://whc.unesco.org/en/soc/2867>.
- World Heritage Committee. 2015.** *Committee Decision 39 COM 7B.7*, UNESCO. Available at <http://whc.unesco.org/en/soc/3234>.
- Yoccoz NG, Nichols JD, Boulinier T. 2001.** Monitoring of biological diversity in space and time. *Trends in Ecology & Evolution* **16(8)**:446–453 DOI [10.1016/s0169-5347\(01\)02205-4](https://doi.org/10.1016/s0169-5347(01)02205-4).
- Yokomizo H, Coutts SR, Possingham HP. 2014.** Decision science for effective management of populations subject to stochasticity and imperfect knowledge. *Population Ecology* **56(1)**:41–53 DOI [10.1007/s10144-013-0421-2](https://doi.org/10.1007/s10144-013-0421-2).