




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



The impact of COVID-19 on cardiovascular health behaviors in student veterans



Matthew R. Canjar, Dylan L. Richard, Rebecca M. Kappus*

Cardiovascular Research Laboratory, Department of Health and Exercise Science, Appalachian State University, Boone, NC, USA

Received 6 October 2021; received in revised form 16 December 2021; accepted 17 December 2021

Handling Editor: P. Russo

Available online 22 December 2021

KEYWORDS

Student Veteran;
COVID-19;
Cardiovascular
disease;
Anxiety;
Sleep;
Stress

Abstract *Background and aims:* Pandemics have previously resulted in increased cardiovascular morbidity and mortality. It is unclear if the effects of the COVID-19 pandemic will be amplified in individuals at high risk for cardiovascular disease, such as military populations, resulting in augmented cardiovascular events in Veterans. The purpose of this study was to determine if traditional behavioral risk factors for cardiovascular disease are amplified due to the COVID-19 pandemic and if risk factors are more prevalent in Veterans compared to non-Veterans.

Methods and results: Thirty-two student Veterans and 46 non-Veteran students between the ages of 18 and 35 completed a Qualtrics self-report questionnaire assessing health behaviors, physical activity, and mental health both before and during COVID-19. Veterans displayed worse pre-COVID cardiovascular health behaviors such as poor sleep habits, greater use of tobacco, alcohol, and energy drinks, and lower values of social engagement compared to non-Veterans. Many health behaviors remained unchanged in student Veterans during the pandemic. The non-Veteran group exhibited augmentation of cardiovascular health behaviors during COVID-19, shown through the worsening sleep habits, increased anxiety, and reduced physical activity.

Conclusion: Student Veterans demonstrate heightened risk for cardiovascular disease based on the pre-COVID elevation of behavioral risk factors. These behavioral factors continued to remain elevated during the COVID-19 pandemic. Non-Veteran students displayed amplification of behavioral risk factors for cardiovascular disease due to the COVID-19 pandemic. These results highlight the need for resources and interventions for our student veterans and suggest long-term cardiovascular consequences for all students who suffered through the COVID-19 pandemic.

© 2021 The Italian Diabetes Society, the Italian Society for the Study of Atherosclerosis, the Italian Society of Human Nutrition and the Department of Clinical Medicine and Surgery, Federico II University. Published by Elsevier B.V. All rights reserved.

1. Introduction

Previous pandemics, such as the Spagnola pandemic in 1918, have been followed by an increase in cardiovascular

events and deaths from coronary heart disease [1–4]. This may be due to the development and amplification of cardiovascular disease (CVD) risk factors during the pandemic. The COVID-19 pandemic has altered lifestyles and habits, leading to decreased physical activity, poor diet, and increased depression, loneliness, and stress [5–8], which are established risk factors for CVD [9–11]. Not surprisingly, countries significantly affected by COVID-19 are already displaying increased cardiovascular mortality [12].

* Corresponding author. Appalachian State University, Department of Health and Exercise Science, 1179 State Farm Rd, Boone, NC 28608-2071, USA.

E-mail address: kappusrm@appstate.edu (R.M. Kappus).

The burden of COVID-19 may be augmented in those who suffer a heightened risk of CVD, such as the military population. It is well established that Veterans display inferior cardiovascular health and a higher prevalence of CVD risk factors such as hypertension, sleep disorders, binge drinking, and mental health issues that far outweigh the prevalence in the civilian population [13,14]. Currently, the majority of Veteran-based cardiovascular studies are completed by The Department of Veterans Affairs and focus on older individuals and those with severe post-traumatic stress disorder. It is possible that even young Veterans will display heightened risk factors for CVD prior to the negative cardiovascular effects of aging [13]. Understanding onset of CVD risk may assist in the development of preventative measures in this vulnerable population, however, young Veterans remain unstudied. Further, response to a stressor, such as a global pandemic, may provide insight into future CVD risk [10].

This study assessed health habits linked with CVD in student Veterans and their non-Veteran student counterparts before and during COVID-19, including sleeping habits, social engagement, physical activity, and use of alcohol, energy drinks, and tobacco [15–20]. The purpose of this study was to determine if individuals with increased risk of CVD, such as Veterans, were more affected by the COVID-19 pandemic compared to the civilian population. Our hypothesis was that young Veterans would be more affected by the COVID-19 pandemic, shown through decreased physical activity, poorer mental health, and inadequate sleep and nutritional habits, more so than the non-Veteran population.

2. Methods

Data were obtained via Qualtrics (Provo, Utah) self-report questionnaire and responses were collected between February and March 2021. This study was determined to be exempt by the Appalachian State University Institutional Review Board. All participants provided electronic informed consent and data was collected anonymously without recording IP addresses.

2.1. Study population

Participants were recruited amongst the student population of Appalachian State University and qualified for the study if between the ages of 18 and 35, a current university student, and either had never served time in the military (non-Veteran) or had prior active duty service for a minimum of 3 years (student Veteran). Exclusion criteria included a history of cardiovascular, pulmonary, metabolic, or renal disease.

2.2. Self-report questionnaire

Participants self-reported demographics such as age, height, and weight. Race and ethnicity were reported using US Census questions, with multiple responses allowed.

Primary outcome measures were assessed utilizing a Likert scale. Participants were asked to reflect upon health and habit status prior to March 2020, considered the “Pre-COVID” timepoint, and their current health and habit status (February/March 2021), considered the “Post-COVID” timepoint. Health habits, such as obtaining 6–8 h of sleep, trouble sleeping, and use of alcohol, energy drinks, and tobacco were scored from 1 (never) to 5 (always). Social engagement was assessed utilizing questions regarding campus involvement and close friends. Responses were scored from 1 (disagree) to 5 (agree). Participants were asked to self-report their physical activity level from 1 (rarely or never perform physical activity) to 5 (vigorous physical activity weekly) and respond “Yes” or “No” if they engaged in resistance training and aerobic training. Anxiety symptoms were assessed using a Generalized Anxiety Disorder Questionnaire (GAD-7). The GAD-7 is a validated survey [21] which utilizes 7 questions to measure anxiety; for example, how often students were bothered by feelings of anxiety, worry, or annoyance. Answers are self-rated ranging from “Not at all” to “Most days.”

2.3. Statistical analysis

Descriptive characteristics were compared between groups using an independent *t*-test. A repeated measures ANOVA was conducted to determine the main effects of COVID (time) and Veteran status (group), and interaction (time x group). Any significant findings were followed up with two-tailed independent and dependent *t* tests. Because of sex difference in alcohol consumption guidelines, a post-hoc ANCOVA was performed controlling for sex in reported alcohol use. The relationship between anxiety severity and veteran status was assessed utilizing chi-squared test. Statistical significance was set a priori at $p < 0.05$. All data are reported as mean \pm standard deviation. Data were analyzed using SPSS version 26 (IBM Corp, Armonk, NY, USA).

3. Results

Seventy-eight volunteers qualified for and completed the Qualtrics survey. Participant characteristics are described in Table 1. The student Veterans ($n = 32$) were comprised of 27 male and 5 females while the non-Veteran students ($n = 46$) consisted of 21 male and 25 female participants. The Veteran group performed an average of 5 years of active service and was significantly older, taller, and heavier than the non-Veteran group.

The health habits of Veteran and non-Veteran students are reported in Table 2. At pre-COVID the Veteran group reported more trouble sleeping and fewer hours of sleep compared to the non-Veteran group. The Veteran group saw no changes in their sleep habits due to COVID and continued to have worse sleep habits when compared to the non-Veteran group. The non-Veteran group displayed an increase in trouble sleeping and a decrease in hours of sleep during the pandemic. The Veteran group displayed higher alcohol use both pre- and post-COVID compared to

Table 1 Participant characteristics.

	Veterans(n = 32)	Non-Veterans(n = 46)	p-value
Male (%)	84	46	
Age (years)*	27 ± 3	21 ± 2	0.001
Height (cm)*	178.8 ± 7.7	171.6 ± 11.0	<0.001
Weight (kg)*	93.1 ± 19.5	72.4 ± 14.3	<0.001
BMI (kg/m ²)*	29.0 ± 5.1	24.5 ± 4.1	<0.001
White (%)	91	89	
Hispanic or Latino (%)	9	4	
Black or African American (%)	3	4	
Asian/Pacific Islander (%)	3	3	
Multiple races (%)	3	4	

*p < 0.05, significant differences between student Veterans and Non-Veterans. BMI, Body Mass Index.

non-Veterans (Cohen's d = 0.92). Neither group had changes in alcohol use during COVID. Because the non-Veteran group had more female respondents, we performed a post-hoc analysis and found that Veterans continued to have significantly higher alcohol use, both pre- and post-COVID, compared to non-Veterans even after controlling for sex. Finally, veterans reported more energy drink usage both pre- and post-COVID, and more tobacco use pre-COVID when compared to the non-Veteran group. The Veteran group showed a significant decrease in both energy drink and tobacco use during COVID while the non-Veteran group displayed no change.

Social engagement parameters are displayed in Table 3. There were no group differences in campus involvement or school stress pre- or post-COVID and both groups decreased campus involvement and increased school

stress during the pandemic. The Veteran group displayed less school excitement pre-COVID and both groups decreased school excitement during COVID. The Veteran group reported having a less enjoyable major both before and during COVID and neither group demonstrated changes in their major's enjoyability during COVID. Veterans reported fewer close friends pre-COVID when compared to the Non-Veterans and both groups reported a decrease in close friends during COVID.

Participants were asked about their general physical activity level prior to and during COVID. Both groups self-reported engaging in moderate physical activity weekly prior to COVID, and the non-Veteran group reduced their physical activity level to "light" during COVID (p = 0.007) while Veterans maintained their level of weekly physical activity (p = 0.195), resulting in group differences during COVID (p = 0.01). Further, participants were asked if they performed resistance and/or aerobic training prior to and during COVID. The Veteran and non-Veteran group reported similar engagement in resistance training pre-COVID (81% versus 63%, respectively), however, the non-Veteran group reported a significant decrease (p < 0.001) in resistance training during COVID with only 30% engaging in resistance training, while Veterans maintained training (72%), resulting in group differences post-COVID (p < 0.001). Both the Veteran and non-Veteran groups reported similar prevalence of aerobic training prior to COVID (78% versus 73%), and both decreased aerobic training during COVID (53% for both groups).

There were no group differences pre- or post-COVID for overall generalized anxiety scores (p = 0.09; data not shown), however the non-Veteran group experienced a significant increase in anxiety levels from pre- to post-COVID (p = 0.02) while Veterans had no change

Table 2 Health habits.

	Veterans		Non-Veterans		Time p-value	Interaction	Group
	Pre-COVID	Post-COVID	Pre-COVID	Post-COVID			
6–8 Hours of Sleep	3.1 ± 1.0*	3.0 ± 1.0*	3.9 ± 0.7	3.6 ± 1.0#	0.021	0.184	0.001
Trouble Sleeping	3.7 ± 0.9*	3.7 ± 0.8*	2.6 ± 0.9	3.2 ± 1.0#	0.004	0.009	0.001
Alcohol Use (>3 per sitting)	3.5 ± 1.0*	3.2 ± 1.1*	2.4 ± 1.2	2.4 ± 1.2	0.390	0.288	0.001
Energy Drink Use	3.2 ± 1.2*	2.8 ± 1.3*#	1.8 ± 1.1	1.7 ± 1.3	0.065	0.155	0.001
Tobacco Use	2.4 ± 1.4*	2.0 ± 1.4#	1.6 ± 1.1	1.8 ± 1.3	0.318	0.023	0.053

Scale: 1 = never; 2 = hardly ever; 3 = sometimes; 4 = often; 5 = always. *p < 0.05, significant differences between student Veterans and Non-Veteran students; #p < 0.05, significantly different from Pre-COVID; significant p values displayed in bold font.

Table 3 Social engagement.

	Veterans		Non-Veterans		Time p-value	Interaction	Group
	Pre-COVID	Post-COVID	Pre-COVID	Post-COVID			
Campus Involvement	3.0 ± 1.5	2.2 ± 1.4#	3.5 ± 1.3	2.8 ± 1.4#	0.001	0.851	0.044
School Stress	3.5 ± 1.0	4.0 ± 1.1#	3.5 ± 1.1	4.1 ± 1.2#	0.001	0.634	0.733
School Excitement	3.1 ± 1.4*	2.6 ± 1.5#	3.7 ± 1.2	2.7 ± 1.4#	0.001	0.110	0.184
Enjoyable Major	3.7 ± 1.3*	3.6 ± 1.4*	4.3 ± 0.9	4.1 ± 1.1	0.416	0.762	0.013
Close Friends	3.0 ± 1.4*	2.4 ± 1.4#	3.7 ± 1.2	3.0 ± 1.5#	0.001	0.926	0.024

Scale: 1 = disagree; 2 = disagree somewhat; 3 = neutral; 4 = agree somewhat; 5 = agree *p < 0.05, significant differences between student Veterans and Non-Veteran students; #p < 0.05, significantly different from Pre-COVID; significant p values displayed in bold font.

($p = 0.20$). There were no group differences pre- ($p = 0.09$) or post-COVID ($p = 0.42$). The breakdown of anxiety rating (none, mild, moderate, severe) is included in Table 4, highlighting that prior to COVID, a higher percentage of Veterans exhibited some form of anxiety compared to Non-Veterans (94% versus 78%, respectively), although this did not reach statistical significance, $X^2(1) = 0.28$, $p = 0.59$. Veterans were more likely to experience severe anxiety compared to non-Veterans (25% vs 8%) pre-COVID, $X^2(1) = 3.85$, $p = 0.04$.

4. Discussion

This study sought to determine if the COVID-19 pandemic led to a worsening of cardiovascular health behaviors in a student Veteran population more so than a civilian student population. The main findings of this study were 1) several cardiovascular health behaviors, including sleep habits, anxiety, and alcohol use were unaltered by the pandemic in the Veteran cohort; and 2) non-Veteran students appeared to be more affected by the COVID-19 pandemic, shown through the deterioration of cardiovascular health behaviors such as worsening sleep habits, amplified anxiety, and reduced physical activity, when compared to Veteran students. Furthermore, the results of this study highlight that Veterans, even at a young age, are reporting cardiovascular health behaviors associated with CVD, including poor sleep and health habits, elevated anxiety, and low levels of social engagement.

4.1. Sleep habits

Veterans reported higher rates of sleep trouble and fewer hours of sleep, which are associated with an increased risk of CVD such as coronary artery disease, heart failure, and hypertension [22]. Insufficient sleep can alter autonomic modulation, resulting in higher sympathetic and lower parasympathetic tone [23], which are linked to CVD [24]. Post-COVID, Veterans reported the same poor sleep habits as pre-COVID, while non-Veterans, although still reporting better sleep than Veterans, worsened sleep habits. Sleep plays a crucial role in supporting the immune system [25] and sleep loss has been shown to reduce natural killer cell activity, generate inflammatory cytokines, and increase risk of infection [26–28]. Further, because low sleep efficiency is associated with decreased immunity to the common cold [29], this relationship between sleep and immunity not only increases future CVD risk, but may make those with sleep disturbances more susceptible to COVID [30].

Table 4 Generalized anxiety disorder prevalence.

	Veterans		Non-Veterans	
	Pre-COVID	Post-COVID	Pre-COVID	Post-COVID
No anxiety (%)	6	9	22	9
Mild (%)	44	35	46	30
Moderate (%)	25	31	24	26
Severe (%)	25	25	8	35

4.2. Health habits

The Veterans in this study reported they were more likely to consume higher amounts of alcohol compared to non-Veterans, who hardly ever consumed 3 or more alcoholic drinks in one sitting. Although light to moderate alcohol consumption may reduce the risk of CVD, heavy drinkers (2 or more drinks per day for women; 3 or more drinks per day for men) have the highest risk of CVD [31,32]. Heavy drinking is associated with increased cardiovascular mortality, coronary heart disease, hypertension, cerebrovascular events, arrhythmias, and cardiomyopathy [18,33,34]. There were no changes in alcohol usage in either group with COVID, which is in contradiction to preliminary data demonstrating increased alcohol usage during COVID [35]. However, many students were forced to move out of dorms and in with parents/guardians, resulting in fewer social opportunities for drinking. Although there was not a decrease in drinking, the lack of increase in heavy drinking may be partially due to respondents' change in housing. Indeed, there was a reduction in student quantity of drinking during the COVID-19 pandemic due to residing with parents [36].

The Veteran group demonstrated higher energy drink usage, which has many negative cardiovascular implications to include sudden cardiac arrest, myocardial infarction, spontaneous coronary dissection, and increased systolic blood pressure [20]. Energy drink usage is widespread in military personnel and nearly half of deployed troops consume energy drinks daily [37]. Our findings highlight that energy drink usage remains amplified following deployment. Due to the relationship between energy drink use, sleep, and mental health problems [37], Veteran's energy drink usage may also contribute to increased mental health and sleep disorders, further amplifying CVD risk.

Tobacco use, a leading cause of premature death from CVD and the cause of coronary heart disease, cerebrovascular disease, aortic aneurysm, and heart failure [38,39], was more prevalent in the Veteran group pre-COVID compared to non-Veterans. This study's findings are similar to other studies showing increased tobacco use in Veterans compared to civilians [13,40]. It is unclear what caused the decreased use of tobacco products post-COVID; if it was preventative because of the increased risk of severe COVID-19 in smokers [41], due to financial concerns, or some unknown factor; however, it is worth noting that no amount of tobacco use is safe and even one cigarette per day leads to an increased risk of disease [42].

4.3. Social engagement

The decline in social engagement (reduced campus involvement, school excitement, and close friends, along with an increase in school stress) in both groups during the pandemic was expected due to social distancing guidelines and a shift from in-person to online learning [43]. In one study, 89% of university students reported difficulty in concentrating, 82% reported increased concerns about academic performance, and 86% reported

decreased social interactions [44]. The Veteran group reported lower levels of social engagement prior to COVID, shown through less school excitement, the lack of an enjoyable major, and less close friends. Social engagement has a protective effect against cardiovascular mortality [17] and social isolation and a lack of friends and community engagement is associated with an increased risk of CVD and stroke [45,46]. Older Veterans experience more loneliness and social isolation when compared to the general population [47,48] and the current study highlights that these feelings are present in young Veterans.

4.4. Exercise

Aerobic exercise training decreased during COVID in both groups, which is concerning not only due to the link between physical inactivity and CVD incidence and mortality [49,50], but also because exercise is beneficial in addressing risk factors highlighted in this study, including anxiety [51], tobacco use [52], and sleep [53].

4.5. Mental health

Although there were no overall group differences in general anxiety disorder scores, Veterans were more likely to experience severe anxiety pre-COVID. Previous studies have reported increased anxiety during COVID in university students [54] and support our findings of increased school stress and decreased school enjoyment and close friends, which are further linked with anxiety [44]. Poor mental health is associated with an increased risk of CVD, via several mechanisms including increased sympathetic nervous activity, increased blood pressure, and endothelial dysfunction [46]. Additionally, severe anxiety is associated with the development and progression of coronary artery disease and heart failure [55]. The higher frequency of anxiety, specifically severe anxiety, in the student Veteran cohort is in agreement with previous research demonstrating an increased prevalence of mental health disorders in Veterans [13].

Overall, the results of this study suggest that student Veterans have an increased risk of developing CVD, demonstrated through elevated risk factors not displayed in civilian students. These findings support previous literature showing Veterans display heightened risk factors for CVD when compared to their civilian counterparts [14,56], but further contributes that these risk factors are present in college-aged students. Contrary to our hypothesis, the Veteran group did not exhibit an increase in cardiovascular risk factors when compared to the civilian population during the COVID-19 pandemic. Although Veterans and non-Veterans were affected similarly by the pandemic in school stress, social engagement, and aerobic exercise, the non-Veteran group reported several significant changes in sleeping habits, overall anxiety, and resistance exercise training, which remained unchanged in Veterans. The worsening of these CVD risk factors suggest that non-Veteran students demonstrate an increased risk of CVD due to the COVID-19 pandemic.

The lack of effect of the pandemic on the Veteran group's risk factors may be due to a few reasons, the first being a blunted stress response. Veterans demonstrate blunted cortisol concentrations to a social stress test compared to civilians [57] and this finding is similar in individuals with post-traumatic stress disorder [58,59]. These attenuated stress responses may be due to an altered hypothalamic-pituitary-adrenal (HPA) axis and sympathetic nervous system. In individuals who experience chronic stress or those who have experienced trauma, there is a lack of responsiveness to stimuli that would otherwise stimulate the HPA axis and create a stress response, such as increased cortisol [60]. Veterans in the current study displayed a significantly higher prevalence of anxiety pre-COVID, especially severe anxiety, compared to their fellow civilian students. This amplified pre-stimulus (pre-COVID) risk factor may contribute to a chronic stress response. It is possible with the presence of already heightened risk factors, the pandemic may not have been able to evoke substantial changes, resulting in a blunted response.

Another possible explanation could be due to Veterans' maintenance of resistance training. During times of high stress and change, such as on deployment, Veterans display increases in muscle strength and power, reflective of resistance training, while decreasing aerobic performance [61]. It is possible that the continuation of resistance training during this time of stress may have helped to curb the impact of the COVID-19 pandemic and could explain the lack of increase in overall anxiety, as resistance exercise training is associated with improved anxiety levels and symptoms [15].

Finally, the Veteran group may have better psychological resilience. Psychological resilience is the ability to adapt well to a crisis, and traumatic events and major life stressors are required for the development of resilience [62]. Veterans are exposed to substantial traumatic events while deployed and reintegrating back into civilian life, as well as transitioning to college, results in significant stress [63]. Indeed, studies show that Veterans who are exposed to trauma display higher levels of psychological resilience and emotional stability [64,65]. Therefore, student Veterans may be better able to deal with a stressor such as a global pandemic more so than the non-Veteran population. While further studies are needed, psychological resilience may help explain why the Veteran group demonstrated little to no amplification of CVD risk factors during the COVID-19 pandemic.

4.6. Limitations

These are several limitations of the present study. The study's main findings were assessed via Qualtrics self-report questionnaire. Although widely used and valid, the responses could be reflective of bias in self-reporting [66]. Furthermore, health status and behaviors from pre-COVID were assessed approximately one year later and it is possible participants did not accurately recall health habits prior to COVID. The number of respondents was low, due

to the accessibility of the targeted population of student Veterans and the limited data collection period due to the changing pandemic. However, this is the first study to assess CVD risk factors in student Veterans and determine the effects of COVID on a young, student population, although future studies should utilize a larger group of participants. A final limitation was a significant difference in age between groups. Psychological resilience does have an age-related component in which older adults display increased levels of resilience when compared to young adults [67]. However, it is more likely any discrepancies in resilience between groups is due to the trauma endured during deployment and in transitioning back to civilian life, and further study is needed in this area.

In conclusion, this study highlights that young, otherwise healthy student Veterans are displaying behavioral risk factors for cardiovascular disease even prior to the detrimental effects of aging on cardiovascular health. Further, following a stressor such as a global pandemic (COVID-19), civilian students worsened their cardiovascular health behaviors more so than student Veterans. The lack of response in student Veterans due to COVID-19 may be due to the presence of amplified risk factors and chronic stressors prior to the pandemic, or their psychological resilience due to previously experienced trauma. Because limited data exists assessing the cardiovascular health of student Veterans, this knowledge may help to improve resources, both at the University and the government level, for our student Veterans. Further, it is important to understand the response to stressors, such as COVID-19, as this provides insight into future cardiovascular disease risk, not only in student Veterans, but also civilian students.

Source of support

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

The authors declared no conflicts of interest to disclose.

References

- [1] Azambuja MI. Spanish flu and early 20th-century expansion of a coronary heart disease-prone subpopulation. *Tex Heart Inst J* 2004; 31:14–21.
- [2] Myrskylä M, Mehta NK, Chang VW. Early life exposure to the 1918 influenza pandemic and old-age mortality by cause of death. *Am J Publ Health* 2013;103:e83–90.
- [3] Mazumder B, Almond D, Park K, Crimmins EM, Finch CE. Lingering prenatal effects of the 1918 influenza pandemic on cardiovascular disease. *J Dev Orig Health Dis* 2010;1:26–34.
- [4] Madjid M, Miller CC, Zarubaev VV, Marinich IG, Kiselev OI, Lobzin YV, et al. Influenza epidemics and acute respiratory disease activity are associated with a surge in autopsy-confirmed coronary heart disease death: results from 8 years of autopsies in 34,892 subjects. *Eur Heart J* 2007;28:1205–10.
- [5] Meyer J, McDowell C, Lansing J, Brower C, Smith L, Tully M, et al. Changes in physical activity and sedentary behavior in response to COVID-19 and their associations with mental health in 3052 US adults. *Int J Environ Res Publ Health* 2020;17.
- [6] Killgore WDS, Cloonan SA, Taylor EC, Dailey NS. Loneliness: a signature mental health concern in the era of COVID-19. *Psychiatr Res* 2020;290:113117.
- [7] Ettman CK, Abdalla SM, Cohen GH, Sampson L, Vivier PM, Galea S. Prevalence of depression symptoms in US adults before and during the COVID-19 pandemic. *JAMA Netw Open* 2020;3:e2019686.
- [8] Mattioli AV, Ballerini Puviani M, Nasi M, Farinetti A. COVID-19 pandemic: the effects of quarantine on cardiovascular risk. *Eur J Clin Nutr* 2020;74:852–5.
- [9] Tzoulaki I, Elliott P, Kontis V, Ezzati M. Worldwide exposures to cardiovascular risk factors and associated health effects: current knowledge and data gaps. *Circulation* 2016;133:2314–33.
- [10] Steptoe A, Kivimäki M. Stress and cardiovascular disease. *Nat Rev Cardiol* 2012;9:360–70.
- [11] Valtorta NK, Kanaan M, Gilbody S, Ronzi S, Hanratty B. Loneliness and social isolation as risk factors for coronary heart disease and stroke: systematic review and meta-analysis of longitudinal observational studies. *Heart* 2016;102:1009–16.
- [12] Brant LCC, Nascimento BR, Teixeira RA, Lopes M, Malta DC, Oliveira GMM, et al. Excess of cardiovascular deaths during the COVID-19 pandemic in Brazilian capital cities. *Heart* 2020;106:1898–905.
- [13] Meadows SO, Engel CC, Collins RL, Beckman RL, Cefalu M, Hawes-Dawson J, et al. 2015 Department of defense health related behaviors survey (HRBS). *Rand Health Q* 2018;8:5.
- [14] Shrestha A, Ho TE, Vie LL, Labarthe DR, Scheier LM, Lester PB, et al. Comparison of cardiovascular health between US army and civilians. *J Am Heart Assoc* 2019;8:e009056.
- [15] Gordon BR, McDowell CP, Lyons M, Herring MP. The effects of resistance exercise training on anxiety: a meta-analysis and meta-regression analysis of randomized controlled trials. *Sports Med* 2017;47:2521–32.
- [16] Nagai M, Hoshida S, Kario K. Sleep duration as a risk factor for cardiovascular disease— a review of the recent literature. *Curr Cardiol Rev* 2010;6:54–61.
- [17] Ramsay S, Ebrahim S, Whincup P, Papacosta O, Morris R, Lennon L, et al. Social engagement and the risk of cardiovascular disease mortality: results of a prospective population-based study of older men. *Ann Epidemiol* 2008;18:476–83.
- [18] Ronsley PE, Brien SE, Turner BJ, Mukamal KJ, Ghali WA. Association of alcohol consumption with selected cardiovascular disease outcomes: a systematic review and meta-analysis. *BMJ* 2011;342:d671.
- [19] Roy A, Rawal I, Jabbour S, Prabhakaran D. Tobacco and cardiovascular disease: a summary of evidence. In: Prabhakaran D, Anand S, Gaziano TA, Mbanya JC, Wu Y, Nugent R, editors. *Cardiovascular, respiratory, and related disorders*. 3rd. Washington (DC): The International Bank for Reconstruction and Development / The World Bank; 2017.
- [20] Shah SA, Szeto AH, Farewell R, Shek A, Fan D, Quach KN, et al. Impact of high volume energy drink consumption on electrocardiographic and blood pressure parameters: a randomized trial. *J Am Heart Assoc* 2019;8:e011318.
- [21] Lowe B, Decker O, Muller S, Brahler E, Schellberg D, Herzog W, et al. Validation and standardization of the generalized anxiety disorder screener (GAD-7) in the general population. *Med Care* 2008;46:266–74.
- [22] Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Sleep duration and all-cause mortality: a systematic review and meta-analysis of prospective studies. *Sleep* 2010;33:585–92.
- [23] Castro-Diehl C, Diez Roux AV, Redline S, Seaman T, McKinley P, Sloan R, et al. Sleep duration and quality in relation to autonomic nervous system measures: the multi-ethnic study of atherosclerosis (MESA). *Sleep* 2016;39:1927–40.
- [24] Curtis BM, O'Keefe Jr JH. Autonomic tone as a cardiovascular risk factor: the dangers of chronic fight or flight. *Mayo Clin Proc* 2002; 77:45–54.
- [25] Lange T, Dimitrov S, Fehm HL, Westermann J, Born J. Shift of monocyte function toward cellular immunity during sleep. *Arch Intern Med* 2006;166:1695–700.
- [26] Irwin M, Mascovich A, Gillin JC, Willoughby R, Pike J, Smith TL. Partial sleep deprivation reduces natural killer cell activity in humans. *Psychosom Med* 1994;56:493–8.

- [27] Imai K, Matsuyama S, Miyake S, Suga K, Nakachi K. Natural cytotoxic activity of peripheral-blood lymphocytes and cancer incidence: an 11-year follow-up study of a general population. *Lancet* 2000;356:1795–9.
- [28] Irwin MR, Wang M, Campomayor CO, Collado-Hidalgo A, Cole S. Sleep deprivation and activation of morning levels of cellular and genomic markers of inflammation. *Arch Intern Med* 2006;166:1756–62.
- [29] Cohen S, Doyle WJ, Alper CM, Janicki-Deverts D, Turner RB. Sleep habits and susceptibility to the common cold. *Arch Intern Med* 2009;169:62–7.
- [30] Kim H, Hegde S, LaFiura C, Raghavan M, Luong E, Cheng S, et al. COVID-19 illness in relation to sleep and burnout. *BMJ Nutr Prev Health* 2021;4:132–9.
- [31] Di Castelnuovo A, Costanzo S, Bagnardi V, Donati MB, Iacoviello L, de Gaetano G. Alcohol dosing and total mortality in men and women: an updated meta-analysis of 34 prospective studies. *Arch Intern Med* 2006;166:2437–45.
- [32] O’Keefe JH, Bybee KA, Lavie CJ. Alcohol and cardiovascular health: the razor-sharp double-edged sword. *J Am Coll Cardiol* 2007;50:1009–14.
- [33] Mukamal KJ, Tolstrup JS, Friberg J, Jensen G, Gronbaek M. Alcohol consumption and risk of atrial fibrillation in men and women: the Copenhagen City Heart Study. *Circulation* 2005;112:1736–42.
- [34] Fernandez-Sola J. Cardiovascular risks and benefits of moderate and heavy alcohol consumption. *Nat Rev Cardiol* 2015;12:576–87.
- [35] Pollard MS, Tucker JS, Green Jr HD. Changes in adult alcohol use and consequences during the COVID-19 pandemic in the US. *JAMA Netw Open* 2020;3:e2022942.
- [36] White HR, Stevens AK, Hayes K, Jackson KM. Changes in alcohol consumption among college students due to COVID-19: effects of campus closure and residential change. *J Stud Alcohol Drugs* 2020;81:725–30.
- [37] Centers for Disease C, Prevention. Energy drink consumption and its association with sleep problems among U.S. service members on a combat deployment - Afghanistan, 2010. *MMWR Morb Mortal Wkly Rep* 2012;61:895–8.
- [38] Rigotti NA, Clair C. Managing tobacco use: the neglected cardiovascular disease risk factor. *Eur Heart J* 2013;34:3259–67.
- [39] Roth GA, Forouzanfar MH, Moran AE, Barber R, Nguyen G, Feigin VL, et al. Demographic and epidemiologic drivers of global cardiovascular mortality. *N Engl J Med* 2015;372:1333–41.
- [40] Odani S, Agaku IT, Graffunder CM, Tynan MA, Armour BS. Tobacco product use among military Veterans - United States, 2010-2015. *MMWR Morb Mortal Wkly Rep* 2018;67:7–12.
- [41] Reddy RK, Charles WN, Sklavounos A, Dutt A, Seed PT, Khajuria A. The effect of smoking on COVID-19 severity: a systematic review and meta-analysis. *J Med Virol* 2021;93:1045–56.
- [42] Hackshaw A, Morris JK, Boniface S, Tang JL, Milenkovic D. Low cigarette consumption and risk of coronary heart disease and stroke: meta-analysis of 141 cohort studies in 55 study reports. *BMJ* 2018;360:j5855.
- [43] Wester ER, Walsh LL, Arango-Caro S, Callis-Duehl KL. Student engagement declines in STEM undergraduates during COVID-19-driven remote learning. *J Microbiol Biol Educ* 2021:22.
- [44] Son C, Hegde S, Smith A, Wang X, Sasangohar F. Effects of COVID-19 on college students’ mental health in the United States: interview survey study. *J Med Internet Res* 2020;22:e21279.
- [45] Kawachi I, Colditz GA, Ascherio A, Rimm EB, Giovannucci E, Stampfer MJ, et al. A prospective study of social networks in relation to total mortality and cardiovascular disease in men in the USA. *J Epidemiol Community Health* 1996;50:245–51.
- [46] Mensah GA, Collins PY. Understanding mental health for the prevention and control of cardiovascular diseases. *Glob Heart* 2015;10:221–4.
- [47] Kuwert P, Knaevelsrud C, Pietrzak RH. Loneliness among older Veterans in the United States: results from the national health and resilience in Veterans study. *Am J Geriatr Psychiatr* 2014;22:564–9.
- [48] Carr DC, Urena S, Taylor MG. Adjustment to widowhood and loneliness among older men: the influence of military service. *Gerontologist* 2018;58:1085–95.
- [49] Kohl 3rd HW. Physical activity and cardiovascular disease: evidence for a dose response. *Med Sci Sports Exerc* 2001;33: S472–83. discussion S93-S94.
- [50] Reddigan JI, Ardern CI, Riddell MC, Kuk JL. Relation of physical activity to cardiovascular disease mortality and the influence of cardiometabolic risk factors. *Am J Cardiol* 2011;108:1426–31.
- [51] Goodwin RD. Association between physical activity and mental disorders among adults in the United States. *Prev Med* 2003;36: 698–703.
- [52] Ussher M, Nunziata P, Cropley M, West R. Effect of a short bout of exercise on tobacco withdrawal symptoms and desire to smoke. *Psychopharmacology (Berl)* 2001;158:66–72.
- [53] Jurado-Fasoli L, De-la OA, Molina-Hidalgo C, Migueles JH, Castillo MJ, Amaro-Gahete FJ. Exercise training improves sleep quality: a randomized controlled trial. *Eur J Clin Invest* 2020;50: e13202.
- [54] Marelli S, Castelnuovo A, Somma A, Castronovo V, Mombelli S, Bottoni D, et al. Impact of COVID-19 lockdown on sleep quality in university students and administration staff. *J Neurol* 2021;268: 8–15.
- [55] Celano CM, Daunis DJ, Lokko HN, Campbell KA, Huffman JC. Anxiety disorders and cardiovascular disease. *Curr Psychiatr Rep* 2016;18:101.
- [56] Assari S. Veterans and risk of heart disease in the United States: a cohort with 20 years of follow up. *Int J Prev Med* 2014;5:703–9.
- [57] Pierce ME, Pritchard LM. Lower stress-reactive cortisol in female Veterans associated with military status but not PTSD. *Stress* 2016; 19:486–91.
- [58] Metz S, Duesenberg M, Hellmann-Regen J, Wolf OT, Roepke S, Otte C, et al. Blunted salivary cortisol response to psychosocial stress in women with posttraumatic stress disorder. *J Psychiatr Res* 2020;130:112–9.
- [59] Geraciotti Jr TD, Baker DG, Kasckow JW, Strawn JR, Jeffrey Mulchahey J, Dashevsky BA, et al. Effects of trauma-related audiovisual stimulation on cerebrospinal fluid norepinephrine and corticotropin-releasing hormone concentrations in post-traumatic stress disorder. *Psychoneuroendocrinology* 2008;33:416–24.
- [60] Tsigos C, Chrousos GP. Hypothalamic-pituitary-adrenal axis, neuroendocrine factors and stress. *J Psychosom Res* 2002;53: 865–71.
- [61] Lester ME, Knapik JJ, Catrambone D, Antczak A, Sharp MA, Burrell L, et al. Effect of a 13-month deployment to Iraq on physical fitness and body composition. *Mil Med* 2010;175:417–23.
- [62] Bonanno GA, Westphal M, Mancini AD. Resilience to loss and potential trauma. *Annu Rev Clin Psychol* 2011;7:511–35.
- [63] Reyes AT, Kearney CA, Isla K, Bryant R. Student Veterans’ construction and enactment of resilience: a constructivist grounded theory study. *J Psychiatr Ment Health Nurs* 2018;25:37–48.
- [64] Isaacs K, Mota NP, Tsai J, Harpaz-Rotem I, Cook JM, Kirwin PD, et al. Psychological resilience in U.S. military Veterans: a 2-year, nationally representative prospective cohort study. *J Psychiatr Res* 2017;84:301–9.
- [65] Pietrzak RH, Cook JM. Psychological resilience in older U.S. Veterans: results from the national health and resilience in Veterans study. *Depress Anxiety* 2013;30:432–43.
- [66] Devaux M, Sassi F. Social disparities in hazardous alcohol use: self-report bias may lead to incorrect estimates. *Eur J Publ Health* 2016;26:129–34.
- [67] Gooding PA, Hurst A, Johnson J, Tarrier N. Psychological resilience in young and older adults. *Int J Geriatr Psychiatr* 2012;27:262–70.