


Magnetic compression anastomosis using a double-balloon enteroscope for complete obstruction of Roux-en-Y hepaticojejunostomy anastomosis


Standard treatment for most post-cholecystectomy bile duct injuries is Roux-en-Y hepaticojejunostomy [1]. Conventional treatment of complete anastomotic obstruction after this procedure is revision surgery, which has high morbidity and mortality. Therefore, less invasive treatments are preferred, such as guidewire tapping, needle knife puncture, and magnetic compression anastomosis [2, 3].

A 72-year-old woman developed obstructive cholangitis 3 months after Roux-en-Y hepaticojejunostomy was performed for a post-cholecystectomy bile duct injury. Complete anastomotic obstruction was observed on percutaneous transhepatic cholangiography and the guidewire could not be passed through the anastomosis (► Fig. 1 a). We decided to apply magnetic compression anastomosis by using a double-balloon enteroscope (EN 580; Fujifilm, Tokyo, Japan) because the patient was at a high risk for surgery owing to comorbid diseases (► Fig. 2, ► Video 1).

First, the bowel site of the anastomosis was reached with the double-balloon enteroscope (► Fig. 1 b), which was then removed, leaving the overtube in place. The neodymium magnet (4×10 mm), which was held by a snare at the tip of the endoscope without the balloon, was introduced into the afferent jejunal limb through the overtube (► Fig. 3, ► Fig. 4 a). Simultaneously, the neodymium magnet (3×10 mm) was inserted into the hepatic side of the obstruction through the percutaneous transhepatic cholangiography catheter (► Fig. 4 a). After confirming that the magnets were aligned in the fluoroscopy, the magnets were released. After 8 days, the magnets were seen to be coupled on fluoroscopy (► Fig. 4 b) and were removed by re-entering with the double-balloon endoscope. In the same session, the percutaneous transhepatic cholangiography catheter was advanced through the new



► Fig. 1 Complete obstruction of the Roux-en-Y hepaticojejunostomy anastomosis (arrow). a Percutaneous transhepatic cholangiography view. b Double-balloon enteroscope view.



► Fig. 2 An illustration of magnetic compression anastomosis using a double-balloon enteroscope (asterisk) and percutaneous transhepatic cholangiography (arrow) performed for complete obstruction of Roux-en-Y hepaticojejunostomy anastomosis. The arrowhead represents the alignment of the magnets.



► **Video 1** Magnetic compression anastomosis using a double-balloon enteroscope for complete obstruction of Roux-en-Y hepaticojejunostomy anastomosis after bile duct injury due to laparoscopic cholecystectomy.



► **Fig. 3** Endoscopic view of the magnet brought to the anastomosis line by snare at the tip of the double-balloon enteroscope.



► **Fig. 4** Fluoroscopic images showing: **a** The magnets inserted percutaneously and endoscopically. **b** The coupled magnets.



► **Fig. 5** Successful recanalization is observed under percutaneous transhepatic cholangiogram.

anastomosis and passed into the intestine. The cholangiography 60 days later showed that the radiopaque material had passed into the jejunum very easily, and the drainage catheter was removed (► **Fig. 5**). There was no recurrence of the stenosis on clinical and radiological follow-ups at 1 year.

In conclusion, magnetic compression anastomosis using double-balloon endoscopy may be used as a viable alternative to surgery for the treatment of anastomotic complete obstruction after Roux-en-Y hepaticojejunostomy.

Endoscopy_UCTN_Code_TTT_1AR_2AK

Competing interests

The authors declare that they have no conflict of interest.

Funding

Kuzey Medical Company (Distributor of Fujifilm Endoscopy), Ankara, Turkey
N/A

The authors

Dilara Turan Gökçe **Bülent Ödemiş, Çağdaş Erdoğan**

Ankara City Hospital, Department of Gastroenterology, Ankara, Turkey

Corresponding author

Dilara Turan Gökçe, MD

Department of Gastroenterology, Ankara City Hospital, 06100, Çankaya, Ankara, Turkey
dilaraturan89@yahoo.com

References

- [1] Benkabbou A, Castaing D, Salloum C et al. Treatment of failed Roux-en-Y hepaticojejunostomy after post-cholecystectomy bile ducts injuries. *Surgery* 2013; 153: 95–102
- [2] Rhee K, Jang S, Lee D. Recanalization of completely obstructed bilioenteric anastomoses using a needle knife puncture. *Gastrointest Interv* 2013; 2: 68–71
- [3] Suyama K, Takamori H, Yamanouchi E et al. Recanalization of obstructed choledochojejunostomy using the magnet compression anastomosis technique. *Am J Gastroenterol* 2010; 105: 230–231

Bibliography

Endoscopy 2022; 54: E945–E947

DOI 10.1055/a-1841-5734

ISSN 0013-726X

published online 14.7.2022

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>