

## Research Article

# The Role of *Rhodomyrtus tomentosa* (Aiton) Hassk. Fruits in Downregulation of Mast Cells-Mediated Allergic Responses

Thanh Sang Vo <sup>1</sup>, Young-Sang Kim,<sup>2</sup> Dai-Nghiep Ngo,<sup>3</sup> and Dai-Hung Ngo <sup>4</sup>

<sup>1</sup>NTT Hi-Tech Institute, Nguyen Tat Thanh University, Ho Chi Minh City, Vietnam

<sup>2</sup>Department of Chemistry, Pukyong National University, Busan 608-737, Republic of Korea

<sup>3</sup>Faculty of Biology and Biotechnology, University of Science, Vietnam National University, Ho Chi Minh City, Vietnam

<sup>4</sup>Faculty of Natural Sciences, Thu Dau Mot University, Thu Dau Mot City, Binh Duong Province, Vietnam

Correspondence should be addressed to Thanh Sang Vo; [vtsang@ntt.edu.vn](mailto:vtsang@ntt.edu.vn) and Dai-Hung Ngo; [hungnd@tdmu.edu.vn](mailto:hungnd@tdmu.edu.vn)

Received 16 March 2019; Revised 29 April 2019; Accepted 26 May 2019; Published 9 June 2019

Academic Editor: Oscar Palomares

Copyright © 2019 Thanh Sang Vo et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Rhodomyrtus tomentosa*, a flowering plant of Myrtaceae family from southern and southeastern Asia, was known to possess a rich source of structurally diverse and various biological activities. In this study, the inhibitory effect of *R. tomentosa* fruit extract (RFE) on allergic responses in calcium ionophore A23187-activated RBL-2H3 mast cells was investigated. The result showed that RFE was able to inhibit mast cell degranulation via decreasing  $\beta$ -hexosaminidase release and intracellular  $\text{Ca}^{2+}$  elevation at the concentration of 400  $\mu\text{g/ml}$ . Moreover, the suppressive effects of RFE on the production of interleukin-1 $\beta$  (IL-1 $\beta$ ) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) were evidenced. In addition, RFE effectively scavenged DPPH radical and suppressed the reactive oxygen species generation in a dose-dependent manner. Notably, the pretreatment of RFE caused the downregulation of tyrosine kinase Fyn phospholipid enzyme phospholipase C $\gamma$  (PLC $\gamma$ ), extracellular-signal-regulated kinase (ERK), and nuclear factor kappa B (NF- $\kappa$ B) phosphorylation. These results indicated that RFE could be a promising inhibitor of allergic responses and may be developed as bioactive ingredient for prevention or treatment of allergic diseases.

## 1. Introduction

Allergy is a hypersensitivity disorder that relates to exaggerated reaction of the immune system to harmless environmental substances such as animal dander, house dust mites, foods, pollen, insects, and chemical agents [1]. Allergic rhinitis, asthma, and atopic eczema are among the most common allergic diseases [2]. The prevalence, severity, and complexity of these diseases in the population are rapidly rising. So far, various medicines such as antihistamine, mast cell stabilizers, and immune suppressors have been applied for ameliorating allergic symptoms and reducing the suffering of anaphylaxis [3]. However, these medicines were not available to cure allergic diseases completely and exhibited several side effects [4]. Therefore, the discovery of safe and efficient therapeutics derived from natural products for prevention and treatment of allergic diseases is necessary.

*Rhodomyrtus tomentosa* is a flowering plant that belongs to the family Myrtaceae and is native to southern and

southeastern Asia. It has been used in traditional Vietnamese, Chinese, and Malaysian medicine for a long time for treatment of diarrhea, dysentery, gynecopathy, stomachache, and wound-healing [5]. Moreover, *R. tomentosa* has been known to contain a rich source of structurally diverse and biologically active metabolites such as triterpenes, steroids, and phenolic compounds [6, 7]. In particular, various biological activities of *R. tomentosa* have been evaluated and reported recently [8]. Hence, it is considered as a potential source for exploring novel therapeutic agents. In the present study, the biological activity of *R. tomentosa* was further evaluated via investigating its inhibitory capacity on allergic responses *in vitro*.

Rat basophilic leukemia (RBL-2H3) cells display properties of mucosal-type mast cells. The activation of these cells leads to the release and generation of several inflammatory mediators [9]. Thus, RBL-2H3 cells have been commonly and successfully used in *in vitro* studies for screening antiallergic agents. Herein, RBL-2H3 cells were used as an *in vitro* model

for evaluation of antiallergic activity of *R. tomentosa* fruit extract.

## 2. Materials and Methods

**2.1. Materials.** *R. tomentosa* fruits were purchased from Duong Dong Town, Phu Quoc district, Kien Giang province. ELISA kits were purchased from R&D Systems (Minneapolis, MN, USA). The antibodies were purchased from Santa Cruz Biotechnology Inc. (Santa Cruz, CA, USA). All other reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA).

**2.2. Extraction.** *R. tomentosa* fruits were air-dried under shade and powdered using a grinder. The powder was soaked with ethanol 80% under the extract conditions of ratio (1/4, w/v), time (4 h), and temperature (60°C). The *R. tomentosa* fruit extract (RFE) was kept at 4°C for further investigation.

**2.3. Cell Culture.** RBL-2H3 cells (Korean Cell Line Bank, Seoul, Korea) were maintained in an incubator containing 5% CO<sub>2</sub> at 37°C. The culture medium contains DMEM, 10% heat-inactivated FBS, 2 mM L-glutamine, 10 mM HEPES, 100 U/ml of penicillin G, and 100 mg/ml of streptomycin.

**2.4. Cell Viability.** The cytotoxic effect of the extract on RBL-2H3 cells was examined by MTT assay as previously described [10]. The cells (1x10<sup>5</sup> cells/ml) were treated with the extract (100, 200, or 400 µg/ml) for 24 h before incubation with MTT solution (1 mg/ml, final concentration) for 4 h. The supernatant was then removed, and DMSO (100 µl) was added to solubilize the formed formazan salt. The absorbance was measured at 540 nm using a microplate reader (GENios® Tecan Austria GmbH, Austria). The cell viability was shown as a percentage compared to blank.

**2.5. Degranulation Assay.** RBL-2H3 cells (2 × 10<sup>5</sup> cells/ml) were pretreated with the extract (100, 200, or 400 µg/ml) for 24 h. The culture medium was replaced by Tyrode buffer before stimulation with calcium ionophore A23187 (1 µM) for 30 min at 37°C. The level of β-hexosaminidase release was measured as previously described [11]. The β-hexosaminidase releases were calculated as a percentage compared to control: release ratio (%) = (T – B)/(C – B) × 100, where B is blank group, C is control group, and T is the tested group.

**2.6. The Intracellular Ca<sup>2+</sup> Elevation Assay.** RBL-2H3 cells (5 × 10<sup>4</sup> cells/ml) were pretreated with the extract (400 µg/ml) for 24 h and subsequently incubated with Fura-3/AM (2 µM) for 60 min at 37°C. The culture medium was then replaced by Tyrode buffer before exposure to calcium ionophore A23187 (1 µM) for 5 min at 37°C. The Fura-3/AM fluorescence intensity was measured by a microplate reader (GENios Tecan, Austria, Grodigl/Salzburg, Austria) at 360 nm of excitation wavelength and 528 nm of emission wavelength [12].

**2.7. Measurement of Cytokine Production.** Different concentration of the extract (100, 200, or 400 µg/ml) was

added to the culture medium of RBL-2H3 cells (2 × 10<sup>5</sup> cells/ml) for 24 h. The culture medium was then removed and Tyrode buffer was replaced before stimulation with calcium ionophore A23187 (1 µM) for 1 h at 37°C. The supernatants were collected, and the amount of IL-1β or TNF-α was measured by ELISA kit.

**2.8. 1,1-Diphenyl-2-Picryl-Hydrazyl Assay.** The 1,1-diphenyl-2-picryl-hydrazyl (DPPH) assay was conducted as previously described [13]. Briefly, the mixture containing 100 µl of the extract (400 µg/ml) and 100 µl of DPPH solution was incubated in the dark at room temperature for 30 min. The absorbance was then measured at 490 nm by Genova Nano (Jenway, UK). The DPPH radical scavenging ability was determined following the following formula:

DPPH scavenging ability

$$= \left[ \frac{(\text{OD}_{\text{control}} - \text{OD}_{\text{sample}})}{\text{OD}_{\text{control}}} \right] \times 100\% \quad (1)$$

**2.9. Measurement of Reactive Oxygen Species Production (ROS).** The extract (400 µg/ml) was added to the culture medium of RBL-2H3 cells (1 × 10<sup>3</sup> cells/ml) for 24 h before incubation with dihydroethidium (2 µM) for 60 min at 37°C. The culture medium was then removed and Tyrode buffer was replaced before stimulation with calcium ionophore A23187 (1 µM) for 30 min at 37°C. Paraformaldehyde 3% was used to fix the cells and the fluorescence intensity was conducted under a fluorescence microscope (CTR 6000, Leica, Wetzlar, Germany).

**2.10. Western Blot Analysis.** The protein expression level was measured by Western blot method. Various doses of the extract (100, 200, or 400 µg/ml) were introduced to RBL-2H3 cells for 24 h prior to stimulation with calcium ionophore A23187 (1 µM) for 30 min at 37°C. The procedure of protein detection was performed as previously described [12]. The protein band was visualized using LAS3000® Luminescent image analyzer (Fujifilm Life Science, Tokyo, Japan).

**2.11. Statistical Analysis.** Statistical analysis was performed by using the analysis of variance (ANOVA) test of Statistical Package for the Social Sciences (SPSS). The statistical significance of differences among groups was analyzed using Duncan's multiple range tests, wherein p < 0.05 was considered significant.

## 3. Results and Discussion

**3.1. Effect of RFE on Mast Cell Degranulation.** Mast cells play an important role in the development of allergic diseases and inflammatory processes [14]. Activation of mast cells triggers a cascade of intracellular events, especially degranulation [15]. Mast cell degranulation is considered to be one of the critical steps in allergic responses, causing the elevation of intracellular Ca<sup>2+</sup> level and the subsequent




FIGURE 1: The inhibitory effect of RFE on mast cell degranulation in calcium ionophore A23187-activated RBL-2H3 cells ((a) and (b)) and cell viability (c). (a) The cells were treated with different concentrations of RFE for 24 h before stimulation with calcium ionophore A23187 for 30 min. The level of degranulation was measured via a spectrofluorometric assay. (b) The cells were pretreated with RFE for 24 h and incubated with Fura-3/AM for 1 h before exposure to calcium ionophore A23187 for 5 min. The Fura-3/AM fluorescence intensity was measured. (c) The cells were treated with various concentrations of RFE for 24 h. Cell viability was assessed by MTT method, and the results were expressed as percentage of surviving cells over blank cells (no addition of RFE). Each determination was made in three independent experiments, and the data are shown as means  $\pm$  SD. Different letters A–E indicate significant difference among groups at  $p < 0.05$  by Duncan’s multiple-range test. The same letter indicates that the difference between the means is not statistically significant. If two groups have different letters, they are significantly different from each other at  $p < 0.05$ .

release of various preformed mediators. These mediators are the origination of various pathophysiologic events in acute allergic responses [16]. Therefore, various antiallergic drugs have been developed so far, which are able to inhibit degranulation of mast cells. In this study, the inhibitory effect of RFE on mast degranulation was evaluated via measuring  $\beta$ -hexosaminidase release and intracellular  $Ca^{2+}$  elevation in the activated RBL-2H3 cells (Figure 1). It was observed that RFE was able to reduce  $\beta$ -hexosaminidase release to 79, 63, and 37% at the concentrations of 100, 200, and 400  $\mu\text{g/ml}$  (Figure 1(a)). Moreover, the increase in intracellular  $Ca^{2+}$  level induced by calcium ionophore A23187 was remarkably alleviated by RFE at concentration of 400  $\mu\text{g/ml}$  (Figure 1(b)).

In particular, the inhibitory effect of RFE on mast cell degranulation was not due to cytotoxicity (Figure 1(c)). Notably, RFE possessed the similar inhibitory activity as compared with *Smilax glabra* [17], *Morinda citrifolia* [18], and *grapeseed extract* [19]. Evidently, calcium ionophore A23187 induces mast cell degranulation by increasing cell-membrane permeability [20, 21]. Thus, antiallergic agents having a membrane-stabilizing action may be desirable such as disodium cromoglycate or sodium hydroxypropylchromate. As a result, RFE may be suggested to stabilize the lipid bilayer membrane, thus reducing the degranulation in RBL-2H3 mast cells. Indeed, various phytochemical components of medicinal plants such as saponins, glycosides, flavonoids,




FIGURE 2: The suppressive effect of RFE on cytokine productions in calcium ionophore A23187-activated RBL-2H3 mast cells. Cells were pretreated with various concentrations of RFE for 24 h before exposure to A23187 for 1 h. The production level of IL-1 $\beta$  (a) and TNF- $\alpha$  (b) was quantified in culture media using commercial ELISA kits. Each determination was made in three independent experiments, and the data are shown as means  $\pm$  SD. Different letters A–D indicate significant difference among groups ( $p < 0.05$ ) by Duncan's multiple-range test. The same letter indicates that the difference between the means is not statistically significant. If two groups have different letters, they are significantly different from each other at  $p < 0.05$ . If a group is similar to A and B or B and C, it was addressed by letters "AB" or "BC," respectively.

tannins, and phenolic compounds have been evidenced as mast cell stabilizing agents [22]. Meanwhile, RFE has been reported to contain various tannins and phenolic compounds that may contribute to stabilizing mast cell membrane and preventing degranulation [23].

**3.2. Effect of RFE on Cytokine Production.** The allergic reactions are also characterized by the production of various cytokines. The activated mast cells are well established as an important source of several cytokines such as tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-1 $\beta$  (IL-1 $\beta$ ), IL-4, IL-6, IL-8, and IL-13 [24]. The excessive production of these cytokines leads to the recruitment of inflammatory cells such as neutrophils and eosinophils, which increase the inflammatory responses [25]. Therefore, the decrease in cytokine production from the activated mast cells is one of the key indicators of the ameliorated allergic symptoms. Herein, the suppressive effect of RFE on IL-1 $\beta$  and TNF- $\alpha$  productions was investigated in RBL-2H3 mast cells. It was shown that the production levels of IL-1 $\beta$  and TNF- $\alpha$  were increased in the culture supernatants of A23187-exposed RBL-2H3 cells (Figure 2). The amount of IL-1 $\beta$  and TNF- $\alpha$  from the exposed cells was  $167 \pm 4.9$  and  $216 \pm 6.4$  pg/ml, respectively. Conversely, this increase was significantly reduced in a concentration-dependent manner by RFE pretreatment. At the concentration of  $400 \mu$ g/ml, RFE reduced IL-1 $\beta$  and TNF- $\alpha$  levels to  $93 \pm 5.7$  and  $80 \pm 6$  pg/ml, respectively.

**3.3. Radical Scavenging Activity of RFE.** A free radical is considered as a molecule that contains one or more unpaired electrons in its outermost atomic or molecular orbital. It is generated from endogenous sources such as intracellular autooxidation and inactivation of small molecules or from

exogenous sources such as tobacco smoke, certain pollutants, organic solvents, and pesticides [26]. The free radicals are recognized as agents involved in the pathogenesis of allergic diseases [27]. Meanwhile, the antioxidant agents from natural products have been proposed as an approach to reduce the allergic diseases [28]. Herein, RFE was determined to be effective in scavenging radicals (Figure 3). RFE was able to scavenge 85% DPPH radical at the concentration of  $400 \mu$ g/ml (Figure 3(a)). Moreover, the result from light microscope assay showed that the fluorescence density of ROS was markedly decreased in RFE-pretreated group as compared to A23187-stimulated group. It evidenced radicals as an inducer of  $Ca^{2+}$  elevation and subsequent degranulation in the mast cells [29]. Meanwhile, the blockade of ROS production by DPI or antioxidants such as (-)-epigallocatechin gallate suppressed mast cell degranulation [30, 31]. These results suggested that the antioxidant activity of RFE may contribute to the depression of mast cell degranulation.

**3.4. Effect of RFE on the Intracellular Signaling Molecules in the Activated RBL-2H3 Mast Cells.** Evidently, the allergic responses were also indicated by the activation of a cascade of intracellular signaling molecules such as Fyn, PLC $\gamma$ , MAPKs, and NF- $\kappa$ B [15, 32]. It was reported that the activation of Fyn and PLC $\gamma$  leads to microtubule polymerization, intracellular  $Ca^{2+}$  elevation, and subsequent mast cell degranulation [33, 34]. Meanwhile, the activation of NF- $\kappa$ B triggers gene expression and production [32]. Thus, the inhibition of mast cell degranulation and cytokine production may relate to the inactivation of these intracellular signaling molecules. As shown in Figure 4, the phosphorylation of Fyn, PLC $\gamma$ , ERK MAPK, and NF- $\kappa$ B was increased in the control group exposed to A23187 alone. Conversely, the RFE




FIGURE 3: The radical scavenging activity of RFE. (a) 100 µl of RFE (400 µg/ml) was mixed with 100 µl of DPPH solution for 30 min. The absorbance of the mixture was then measured at 490 nm. Each determination was made in three independent experiments, and the data are shown as means ± SD. Different letters A–C indicate significant difference among groups ( $p < 0.05$ ) by Duncan’s multiple-range test. If two groups have different letters, they are significantly different from each other at  $p < 0.05$ . (b) The cells were pretreated with RFE (400 µg/ml) for 24 h and incubated with dihydroethidium for 1 h before exposure to calcium ionophore A23187 for 30 min. The level of ROS production was monitored by a light microscope with 10× magnification. Each determination was made in three independent experiments.




FIGURE 4: The downregulative effect of RFE on the intracellular signaling molecules in the activated RBL-2H3 mast cells. (a) The protein expression level was assessed by Western blotting. β-Actin was used as internal controls. Each determination was made in three independent experiments. (b) Densitometrically calculated expression levels of the intracellular signaling molecules were given as a percentage compared to the control group.

pretreatment caused significant suppression on Fyn, PLCγ, ERK MAPK, and NF-κB phosphorylation at the concentration of 400 µg/ml. As a result, this suppressive effect of RFE may contribute to the inhibition of mast cell degranulation.

#### 4. Conclusion

In conclusion, this study has evidenced the inhibitory effect of *R. tomentosa* fruit extract on allergic responses in

A23187-activated RBL-2H3 mast cells. The inhibitory effect has been found due to decreasing β-hexosaminidase release and cytokine productions, suppressing ROS production, and downregulating the phosphorylation of tyrosine kinase Fyn and phospholipid enzyme phospholipase Cγ. Therefore, *R. tomentosa* fruits could offer an attractive strategy for the control of allergic diseases. However, further studies related to safety and efficacy need to be evaluated.

## Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

## Conflicts of Interest

The authors declare that there are no conflicts of interest.

## Acknowledgments

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under Grant no. 106-NN.02-2016.68.

## References

- [1] T.-S. Vo, D.-H. Ngo, and S.-K. Kim, "Marine algae as a potential pharmaceutical source for anti-allergic therapeutics," *Process Biochemistry*, vol. 47, no. 3, pp. 386–394, 2012.
- [2] R. Shokouhi Shoormasti, Z. Pourpak, M. R. Fazlollahi et al., "The prevalence of allergic rhinitis, allergic conjunctivitis, atopic dermatitis and asthma among adults of Tehran," *Iranian Journal of Public Health*, vol. 47, no. 11, pp. 1749–1755, 2018.
- [3] G. J. Molderings, B. Haenisch, S. Brettner et al., "Pharmacological treatment options for mast cell activation disease," *Naunyn-Schmiedeberg's Archives of Pharmacology*, vol. 389, no. 7, pp. 671–694, 2016.
- [4] H. Nagai, H. Teramachi, and T. Tuchiya, "Recent advances in the development of anti-allergic drugs," *Allergology International*, vol. 55, no. 1, pp. 35–42, 2006.
- [5] V. T. Sang, N. D. Hung, L. P. Uyen, N. D. Nghiep, and K. Se-Kwon, "Investigation of the biological activities of Phu Quoc Sim fruits *Rhodomyrtus tomentosa* (Aiton) Hass," *EurAsian Journal of BioSciences*, vol. 13, no. 1, pp. 49–55, 2019.
- [6] Y.-B. Zhang, W. Li, Z.-M. Zhang et al., "Two new triterpenoids from the roots of *rhodomyrtus tomentosa*," *Chemistry Letters*, vol. 45, no. 3, pp. 368–370, 2016.
- [7] L. Zhao, H. Liu, L. Wang, Z. Xu, H. Tan, and S. Qiu, "Rhodomyrtosone B, a membrane-targeting anti-MRSA natural acylphloroglucinol from *Rhodomyrtus tomentosa*," *Journal of Ethnopharmacology*, vol. 228, pp. 50–57, 2019.
- [8] T. Vo and D. Ngo, "The Health Beneficial Properties of *Rhodomyrtus tomentosa* as Potential Functional Food," *Biomolecules*, vol. 9, no. 2, p. 76, 2019.
- [9] E. Passante and N. Frankish, "The RBL-2H3 cell line: its provenance and suitability as a model for the mast cell," *Inflammation Research*, vol. 58, no. 11, pp. 737–745, 2009.
- [10] T.-S. Vo, D.-H. Ngo, Q. V. Ta, I. Wijesekara, C.-S. Kong, and S.-K. Kim, "Protective effect of chitin oligosaccharides against lipopolysaccharide-induced inflammatory response in BV-2 microglia," *Cellular Immunology*, vol. 277, no. 1-2, pp. 14–21, 2012.
- [11] T.-S. Vo, C.-S. Kong, and S.-K. Kim, "Inhibitory effects of chitooligosaccharides on degranulation and cytokine generation in rat basophilic leukemia RBL-2H3 cells," *Carbohydrate Polymers*, vol. 84, no. 1, pp. 649–655, 2011.
- [12] T.-S. Vo, D.-H. Ngo, and S.-K. Kim, "Gallic acid-grafted chitooligosaccharides suppress antigen-induced allergic reactions in RBL-2H3 mast cells," *European Journal of Pharmaceutical Sciences*, vol. 47, no. 2, pp. 527–533, 2012.
- [13] T. S. Vo, P. U. Le, and D. H. Ngo, "The increased gamma-aminobutyric acid content by optimizing fermentation conditions of bacteria from kimchi and investigation of its biological activities," *EurAsian Journal of BioSciences*, vol. 12, no. 2, pp. 369–376, 2018.
- [14] K. Amin, "The role of mast cells in allergic inflammation," *Respiratory Medicine*, vol. 106, no. 1, pp. 9–14, 2012.
- [15] A. M. Gilfillan and C. Tkaczyk, "Integrated signalling pathways for mast-cell activation," *Nature Reviews Immunology*, vol. 6, no. 3, pp. 218–230, 2006.
- [16] S. J. Galli, M. Tsai, and A. M. Piliponsky, "The development of allergic inflammation," *Nature*, vol. 454, no. 7203, pp. 445–454, 2008.
- [17] A. Itharat, K. Srikwan, S. Ruangnoo, and P. Thongdeeying, "Anti-allergic activities of *Smilax glabra* rhizome extracts and its isolated compounds," *Journal of the Medical Association of Thailand*, vol. 98, pp. S66–S74, 2015.
- [18] K. Murata, Y. Abe, K. Shinohara et al., "Anti-allergic activity of the *Morinda citrifolia* extract and its constituents," *Pharmacognosy Research*, vol. 6, no. 3, pp. 260–265, 2014.
- [19] B.-H. Chen, M.-H. Hung, J. Y.-F. Chen et al., "Anti-allergic activity of grapeseed extract (GSE) on RBL-2H3 mast cells," *Food Chemistry*, vol. 132, no. 2, pp. 968–974, 2012.
- [20] B. K. De and I. Friedberg, "Effect of ionophore A23187 on the membrane permeability in mouse fibroblasts," *Biochemical and Biophysical Research Communications*, vol. 178, no. 3, pp. 830–841, 1991.
- [21] M. G. Valle-Dorado, C. E. Santana-Gómez, S. A. Orozco-Suárez, and L. Rocha, "The mast cell stabilizer sodium cromoglycate reduces histamine release and status epilepticus-induced neuronal damage in the rat hippocampus," *Neuropharmacology*, vol. 92, pp. 49–55, 2015.
- [22] D. J. Taur and R. Y. Patil, "Mast cell stabilizing and antiallergic activity of *Abrus precatorius* in the management of asthma," *Asian Pacific Journal of Tropical Medicine*, vol. 4, no. 1, pp. 46–49, 2011.
- [23] H. Hazrulrizawati and S. S. Zeyohannes, "Rhodomyrtus tomentosa: a phytochemical and pharmacological review," *Asian Journal of Pharmaceutical and Clinical Research*, vol. 10, no. 1, pp. 1–7, 2017.
- [24] T. C. Theoharides, K.-D. Alysandratos, A. Angelidou et al., "Mast cells and inflammation," *Biochimica et Biophysica Acta*, vol. 1822, no. 1, pp. 21–33, 2012.
- [25] N. Sismanopoulos, D.-A. Delivanis, K.-D. Alysandratos et al., "Mast cells in allergic and inflammatory diseases," *Current Pharmaceutical Design*, vol. 18, no. 16, pp. 2261–2277, 2012.
- [26] S. Perrone, A. Santacroce, M. Longini, F. Proietti, F. Bazzini, and G. Buonocore, "The free radical diseases of prematurity: from cellular mechanisms to bedside," *Oxidative Medicine and Cellular Longevity*, vol. 2018, Article ID 7483062, 14 pages, 2018.
- [27] J. M. Matés, C. Pérez-Gómez, and M. Blanca, "Chemical and biological activity of free radical 'scavengers' in allergic diseases," *Clinica Chimica Acta*, vol. 296, no. 1-2, pp. 1–15, 2000.
- [28] H. Moreno-Macias and I. Romieu, "Effects of antioxidant supplements and nutrients on patients with asthma and allergies," *The Journal of Allergy and Clinical Immunology*, vol. 133, no. 5, pp. 1237–1244, 2014.
- [29] Y. Suzuki, T. Yoshimaru, T. Inoue, O. Niide, and C. Ra, "Role of oxidants in mast cell activation," *Chemical Immunology and Allergy*, vol. 87, pp. 32–42, 2005.

- [30] K. Yamashita, Y. Suzuki, T. Matsui et al., "Epigallocatechin gallate inhibits histamine release from rat basophilic leukemia (RBL-2H3) cells: role of tyrosine phosphorylation pathway," *Biochemical and Biophysical Research Communications*, vol. 274, no. 3, pp. 603–608, 2000.
- [31] T. Yoshimaru, Y. Suzuki, T. Matsui et al., "Blockade of superoxide generation prevents high-affinity immunoglobulin E receptor-mediated release of allergic mediators by rat mast cell line and human basophils," *Clinical & Experimental Allergy*, vol. 32, no. 4, pp. 612–618, 2002.
- [32] S. Klemm and J. Ruland, "Inflammatory signal transduction from the FcεRI to NF-κB," *Immunobiology*, vol. 211, no. 10, pp. 815–820, 2006.
- [33] S. Marin-Vardeaux, I. Pombo, B. Iannascoli et al., "Evidence of a role for Munc 18-2 and microtubules in mast cell granule exocytosis," *Journal of Cell Science*, vol. 116, no. 2, pp. 325–334, 2003.
- [34] K. Nishida, S. Yamasaki, Y. Ito et al., "FcεRI-mediated mast cell degranulation requires calcium-independent microtubule-dependent translocation of granules to the plasma membrane," *The Journal of Cell Biology*, vol. 170, no. 1, pp. 115–126, 2005.