




ORIGINAL ARTICLE

Prevalence of gastrointestinal parasites in three groups of domestic poultry managed under backyard system in the Savanna subregion, Department of Sucre, Colombia

Donicer Eduardo Montes-Vergara¹ , José Cardona-Alvarez² , Alexander Pérez-Cordero¹ 

¹Faculty of Agricultural Sciences, University of Sucre, Sincelejo, Colombia

²Department of Veterinary Medicine and Zootechnics, University of Córdoba, Montería, Colombia

ABSTRACT

Objective: To identify the prevalence of gastrointestinal parasites that affect the backyard poultry system in the Savanna region, Department of Sucre, Colombia.

Materials and Methods: Fecal samples were collected from 860 native birds, both hens (*Gallus domesticus*), ducks (*Anas platyrhynchos domesticus*), and turkeys (*Meleagris gallopavo*), regardless of age and sex. Samples were processed using direct techniques with ZnSO₄ and indirect methods such as modified Sloss. Data were presented as frequencies, and the nonparametric odds ratio test was used for two independent samples.

Results: A total of 77.3% (665/860) of the birds were infected with one or more species of gastrointestinal parasites. Among the nematodes, *Capillaria* spp. (45.6%), *Ascaridia galli* (18.4%), *Heterakis gallinarum* (59.4%), *Syngamus trachea* (38.9%), *Tetrameres* spp. (25.2%), and *Strongylus* spp. (12.2%) were recorded. The cestodes were *Choanotaenia infundibulum* (22.6%), *Davainea proglottina* (42.3%), *Raillietina* spp. (58.3%), and *Hymenolepis* spp. (54.7%), while only *Eimeria* spp. (90%) was recorded as protozoa.

Conclusions: The study showed a high incidence of gastrointestinal parasite infestations, the most common species being *Hymenolepis* spp., *Eimeria* spp., *Raillietina* spp., and *Heterakis gallinarum*.

ARTICLE HISTORY

Received August 06, 2021

Revised September 08, 2021

Accepted September 16, 2021

Published November 01, 2021

KEYWORDS

Gastrointestinal parasites; prevalence; backyard poultry; *Ascaridia*; *Heterakis*; *Capillaria*; cestodes



© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

Introduction

Backyard or rustic “family poultry farming” is a form of traditional domestic breeding that requires few inputs and includes a variety of bird species such as chickens, turkeys, ducks, geese, and quails [1], and is the most traditional and widespread livestock activity in rural communities, as it benefits rural families by providing high-nutrient products such as meat and eggs, as well as revenue from surpluses [2]. The poultry population in Colombia is distributed among 440,381 farms, of which 98.7% are backyard farms, and the remaining 1.3% correspond to technological farms. Of the total number of birds in Colombia (210,541,160), 95.8% are technified poultry and 4.2% are backyard poultry. The backyard poultry population in 2021 was more than 8 million,

of which 856,764 were registered in the department of Sucre, Colombia [3].

Backyard poultry farming is an important economic activity for the rural population as a source of income and as a way to guarantee food security in unprotected communities [4]. In this sense, support for backyard poultry farming has been widely used since it is considered that small-scale livestock production represents an effective alternative to achieving food security [5]. Farm animals constitute an essential element of subsistence for the rural poor, performing multiple functions: food production, fertilizer, and income generation [6].

Birds managed in BPS are vulnerable to parasite attacks [7]. The occurrence of parasitic infestations has a high

Correspondence Donicer Eduardo Montes-Vergara ✉ donicer.montes@unisucra.edu.co 📧 Faculty of Agricultural Sciences, University of Sucre, Sincelejo, Colombia.

How to cite: Montes-Vergara DE, Cardona-Alvarez J, Pérez-Cordero A. Prevalence of gastrointestinal parasites in three groups of domestic poultry managed under the backyard system in the Savanna subregion, Department of Sucre, Colombia. J Adv Vet Anim Res 2021; 8(4):606–611.

prevalence, causing low economic conditions, increased mortality and prophylaxis, leading to low production, death of animals, and limited productivity. Improved poultry management practices are responsible for the reduction in the incidence of parasitic infections [8]. Thus, the objective of this work was to identify the prevalence of gastrointestinal parasites in *Gallus domesticus*, *Anas platyrhynchos domesticus*, and *Meleagris gallopavo* managed under the backyard poultry system in the Savanna subregion, Department of Sucre, Colombia.

Materials and Methods

The study was conducted in the Savanna subregion (Sincedejo, Sincé, El Roble, San Pedro, Sampués, Los Palmitos, Galeras, Buenavista, Corozal and San Juan de Betulia) of the Department of Sucre, Colombia (Fig. 1) between January 2019 and March 2020. The climate of the study area is characteristic of tropical dry forest (bs-t) zones, with an average annual temperature of 27.2°C,

yearly rainfall between 990 and 1,275 mm, and relative humidity of 80% [9]. The region is characterized by commercial poultry farms and numerous backyard poultry operations.

The Savannas subregion in the Department of Sucre, Colombia, includes 10 municipalities and has a backyard poultry census of 42,624, distributed among 794 farms [3]. The sample size was determined according to Suarez and Tapia [10], using a confidence level of 95% ($Z = 1.96$) and a maximum margin of error of 10%, resulting in a total of 85.6 farms. Thus, an average of 8.6 farms per municipality was visited, using stratified random sampling, in which 860 adult criollo birds were characterized, considering environmental, sanitary, and management factors.

We worked with four geographical groups, which were characterized by their relative proximity (Group 1 = Sincedejo, Los Palmitos, Corozal; Group 2 = Sincé, Galeras, S-Betulia; Group 3 = El Roble, Sampués; Group 4 = San Pedro, Buenavista), in order to establish prevalence levels by geographical group.



Figure 1. Location of the study area. Savanna sub-region, Department of Sucre Colombia. Wikipedia, the free encyclopedia. Accessed on: November 1, 2021, from [https://es.wikipedia.org/w/index.php?title=Subregi%C3%B3n_de_La_Sabana_\(Sucre\)&oldid=135872197](https://es.wikipedia.org/w/index.php?title=Subregi%C3%B3n_de_La_Sabana_(Sucre)&oldid=135872197).

A cross-sectional coprological study of feces from hens (*G. domesticus*), ducks (*A. platyrhynchos domesticus*), and turkeys (*M. gallopavos*) was carried out. The samples were taken in the morning hour before feeding. The animals evaluated on each farm were placed in a portable cage according to the group under study. Black plastic was placed on the floor at the bottom of the cage to avoid feces contamination with the ground. Once the birds had deposited their feces, a swab was taken, and the sample was collected in a sterile fecal matter collection container. The sample was kept cold at 5°C for subsequent analysis in the Microbiology I laboratory of the University of Sucre.

A total of 860 fecal samples were processed and analyzed using direct techniques with ZnSO₄ and indirect methods such as modified Sloss [11], quantified with MacMaster chamber [8,12]. To calculate the frequency of gastrointestinal parasite infection, descriptive statistics using absolute and relative frequency were used. The nonparametric odds ratio test for two independent samples was used to determine the risk factors with a significance level of 5% in each geographical group. The variables were subjected to the Shapiro–Wilk test as a normality distribution test ($p < 0.05$). The presence of parasites by the geographic group was compared using the Kruskal–Wallis test for nonparametric variables and Dunn’s test to determine differences between groups [13]. Statistical analysis was carried out with the R program.

Results and Discussion

A total of 77.3% (665/860) of the samples were positive for gastrointestinal parasites, with one protozoan species, four cestodes species, and six nematodes species. The latter were *Capillaria* ssp. (45.6%), *Ascaridia galli* (18.4%), *Heterakis gallinarum* (59.4%), *Syngamus trachea* (38.9%), *Tetrameres* spp. (25.2%), and *Strongylus* (12.2%). The cestodes corresponded to *Choanotaenia infundibulum* (22.6%), *Davainea proglottina* (42.3%), *Raillietina* spp. (58.3%), and *Hymenolepis* spp. (54.7%). The protozoan recorded was *Eimeria* spp. (90%) (Table 1).

The high prevalence rate could be due to the fact that birds are kept in absolute freedom, feeding on many agricultural byproducts, pastures, and intermediate hosts of parasites (beetle, grasshoppers, cockroaches, crustaceans, earthworm, and snail) available in the environment [11,14].

The prevalence of gastrointestinal parasite infections varies between regions and countries [15]. For example, prevalences between 97.6% and 99.2% were reported in Germany [16], 99.3% in Italy [17], 72% in Iran [18,19], 73.9% in Thailand [20], and 95.8% in Zambia [21], with

Table 1. Frequency of gastrointestinal parasites in domestic poultry ($n = 860$) managed under the backyard system in the Savanna subregion, Department of Sucre, Colombia.

GP	Species	Frequency (%)	IC 95%
Protozoa	<i>Eimeria</i> spp.	90.0	82.5–94.4
	<i>C. infundibulum</i>	22.6	15.5–31.7
	<i>D. proglottina</i>	42.3	33.0–52.0
Cestodes	<i>Raillietina</i> spp.	58.3	48.5–67.4
	<i>Hymenolepis</i> spp.	54.7	44.9–64.1
	<i>Capillaria</i> ssp.	45.6	36.1–55.3
	<i>A. galli</i>	18.4	12.0–27.1
Nematodes	<i>H. gallinarum</i>	59.4	49.6–68.5
	<i>S. trachea</i>	38.9	29.9–48.6
	<i>Tetrameres</i> spp.	25.2	17.7–34.5
	<i>Strongylus</i> spp.	12.2	7.14–20.0

GP = parasitic group.

environmental conditions, age of birds, and availability of intermediate hosts being the main determinants for the variability of parasite prevalence [22].

The most prevalent parasite in the three groups was *Eimeria* spp, with a prevalences of 87.4%, 91.2%, and 90.2% in hens, ducks, and turkeys, respectively. The frequency of parasites according to avian species is shown in Table 2. The high prevalence of *Eimeria* spp. is a product of poor sanitation and poor management of the poultry habitat. Studies in Nigeria report the presence of *Eimeria* spp. as the most prevalent protozoan among gastrointestinal parasites infecting poultry [23–25].

Raillietina spp. was the most prevalent among the cestodes in the studied population; however, it was not observed in turkeys. *Hymenolepis* spp. predominated (Table 2) the most common type of parasite infecting domestic fowl [26–29].

The prevalence of nematodes ranged from 10.2% to 68.6% in the study birds (Table 2). Reports of nematode prevalence in domestic poultry managed under the SAT scheme have been reported [30–33] in various countries. Among the nematodes encountered, *H. gallinarum* is not pathogenic. Still, it is a vector of *Histomonas meleagridis*, which is highly pathogenic and involved in “black head” disease, which is lethal to many domestic fowl [34].

Nematode parasitism changes the gastrointestinal system of chickens, resulting in decreased performance and, in some cases, mortality [35,33]. While there is no information on the direct economic costs of nematode infestation in the Savanna, Sucre sub-region, the prevalence discovered warrants the use of mitigation techniques to offset the detrimental impact on SAT productivity.

Table 2. Frequency of gastrointestinal parasites in domestic poultry managed in the backyard systems in the Savanna subregion, Department of Sucre, Colombia.

Poultry	Samples		Parasites		Frequency (%)			
	n	Positive	Groups	Species				
Hens	535	414	Protozoa	<i>Eimeria</i> spp.	87.4			
				Cestodes	<i>Raillietina</i> spp.	45.5		
			<i>Hymenolepis</i> spp.		28.3			
			<i>D. proglottina</i>		55.1			
			Nematodes		<i>Capillaria</i> spp.	31.2		
				<i>A. galli</i>	26.8			
				<i>H. gallinarum</i>	39.6			
				<i>S. trachea</i>	24.2			
			Ducks	179	138	Protozoa	<i>Eimeria</i> spp.	91.2
							Cestodes	<i>Raillietina</i> spp.
<i>Hymenolepis</i> spp.	26.7							
Nematodes	<i>Capillaria</i> ssp.	22.3						
	<i>A. galli</i>	36.2						
	<i>H. gallinarum</i>	28.4						
	<i>S. trachea</i>	38.2						
Turkeys	146	113				Protozoa	<i>Eimeria</i> spp.	90.2
							Cestodes	<i>C. infundibulum</i>
						<i>Hymenolepis</i> spp.		76.1
			Nematodes	<i>Capillaria</i> ssp.	68.2			
				<i>A. galli</i>	10.2			
				<i>H. gallinarum</i>	68.6			
				<i>S. trachea</i>	45.9			
						<i>Tetrameres</i> spp.	25.2	
			<i>Strongylus</i> spp.	12.3				
Total	860	665						

The high prevalence rate in the study area may be a result of poor sanitary conditions, high poultry population density, uncontrolled feeding, and a lack of attention to treatment and disease control and prevention measures, all of which expose birds to poor hygiene on farms and in poultry houses, allowing them to contract a wide variety of harmful parasites [36].

Chemical control of parasites is simple, inexpensive, and can be used both therapeutically and prophylactically. However, chemical treatment has several drawbacks, such as weakening natural immunity and the presence of residues in food and the environment. In addition, chemical anthelmintics can stimulate resistance, so alternative forms of control are needed [8].

Table 3. Distribution and prevalence of gastrointestinal parasites according to geographic location in the Savanna subregion, Department of Sucre, Colombia.

Grups	Municipality	Total	Positive		OR	IC -95%	p-value
			%	%			
1	Sincelejo	85	72.9	68.2	0.87	0.66–1.14	0.34
	Los Palmitos	90	64.4				
	Corozal	92	67.4				
2	Sincé	90	72.2	78.1	0.97	0.71–1.33	0.93
	Galeras	83	88.0				
	SJ de Betulia	73	74.0				
3	El Roble	82	84.1	82.1	0.97	0.71–1.33	0.93
	Sampués	87	81.6				
4	San Pedro	94	84.0	84.8	0.97	0.71–1.33	0.93
	Buenavista	84	85.7				
Total		860	77.3				

The prevalence of gastrointestinal parasites was high in all the study municipalities, regardless of geographic location (Table 3), without significant differences. Education and motivation of farmer producers on biosecurity measures may aid in mitigating the negative effects of parasitic infection on poultry response effectiveness [33,37].

Conclusion

Gastrointestinal parasites are endemic among the domestic poultry managed under the backyard system in the Savanna region, Sucre, Colombia, showing a high prevalence of gastrointestinal parasites, the most frequent being *Hymenolepis* spp., *Eimeria* spp., *Raillietina* spp., and *H. gallinarum*, in the period studied. It is essential to know the conditions of the farm to develop the best prevention program, allowing the recognition of the factors that influence the possibility of disease incidence.

List of abbreviations

BPS, backyard poultry system; GP, parasitic group; IC, confidence interval; *n*, sample size; OR, Odds ratio; *p*, probability value; spp., species.

Acknowledgments

The authors thank the farmers interviewed for their cooperation, willingness, and access to backyard poultry systems and also the students of the Animal Breeding 2020 course at the University of Sucre for their support in the field.

Conflict of interest

The authors declare that they have no conflict of interest

Authors' contributions

MVDE: research, data analysis, writing of original draft.

CAJ: research, conceptualization, writing, revision, and editing

PCA: data analysis, review, and editing.

All authors made an effort to edit the final draft of the manuscript.

References

- [1] Centeno-Bautista S, López-Díaz C, Juárez-Estrada M. Producción avícola familiar en una comunidad del Municipio de Ixtacamaxitlán, Puebla-México. *Téc Pecu Méx* 2007; 45(1):41–60.
- [2] Toapanta M, Avilés-Esquivel DF, Montero-Recalde M, Pomboza P. Caracterización del sistema de producción de aves de traspatio del cantón Cevallos. *Actas Iberoam Conserv Anim* 2019; 13:1–5.
- [3] Colombia, Instituto Colombiano Agropecuario (ICA). Censo agropecuario, Bogotá, Distrito Capital-Colombia, América del Sur; 2021; <https://www.ica.gov.co/areas/pecuaria> (Accessed on May 01, 2021)
- [4] Luka SA, Ndams IS. Gastrointestinal parasites of domestic chicken *Gallus gallus domesticus* Linnaeus 1758 in Samary, Zaria Nigeria. *Sci World J* 2007; 2(1):27–30; <https://doi.org/10.4314/swj.v2i1.51723>
- [5] Ara I, Khan H, Syed T, Bhat B. Prevalence and seasonal dynamics of gastrointestinal nematodes of domestic fowls (*Gallus gallus domesticus*) in Kashmir, India. *J Adv Vet Anim Res* 2021; 8(3):448–53; <http://doi.org/10.5455/javar.2021.h533>
- [6] Rodrigues Fortes A, Ferreira V, Barbosa Simões E, Baptista I, Grando S, Sequeira E. Food systems and food security: the role of small farms and small food businesses in Santiago Island, Cabo Verde. *Agriculture* 2020; 10:216; <https://doi.org/10.3390/agriculture10060216>
- [7] Suhaila AH, Sabrina DL, Ahmad N, Irwan Izzauddin NH, Hamdan A, Khadijah S. Study of parasites in commercial free-range chickens in northern peninsular Malaysia. *Malays J Vet Res* 2015; 6(2):53–64.
- [8] Jaiswal K, Mishra S, Bee A. Prevalence of gastrointestinal helminth parasites in *Gallus gallus domesticus* in Lucknow, UP, India. *Adv Zool Bot* 2020; 8(5):422–30; <https://doi.org/10.13189/azb.2020.080506>
- [9] Ossa-Lacayo A, Ballut-Dajud G, Monroy-Pineda M. Análisis temporal de la cobertura en sabanas antrópicas de Sucre, Colombia. *Rev Colomb Cienc Anim* 2017; 9(26):26–30; <https://doi.org/10.24188/recia.v9.nS.2017.517>
- [10] Suarez M, Tapia F. Interaprendizaje de Estadística Básica. Universidad Técnica del Norte, Facultad de Ciencias Administrativas y económicas, Ibarra, Ecuador; pp 130, 2014.
- [11] Ensuncho C, Herrera Y, Montalvo A, Almanza M, Vergara J, Pardo E, et al. Frecuencia de parásitos gastrointestinales en *Gallinas criollas* (*Gallus domesticus*) en el departamento de Córdoba, Colombia. *Ret Vet* 2015; 16(6):1–9.
- [12] Singla LD, Gupta SK. Advances in diagnosis of coccidiosis in poultry. In: Gupta RP, Garg SR, Nehra V, Lather D (eds.). *Veterinary Diagnostics: Current Trends*, Satish Serial Publishing House, Delhi, India, pp 615–2, 2012.
- [13] Da Silva GS, Romera DM, Da Silva CG, Soares VE, Meireles MV. Helminth infections in chickens (*Gallus domesticus*) raised in different production systems in Brazil *Vet Parasitol Reg Stud Rep* 2018; 12:55–60; <https://doi.org/10.1016/j.vprsr.2018.02.003>
- [14] Yonairo HB, Michael AP, Luis GM. Frecuencia de parásitos gastrointestinales en patos domésticos (*Anas platyrhynchos domesticus*) en el departamento de Córdoba, Colombia. *Rev Elect Vet* 2016; 17(9):1–7.
- [15] Shifaw A, Feyera T, Walkden-Brown SW, Sharpe B, Elliott T, Ruhnke I. Global and regional prevalence of helminth infection in chickens over time: a systematic review and meta-analysis. *Poult Sci* 2021; 100(5):1–11; <https://doi.org/10.1016/j.psj.2021.101082>
- [16] Wongrak K, Das G, Moors E, Sohnrey B, Gauly M. Establishment of gastrointestinal helminth infections in free-range chickens: a longitudinal farm study. *Berl Munch Tierarztl Wochenschr* 2014; 127:305–13.
- [17] Wuthijaree K, Lambertz C, Gauly M. Prevalence of gastrointestinal helminth infections in free-range laying hens under mountain farming production conditions. *Br Poult Sci* 2017; 58:649–55; <https://doi.org/10.1080/00071668.2017.1379049>
- [18] Ebrahimi M, Asadpour M, Khodaverdi M, Borji H. Prevalence and distribution of gastrointestinal helminths in free range chickens in Mashhad, northeast of Iran. *Sci Parasitol* 2014; 15:38–42.
- [19] Larki S, Alborzi A, Chegini R, Amiri R. A preliminary survey on gastrointestinal parasites of domestic ducks in Ahvaz, Southwest Iran. *Iran J Parasitol* 2018; 13(1):137–44.
- [20] Wuthijaree K, Lambertz C, Veerasilp T, Anusatsananun V, Gauly M. Prevalence of gastrointestinal helminths in Thai indigenous chickens raised under backyard conditions in Northern Thailand. *J Appl Poult Res* 2019; 28:221–9; <https://doi.org/10.3382/japr/pfy062>
- [21] Chilinda I, Lungu JCN, Phiri IK, Chibinga OC, Simbaya J. Prevalence of helminths infestation in indigenous free-ranging chickens in different ecological zones in Zambia. *Livest Res Rural Dev* 2020; 32(9):147.
- [22] Sahu S, Anand A, Sinha K P. Studies on the prevalence of helminthic infection in broiler poultrybirds from Darbhanga region of North Bihar, India. *Int J Fauna Biol Stud* 2016; 3:15–8.
- [23] Jatau ID, Sulaimon NH, Musa IW, Lawal AI, Okubanjo OO, Isah I, et al. Prevalence of coccidian infection and preponderance *Eimeria* species in free range indigenous and intensively managed exotic chickens during hot-wet season in Zaria, Nigeria. *Asian J Poult Sci* 2012; 6(3):79–88; <https://doi.org/10.4314/sokjvs.v13i3.5>
- [24] Jegede OC, Asadu IA, Opara M, Obeta SS, Olayemi DO. Gastrointestinal parasitism in local and exotic breeds of chickens reared in Gwagwalada guinea Savannah zone of Nigeria. *Sokoto J Vet Sci* 2015; 13(3):25–30; <https://doi.org/10.4314/sokjvs.v13i3.5>
- [25] Adeyemi OO, Idowu ET, Otubanjo OA, Ajayi MB. Blood and gastrointestinal tract protozoa infections of domesticated chicken slaughtered in Lagos central, southwestern Nigeria. *Unilag J Med Sci Technol* 2019; 7(1):41–52.
- [26] Permin A, Esmann JB, Hoj CH, Hove T, Mukaratirwa S. Ecto-, endo- and haemoparasites in free-range chickens in the Goromonzi District in Zimbabwe. *Prev Vet Med* 2002; 54:213–24; [https://doi.org/10.1016/S0167-5877\(02\)00024-7](https://doi.org/10.1016/S0167-5877(02)00024-7)
- [27] Hussen H, Chaka H, Deneke Y, Bitew M. Gastrointestinal helminths are highly prevalent in scavenging chickens of selected districts of eastern Shewa zone, Ethiopia. *Pak J Biol Sci* 2012; 15:284–9; <https://doi.org/10.3923/pjbs.2012.284.289>
- [28] Divyamery R, Subramanian N, Soundhararajan C, Muthu M. Studies on gastrointestinal parasites of chicken in and around Cheyyar Taluk, Thiruvannamalai district. *Int J Recent Adv Multidiscipl Res* 2016; 3:2024–30.
- [29] Rukambile EJ, Chengula A, Swai ES, Jongejan F. Poultry ecto-, endo- and haemoparasites in Tanzania: a review. *Austin J Vet Sci Anim Husb* 2020; 7(1):1–9.
- [30] Fink M, Permin A, Magwisha HB, Jensen KMV. Prevalence of the proventricular nematode *Tetrameres americana* Cram (1927) in different age groups of chickens in the Morogoro Region, Tanzania. *Trop Anim Health Prod* 2005; 37:133–7; <https://doi.org/10.1023/B:TROP.0000048512.83379.0e>
- [31] Adejinmi JO, Oke M. Gastrointestinal parasites of domestic ducks (*Anas platyrhynchos*) in Ibadan Southwestern Nigeria. *Asian J Poult Sci* 2011; 5:46–50; <https://doi.org/10.3923/ajpsaj.2011.46.50>
- [32] Komba EVG, Mkupasi EM, Mwesiga GK, Mbyuzi AO, Busagwe Z, Mzula A, et al. occurrence of helminths and coccidia in apparently healthy free range local chickens slaughtered at Morogoro live bird market. *Tanzan Vet J* 2013; 28(2):55–61.
- [33] Singh M, Kaur P, Singla LD, Kashyap N, Bal MS. Assessment of risk factors associated with prevalence of gastrointestinal parasites

- in poultry of central plain zone of Punjab, India. *Vet World* 2021; 14(4):972–7; <https://doi.org/10.14202/vetworld.2021.972-977>
- [34] Adang KL, Oniye SJ, Ajanusi JO, Ezealor AU, Abdu PA. Helminths gastrointestinales de las palomas domésticas. *Columba livia domestica* Gmelin, 1789 Aves: Columbidae en Zaria, norte de Nigeria. *Sci World J* 2008; 3:33–7; <https://doi.org/10.4314/swj.v3i1.51769>
- [35] Shaikh AA. Pathology of *Heterakis gallinarum* in the ceca of naturally infected chicken (*Gallus Domesticus*). *Pure Appl Biol* 2016; 5:1–7; <https://doi.org/10.19045/bspab.2016.50102>
- [36] Wangelu K, Walkite F, Debela A. Investigation of major ecto-parasite affecting backyard chicken in Bishoftu Town, Ethiopia. *J Med Health* 2021; 3(1):1–9; <https://doi.org/10.1155/2021/5591932>
- [37] Shrestha D, Subedi JR, Chhetri B. Gastrointestinal parasites of domesticated duck (*Anas platyrhynchos* Linnaeus, 1758) in Chandragiri municipality, Kathmandu, Nepal. *Ife J Sci* 2020; 22(2):15–22; <https://doi.org/10.4314/ij.s.v22i2.2>