

Research Article

Influence of Treadmill Speed and Perturbation Intensity on Selection of Balancing Strategies during Slow Walking Perturbed in the Frontal Plane

Zlatko Matjačić , Matjaž Zadavec, and Andrej Olenšek

University Rehabilitation Institute, Republic of Slovenia, Linhartova 51, SI-1000 Ljubljana, Slovenia

Correspondence should be addressed to Zlatko Matjačić; zlatko.matjacic@ir-rs.si

Received 22 February 2019; Revised 8 May 2019; Accepted 16 May 2019; Published 2 June 2019

Academic Editor: Craig P. McGowan

Copyright © 2019 Zlatko Matjačić et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Common understanding is that adequate foot placement (stepping strategy) is crucial in maintaining stability during walking at normal speed. The aim of this study was to investigate strategies that humans use to cope with lateral perturbations during very slow walking. **Methods.** Ten healthy individuals underwent an experimental protocol whereby a set of perturbations directed inward (medially to a stance leg) and outward (laterally to a stance leg) of three intensities ($F_1 = 5\%$, $F_2 = 10\%$, and $F_3 = 15\%$ of body weight), applied at three instances of a stance phase, were delivered in random order to the pelvis using a balance assessment robot while walking on a treadmill at three walking speeds ($S_1 = 0.4$, $S_2 = 0.6$, and $S_3 = 0.8$ m/s). We analyzed the peak center of mass displacements; step length, step width, and step times; and the lateral component of ground reaction force for perturbations that were delivered at the beginning of the gait cycle. **Results.** Responses after inward perturbations were similar at all tested speeds and consistently employed stepping strategy that was further facilitated by a shortened stance. Wider and shorter steps were applied with increased perturbation intensity. Responses following outward perturbations were more complex. At S_1 , hip strategy (impulse-like increase of mediolateral ground reaction force) augmented with ankle strategy (mediolateral shift of the center of pressure) mainly contributed to responses already during the stance phase. The stance duration was significantly longer for all perturbation intensities. At S_2 , the relative share of hip strategy was reduced while with increased perturbation intensity, stepping strategy was gradually added. The stance duration was significantly longer for F_1 and F_2 . At S_3 , stepping strategy was mainly used while the duration of stance was similar to the one in unperturbed walking. Responses following both inward and outward perturbations at all speeds were characterized by temporary slowing down movement in a sagittal plane that was more pronounced with increased perturbation intensity. **Conclusions.** This study provides novel insights into balancing strategies used at slower walking speeds which may be more relevant to understand the challenges of gait stability following perturbations in the frontal plane in clinical populations.

1. Introduction

An essential component of bipedal walking is maintenance of dynamic balance, particularly in the frontal plane [1]. The main mechanism used in normal unperturbed human walking has been explained through the inverted pendulum model and is related to adequate placement of the swinging limb onto a new stance location [2]. This changes the base of support (BOS) and provides appropriate development of the lateral component of ground reaction force to ensure stable side-to-side movement of the center of mass (COM) [3].

Likewise, following a perturbation that may be imposed by various perturbation modalities, for example, (i) as a push at the waist level, mimicking a sudden bump into another person in a crowd [4–7], (ii) as a movement of the support surface, mimicking a slip [8], and (iii) as a pull on the foot of the swinging leg, mimicking a trip [9], the main balancing strategy used was related to the placement of the swinging limb onto an adequate location [3, 10, 11]. Stepping was additionally augmented by the “ankle strategy,” which is related to the activity of ankle musculature to displace the center of pressure (COP) under the stance leg in the direction

of the action of perturbation [12, 13]. Additionally, for perturbations acting in the inward direction (medially relative to the stance leg in case of perturbing pushes to the waist), the swing time was shortened to facilitate earlier application of balance correction in the next step [4, 6]. However, for the perturbations acting in the outward direction (laterally relative to the stance leg in case of perturbing pushes to the waist), shortening of the swing phase has not been observed [4, 6]. Several studies have also pointed out that in sessions where perturbations in the frontal plane were delivered subjects adopted wider stepping as an additional stabilizing measure, compared to sessions without perturbations [14, 15]. The above referenced studies examined dynamic reactions to perturbations that were imposed during walking in a range of speeds that are normally used (0.8–1.2 m/s).

Various diseases or injuries to the central nervous system (CNS) result in substantially reduced motor capabilities in clinical cases. For example, after completion of clinical rehabilitation, the majority of stroke survivors walk with speeds that range from 0.4 to 0.8 m/s [16]. Our knowledge of balancing mechanisms used following perturbations at these lower speeds of walking is scarce. One consequence of slower walking is that swing times are longer. Thus, for example, if a perturbation is imposed during a double support phase, which would resemble a situation of a slip on the floor [8], it may be the case that a corrective action coming from a wider/narrower next step, which inevitably acts with considerable delay against the induced instability [4], would not be sufficient to successfully correct for the perturbation. Thus, corrective actions may be required to start already during the stance phase. Apart from using ankle strategy under the stance leg that can act fast against perturbation but has limited stabilizing effect due to a narrow foot width [4, 13], additional strategy related to counter-rotation of body segments, termed as “inertial strategy” [12], which is frequently used during one-leg standing [17], may be utilized during slow walking. The most notable example of inertial strategy is related to the pelvis and trunk rotation and has been termed as “hip strategy” [12, 17]. In our previous work with a selected neurologically intact subject walking at the speed of 0.4 m/s, we observed that an important contribution to the balancing response after an outward perturbing push was a hip strategy related to the activity of hip abductors of the stance leg [18]. Vlutters et al. [19] have also observed important activity of the gluteus medius muscle of the stance leg following outward pushes at walking speed of 0.6 m/s. On the other hand, studies from Hof et al. [4] and Vlutters et al. [19] where pelvis perturbations of similar intensity were applied in the frontal plane at walking speed of 1.2 m/s have not observed use of hip strategy. This indicates that walking speed may have a considerable influence on the selection of a suitable balancing strategy or a synergy of balancing strategies following perturbations applied in the frontal plane.

The aim of this study was to systematically investigate the kinematics and kinetics of reactive dynamic balancing at various speeds of slower walking and at various intensities of inward- and outward-directed perturbing pushes applied at the waist at the beginning of the stance phase, to elucidate

the interplay of strategies that humans use to cope with the consequences of an unexpected lateral perturbation.

2. Methods

2.1. Subjects. Ten healthy males without known history of neuromuscular or orthopedic problems (age: 31 ± 5 years, height: 180 ± 3.9 cm, and mass: 78.7 ± 6.5 kg) participated in this study after signing informed consent forms. The subjects represent a sample of convenience. The study was approved by the Slovenian National Ethics Committee.

2.2. Instrumentation. Figure 1 shows the experimental environment, which consisted of a balance assessment robot and an instrumented treadmill (BART). Here, only a brief description of the experimental setup is given, as a more detailed description is provided elsewhere [6, 7]. The BART interfaces with the pelvis of a walking participant with six degrees of freedom (DOF). Five of the DOFs (translation of the pelvis in the sagittal, lateral, and vertical directions; pelvic rotation; and pelvic list) are actuated and admittance-controlled, providing transparent haptic interaction with negligible power transfer [7]. The sixth DOF (pelvic tilt) is passive. The BAR-TM is capable of delivering perturbations in the forward/backward and left/right directions. In this study, we only considered inward and outward perturbations delivered in the frontal plane as depicted in Figure 1.

COM movement was estimated from the translational movement of the subjects' pelvis and assessed from the movement of the BAR-TM, similarly as in our previous studies [7, 18]. Recordings of the ground reaction force (GRF) and COP in the transversal plane during walking were obtained by means of four force transducers (K3D120, ME Systeme GmbH) placed underneath the treadmill. Spatiotemporal data were assessed by means of an OptiTrack camera (NaturalPoint Inc.). Passive reflective markers were placed on the participants' feet (on the medial malleoli and the first and fourth metatarsal joints) [7, 18]. Sampling frequency for the kinematic and kinetic data was 50 Hz which is considered to be adequate for this type of study [20].

2.3. Experimental Protocol. The experimental protocol is shown schematically in Figure 2. First, subjects walked at a treadmill speed set to 0.4 m/s for a period of three minutes—unperturbed walking session. This was followed by a period of around half an hour of perturbed walking—perturbed walking session. These two experimental blocks were then repeated for treadmill speeds of 0.6 m/s and 0.8 m/s. The whole protocol was done in a single day and took around 2 hours. Perturbations were delivered with a randomly varied pause that ranged from six to eight seconds in order to avoid predictability of the perturbation occurrence. Four perturbation directions (outward RR and LL and inward RL and LR), three perturbation onsets (at 0%, 30%, and 60% of the stance phase of a gait cycle), and three perturbation amplitudes (5%, 10%, and 15% of body weight) were varied. Each combination of perturbation parameters was repeated seven times.




FIGURE 1: Photo of a subject walking on an instrumented treadmill while being embraced by the BAR-TM perturbing device; projection on the wall shows the middle of the BAR-TM working space as well as the current position and orientation of the pelvis in a transverse plane—the subjects were instructed to return to the middle of the BAR-TM working space after they rejected perturbation (a). Top view illustration of perturbation directions: outward RR: perturbation to the right triggered at right-foot contact; inward RL: perturbation to the left triggered at right-foot contact; outward LL: perturbation to the left triggered at left-foot contact; inward LR: perturbation to the right triggered at left-foot contact (b).

Unperturbed walking	Perturbed walking	Unperturbed walking	Perturbed walking	Unperturbed walking	Perturbed walking
(3 min)	252 random perturbations (30 min)	Rest (5 min)	252 random perturbations (30 min)	Rest (3 min)	252 random perturbations (30 min)
$S_1 = 0.4 \text{ m/s}$			$S_2 = 0.6 \text{ m/s}$		$S_3 = 0.8 \text{ m/s}$

FIGURE 2: Schematic diagram of the experimental protocol.

This yielded a total of 252 perturbing pushes at each walking speed that were block-randomized. Perturbations took the form of a force impulse lasting 150 ms [6, 7, 18]. Prior to this study, all subjects visited our laboratory where they practiced unperturbed and perturbed walking on the BAR-TM system for approximately half an hour.

2.4. Measurements and Data Analysis. The COM, COP, and GRF were first segmented into strides with the gait cycle defined as the period between two consecutive left (for LL and LR responses) or right (for RR and RL responses) heel strikes, as detected from COP_{ML} and COP_{AP} signals. Two full gait cycles, half of a cycle prior to and one and a half cycles after the onset of perturbation, were analyzed. Spatiotemporal responses were investigated in terms of step length, step width, and step time where left (right) step length was taken to be the antero-posterior distance between ankle markers at the moment of left (right) foot strike while left (right) step width was defined as the mediolateral distance between the same markers. Step times were defined as the time elapsed between two consecutive left (right) and right (left) foot strikes. In each combination of perturbation parameters, COM, COP, and GRF trajectories and spatiotemporal parameters were averaged across

seven repetitions. We also averaged spatiotemporal parameters for unperturbed walking in unperturbed walking sessions and unperturbed walking (the periods between the complete recoveries from previous perturbation until the onset of the next perturbation) in the perturbed walking sessions at each tested treadmill speed.

Although we assessed postural responses at three levels of perturbation onset, we included in further analysis only perturbations that commenced at 0% of a gait cycle.

The following data were used as outcome measures: step lengths, step widths, and step times for perturbed (we analyzed the first step after the perturbation onset which determines the “stepping” response) and unperturbed experimental conditions; peak displacements of COM within the first stride (from 0% to 100% of the gait cycle) in sagittal ($COM_{AP \text{ peak}}$) and frontal planes ($COM_{ML \text{ peak}}$); and integral of the lateral component of GRF ($GRF_{ML \text{ impulse}}$) for the period of the first stance phase (from 0% to approx. 50% of a gait cycle) (“in-stance response”) and for the period of the second stance phase (from approx. 50% to 100% of a gait cycle) (“stepping response”). Thus, the “in-stance response” period encompassed the balancing activity prior to the first step after the onset of perturbation, while the “stepping response” period encompassed the balancing

activity between the first and the second steps after the onset of perturbation. Since GRF_{ML} determines the acceleration of COM_{ML} , the $GRF_{ML \text{ impulse}}$ provides a measure of the overall balancing activity in both “in-stance” and “stepping” periods of balance responses.

2.5. Statistical Analysis. For unperturbed walking, a two-way repeated measures analysis of variance (rmANOVA) was used to test for the main effects and interactions on step length, step width, and step time between walking speed (3 levels: 0.4, 0.6, and 0.8 m/s) and walking condition (2 levels: unperturbed walking during unperturbed walking sessions and unperturbed walking during perturbed walking sessions). When a significant main effect or interaction was found, we performed post hoc pairwise comparisons for each of the walking speeds separately. A significance level of 0.05 was used.

For perturbed walking, a two-way rmANOVA was used to test for the main effects and interactions on step length, step width, step time, $COM_{ML \text{ peak}}$, $COM_{AP \text{ peak}}$, and $GRF_{ML \text{ impulse}}$ between walking speed (3 levels: 0.4, 0.6, and 0.8 m/s) and perturbation amplitude (4 levels: 0% (unperturbed strides from perturbed sessions), 5%, 10%, and 15% of body weight). When a significant main effect or interaction was found, we performed post hoc pairwise comparisons versus unperturbed walking for each of the walking speeds separately. A significance level of 0.05 was used, and a Bonferroni correction was applied to correct for multiple comparisons (0.016).

3. Results

The results for pushes RR (outward perturbation) and RL (inward perturbation) are presented in this section. The effects of pushes to both outward directions (LL and RR) were comparable. Likewise, the effects of pushes to both inward directions (LR and RL) were comparable.

3.1. Dynamic Balancing Responses following Perturbations

3.1.1. Outward Perturbations. Figure 3 shows COP, COM, and GRF responses to outward perturbations (RR) for all three tested walking speeds and for all three tested perturbation intensities for a representative subject.

(1) Frontal Plane. At a walking speed of 0.4 m/s, we can observe increased lateral displacement of COP_{ML} in the “in-stance” period of the response (from 0% to approx. 50% of a gait cycle) in relation to unperturbed walking. An impulse-like rise in the GRF_{ML} can be seen in the first half of the stance that is similar for all three intensities and acts in the direction opposite to the perturbation. Perturbation was fully contained during the “in-stance” period for perturbation intensities of 5% and 10% while following a perturbation intensity of 15%, there was medial displacement of COP_{ML} and related decrease in GRF_{ML} in the “stepping period” (from approx. 50% to approx. 100% of a gait cycle) that finally contained the instability.

At a walking speed of 0.6 m/s, we can observe increased lateral displacement of COP_{ML} in the “in-stance” period while the impulse-like rise in GRF_{ML} in the first half of the “in-stance” period was smaller in comparison to those at walking speed 0.4 m/s. Medial displacement of COP_{ML} and related decrease in GRF_{ML} were observed in the “stepping” period for the perturbation intensity of 15%.

At a walking speed of 0.8 m/s, increased lateral displacement of COP_{ML} in the “in-stance” period was observed while the impulse-like rise in the GRF_{ML} in the first half of the “in-stance” period was not present. In the second half of the same period, there was a gradual decrease of GRF_{ML} with increasing perturbation intensity followed by a progressively larger medial displacement of COP_{ML} and related decrease in GRF_{ML} in the “stepping” period.

(2) Sagittal Plane. At walking speed of 0.4 m/s, COP_{AP} was displaced increasingly forward in the first half of the stance with increasing intensity of perturbation while GRF_{AP} showed increased braking action that decelerated COM_{AP} in relation to unperturbed walking. Slowing down of COM_{AP} and associated changes in COP_{AP} were progressively smaller at walking speeds of 0.6 m/s and 0.8 m/s compared to those observed at the speed of 0.4 m/s.

3.1.2. Inward Perturbations. Figure 4 shows COP, COM, and GRF responses to inward perturbations (RL) for all three tested walking speeds and for all three tested perturbation intensities for a representative subject. The responses look similar across the tested walking speeds.

(1) Frontal Plane. In the “in-stance” period, no noticeable difference can be observed in COP_{ML} , COM_{ML} , and GRF_{ML} in relation to unperturbed walking except for a shortened duration of the stance phase. The dominant balancing response can be observed in the “stepping” period where depending on the perturbation intensity COP_{ML} was shifted laterally which was accompanied with a progressively increased GRF_{ML} .

(2) Sagittal Plane. In the second part of the “in-stance” period, a shortened posterior displacement of COP_{AP} can be observed. Consequently, GRF_{AP} was also reduced thus slowing down movement of COM_{AP} . Throughout the “stepping” period, a smaller anterior displacement of COP_{AP} can be seen with accompanying reduction of GRF_{AP} which enabled COM_{AP} to catch up with the relative position of COM_{AP} on the treadmill that the subject assumed before the action of perturbation.

3.2. Peak COM Displacements. Figure 5 shows peak excursions of COM_{ML} and COM_{AP} for both outward (RR) and inward (RL) perturbations. $COM_{ML \text{ peak}}$ following outward perturbation was significantly affected by the speed of walking ($F(2, 18) = 10.015$, $p = 0.001$), the perturbation intensity ($F(3, 27) = 274.194$, $p < 0.001$), and their interaction ($F(6, 54) = 9.790$, $p < 0.001$). Post hoc analysis has shown significantly larger peak COM_{ML} displacements for all intensities and at all speeds in comparison to




FIGURE 3: Kinematics and kinetics of balancing responses following outward RR perturbation assessed in a representative subject. The first row shows the trajectories of COP_{ML} (solid lines) and COM_{ML} (dotted lines), while the second row shows GRF_{ML} trajectories. The third row shows COP_{AP} (solid lines) and COM_{AP} (dotted lines) trajectories, while the fourth row shows GRF_{AP} trajectories. Each graph contains responses to all three perturbation intensities along with the trajectories assessed during the unperturbed walking sessions. The left, middle, and right columns show the balancing responses at speeds S_1 , S_2 , and S_3 , respectively. Half a stride prior to and one and a half strides following the perturbation commencement are shown. A stride is defined as the period between two consecutive right-foot contacts. The trajectories displayed show mean values of seven balancing responses.

unperturbed walking. $COM_{AP\ peak}$ following outward perturbation was significantly affected by the speed of walking ($F(2, 18) = 69.523, p < 0.001$), the perturbation intensity ($F(3, 27) = 88.255, p < 0.001$), and their interaction ($F(6, 54) = 5.551, p < 0.001$). Post hoc analysis has shown significantly larger peak COM_{AP} displacements for all intensities and at all speeds in comparison to unperturbed walking. $COM_{ML\ peak}$ following inward perturbation was significantly affected by the intensity of perturbation ($F(3, 27) = 53.150, p < 0.001$) and interaction between intensity and speed ($F(6, 54) = 3.556, p = 0.005$) but not by the speed of walking ($F(2, 18) = 3.088, p = 0.070$). Post hoc analysis has shown significantly larger peak COM_{ML} displacements for all intensities and at all speeds in comparison to unperturbed walking. $COM_{AP\ peak}$ following inward perturbation was significantly affected only by the perturbation intensity ($F(3, 27) = 18.892, p < 0.001$) but not by the walking speed ($F(2, 18) = 1.620, p = 0.225$) nor the interaction between the intensity and walking speed ($F(6, 54) = 1.140,$

$p = 0.352$). Post hoc analysis has shown significantly larger peak COM_{ML} displacements for majority of intensities at all speeds in comparison to unperturbed walking.

3.3. Spatiotemporal Parameters. Figure 6 shows spatiotemporal parameters for unperturbed walking during the unperturbed walking sessions (UWS) and for unperturbed walking during the perturbed walking sessions (PWS). Step lengths and step times were significantly affected only by the walking speed (step length $F(2, 18) = 354.221, p < 0.001$; step time $F(2, 18) = 238.195, p < 0.001$) but not by the walking condition (step length $F(1, 9) = 3.089, p = 0.113$; step time $F(1, 9) = 4.822, p = 0.056$) nor the interaction between the walking condition and the walking speed (step length $F(2, 18) = 1.304, p = 0.296$; step time $F(2, 18) = 2.866, p = 0.083$). Step lengths increased with increased walking speed while the step times decreased with increased walking speed. Step widths were significantly affected by walking speed ($F(2, 18) = 4.996, p = 0.019$) and walking condition




FIGURE 4: Kinematics and kinetics of balancing responses following inward RL perturbation assessed in a representative subject. The first row shows the trajectories of COP_{ML} (solid lines) and COM_{ML} (dotted lines), while the second row shows GRF_{ML} trajectories. The third row shows COP_{AP} (solid lines) and COM_{AP} (dotted lines) trajectories, while the fourth row shows GRF_{AP} trajectories. Each graph contains responses to all three perturbation intensities along with the trajectories assessed during the unperturbed walking sessions. The left, middle, and right columns show the balancing responses at speeds S_1 , S_2 , and S_3 , respectively. Half a stride prior to and one and a half strides following the perturbation commencement are shown. A stride is defined as the period between two consecutive right-foot contacts. The trajectories displayed show mean values of seven balancing responses.

($F(1, 9) = 70.489$, $p < 0.001$) but not by interaction of the two ($F(2, 18) = 0.830$, $p = 0.452$). Post hoc analysis has shown that the difference between step widths of unperturbed walking during unperturbed walking sessions and during perturbed walking sessions was on average 4 cm.

Figure 7 shows spatiotemporal parameters for perturbed walking following outward (RR) and inward (RL) perturbations. Step lengths, step widths, and step times following outward perturbation were significantly affected by the walking speed (step length $F(2, 18) = 38.259$, $p < 0.001$; step width $F(2, 18) = 8.869$, $p = 0.002$; and step time $F(2, 18) = 153.168$, $p < 0.001$), by intensity (step length $F(3, 27) = 6.721$, $p = 0.002$; step width $F(3, 27) = 51.945$, $p < 0.001$; and step time $F(3, 27) = 12.214$, $p < 0.001$), and by interaction of both factors (step length $F(6, 54) = 5.407$, $p < 0.001$; step width $F(6, 54) = 12.023$, $p < 0.001$; and step time $F(6, 54) = 13.333$, $p < 0.001$). Post hoc analysis showed no significant differences in step lengths at 0.4 m/s; at 0.6 m/s, significantly longer steps were taken at perturbation intensities of

5% and 10% while at 0.8 m/s, significantly shorter steps were made at a perturbation intensity of 15% in comparison to unperturbed walking. Post hoc analysis further revealed that step widths at a walking speed of 0.4 m/s were not statistically different; at the walking speed of 0.6 m/s, step width at the strongest perturbation was significantly smaller in comparison to unperturbed walking while at a walking speed of 0.8 m/s, step widths for all intensities were significantly smaller in comparison to unperturbed walking. Post hoc analysis for step times has shown significantly longer steps at 0.4 m/s for all intensities in comparison to unperturbed walking; at 0.6 m/s, this was the case for intensities of 5% and 10% while at 0.8 m/s, the step time for an intensity of 10% was significantly longer in comparison to unperturbed walking.

Step lengths and step times following inward perturbation were significantly affected by the walking speed (step length $F(2, 18) = 43.703$, $p < 0.001$; step time $F(2, 18) = 75.724$, $p < 0.001$) but not the step width ($F(2, 18) = 3.085$,




FIGURE 5: Group average (\pm standard deviation) of peak COM_{ML} and COM_{AP} excursions across the three walking speeds during unperturbed walking and perturbed walking is shown for outward RR (a) and inward RL (b) perturbations along with the p values of 2-way rmANOVA. Asterisks (*) indicate significant difference from unperturbed walking in Bonferroni post hoc pairwise comparisons ($p < 0.016$).

$p = 0.071$). On the other hand, perturbation intensity had significant effect on step length, step width, and step time (step length $F(3, 27) = 195.329$, $p < 0.001$; step width $F(3, 27) = 31.060$, $p < 0.001$; and step time $F(3, 27) = 361.699$, $p < 0.001$). Only step width and step time were significantly affected by interaction of both factors (step width $F(6, 54) = 2.758$, $p = 0.021$; step time $F(6, 54) = 44.580$, $p < 0.001$) and not the step length ($F(6, 54) = 1.103$, $p = 0.373$).

Post hoc analysis has shown a decrease of step lengths and step times and an increase of step widths with increased intensity at all tested speeds in comparison to unperturbed walking.

3.4. Lateral GRF Impulses. Figure 8 shows the integral of GRF_{ML} over the “in-stance” and “stepping” periods of dynamic responses following outward (RR) and inward




FIGURE 6: Group average (\pm standard deviation) of step lengths, step widths, and step times across the three walking speeds during unperturbed walking in unperturbed walking session (UWS) and during unperturbed walking in perturbed walking sessions (PWS) is shown along with the p values of 2-way rmANOVA. Asterisks (*) indicate significant difference from unperturbed walking in post hoc pairwise comparisons ($p < 0.05$).

(RL) perturbations. $GRF_{ML\ impulse}$ following outward perturbation was significantly affected by the walking speed (“in-stance” $F(2, 18) = 37.079$, $p < 0.001$; “stepping” $F(2, 18) = 82.463$, $p < 0.001$), by intensity (“in-stance” $F(3, 27) = 26.849$, $p < 0.001$; “stepping” $F(3, 27) = 87.569$, $p < 0.001$), and by interaction of both factors (“in-stance” $F(6, 54) = 10.037$, $p < 0.001$; “stepping” $F(6, 54) = 24.318$, $p < 0.001$) during both periods. Post hoc analysis for the “in-stance” period has shown significant increases for intensities of 5% and 10% at a walking speed of 0.4 m/s while $GRF_{ML\ impulse}$ value at an intensity of 15% was not significantly different from unperturbed walking. At 0.6 m/s, $GRF_{ML\ impulse}$ values at intensities of 5% and 10% were significantly increased while $GRF_{ML\ impulse}$ value at an intensity of 15% was significantly decreased in comparison to unperturbed walking. At 0.8 m/s, only $GRF_{ML\ impulse}$ value at an intensity of 5% was significantly increased while $GRF_{ML\ impulse}$ values at intensities of 10% and 15% were significantly decreased. Post hoc analysis for the “stepping” period has shown no statistically significant differences among intensity factors at a walking speed of 0.4 m/s. At walking speeds 0.6 m/s and 0.8 m/s, the $GRF_{ML\ impulse}$ shows increasingly smaller values with increasing intensity of perturbation in comparison to unperturbed walking.

Following inward perturbation, $GRF_{ML\ impulse}$ was significantly affected in both periods by the perturbation intensity (“in-stance” $F(3, 27) = 53.116$, $p < 0.001$; “stepping” $F(3, 27) = 28.598$, $p < 0.001$) and by interaction of both factors (“in-stance” $F(6, 54) = 4.832$, $p = 0.001$; “stepping” $F(6, 54) = 7.391$, $p < 0.001$) but was not significantly affected by walking speed (“in-stance” $F(2, 18) = 1.289$, $p = 0.300$; “stepping” $F(2, 18) = 1.319$, $p = 0.292$). Post hoc analysis for the “in-stance” period at all speeds showed increasingly smaller values of $GRF_{ML\ impulse}$ with increasing intensity of perturbation in comparison to unperturbed walking. Post hoc analysis

for the “stepping” period at speeds of 0.4 m/s and 0.6 m/s showed significantly higher values of $GRF_{ML\ impulse}$ for intensities of 10% and 15% while at 0.8 m/s, significantly higher values were observed for all intensities in comparison to unperturbed walking.

4. Discussion

The main aim of this study was to investigate how slower walking speeds and various intensities of perturbing pushes delivered at the heel strike influence the selection of dynamic balancing responses.

Responses after inward perturbations were similar at all tested speeds and consistently employed a predominantly stepping strategy facilitated by a shortened stance. Wider steps and shorter stances were applied with increasing perturbation strengths. The role of hip/inertial balancing strategies was not observed. These observations are well in line with the observations of other studies which were mostly performed at higher walking speeds [4, 5, 7, 11].

On the contrary, when subjects were faced with outward perturbations, additional inertial balancing strategies were used. The predominant inertial strategy associated with the counter-rotation of body segments changing GRF is the hip strategy [12, 17, 21]. Depending on the walking speed and perturbation intensity, the relative contribution of individual balancing strategy also varied. At the slowest walking speed, we consistently observed a significant contribution of hip strategy in the first half of the “in-stance” period. The hip strategy was augmented with an ankle strategy, displacing the COP in the lateral direction away from the direction of perturbation [13]. The stance duration was significantly longer for all perturbation intensities. At medium walking speed, the hip strategy augmented with an ankle strategy was sufficient at the weakest perturbation magnitude while a stepping




FIGURE 7: Group average (\pm standard deviation) of step lengths, step widths, and step times across the three walking speeds during unperturbed walking and perturbed walking is shown for outward RR (a) and inward RL (b) perturbations along with the p values of 2-way rmANOVA. Asterisks (*) indicate significant difference from unperturbed walking in Bonferroni post hoc pairwise comparisons ($p < 0.016$).

strategy was gradually added with increasing perturbation intensity. At the highest walking speed, stepping was the main strategy used to counteract the effects of perturbation while the duration of stance was similar to those in unperturbed walking.

4.1. Synergy of Balancing Strategies following Outward Perturbations. This study provides an important insight into the balancing strategies used at walking speeds that are well below those normally used and which may be more relevant for understanding the challenges of gait stability following




FIGURE 8: Group average (\pm standard deviation) of integrals of GRF_{ML} over the “in-stance” and “stepping” periods across the three walking speeds during unperturbed walking and perturbed walking is shown for outward RR (a) and inward RL (b) perturbations along with the p values of 2-way rmANOVA. Asterisks (*) indicate significant difference from unperturbed walking in Bonferroni post hoc pairwise comparisons ($p < 0.016$).

perturbations in the frontal plane in clinical populations. This study complements the existing body of knowledge on the organization of balancing responses during walking following perturbations acting in the frontal plane. The results of this study to some extent challenge the currently accepted

opinion that control of human gait is predominantly achieved through foot placement [3–5, 7, 10]. Stepping strategy may well be the primary and energetically wise optimal coping option following perturbations at normal speeds of walking; however, the inherent time-delay associated with

swing time needed for a stepping strategy to start acting against a perturbation after an outward perturbation is reciprocal to the walking speed. This becomes critical at lower speeds of walking in particular when the perturbation is applied in early stance thus maximizing the time for instability to develop. Wang and Srinivasan [22] have indicated that as much as 80% of the variance in deviations of foot placement from the average during unperturbed walking at speeds ranging from 1 to 1.4 m/s could be explained by deviations from the average in pelvis position and speed at midstance. Vlutters et al. [23] have also shown a similar correlation for perturbed walking at speed of 1.2 m/s. However, a study from Stimpson et al. [24] that examined step-by-step control of step width during unperturbed walking at speeds ranging from 0.2 to 1.2 m/s has shown that the strength of the relationship between the step width and pelvis mechanics is significantly reduced at lower speeds. These findings imply that utilization of adequate foot placement (stepping strategy) as the main strategy to maintain dynamic stability in the frontal plane depends substantially on walking speed. Therefore, during very slow walking, which is characteristic for clinical populations, other balancing mechanisms, primarily the inertial strategy in a form of hip strategy, which are considered to have a limited control ability to counteract perturbations applied in the frontal plane during walking at normally used speeds [2], should be employed earlier in the stance to impede the development of instability and thus decisively contribute to successful correction of an outward perturbation.

Previous studies have shown that mediolateral ankle strategy is employed following an outward push regardless of walking speed and intensity of perturbation [4, 5, 13] probably because it can, according to the inverted pendulum model, act fast against the developing instability by increasing GRF_{ML} in the direction opposite to the action of a perturbation. This is also what we observed in this study. No visually noticeable movement of the trunk or the arms was observed following perturbations in our study, which is consistent with observations from other studies using similar perturbation intensities and at walking speeds of 1.2 m/s [4] and 0.6 m/s [5]. However, the observed impulse-like increase of GRF_{ML} immediately after the perturbation commencement at low walking speed is not consistent with the inverted pendulum model [1, 12] and implies movement of other body segments. Balancing activity during the “in-stance” period following the commencement of outward perturbation seems to be similar to balancing while standing on one leg [12, 17] where visually notable counter-rotation of body segments resulting in an increase of GRF_{ML} that acts in the opposite direction of the COM_{ML} movement is readily used to maintain balance. Our recent study [18] and the study from Vlutters et al. [19] have shown pronounced activity of hip abductors of the stance leg following an outward perturbation which seems to be the cause of the observed impulse-like increase in GRF_{ML} constituting a hip strategy. Since this GRF_{ML} impulse was rather small, it has limited capacity to substantially move the relatively heavy trunk. Future studies should explore the neuromechanics of the “in-stance” balancing responses (consisting of ankle and hip strategies) follow-

ing an outward perturbation at low walking speed in more details.

4.2. Shortening and Prolongation of the Stance Duration as a Balancing Strategy. It was shown that humans while walking at normal speeds and when subjected to larger perturbations of inward direction react more quickly by shortening their stance duration so that the next step which is also a corrective step is made earlier [4, 11]. Our results show that this is also the case for lower speeds.

However, when reacting to outward perturbations at lower speeds, the stance phase was in most cases substantially prolonged, thus prolonging utilization of the “in-stance” hip strategy resulting in an increase of GRF_{ML} in the first half of the stance that acted in the direction opposite to the movement of COM.

4.3. Braking of the Movement in the Plane of Progression as a Balancing Strategy. Both types of perturbation lead to temporary slowing down of progression in the sagittal plane, which was more pronounced for outward perturbations at lower speeds and stronger perturbations. Similar results were also observed in the studies of Hof et al. [4, 11, 13], however, at higher walking speeds. It seems that slowing down of the movement in the plane of progression following perturbation acting in the frontal plane is related to stiffening of the ankle [13] and also knee and hip joints [19] of the stance leg.

4.4. Widening of Steps as a General Strategy to Increase Stability when Faced with the Prospect of a Period of Perturbations. Several studies have identified a common precautionary strategy of widening steps when faced with prospective perturbations [14, 15]. The results of our study show that the subjects consistently adopted wider steps during perturbed walking sessions. In our opinion, this further stresses the importance of utilizing the hip balancing strategies in the first half of the stance at the lowest speed of walking as seen in this study, since in a real-life situation, the occurrence of perturbation cannot be expected in advance. Therefore, the narrower stepping that is normally exercised during walking would facilitate an even larger destabilizing effect of a perturbation applied in the frontal plane as compared to walking with adopted wider stepping.

4.5. Relevance of COM-Based Pushes to Real-Life Situations. Hof and Duysens [13] compared the results of ankle muscle activity responses obtained in their study, where the pushes at the level of COM were applied, to the ankle muscle activity obtained in studies that used walking surface translations as a source of perturbation. They concluded that the lateral translation of the floor is comparable to inward (medial) perturbation at the level of the waist while the medial translation of the floor is similar to outward (lateral) perturbation at the waist level. Oddsson et al. [8] applied surface translations and observed that lateral translation of the standing foot during the midstance caused the upcoming step to be wider while a medially directed translation caused the upcoming step to be narrower. This is also in agreement with the notion of Hof and Duysens [13] with respect to the similarity of surface translation-based and waist push-based perturbations.

4.6. Methodological Considerations and Study Limitations. We limited our analysis to single perturbation timing. The instant of perturbation commencement at the beginning of the stance was selected because this particular timing gives the perturbation opportunity to develop instability for the longest period until the leg in swing can enter stance onto a new location. For example, Oddsson et al. [8] have applied perturbations during the midstance and not at the beginning of the stance phase when one would normally expect a real slip to occur. They did that in order to avoid possible tripping which is clearly associated with the stepping response following an outward perturbation. Therefore, a perturbation applied at the beginning of the stance seems to be the most challenging to cope with. Additionally, at this perturbation timing, the responses throughout the varied speeds and intensities finished at the end of the next step (within one stride—0–100% of a gait cycle). This enabled consistent treatment of all responses. However, from a methodological point of view, the inclusion of two additional instances of perturbation occurrence (at 30% and 60% of stance) increased the level of unpredictability which increases the strength of our findings.

The largest perturbation intensity used in this study was 15% of body weight which was also the value used in our previous study [7]. At this perturbation intensity, no noticeable trunk movement or arm movement was observed; however, at the lowest treadmill speed, a perturbation of 15% elicited responses in some subjects that, beside hip strategy, also required utilization of stance foot repositioning through pivoting on the toes and heel which has also been shown in our previous study [14]. This response indicates that if we increased the perturbation magnitude even further, additional balancing strategies such as trunk rotation, rapid arm and leg movements, and possibly also hopping on the stance leg could be employed, possibly resulting in inconsistent and variable within-subject responses.

The balance assessment robot was controlled such that the interaction forces between the walking subject and pelvis link were as low as possible. We have assessed the peak interaction forces in a previous study and found that the influence of these forces on COP and GRF in sagittal and frontal planes as well as on EMGs of major lower limb muscles during unperturbed walking had negligible effects in the range of walking speed from 0.4 to 0.8 m/s [25]. In another study, we have demonstrated that the interaction between the balance assessment robot and the pelvis of a walking subject is purely passive; thus, the interaction forces can be perceived as reflected inertia which was estimated for the balance assessment robot to be below 5 kg at walking speed of 0.85 m/s [7]. This is below a value of 6 kg identified in the study of Meuleman et al. [26] that can be added to the pelvis without significantly affecting the gait. Therefore, we may conclude that the interaction forces between the balance assessment robot and the walking subject had minimal influence on the dynamic balancing responses observed in this study.

A valid question one can pose is how relevant may be the balancing responses assessed in healthy people to indi-

viduals with gait pathology. Able-bodied and neurologically impaired subjects have in general different repertoires of available muscle actions to react to unexpected mechanical perturbations. Balancing responses as assessed in able-bodied subjects can be regarded as optimal solution for a given speed of walking and a given perturbation strength. In this study, we found differences in a way how the able-bodied population reacts to perturbations of the same intensity at different speeds of walking. A significant balancing activity must commence already during the “in-stance period,” which is not the case for higher walking speeds. This implies that at low speed of walking, the task of balancing for neurologically impaired will be substantially the same as for able-bodied, and if they will not react appropriately already in the “in-stance period,” this will have consequences for subsequent phases of response. Therefore, understanding balancing responses in able-bodied individuals at speeds that are similar to those used by neurologically impaired individuals enables us to better understand what the task at hand is for neurologically impaired when subjected to mechanical perturbation.

5. Conclusion

In conclusion, our findings reveal reactive dynamic balancing responses following perturbations delivered at the waist at very slow walking speeds. The responses to inward perturbations predominantly consist of a stepping strategy at all walking speeds and all tested perturbation intensities. The responses to outward perturbations are highly dependent on the walking speed and perturbation intensity. Our interpretation is that at the slowest speed and lowest intensity, the hip strategy is dominant while at the greatest speed and highest intensity, the stepping strategy dominates. Between these two extremes, a synergy of both strategies is used with the relative share of each strategy depending on the walking speed and perturbation intensity. Further studies should explore in more details the neuromechanics of the “in-stance” balancing responses (consisting of ankle and hip strategies) following an outward perturbation at low walking speeds.

The results related to balancing responses following outward perturbations have implications for the development of a screening method which could identify potential fallers among either elderly or neurologically impaired populations. Inability to generate an adequate “in-stance” response at slower speeds of walking could be an indication of diminished balancing abilities. The results of this study may also be very relevant to the developers of control approaches applied in robot exoskeletons that are used to support walking and balance functions.

Data Availability

The datasets used and/or analyzed during the current study are available from the corresponding author on request for reasonable use.

Ethical Approval

Ethical approval for this study was obtained from Republic of Slovenia National Medical Ethics Committee, decision number 80/03/15.

Consent

Participants gave consent to use and publish data in such way that anonymity is assured. All participants gave signed, written, informed consent.

Conflicts of Interest

The authors declare that they have no competing interests.

Authors' Contributions

ZM conceived the draft structure and wrote the final version of the paper. MZ and AO critically revised the draft paper and contributed to all sections. ZM, MZ, and AO all contributed substantially to the analysis of the data. MZ did most of the data processing and prepared the figures. All authors read and approved the final manuscript.

Acknowledgments

This research was supported by the Slovenian Research Agency under research project J2-8172 and research program number P2-0228.

References

- [1] C. D. MacKinnon and D. A. Winter, "Control of whole body balance in the frontal plane during human walking," *Journal of Biomechanics*, vol. 26, no. 6, pp. 633–644, 1993.
- [2] C. E. Bauby and A. D. Kuo, "Active control of lateral balance in human walking," *Journal of Biomechanics*, vol. 33, no. 11, pp. 1433–1440, 2000.
- [3] S. M. Bruijn and J. H. van Dieën, "Control of human gait stability through foot placement," *Journal of The Royal Society Interface*, vol. 15, no. 143, article 20170816, 2018.
- [4] A. L. Hof, S. M. Vermerris, and W. A. Gjaltema, "Balance responses to lateral perturbations in human treadmill walking," *The Journal of Experimental Biology*, vol. 213, no. 15, pp. 2655–2664, 2010.
- [5] M. Vlutters, E. H. F. van Asseldonk, and H. van der Kooij, "Center of mass velocity based predictions in balance recovery following pelvis perturbations during human walking," *The Journal of Experimental Biology*, vol. 219, no. 10, pp. 1514–1523, 2016.
- [6] A. Olenšek, M. Zadavec, and Z. Matjačić, "A novel robot for imposing perturbations during overground walking: mechanism, control and normative stepping responses," *Journal of Neuroengineering and Rehabilitation*, vol. 13, no. 1, p. 55, 2016.
- [7] Z. Matjačić, M. Zadavec, and A. Olenšek, "An effective balancing response to lateral perturbations at pelvis level during slow walking requires control in all three planes of motion," *Journal of Biomechanics*, vol. 60, pp. 79–90, 2017.
- [8] L. I. E. Oddsson, C. Wall III, M. D. McPartland, D. E. Krebs, and C. A. Tucker, "Recovery from perturbations during paced walking," *Gait & Posture*, vol. 19, no. 1, pp. 24–34, 2004.
- [9] Z. Potocanac, M. Pijnappels, S. Verschuereen, J. van Dieën, and J. Duysens, "Two-stage muscle activity responses in decisions about leg movement adjustments during trip recovery," *Journal of Neurophysiology*, vol. 115, no. 1, pp. 143–156, 2016.
- [10] B. L. Rankin, S. K. Buffo, and J. C. Dean, "A neuromechanical strategy for mediolateral foot placement in walking humans," *Journal of Neurophysiology*, vol. 112, no. 2, pp. 374–383, 2014.
- [11] A. L. Hof and J. Duysens, "Responses of human hip abductor muscles to lateral balance perturbations during walking," *Experimental Brain Research*, vol. 230, no. 3, pp. 301–310, 2013.
- [12] A. L. Hof, "The equations of motion for a standing human reveal three mechanisms for balance," *Journal of Biomechanics*, vol. 40, no. 2, pp. 451–457, 2007.
- [13] A. L. Hof and J. Duysens, "Responses of human ankle muscles to mediolateral balance perturbations during walking," *Human Movement Science*, vol. 57, pp. 69–82, 2018.
- [14] L. Hak, H. Houdijk, F. Steenbrink et al., "Speeding up or slowing down?: gait adaptations to preserve gait stability in response to balance perturbations," *Gait & Posture*, vol. 36, no. 2, pp. 260–264, 2012.
- [15] H. E. Stokes, J. D. Thompson, and J. R. Franz, "The neuromuscular origins of kinematic variability during perturbed walking," *Scientific Reports*, vol. 7, no. 1, p. 808, 2017.
- [16] B. Raja, R. R. Neptune, and S. A. Kautz, "Quantifiable patterns of limb loading and unloading during hemiparetic gait: relation to kinetic and kinematic parameters," *The Journal of Rehabilitation Research and Development*, vol. 49, no. 9, pp. 1293–1304, 2012.
- [17] E. Otten, "Balancing on a narrow ridge: biomechanics and control," *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, vol. 354, no. 1385, pp. 869–875, 1999.
- [18] Z. Matjačić, M. Zadavec, and A. Olenšek, "Feasibility of robot-based perturbed-balance training during treadmill walking in a high-functioning chronic stroke subject: a case-control study," *Journal of Neuroengineering and Rehabilitation*, vol. 15, no. 1, p. 32, 2018.
- [19] M. Vlutters, E. H. F. van Asseldonk, and H. van der Kooij, "Lower extremity joint-level responses to pelvis perturbation during human walking," *Scientific Reports*, vol. 8, no. 1, article 14621, 2018.
- [20] C. K. Rhea, A. W. Kiefer, W. G. Wright, L. D. Raisbeck, and F. J. Haran, "Interpretation of postural control may change due to data processing techniques," *Gait & Posture*, vol. 41, no. 2, pp. 731–735, 2015.
- [21] R. R. Holt, D. Simpson, J. R. Jenner, S. G. B. Kirker, and A. M. Wing, "Ground reaction force after a sideways push as a measure of balance in recovery from stroke," *Clinical Rehabilitation*, vol. 14, no. 1, pp. 88–95, 2000.
- [22] Y. Wang and M. Srinivasan, "Stepping in the direction of the fall: the next foot placement can be predicted from current upper body state in steady-state walking," *Biology Letters*, vol. 10, no. 9, article 20140405, 2014.
- [23] M. Vlutters, E. H. F. van Asseldonk, and H. van der Kooij, "Foot placement modulation diminishes for perturbations near foot contact," *Frontiers in Bioengineering and Biotechnology*, vol. 6, p. 48, 2018.

- [24] K. H. Stimpson, L. N. Heitkamp, J. S. Horne, and J. C. Dean, "Effects of walking speed on the step-by-step control of step width," *Journal of Biomechanics*, vol. 68, pp. 78–83, 2018.
- [25] A. Olenšek, M. Zadavec, and Z. Matjačić, "The effects of haptic interaction between balance assessment robot and pelvis on muscle activation of leg muscles," in *2017 International Conference on Rehabilitation Robotics (ICORR)*, pp. 234–239, London, UK, July 2017.
- [26] J. H. Meuleman, E. H. F. van Asseldonk, and H. van der Kooij, "The effect of directional inertias added to pelvis and ankle on gait," *Journal of Neuroengineering and Rehabilitation*, vol. 10, no. 1, p. 40, 2013.