

Review Article

Effects of Lycopene Attenuating Injuries in Ischemia and Reperfusion

Sijia Wu,¹ Xiajun Guo,¹ Jia Shang,² Yuanyuan Li,¹ Wanglin Dong,¹ Qianwen Peng,¹ Zhenxing Xie^{ID},³ and Chaoran Chen^{ID}¹

¹Institute of Nursing and Health, College of Nursing and Health, Henan University, Kaifeng, Henan, China

²Arts Department, School of Kaifeng Culture and Tourism, Kaifeng, Henan, China

³School of Basic Medical Sciences, Henan University, Jinming Avenue, Kaifeng, Henan 475004, China

Correspondence should be addressed to Zhenxing Xie; pet213@163.com and Chaoran Chen; kfccr@126.com

Received 2 May 2022; Revised 25 July 2022; Accepted 15 September 2022; Published 7 October 2022

Academic Editor: Pasquale Pagliaro

Copyright © 2022 Sijia Wu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Tissue and organ ischemia can lead to cell trauma, tissue necrosis, irreversible damage, and death. While intended to reverse ischemia, reperfusion can further aggravate an ischemic injury (ischemia-reperfusion injury, I/R injury) through a range of pathologic processes. An I/R injury to one organ can also harm other organs, leading to systemic multiorgan failure. A type of carotenoid, lycopene, has been shown to treat and prevent many diseases (e.g., rheumatoid arthritis, cancer, diabetes, osteoporosis, male infertility, neurodegenerative diseases, and cardiovascular disease), making it a hot research topic in health care. Some recent researches have suggested that lycopene can evidently ameliorate ischemic and I/R injuries to many organs, but few clinical studies are available. Therefore, it is essential to review the effects of lycopene on ischemic and I/R injuries to different organs, which may help further research into its potential clinical applications.

1. Introduction

Ischemia refers to insufficient blood supply to the tissues or organs of the body [1]. It is usually caused by events such as thromboembolism, trauma, atherosclerosis, tachycardia, and organ transplantation [2]. Ischemia can lead to the lack of oxygen and nutritional elements required for cell metabolism and insufficient clearance of metabolic waste [3, 4]. The direct outcomes of ischemia are cell damage, tissue necrosis, irreversible pathologies, and death [5]. The severity of ischemic damage rests with the degree and duration of blood flow restriction. In cases of the short-term or mild ischemia, a new balance formed between energy supply and utilization results in less severe injuries and blood supply can be recovered spontaneously [6, 7]. However, severe or long-term ischemia can lead to infarctions at the ischemic site. Although reperfusion can restore blood flow to the ischemic tissues, it often leads to further damage resulting in molecular (lipid, protein,

and DNA) damage, cell necrosis, and tissue dysfunction [8–10]. Ischemia-reperfusion (I/R) injuries often occur following postischemia treatments (e.g., relieving coronary artery spasm, dredging the microcirculation in shock, and cardiopulmonary cerebral resuscitation) or surgery (arterial bypass grafting, thrombolytic therapy [11], and organ transplantations [12]). Furthermore, I/R injuries not only harm the I/R site but can also injure other organs and systems. For example, a liver I/R injury not only damages the liver itself but also triggers remote organ dysfunction to the lungs, kidneys, and myocardium [13–15]. Therefore, it is necessary to identify natural active substances that can reduce the impact of ischemic and I/R injuries. Prior studies have shown that several bioactive substances (valsartan [16], carbon monoxide [17], hydrogen sulfide [18], oleuropein [19], and necrostatin-1 [20]) can be used as adjunct therapies for ischemic and I/R injuries. Besides, lycopene has been found to be protective effects on ischemic and I/R injuries.

2. Lycopene

2.1. Sources and Structure. Lycopene has been isolated and identified for more than one hundred years. As a natural pigment, lycopene is mainly present in red fruits and vegetables, for example, watermelon, papaya, red carrot, and pink guava [21]. Photosynthetic bacteria can also produce a certain amount of lycopene. Lycopene consists of eight isoprene units that contains eleven conjugated double bonds and two nonconjugated double bonds. As a member of the carotenoid family, lycopene exists in the all-*trans* form and can be *cis*-isomerized by thermal energy, light energy, and chemical reactions [22]. 5-*cis*, 9-*cis*, 13-*cis*, and 15-*cis* are the most common *cis* isomers of lycopene. Of these, 5-*cis* lycopene has the highest stability and antioxidation. Although the stability of all-*trans* lycopene is second only to 5-*cis* lycopene, it has the lowest antioxidant properties [23].

The bioavailability and biological properties of lycopene are affected by the conversion of *trans*-lycopene to *cis*-lycopene during food processing (e.g., cooking), digestion, and absorption (Figure 1) [24]. For instance, dietary fiber and β -carotene impede lycopene absorption, but dietary fats favor it [25]. In micelles, the *cis* isomer of lycopene is superior to the *trans* isomer in solubility, and after entering the intestinal cells through passive diffusion and cholesterol membrane transporters (scavenger receptor class B type I, SR-BI) [26, 27], the *cis*-lycopene isomer preferentially merges into chylomicron particles, enters the lymphatic system, and is released into the bloodstream [28, 29]. Due to the lipophilic nature of lycopene, the concentration of lycopene in adipose tissue is higher than in the serum [30]. In the serum, lycopene is concentrated in low-density lipoproteins (LDL) and very low-density lipoproteins (VLDL) [31]. The testis, adrenal glands, liver, and prostate are also storage sites for lycopene [32, 33]; in these tissues and organs, the concentration of *cis*-lycopene is mildly higher than that of the *trans* isomer.

2.2. Biological Functions of Lycopene

2.2.1. Biological Effects of Lycopene. Although lycopene is not an indispensable nutrient for humans, it is still a hot topic in health care. This is mainly because lycopene is ubiquitous in our daily diet and has a significant biological effect on the human body. Multiple kinds of studies (e.g., epidemiological studies, animal studies, human clinical studies, and intervention studies) have substantiated that lycopene has a biological influence on almost every organ and tissue. Moreover, many studies have shown that lycopene administration can prevent and treat many diseases, including cancer [34], cardiovascular disease [35], osteoporosis [36], male infertility [37], obesity and diabetes [38], neurodegenerative diseases [39], rheumatoid arthritis [40], skin [41], and retinal diseases [42] (Figure 2). Because lycopene does not have a β -ion ring structure, it does not have provitamin A activity [43]. Instead, lycopene exerts its biological effects primarily through antioxidation and non-antioxidation. As a highly effective antioxidant, lycopene can decrease the risk of disease by neutralizing reactive oxy-

gen species (ROS) and reducing oxidative damage of some biomolecules (DNA, proteins, and lipids) [44, 45]. In addition, lycopene has also acted through other mechanisms, including anti-inflammation [40], autophagy [46, 47], apoptosis [48, 49], proliferation [50, 51], and pyroptosis [52].

2.2.2. Dietary and Safety. Apart from dietary intake, lycopene had been sold over the counter as a health-care product to prevent diseases. There are different opinions on the recommended dosage and administration time of lycopene for different health purposes. The 67th meeting of the Joint Committee of Experts of the World Health Organization and the Food and Agriculture Organization of the United Nations (FAO) established a permissible daily intake of lycopene of 0–0.5 mg/kg b.w. in healthy people. However, more lycopene may be needed to prevent some diseases. One human study suggested that intake of 6.5 mg lycopene reduced prostate cancer risk in men [53]. Another research reported that increasing the intake of lycopene to 7 mg/d improved the endothelial function of patients with cardiovascular disease [54]. Besides, side effects should be considered when using lycopene to prevent diseases. An animal study testing the safety of two synthetic lycopene products at different concentrations (0, 500, 1500, and 3000 mg/kg b.w./d) found that rats ingesting lycopene at the highest concentration (3000 mg/kg b.w./d) for 13 weeks did not develop any adverse effects [55]. Few side effects have been reported in humans unless a large quantity of tomatoes or lycopene are ingested. Meanwhile, one study reported that after ingesting two liters of tomato juice per day for several consecutive years, one 61-year-old woman developed lycopenemia, but no other adverse events were observed [56]. Three weeks after she stopped ingesting lycopene, the orange color of her skin faded. Although supplementation with a low-dose of lycopene might yield beneficial effects (such as preventing preeclampsia [57]), it should be used carefully by women who are pregnant or lactating. In summary, lycopene is a relatively safe functional ingredient at most doses.

3. The Effects of Lycopene on Ischemic and I/R Injuries

3.1. The Effects of Lycopene on Myocardial Ischemic and I/R Injuries. The primary cause of global human death is ischemic heart disease which has attracted significant research attention [58]. Ischemic heart disease is brought about by several reasons, including thrombosis, coronary artery stenosis, and excessive contraction of the coronary artery [59]. Myocardial ischemia leads to inadequate oxygen and energy for supporting normal heart function. Besides, long-term or severe cardiac ischemia can result in arrhythmia, myocardial infarction, and sudden death. Mechanical and drug interventions, such as angioplasty, thrombolytic therapy, and coronary artery bypass grafting, can quickly restore blood flow to the ischemic site [60–63]. However, these treatments not only fail to reduce the area of the infarction but can cause more serious damage to the ischemic site (increasing area of infarction and death such as through I/R injury) [64, 65]. Myocardial I/R injuries are extremely

enzymes) by the presence of comorbidities (e.g., myocardial hypertrophy, diabetes, and metabolic syndrome [82–86]). Increased doses of antioxidants should be considered to achieve a better therapeutic effect in these patients. Thus, when lycopene is given as an adjunctive therapeutic measure, it is essential to identify the best dose and time of administration for this drug.

3.1.1. Lycopene Attenuates Myocardial Ischemic Injury. Prior epidemiological studies reported that lycopene might exert a protective effect against myocardial infarctions (MI). According to a cross-sectional study [87], the level of lycopene in serum was negatively related to the risk of an acute MI in men. Meanwhile, according to the case-control study, it could be found that in the adipose tissues, higher levels of lycopene were significantly correlated with a lower risk of MI [88]. Because lycopene concentration in adipose tissues can better reflect its long-term accumulation than serum concentration, this result more reliably and accurately reflects the correlation between high levels of lycopene and a low risk of MI. Further, it also hints that maintaining high levels of lycopene *in vivo* may prevent myocardial ischemia. In addition, lycopene may also lower the risk factors for myocardial ischemia. An epidemiological study found that in the early stages of atherosclerosis, serum lycopene concentration was negatively associated with common carotid intima artery media thickness [89]. Based on these studies, it can be found that high levels of lycopene in the body appear to not only directly prevent MI but also reduce the impact of one of its primary risk factors (atherosclerosis). However, it is worth doing more clinical studies to explore the specific effects and related mechanisms of lycopene in preventing MI.

Some recent studies using animal ischemic models showed that supplying lycopene before and after ischemia could improve the consequences of an ischemic MI (Table 1). It is well known that adrenaline and isoprenaline (ISP) are usually used to induce animal myocardial ischemia. Adrenaline and ISP produce free radicals, inducing MI through autooxidation and positive inotropic and chronotropic effects. Moreover, this MI model better simulates human MI pathophysiology, biochemistry, morphology, functional changes, and histopathology [90–92]. The cardioprotective effects of lycopene in this MI model were related to anti-inflammation (reducing IL-1 β , IL-6, TNF- α , and NF- κ B); antioxidation via increasing antioxidant enzymes activities (SOD, GPx, CAT, and GSH) and decreasing the level of ROS; antiapoptosis by increasing Bcl-2 and reducing Bax, Cyt-c, caspase-3, caspase-8, caspase-9, the ratio of Bax/Bcl-2, and JNK/(ERK1, STAT1,3); promoting repair (inhibiting eight miRNAs and increasing IGF-1 and SIRT-1); and reducing remodeling (inhibiting p38/MMP-9) [93–99]. Besides, other studies have shown that quercetin can be used in combination with other antioxidants to protect against ischemic injury [100, 101]. The combined application of lycopene and quercetin improved myocardial ischemic injuries in a rat model [102]. In contrast with quercetin treatment (100 mg/kg b.w.) alone [103], lower doses of quercetin administration (80 mg/kg) combined with low-

dose lycopene (3 mg/kg) could achieve similar protective effects (restoring the ischemic injury of myocardial to normal status). This combination could also shorten administration time (10 d) compared with quercetin alone (30 d). Therefore, in-depth exploration of additional synergies between lycopene and other agents should be pursued in order to achieve the best therapeutic effects.

3.1.2. Lycopene Attenuates Myocardial I/R Injury. Many studies have discussed lycopene in protective effects on myocardial I/R injury. The only published randomized controlled clinical study reported that after standard MI treatment (giving aspirin, statin, and β -blocker), lycopene administration (30 mg 12 hours before PCI and 15 mg before and 8 hours after the procedure) significantly reduced the serum CK-MB levels of 28 patients [104]. The findings showed that lycopene had the potential to prevent cardiovascular events after a myocardial I/R injury. However, lycopene supplementation had no significant effect against increases in serum TnI and hs-CRP levels. This was counter to the findings of other studies, which showed that lycopene could lower CRP levels in both heart failure patients and healthy individuals [105, 106]. These contrasting results may be because the randomized trial used lower dose of lycopene for a shorter period.

Many studies (*in vivo* and *in vitro*) using animal I/R models showed that lycopene supplementation before and after ischemia could significantly improve myocardial I/R injuries (Table 2). Relevant mechanisms include antioxidation (inhibiting ROS production; increasing GSH, SOD, CAT, and GSH-Px activities; and activating AMPK-ACC pathway), enhanced autophagy (improving p-AMPK, MAP1LC3B, and Beclin 1), anti-inflammatory effects (reducing IL-6, TNF- α , and IL-1 β and weakening JNK pathway), ER stress inhibition (reducing the mRNA expression of eIF2 α , GRP78, CHOP, ATF6, caspase-12, and sXbp-1 and the phosphorylation of AMPK and eIF2 α), and reduced mitochondrial apoptosis (increasing Tfam, Bcl-2, and the phosphorylation of Akt and ERK1/2; inhibiting the opening of mPTP; and decreasing the level of Cyt-c, Bax, APAF-1, cleaved caspase-3 and caspase-9 and p-GSK-3 β , and the ratio of Bax/Bcl-2,). Besides, it could be found that these protective effects were produced by long-term low-dose administration or a single high-dose lycopene. Although lycopene dosage in this study [107] was too high to unsuitable for clinical applications compared with the international intake standard of lycopene for healthy people (0–0.5 mg/kg/d b.w.) [53, 54], the study did not find any toxicity from the use of higher concentrations of lycopene. However, whether such concentrations could be used clinically requires further investigation. Further, some studies showed that high cholesterol not only exacerbated myocardial damage but also affected the therapeutic effects of PostC [108–112]. The combination of PostC and a certain dose of lycopene could significantly improve severe myocardial I/R injuries in patients with high cholesterol [113], justifying further exploration of potential combination therapies in myocardial I/R injuries under other conditions (e.g., diabetes and hyperlipidemia [114, 115]).

TABLE 1: Effects of lycopene on myocardial ischemic infarction.

Species	Ischemic models	Treatments	Effects and mechanisms	Ref
BALA/c mice	Ligating LAD	Lycopene (10 mg/kg/d, i.g.) for 4 weeks after ischemia	Relieving myocardial apoptosis and inflammation via reducing the expression of TGF- β 1, IL-1 β , NF- κ B TNF- α , collagen I, collagen III, caspase-3, caspase-8, and caspase-9	[93]
Wistar rats	Coronary artery ligation	Lycopene (1 mg/kg/d) for 3 months after ischemia	Attenuating cardiac remodeling and improving diastolic dysfunction via \downarrow Nrf-2, TNF- α and eight miRNAs, \uparrow IGF-1R, and SIRT-1	[94]
Wistar rats	At intervals of 24 h, ISP (200 mg/kg, s.c.) for 2 d	Lycopene (10 mg/kg/d, p.o.) for 30 d after ischemia	Reducing infarction area and preventing DNA damage and oxidation of enzyme SH groups via scavenging ROS, \downarrow caspase-3, iNOS, NO, CRP, and MPO	[95]
Sprague-Dawley rats	Ligating LAD	Lycopene (40 mg/kg/d, i.g.) for 28 d after ischemia	Attenuating ventricular remodeling and LV dilation, improving LV systolic, and cardiac function via inhibiting p38/MMM-9 and the type I collagen	[96]
Wistar rats	ISP (85 mg/kg, s.c.) at 28 and 29 d	Lycopene (0.5, 1.0, and 1.5 mg/kg, i.g.) for 30 d	Improving hemodynamics and disorder of the cell morphology and reducing cardiac dysfunction and via \uparrow SOD, CAT, GSH-Px and GSH, \downarrow MDA, CK-MB, and LDH	[97]
Sprague-Dawley rats	ISP (85 mg/kg, s.c.) for 2 d	Lycopene (4 and 6 mg/kg) for 7 d after ischemia	Reducing disorder of the cell morphology and apoptosis via \uparrow SOD, CAT, GSH and GPx, \downarrow IL-6,1 β , TNF- α , Bax/Bcl-2, JNK/(ERK, STAT1,3), Cyt-c, and caspase-3	[98]
Sprague-Dawley rats	ISP (100 mg/kg, i.p.) at 7 and 8 d	Lycopene 3 mg/kg and quercetin 80 mg/kg p.o. for 1-10 d	Reducing cardiotoxicity (\downarrow CK-MB, LDH, MYO, and TROP) and oxidative stress (\downarrow MDA) via \uparrow GST μ , SOD1, SOD2, CAT, and Bcl-2	[102]

Furthermore, one study found that the cardioprotective effects of IP and PostC were activated through the posttranslational modification of proteins (S-nitrosylation, SNO) mediated by RISK and SAFE [116]. Most of the evidence suggests that as an adjunctive therapeutic measure, lycopene acts on many downstream signal molecules of RISK and SAFE (e.g., activating ERK, Akt and PI3K and inhibiting STAT3 and p-GSK-3 β) to attenuate myocardial I/R injury [113]. These studies implied that lycopene seemed to promote SNO. Moreover, the activity of antioxidant enzymes (GSH and SOD) could be improved by lycopene's cardioprotective effects [97]. These antioxidant enzymes could also limit protein SNO levels through denitrosylation, thereby preventing excessive SNO from inducing cytotoxicity in the pathological environment created by nitrosative stress [117–119]. Therefore, lycopene may promote SNO and inhibit the production of excess SNO to alleviate myocardial I/R injury. Nevertheless, further studies are still required on this topic.

3.2. Lycopene Attenuates Liver I/R Injury. In a variety of clinical settings (e.g., hemorrhagic shock, hepatic resection, liver transplantation, and trauma [128–130]), liver I/R injury is prone to occur. Liver I/R injury not only damages the liver itself but also triggers the remote organ dysfunction to the lungs [13], kidneys [14], myocardium [15], and intestines [131]. This effect has made liver I/R injury prevention a hot research topic in recent years. As the liver is the main storage site of lycopene, it is highly likely that lycopene acts on the liver. Many studies have shown that lycopene could

effectively impact liver-related diseases, including cirrhosis [132], nonalcoholic fatty liver [133], and hepatocellular carcinoma [134].

A recent epidemiological study compared changes in the plasma antioxidant levels (vitamin A, vitamin E, β -carotene, and lycopene) of the 12 patients who underwent liver transplantation (liver I/R) at three stages of their transplant (pre-hepatic, anhepatic, and reperfusion) [135]. The levels of lycopene and β -carotene declined much faster than those of vitamin A and vitamin E. This decrease may be related to the reduction or disappearance of the liver's ability to store lycopene in the setting of liver disease. Lipid peroxidation also increased during reperfusion, possibly due to the almost complete depletion of lycopene in the plasma, and hints that lycopene may be essential to maintaining oxidative balance during liver transplantation. It is therefore possible that lycopene supplementation during hepatic I/R injury (e.g., liver transplantation) could significantly reduce oxidative damage during I/R.

Prior animal studies showed that lycopene pretreatment could effectively mitigate hepatic I/R injury by enhancing the Kupffer cell autophagy (activating Nrf-2/HO-1) and inhibiting inflammation (decreasing IL-6, TNF- α , IL-1 β , and NLRP3 inflammasomes and increasing IL-10), oxidative stress (reducing ROS and increasing CAT), and apoptosis (increasing Bcl-2 and decreasing caspase-1 and caspase-3) (Table 3). One study showed that apart from CAT, ALT, AST, LDH, and MDA levels were reduced by lycopene pretreatment in a dose-dependent manner [136]. According to

TABLE 2: Effects of lycopene on myocardial I/R injury.

Species	I/R Models	Treatments	Effects and mechanisms	Ref
Wistar rats	I/R (30 min/120 min)	Lycopene (40 mg/kg/d, i.p.) for 5 d before I/R	Reducing myocardial infarction area, apoptosis, and the opening of mPTP via ↓APAF-1, caspase-3, caspase-9, Bax, Bax/Bcl-2, Cyt-c, and ↑Bcl-2	[120]
Wistar rats	I/R (30 min/60 min) with HCD	Lycopene (20 mg/kg/d, i.p.) for 5 d before I/R	Decreasing myocardial infarction area and apoptosis and improving cardiac functions via ↓GRP78 and CHOP, Cyt-c, caspase-3, caspase-9, p-GSK-3β and ↑p-Akt, and p-ERK1/2	[113]
C57BL/6 mouse; HL-1cell	LAD I/R (20 min/4 h); H/R (2 h/2 h)	Lycopene (33.5 mg/kg* b.w., i.v.) after ischemia; lycopene (1, 2, and 4 μM) after hypoxia	Reducing myocardial infarction area and apoptosis via inhibiting the level of p-JNK and ROS and the activation ERK1/2	[107]
Sprague-Dawley rats	Left coronary artery I/R (30 min/24 h)	Lycopene (1.5 mg/kg/d, i.g.) for 30 d before I/R	Reducing myocardial infarction area, apoptosis, and mtDNA damage via ↓MDA, Cyt-c, caspase-3, ROS level, and ↑Tfam	[121]
Wistar rats	LAD I/R (45 min/1 h)	Lycopene (1 mg/kg, i.g.) for 31 d before I/R	Reducing oxidative stress and disorder of the cell morphology via ↑GSH and GSH-Px level and ↓MDA	[122]
Sprague-Dawley rats	I/R (30 min/2 h)	Lycopene (0.088 mg/d, i.g.) for 3 w before I/R	Improving ventricular functions, reducing myocardial infarction area and apoptosis via ↓MDA and ROS	[123]
H9C2 cardiomyocytes	H/R (4 h/4 h)	Lycopene (10 μM) for 4 h before H/R	Reducing ER stress and apoptosis via inhibiting GRP78, CHOP, caspase-12, p-JNK, and LDH	[124]
H9C2 cardiomyocytes	H/R (16 h/2 h)	Lycopene (2.5 μM) for 48 h before H/R	Increasing autophagy and reducing apoptosis via ↓Bax/Bcl-2, caspase-3, ↑Beclin 1, MAP1LC3B, and p-AMPK	[125]
C57BL/6 mouse cardiomyocytes	H/R (4 h/6 h)	Lycopene (5 μM) for 4 h before H/R	Reducing ER stress and apoptosis via ↑AMPK, ↓caspase-3, caspase-12, CHOP, Bax/Bcl-2, sXbp-1, eIF2α, GRP78, ROS, and ATF6	[126]
C57BL/6 mouse cardiomyocytes	H/R (4 h/8 h)	Lycopene (5 μM) for 4 h before H/R	Reducing apoptosis via inhibiting mPTP opening and decreasing caspase-3 and Cyt-c	[127]

TABLE 3: Effects of lycopene on hepatic I/R injury.

Species	I/R Models	Treatments	Effects and mechanisms	Ref
Sprague-Dawley rats	The portal vein, the bile duct, and the hepatic artery I/R (45 min/60 min)	Lycopene (2.5 and 5 mg/kg, i.p.) at 60 min before I/R	Reducing liver damage via ↑CAT and ↓ROS, ALY, AST, LDH, and MDA	[136]
C57BL/6 mice	The portal vein and hepatic artery I/R (90 min/6 h)	Lycopene (20 mg/kg, i.g.) for 2 weeks before I/R	Inhibiting NLRP3 inflammasome and apoptosis and enhancing autophagy via ↑Bcl-2, LC3B, IL-10, ↓TNF-α, IL-6, MPO, IL-1β, caspase-1, caspase-3, and p62 and activating Nrf2/HO-1	[137]
AML12 hepatic cells	H/R (12 h/4 h)	Lycopene (10 μM) for 4 h before H/R	Reducing apoptosis and oxidative stress via ↓TNF-α, IL-6, MDA, and ROS and activating Nrf2/HO-1	[138]
Wistar rats	H/R (45 min/2 h)	Lycopene (25 mg/kg/d, p.o.) for 14 d before H/R	Relieving edema and reducing oxidative stress via reducing ALT, AST, LDH, and MDA	[139]

these results, the protective effects of lycopene on hepatic I/R injury might be associated with the antioxidant capacity of lycopene or the activation of the Nrf2 pathway [137].

Some studies are worth exploring whether lycopene pre-treatments can improve hepatic I/R injury-induced damage to other organs.

3.3. Lycopene Attenuates Kidney I/R Injury. Some urological procedures (e.g., kidney transplantation, partial nephrectomy, and renal artery surgery) [140] and crisis situations (trauma, hemorrhagic shock, and sepsis) [141] can induce renal ischemia and acute kidney injury. Subsequent reperfusion induces neutrophil activation and the release of ROS and other inflammatory mediators [142, 143], leading to decreased tubular and glomerular function and further exacerbates kidney damage [144]. Exploring effective measures to reduce renal I/R injury is critical.

Encouragingly, some studies have shown that lycopene pretreatment could effectively improve the histopathologic and biochemical changes induced by renal I/R injury via antioxidation and anti-inflammation. A recent study found that before renal I/R injury (30 min/2 h), intake of lycopene (at 30 min before ischemia, 10 mg/kg, i.p.) reduced the levels of urea and creatinine in the serum and improved pathologic changes to renal tissues and cells by inhibiting the Notch2/Hes-1 pathway; decreasing Bax, IL-6, and TLR-2; and increasing Bcl-2 [145]. Another study showed that in the renal I/R injury (45 min/6 h), lycopene pretreatment (at 30 min before ischemia, 100 mg/kg, i.g.) reduced serum MDA levels and histopathologic changes [146]. However, serum BUN and creatinine levels, tissue MDA level, and the GSH levels in the serum and tissues did not significantly change. An additional study showed that the before kidney I/R injury (45 min/24 h) [147], lycopene supplementation (4 mg/kg/d, for 2 d before ischemia) significantly improved sodium, GSH, and CAT levels in addition to pathology scores. However, it did not change the levels of other indices (potassium, creatinine, BUN, MDA, SOD, and GSH-Px). Based on these studies, the duration of ischemia appears to affect lycopene in renal I/R injury protective effects beyond dose and pretreatment length. In view of lower-dose lycopene pretreatment in renal I/R injuries protective effects only following short-term ischemia, higher-dose regimens before and after renal I/R injury following long-term ischemia should be evaluated.

3.4. Lycopene Attenuates Cerebral Ischemic and I/R Injuries. Cerebral ischemia occurs in the setting of brain damage, asphyxia, and cardiac arrest [148]. During cerebral ischemia, the release of cytokines, mitochondrial respiratory enzymes damage, programmed cell death induction, and microglia activation can lead to irreversible brain damage [149–151]. Although timely reperfusion can mitigate these injuries by arresting these pathologic changes, activated inflammation and oxidative stress after reperfusion destroy the blood-brain barrier and induce the death of neuronal [152]. Because lycopene can cross the blood-brain barrier and directly act on neurons [32], lycopene has been used to improve cerebral disease effects (subarachnoid hemorrhage [153], Alzheimer's disease [154], memory impairment [155], and central and peripheral neuropathy [156]). Recent studies have shown that lycopene can not only improve cerebral ischemia but also reduce cerebral I/R injury.

Only one epidemiological investigation and one *in vitro* study have explored the protective effects of lycopene against cerebral ischemic injuries. In a prospective follow-up study,

Karppi et al. [157] found that the high serum concentrations of lycopene were closely associated with a reduction in the risk of ischemic stroke in men. It suggested that lycopene might play a vital role in preventing ischemic stroke. *In vitro*, the trial found that lycopene pretreatment (0.5, 2.0, and 8.0 $\mu\text{mol/L}$, 1 h) could prevent oxygen-glucose deprivation- (OGD-, 3, 6, 12, and 24 h) induced SH-SY5Y cell death by reducing autophagy mediated by activated AMPK/mTOR following the inhibition of oxidative stress (increasing xCT, cysteine, and GSH levels and decreasing the levels of ROS and MDA) [158]. It is worth conducting additional clinical investigations and animal studies to explore the specific protective effects and mechanisms of lycopene against cerebral ischemic and I/R injuries.

Several animal studies showed that lycopene supplementation before and after ischemia and after reperfusion could reduce neuronal apoptosis and cerebral infarction area (Table 4). Lycopene's mechanism of action is related to anti-inflammation (inhibiting IL-6/JAK-STAT3), antioxidation (increasing SOD, GSH, and CAT activities and the mRNA expression of HIF-1 α ; reducing ROS, iNOS, and NO; and activating Nrf-2/HO-1), antiapoptosis (reducing caspase-3, JNK/MAPK, and Bax and increasing Bcl-2), and lowering iron content (raising FNPI and decreasing hepcidin and L-ferritin). Nanoliposome-encapsulated lycopene (L-LYC) was more effective at preventing a brain I/R injury than naked lycopene [159]. This might be because L-LYC more easily crosses the blood-brain barrier, thereby increasing lycopene content in the brain. Among currently known nanoparticles, gold nanoparticles (GNPs) have been of particular interest because of their unique surface chemical properties, excellent biocompatibility, and potential for extensive chemical modification [160, 161]. Some studies found that GNPs played an essential role in treatments (20 nm GNPs remarkably exhibited neuroprotective effects, while 5 nm GNPs aggravated neuronal damage) [162] and diagnosis (GNPs recognized the infarct area of an ischemic stroke by combining with various biological molecules such as hyaluronic acid) [163]. Further, modifying GNPs with some biomolecules could transport nanoparticles to specific locations [164]. To permit the accurate transport of the appropriate dose of lycopene to sites of infarction and improve its therapeutic effects, it may be possible to modify GNPs with some biomolecules (directionally transported to the brain) as carriers instead of using naked GNPs. However, it is important to consider whether modifying GNPs with biomolecules will affect their diagnostic and treatment potential.

3.5. Lycopene Attenuates Reproductive Organ I/R Injuries. The ischemia of the reproductive organs largely occurs in the testicles of males and the ovaries of females. Testicular torsion is the primary cause of testicular ischemia and commonly occurs in neonates and adolescents [169, 170]. Ovarian ischemia is usually caused by ovarian torsion, surgery, and pregnancy [171]. Long-term ischemia of the testis and ovaries can lead to cell damage and declined function, resulting in hormone secretion disorders, low fertility, and infertility [172]. Severe pain due to torsional ischemia requires a

TABLE 4: Effects of lycopene on cerebral I/R injury.

Species	I/R models	Treatments	Effects and mechanisms	Ref
Mongolian gerbils	Bilateral common carotid artery I/R (10 min/3 h)	Lycopene (5 mg/100 g, in diet) after ischemia	Inhibiting apoptosis via increasing Bcl-2 and SOD, reducing caspase-3 and Bax	[165]
C57BL/6 mice	Bilateral common carotid artery I/R (20 min/24 h)	Lycopene (20 mg/kg, i.p.) for 7 d before I/R	Improving neurological function scores, reducing oxidative stress and apoptosis via reducing ROS, and increasing GSH and Nrf2/HO-1	[166]
Sprague-Dawley rats	Middle cerebral artery I/R (2 h/24 h)	Lycopene (5 and 20 mg, i.g.) for 15 d before I/R	Reducing infarct area and nerve function defect via \uparrow SOD, CAT, Bcl-2, HIF-1 α , \downarrow iNOS, and MDA	[167]
Sprague-Dawley rats	Middle cerebral artery I/R (60 min/7 d)	L-LYC (6 mg/d, i.g.) for 14 d before I/R	Reducing infarct area and nerve function defect via reducing iron content (\downarrow hepcidin and L-ferritin and \uparrow FPN1), oxidation (\uparrow SOD, CAT, GSH, \downarrow ROS, iNOS, and NO), apoptosis (\uparrow Bcl-2, \downarrow caspase-3, and JNK/MAPK), and inflammation (\downarrow IL-6 and p-STAT3)	[159]
Wistar rats	Middle cerebral artery I/R (1 h/24 h)	Lycopene (4 mg/kg, i.v.) twice at 15 min before ischemia and reperfusion	Reducing infarct area via inhibiting microglia activation, the production of NO, and lipid peroxidation and scavenging ROS	[168]

timely operation to detorsion of the organ, relieve the patient's pain, and restore blood flow. Although reperfusion is highly necessary to the survival of ischemic tissues, reperfusion also aggravates damage to these reproductive organs. It is therefore equally important to restore blood supply to reverse the ischemic injury and to take measures to reduce reperfusion injury.

Fortunately, current studies had shown that lycopene treatment could protect the reproductive organs from I/R injury. Following testicular I/R injury (1 h/3 h, 24 h, 30 d), lycopene intake (4 mg/kg/d, i.g.) after 5 min of ischemia effectively improved bilateral testicular sperm motility and decrease oxidative stress [173]. However, recent research [174] showed that lycopene supplementation (20 mg/kg/d, i.p.) at 2 h after the ischemia could only reduce germ cell apoptosis following a short-term testicular I/R injury (2 h/3 d) and did not protect against long-term I/R injury (2 h/10 d). According to these studies, it can be found that for short-term ischemia, the supplement with lycopene significantly alleviates testicular I/R injury. However, even increasing lycopene concentration failed to improve the damage caused by long-term ischemia. Furthermore, the longer the testicular reperfusion time, the worse the therapeutic effect of lycopene. This suggests that timely administration of high-dose lycopene can be used as an adjunctive therapy against testicular I/R injury. Moreover, it also suggests that an evaluation about the benefits of lycopene pretreatment against testicular I/R injury is warranted. Further epidemiological investigations should focus on whether patients have milder symptoms of I/R injury after taking more lycopene.

Besides, lycopene has also been shown to improve ovarian I/R injury. One study [175] showed that at 2.5 h after ischemia (3 h), lycopene treatment (100 and 200 mg/kg, i.g.) could effectively reduce tissue congestion, hemorrhage, and polymorphonuclear leukocyte infiltration via increasing GSH and CAT and lowering MDA and MPO. Moreover, the protective effects of lycopene were dose-dependent. Another

study showed that lower doses of lycopene (before 1 h of reperfusion, 2.5 and 5 mg/kg, i.p.) could reduce ovarian tissue degeneration, vascular congestion, inflammatory cell infiltration, and edema [176]. However, there was no difference in other pathological indexes (MDA and p-NF- κ B) between the two different doses. The differing results of these dose-effect studies are likely due to different dose designs. On one hand, the dose of the former was higher than that of the latter. On the other hand, the dose difference in the former study was also relatively larger, which may result in significantly different effects. Further studies are therefore needed to identify the lowest doses of lycopene required to achieve the best therapeutic effects in different I/R circumstances.

3.6. Lycopene Attenuates the Intestinal Tract I/R Injury.

Intestinal ischemia is often seen during the progression of some diseases, for example, volvulus [177], acute mesenteric ischemia [178], neonatal necrotizing enterocolitis [179], trauma [180], and intestinal transplant rejection [181]. As a pathologic process, intestinal ischemia can cause intestinal mucosal barrier and bacterial translocation (BT) [182]. With prolonged ischemic time, intestinal ischemia can lead to irreversible intestinal tissue necrosis and serious systemic metabolic derangement [182]. Although reperfusion can repair some mild injuries and prevent irreversible damage [9], it can aggravate intestinal injuries, leading to shock, multiple organ dysfunction, and sepsis. Although new therapies have been proposed, the morbidity and mortality from intestinal I/R injuries remain high [183]. The study about lycopene in effects on intestinal I/R injury had just begun. A prior research showed that lycopene pretreatment (12 h before ischemia, 0.4 mg/kg, i.g.) could significantly protect the intestinal villi structure from necrosis, degeneration, and spillage of the epithelium, thereby reducing IgA and sIgA loss and limiting BT [184]. These effects were related to anti-inflammation and the maintenance of the tight

junction via activation of retinoic acid receptor- (RAR-) mediated transcriptional pathways. However, lycopene intake (0.4 mg/kg, i.g.) 1 h after reperfusion could not protect the intestine from I/R injury (30 min/12 h). Those results suggest that low-dose preconditioning using lycopene could effectively prevent intestinal I/R injury, but treatment after reperfusion might require higher doses to achieve a therapeutic effect. Moreover, the intestinal IgA, as everyone knows, plays a crucial role in mucosal immunity and infection by binding with pathogens to arrest them from crossing the epithelial barrier [185, 186]. According to the result that lycopene could maintain the intestinal IgA level as shown in that prior study, it may be useful for alleviating gastrointestinal symptoms (anorexia, nausea and vomiting, and diarrhea) and impaired nutrient absorption [187, 188] in patients suffering from intestinal mucosal and gut microbiome dysfunction following COVID-19 infection. However, in order to confirm this hypothesis, more research is needed.

3.7. Lycopene Attenuates Retinal I/R Injury. Retinal ischemia occurs during many pathologic processes, including glaucoma, diabetic retinopathy, and retinal vascular obstruction [189, 190]. As a high oxygen-consuming organ, the retina is extremely sensitive to ischemia [191]. Mild ischemia increases the retina's vascular permeability, leading to edema. With increasing severity, the retinal injury gets worse, causing retinal ganglion cell apoptosis, reduced vision, and blindness [192, 193]. Timely recovery of blood flow is particularly vital to reverse a retinal ischemic injury. However, reperfusion can cause a series of pathologic problems (e.g., oxidative stress, inflammation, and apoptosis) that can exacerbate the damage of ischemic retinal [194–196]. Studies have shown that lycopene may have potential applications in prophylaxis and therapy against cataract and diabetic optic neuropathy through its antioxidative and anti-inflammatory effects [197, 198]. A recent study explored the effects and mechanisms of lycopene on retinal I/R injuries *in vivo* and *in vitro* [42]. *In vivo*, it was found that lycopene treatment (40 μ M, injected into the ocular bulbs) before retinal I/R injury (3 h/24 h) could prevent cell morphologic changes and reduce apoptosis by activating the KEAP1/Nrf2/ARE pathway. Besides, *in vitro*, different doses of lycopene (10 μ M and 20 μ M, for 24 h) culturing cells before OGD/reperfusion (40 min/2 h, 4 h) could reduce ROS and MDA and increase cell viability by enhancing antioxidant enzyme activity (GSH, CAT, and SOD) and activating the KEAP1/Nrf2/ARE/(HO-1, NQO-1) pathways. While low-dose lycopene treatment had no obvious protective effects against long-term retinal I/R injury, high-dose lycopene had significant protective effects against both short- and long-term I/R injuries. Higher doses of lycopene are therefore required to reverse retinal I/R injuries.

3.8. Lycopene Attenuates Spinal Cord I/R Injury. Spinal I/R injury occurs mostly after thoracic aorta, common thoracic, and abdominal artery surgeries [199]. One of the most serious complications of these surgeries is spinal I/R injury, which has a very poor prognosis [200]. Spinal I/R injury can seriously impair nerve function and lead to paraplegia

[200]. One survey showed that the incidence of a spinal I/R injury during thoracic aortic surgeries worldwide was approximately 1 in 10 [201]. However, the therapeutic effects of currently known interventions (e.g., cerebrospinal fluid drainage [202], electro-needle pretreatment [203], drug administration [204], multifunctional hepatocyte use [205], remote ischemia [206], and hyperbaric oxygen pretreatment [207]) are unsatisfactory. Therefore, it is urgent to find better interventions for preventing spinal I/R injury. A recent study showed that after a spinal I/R injury (40 min/48 h) [208], lycopene (25 and 50 mg/kg/d, for 14 d, i.g.) could significantly improve neuroprotection and reduce motor dysfunction, apoptosis (increasing the neuron survival rate), and inflammation (decreasing IL-8, TNF- α , IL-1 β , and IL-6). These effects were related to the downregulation of COX-2 expression via HO-1 activation. Besides, the protective effects of the high-dose lycopene were significantly better than those of low-dose lycopene. Although the study showed that lycopene could lessen spinal I/R injury, lycopene pretreatment was preferred. Further studies should therefore focus on the lycopene pretreatment in protective effects against spinal I/R injuries.

3.9. Lycopene Attenuates Limb I/R Injury. Limb muscle ischemia often occurs in some clinical settings, including thrombosis [209], limb surgical revascularization, trauma [210], and free flap reconstruction [211, 212]. During early ischemia, the limb gets cold, numb, and painful. As the ischemia time continues, the limb's skin color changes to cyan/black and skeletal muscle necrosis occurs. Further worsening of ischemic injuries can lead to amputation and death. Common reperfusion treatments include intravascular interventional surgery [213], bypass surgery [214], hybrid reconstruction [215], and cell therapy [216]. However, reperfusion leads to metabolic waste that accumulated in the ischemic limb reentering the systemic circulation, which may result in additional organ injury, multiple organ failure, and death [217]. One previous study showed that after strenuous exercise, lycopene could alleviate oxidative stress to protect the skeletal muscle by reducing xanthine oxidase and myeloperoxidase activity [218]. It is therefore possible that lycopene could reduce skeletal muscle I/R injury. A previous study showed that lycopene (10 mg/kg/d, for 15 d, p.o.) intake prior to rat limb muscle I/R injury (2 h/2 h) [219] significantly reduced muscular atrophy, perivascular inflammation, and focal central muscle necrosis by decreasing MDA and IMA and increasing GSH-Px and SOD activities. In the serum and muscle tissue, MDA levels both decreased significantly, and the average level of MDA in the muscle was lower than that in the serum. This suggests that the antioxidant effects of lycopene on skeletal muscle were better than in the serum. However, it did not impact lycopene's positive impact on other organs (even distant organs [220, 221]) injured following a limb I/R injury. In a recent study, the results showed that lycopene pretreatment (10 mg/kg/d, for 15 d, p.o.) could prevent lung injury (induced by skeletal muscle I/R injury, 2 h/2 h) via increasing SOD and GPX and reducing MDA [222]. It can be hypothesized that lycopene may prevent single organ I/R

injuries from affecting other organs. However, it is unclear whether the protective effects and mechanisms of lycopene are the same in different organs. Before clinical application, further research is necessary.

3.10. Lycopene Attenuates Ischemic Injury to Mesenchymal Stem Cells. As a kind of multifunctional stem cells, mesenchymal stem cells (MSC) can differentiate into various types of terminal cells, including stem cells, cardiomyocytes, and neurons. So MSC transplantation has been used to treat tissue damage such as limb ischemia [223], acute kidney injury [224], liver fibrosis [225], ischemic heart failure [226], myocardial infarction [227], and the I/R injuries to the heart [228], kidney [229], liver [230], spine [231], and brain [232]. However, the therapeutic effects of MSCs are limited by their low implantation rate, low survival rate, impaired paracrine ability, and delayed administration [233]. Moreover, previous studies showed that the low survival rate of MSCs was due to oxidative stress, apoptosis, and autophagy caused by ischemia [234–236]. Therefore, it seems feasible to improve the implanted cells survival rate through protecting MSC from oxidative stress, apoptosis, and autophagy. A study showed that before hypoxia (6 h), precultured cells with lycopene (20 μ M, for 1 h) could activate the PI3K/Akt pathway to inhibit the production of ROS and reduce apoptosis [237]. Additionally, in a different ischemia model (low glucose and H₂O₂ treatment, 1, 2, 3, and 4 h), lycopene pretreatment (10 μ g/mL, for 30 min) significantly inhibited oxidative stress (reducing ROS levels and activating Akt-MnSOD) and apoptosis (decreasing the levels of p-p38, p-ATM/p53, Bax, and p-JNK and increasing Bcl-2) [238]. According to these studies, it can be found that lycopene may protect MSCs from ischemic injuries. However, there are currently no studies that evaluate lycopene in the protective effects on MSC I/R injury. Therefore, it is essential to further explore the effects and mechanisms of lycopene in the prevention and treatment of MSC I/R injury, in order to improve the MSCs survival rate during transplantation.

3.11. Lycopene Attenuates Bladder I/R Injury. The main functions of the bladder are to store urine under low pressure and discharge urine via coordination with nerves and muscles. However, some conditions (e.g., bladder outlet obstruction, ligation, complications from pregnancy or pelvic surgery, compression by pelvic tumors, and lower urinary tract obstruction [239–241]) can lead to bladder hypoperfusion, resulting in ischemia. Due to ischemia, the structure and function of the bladder will be altered, which can cause noncompliance and hyperreflexia [242, 243]. Therefore, blood flow must be restored as soon as possible. However, the damages occurring in these areas (nerves, synapses, and smooth muscle cells) are further aggravated when oxygen-rich blood quickly enters the ischemic bladder [244, 245]. Reperfusion can lead to neurologic injury, hypercapnia, and irregular and decreased contractile responses [246–248]. Although no studies have directly shown that lycopene could improve bladder I/R injury, a new type of antioxidant complex containing the lycopene chemical structure has been found to reduce detrusor nerve injury in

a guinea pig whole urinary bladder *in vitro* during bladder H/R injury (1 h/2 h) better than a substance containing the chemical structure of vitamin C [249]. Through these results, it can be found that lycopene exerting the neuroprotective effects in the bladder may not relate antioxidant activity, as vitamin C has better antioxidant properties than lycopene. This finding warrants further investigation.

4. Mechanisms

4.1. The Mechanisms behind Lycopene Attenuation of Ischemic Injuries. Ischemia damages organ function by depriving them of nutrition and oxygen. Long-term or severe ischemia can lead to acute pain, infarction, and sudden death. Once the infarct is formed, the currently known interventions such as thrombolytic therapy, angioplasty, and percutaneous coronary intervention can restore blood flow and relieve ischemic pain but have a lesser ability to reduce the infarct area. Adjuvant treatments that can potentially improve the efficacy of the above interventions should therefore be considered. Many studies have shown that lycopene supplementation before and after ischemia can significantly reduce ischemic injuries (e.g., infarct area, cell necrosis, edema, and inflammatory infiltrates) to some organs. Most of these studies focused on myocardial ischemia, while fewer focused on MSC and brain ischemia. The effects of lycopene are made by inhibiting oxidative stress, apoptosis, and inflammation. Moreover, lycopene can reduce remodeling in ischemic pathologies and promote tissue repair (Figure 3).

Firstly, lycopene decreases ROS and lipid peroxidation (MDA) levels primarily due to its strong antioxidant properties. In addition, lycopene can also inhibit oxidative stress via increasing the activity of SOD, GSH, GPx, and CAT. However, the Nrf-2 pathway does not seem to participate in lycopene's inhibition of oxidative stress, and the reduced level of Nrf-2 may be the result of restored redox status by lycopene. At the same time, lycopene also activates the PI3K/Akt pathway and further increases MnSOD levels to inhibit oxidative stress. Lycopene's inhibition of autophagy can also reduce ischemic injuries via decreased oxidative stress by activating AMPK and inhibiting mTOR. In addition, lycopene also reduces ischemic injuries by decreasing apoptosis via inhibition of ROS-dependent and non-ROS-dependent pathways. The former includes decreased (p-p38 or p-ATM)/p-p53 and JNK/(ERK1, or STAT1,3) pathways that are induced by ROS. The latter includes direct reductions in the levels of caspase-3, caspase-8, caspase-9, and Cyt-c via increased Bcl-2 levels and decreased Bax levels. Besides, lycopene can also directly inhibit the JNK/(ERK1, STAT1,3) pathways. Moreover, lycopene's anti-inflammatory properties are the result of inhibiting NF- κ B/inflammatory factors (TNF- α , IL-1 β , and IL-6), MPO, and CRP. Additionally, reduced levels of miRNAs (29c-3p, 194-5p, 503-5p, 20a-5p, 30a-5p, 192-5p, 30e-5p, and 126-3p) or increased IGF-1 and SIRT-1 levels also affect lycopene's reparative properties. Furthermore, inhibition of the p38/MMP-9 pathway by lycopene also impedes remodeling after infarcts.




FIGURE 3: The mechanisms of lycopene attenuating ischemic injury.




FIGURE 4: The mechanisms of lycopene attenuating I/R injury.

4.2. *The Mechanisms behind Lycopene Attenuation of I/R Injuries.* Timely reperfusion is still essential to mitigating ischemic damage to tissues and organs. However, reperfusion can further aggravate pathologic insults (molecules, cells, and organ dysfunctions). Furthermore, compared with ischemic injuries, I/R injuries are more serious due to their longer-lasting effects and incomplete reversal following reperfusion. Therefore, effective adjunctive therapies are urgently needed. So far, many studies have shown that lycopene

administration before and after reperfusion and before ischemia could improve I/R injury to almost all organs (Figure 4). Meanwhile, several mechanisms behind this have been identified, including inhibiting inflammation, oxidative stress, apoptosis, and promoting autophagy.

At first, lycopene inhibits oxidative stress during I/R injuries by directly enhancing the activity of SOD, CAT, GSH, and GSH-Px and reducing the levels of ROS, MDA, and F2IsoP. Lycopene can also indirectly increase NQO-1;

decrease ROS, MDA, and F2IsoP levels; and improve the activity of antioxidant enzymes by activating KEAP1/Nrf2/ARE. Upregulated expression of HIF-1 α mRNA, AMPK/ACC pathway activation and iNOS/NO pathway inhibition by lycopene can also account for its effects on oxidative stress. Secondly, lycopene can also improve the effects of I/R injuries by reducing apoptosis. On one hand, lycopene restores normal mitochondrial function via lowering ROS and raising Tfam. On the other hand, lycopene also directly inhibits caspase-mediated apoptosis and ER stress-induced apoptosis. The former is inhibited not only by lycopene directly but also through reduced ER stress indirectly. Lycopene reduces apoptosis by decreasing p-GSK-3 β and the ratio of Bax/Bcl-2 (increased Bcl-2 and decreased Bax), thereby increasing p-Akt and p-ERK1/2 while inhibiting the opening of mPTP and preventing Cyt-c from being released into the cytoplasm for APAF-1 binding. The latter is inhibited through reduced ER stress by AMPK activation and JNK inactivation.

Besides, lycopene suppression of JNK can lead to reduced inflammation and inflammatory signaling factors (decreasing TNF- α , TLR-2, IL-1 β , and IL-8 and increasing IL-10). Lycopene can also mitigate inflammation by directly inhibiting the Notch2/Hes-1 and IL-6/JAK-STAT3 pathways. Furthermore, HO-1 activated by Nrf-2 can weaken inflammation not only by inhibiting COX-2 but by increasing Kupffer cell (KC) autophagy and reducing the production of NLRP3. Additionally, lycopene can also promote autophagy by increasing the levels of p-AMPK, MAP1LC3B, and Beclin 1. Although the antioxidant effects of Nrf2 are not explored in this study [137], it can be hypothesized that Nrf2 is most likely involved in oxidative stress inhibition in the setting of I/R after lycopene administration.

5. Summary and Prospect

In conclusion, lycopene significantly reduces the impact of ischemic (e.g., infarction) and I/R injuries to multiple organs by inhibiting inflammation, oxidative stress, and apoptosis and promoting autophagy. Lycopene also improves repair and remodeling after ischemic injuries. Besides, lycopene may prevent damage to other organs caused by an ischemic or I/R injury to a different organ. Considering the different nature of ischemic and I/R injuries, a certain dose of lycopene is needed to assure its protective effects against both. For instance, in some special cases (such as high cholesterol [113]), higher doses of lycopene are required in order to achieve better effects. Furthermore, based on previous epidemiological studies, regular monitoring to maintain a certain concentration of lycopene *in vivo* may be a feasible strategy to achieve satisfied effects on preventing ischemic and I/R injuries.

While existing evidence is encouraging, additional studies are required before lycopene can be used clinically. Further clinical studies should focus on lycopene administration method, dosage, and time to achieve optimal protective effects. Additional study should also evaluate if lycopene can improve the damage to other multiple organs caused by ischemic or I/R injuries to one organ and the possibility of

using lycopene to simultaneously attenuate ischemic and I/R injuries to multiple organs. Moreover, new drug delivery techniques should be continuously explored to ensure that a certain dose of lycopene can accurately reach the ischemic site, thereby guaranteeing its protective effects.

Abbreviations

ACC:	Acetyl-CoA carboxylase
Akt:	Protein kinase B
ALT:	Alanine aminotransferase
AMPK:	Adenosine monophosphate-activated protein kinase
APAF-1:	Apoptotic peptidase activating factor 1
ARE:	Antioxidant response element
AST:	Aspartate aminotransferase
ATF6:	Activating transcription factor 6
ATM:	Ataxia telangiectasia mutated protein
Bax:	Bcl-2-associated X protein
Bcl-2:	B-cell lymphoma 2
BT:	Bacterial translocation
CAT:	Catalase
CHOP:	C/EBP homologous protein
CK-MB:	Creatine kinase isoenzymes
COX:	Cyclooxygenase
CRP:	C-reactive protein
Cyt-c:	Cytochrome c
eIF2 α :	Eukaryotic translation initiation factor 2 subunit- α
ER:	Endoplasmic reticulum
ERK:	Extracellular regulated protein kinases
F2IsoP:	F2-Isoprostane
GRP78:	Glucose-regulated proteins 78
GSH:	Glutathione
GSH-Px:	Glutathione peroxidase
GSK-3 β :	Glycogen synthase kinase-3 β
GST:	Glutathione S-transferase
HCD:	High-cholesterol diet
Hes-1:	Hairy enhancer of split
HIF-1:	Hypoxia inducible factor-1
HO-1:	Heme oxygenase 1
hs-CRP:	Hypersensitive C-reactive protein
IGF:	Insulin-like growth factors
IL:	Interleukin
JAK:	The Janus kinase
JNK:	c-Jun N-terminal kinase
KEAP1:	Kelch-like erythroid cell-derived protein with CNC homology-associated protein 1
LAD:	Left anterior descending
LDL:	Low-density lipoprotein
LDH:	Lactate dehydrogenase
LV:	Left ventricular
MAP1LC3B:	Microtubule-associated protein 1-light chain 3 β
MDA:	Malonic dialdehyde
MMP-9:	Matrix metalloproteinase-9
MPO:	Myeloperoxidase
mPTP:	Mitochondrial permeability transition pore
mtDNA:	Mitochondrial DNA
mTOR:	Mammalian target of rapamycin

MYO:	Myoglobin
NF- κ B:	Nuclear factor-kappa B
NLRP3:	NOD-like receptor protein 3
NOS:	Nitric oxide synthase
NQO-1:	NADPH quinone oxidoreductase-1
Nrf2:	Nuclear factor-erythroid 2-related factor 2
PCI:	Percutaneous coronary intervention
PI3K:	Phosphoinositide 3-kinase
PKC:	Protein kinase C
ROS:	Reactive oxygen species
SIRT-1:	Sirtuin 1
sIgA:	Secretory immunoglobulin A
SOD:	Superoxide dismutase
STAT:	Signal transducer and activator of transcription
sXbp-:	Splices X-box binding proteins-1
Tfam:	Mitochondrial transcription factor A
TLR-2:	Toll-like receptor 2
TNF:	Tumor necrosis factor
TnI:	Troponin I
TROP:	Troponin T.

Conflicts of Interest

The authors declare that they have no conflict of interest.

Authors' Contributions

Sijia Wu, Qianwen Peng, Jia Shang, and Zhenxing Xie collected the data and composed the manuscript. Wanglin Dong, Xiajun Guo, Yuanyuan Li, and Chaoran Chen helped a lot in designing the graphic abstracts. All authors have reviewed and agreed to the publication of the manuscript. Xiajun Guo and Sijia Wu contributed equally to this work.

Acknowledgments

The study was sponsored by the Graduate Education Reform and Quality Improvement Project of Henan Province (grant number: YJS2021AL074), Graduate Education Innovation and Quality Improvement Project of Henan University (grant number: SYL19060141), Henan Province Science and Technology Development Plan Soft Science Research (grant number: 222400410497), Planning and Decision Consultation Project of Henan Province (grant number: 2018JC38), and Research and Practice Project of Higher Education Teaching Reform in Henan Province (grant number: HDXJJG2020-25).

References

- [1] J. D. Beard, "Chronic lower limb ischemia," *The Western journal of medicine*, vol. 173, no. 1, pp. 60–63, 2000.
- [2] H. W. Kniemeyer, "Mesenteric infarct—when is the vascular surgeon needed?," *Zentralblatt für Chirurgie*, vol. 123, no. 12, pp. 1411–1417, 1998.
- [3] Y. Zhai, H. Petrowsky, J. C. Hong, R. W. Busuttill, and J. W. Kupiec-Weglinski, "Ischaemia-reperfusion injury in liver transplantation—from bench to bedside," *Nature reviews Gastroenterology & hepatology*, vol. 10, no. 2, pp. 79–89, 2013.
- [4] N. Perico, D. Cattaneo, M. H. Sayegh, and G. Remuzzi, "Delayed graft function in kidney transplantation," *The Lancet*, vol. 364, no. 9447, pp. 1814–1827, 2004.
- [5] A. Mastoraki, S. Mastoraki, E. Tziava et al., "Mesenteric ischemia: pathogenesis and challenging diagnostic and therapeutic modalities," *World journal of gastrointestinal pathophysiology*, vol. 7, no. 1, pp. 125–130, 2016.
- [6] G. A. Pantely, S. A. Malone, W. S. Rhen et al., "Regeneration of myocardial phosphocreatine in pigs despite continued moderate ischemia," *Circulation research*, vol. 67, no. 6, pp. 1481–1493, 1990.
- [7] R. Schulz, B. D. Guth, K. Pieper, C. Martin, and G. Heusch, "Recruitment of an inotropic reserve in moderately ischemic myocardium at the expense of metabolic recovery. A model of short-term hibernation," *Circulation research*, vol. 70, no. 6, pp. 1282–1295, 1992.
- [8] H. Haimovici, "Muscular, renal, and metabolic complications of acute arterial occlusions: myoneuropathic-metabolic syndrome," *Surgery*, vol. 85, no. 4, pp. 461–468, 1979.
- [9] D. A. Parks and D. N. Granger, "Contributions of ischemia and reperfusion to mucosal lesion formation," *American Journal of Physiology-Gastrointestinal and Liver Physiology*, vol. 250, no. 6, pp. G749–G753, 1986.
- [10] E. Rhoden, C. Telöken, M. Lucas et al., "Protective effect of allopurinol in the renal ischemia-reperfusion in uninephrectomized rats," *General Pharmacology*, vol. 35, no. 4, pp. 189–193, 2000.
- [11] S. Warach and L. L. Latour, "Evidence of reperfusion injury, exacerbated by thrombolytic therapy, in human focal brain ischemia using a novel imaging marker of early blood-brain barrier disruption," *Stroke*, vol. 35, no. 11, pp. 2659–2661, 2004.
- [12] M. K. Porteous, J. M. Diamond, and J. D. Christie, "Primary graft dysfunction: lessons learned about the first 72 h after lung transplantation," *Current Opinion in Organ Transplantation*, vol. 20, no. 5, pp. 506–514, 2015.
- [13] I. O. Sherif and N. H. Al-Shaalan, "Alleviation of remote lung injury following liver ischemia/reperfusion: Possible protective role of vildagliptin," *International Immunopharmacology*, vol. 91, article 107305, 2021.
- [14] S. J. Han, H. S. Jang, S. Y. Seu et al., "Hepatic ischemia/reperfusion injury disrupts the homeostasis of kidney primary cilia via oxidative stress," *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, vol. 1863, no. 7, pp. 1817–1828, 2017.
- [15] A. A. Weinbroum, E. Hochhauser, V. Rudick et al., "Direct induction of acute lung and myocardial dysfunction by liver ischemia and reperfusion," *Journal of Trauma and Acute Care Surgery*, vol. 43, no. 4, pp. 627–635, 1997.
- [16] J. Yang, H. Jiang, J. Yang et al., "Valsartan preconditioning protects against myocardial ischemia-reperfusion injury through TLR4/NF-kappaB signaling pathway," *Molecular and cellular biochemistry*, vol. 330, no. 1-2, pp. 39–46, 2009.
- [17] M. Correa-Costa, D. Gallo, E. Csizmadia et al., "Carbon monoxide protects the kidney through the central circadian clock and CD39," *Proceedings of the National Academy of Sciences*, vol. 115, no. 10, pp. E2302–E2310, 2018.
- [18] S. Jeddi, S. Gheibi, K. Kashfi, M. Carlström, and A. Ghasemi, "Dose-dependent effects of long-term administration of hydrogen sulfide on myocardial ischemia-reperfusion injury in male Wistar rats: modulation of RKIP, NF- κ B, and

- oxidative stress,” *International Journal of Molecular Sciences*, vol. 21, no. 4, p. 1415, 2020.
- [19] H. Yu, P. Liu, H. Tang et al., “Oleuropein, a natural extract from plants, offers neuroprotection in focal cerebral ischemia/reperfusion injury in mice,” *European Journal of Pharmacology*, vol. 775, pp. 113–119, 2016.
- [20] M. I. Oerlemans, J. Liu, F. Arslan et al., “Inhibition of RIP1-dependent necrosis prevents adverse cardiac remodeling after myocardial ischemia-reperfusion in vivo,” *Basic research in cardiology*, vol. 107, no. 4, p. 270, 2012.
- [21] A. R. Mangels, J. M. Holden, G. R. Beecher, M. R. Forman, and E. Lanza, “Carotenoid content of fruits and vegetables: an evaluation of analytic data,” *Journal of the American Dietetic Association*, vol. 93, no. 3, pp. 284–296, 1993.
- [22] S. Xianquan, J. Shi, Y. Kakuda, and J. Yueming, “Stability of lycopene during food processing and storage,” *Journal of medicinal food*, vol. 8, no. 4, pp. 413–422, 2005.
- [23] G. A. Chasse, M. L. Mak, E. Deretey et al., “An ab initio computational study on selected lycopene isomers,” *Journal of Molecular Structure-Theochem*, vol. 571, no. 1-3, pp. 27–37, 2001.
- [24] A. Agarwal, H. Shen, S. Agarwal, and A. V. Rao, “Lycopene content of tomato products: its stability, bioavailability and in vivo antioxidant properties,” *Journal of medicinal food*, vol. 4, no. 1, pp. 9–15, 2001.
- [25] W. Stahl and H. Sies, “Uptake of lycopene and its geometrical isomers is greater from heat-processed than from unprocessed tomato juice in humans,” *The Journal of nutrition*, vol. 122, no. 11, pp. 2161–2166, 1992.
- [26] M. Moussa, J. F. Landrier, E. Reboul et al., “Lycopene absorption in human intestinal cells and in mice involves scavenger receptor class B type I but not Niemann-Pick C1-like 1,” *The Journal of nutrition*, vol. 138, no. 8, pp. 1432–1436, 2008.
- [27] A. During, H. D. Dawson, and E. H. Harrison, “Carotenoid transport is decreased and expression of the lipid transporters SR-BI, NPC1L1, and ABCA1 is downregulated in Caco-2 cells treated with ezetimibe,” *The Journal of nutrition*, vol. 135, no. 10, pp. 2305–2312, 2005.
- [28] P. Singh and G. K. Goyal, “Dietary lycopene: its properties and anticarcinogenic effects,” *Comprehensive Reviews in Food Science and Food Safety*, vol. 7, no. 3, pp. 255–270, 2008.
- [29] R. S. Parker, “Absorption, metabolism, and transport of carotenoids,” *The FASEB Journal*, vol. 10, no. 5, pp. 542–551, 1996.
- [30] N. E. Moran, J. W. Erdman Jr., and S. K. Clinton, “Complex interactions between dietary and genetic factors impact lycopene metabolism and distribution,” *Archives of biochemistry and biophysics*, vol. 539, no. 2, pp. 171–180, 2013.
- [31] A. V. Rao and S. Agarwal, “Bioavailability and in vivo antioxidant properties of lycopene from tomato products and their possible role in the prevention of cancer,” *Nutrition and Cancer*, vol. 31, no. 3, pp. 199–203, 1998.
- [32] F. Khachik, L. Carvalho, P. S. Bernstein, G. J. Muir, D. Y. Zhao, and N. B. Katz, “Chemistry, distribution, and metabolism of tomato carotenoids and their impact on human health,” *Experimental biology and medicine*, vol. 227, no. 10, pp. 845–851, 2002.
- [33] W. Stahl, W. Schwarz, A. R. Sundquist, and H. Sies, “_cis-trans_ isomers of lycopene and β -carotene in human serum and tissues,” *Archives of biochemistry and biophysics*, vol. 294, no. 1, pp. 173–177, 1992.
- [34] J. Levy, E. Bosin, B. Feldman et al., “Lycopene is a more potent inhibitor of human cancer cell proliferation than either α -carotene or β -carotene,” *Nutrition and Cancer*, vol. 24, no. 3, pp. 257–266, 1995.
- [35] K. Klipstein-Grobusch, L. J. Launer, J. M. Geleijnse, H. Boeing, A. Hofman, and J. C. M. Witteman, “Serum carotenoids and atherosclerosis: The Rotterdam Study,” *Atherosclerosis*, vol. 148, no. 1, pp. 49–56, 2000.
- [36] L. G. Rao, E. S. Mackinnon, R. G. Josse, T. M. Murray, A. Strauss, and A. V. Rao, “Lycopene consumption decreases oxidative stress and bone resorption markers in postmenopausal women,” *Osteoporosis international*, vol. 18, no. 1, pp. 109–115, 2007.
- [37] A. Iwasaki and C. Gagnon, “Formation of reactive oxygen species in spermatozoa of infertile patients*,” *Fertility and sterility*, vol. 57, no. 2, pp. 409–416, 1992.
- [38] R. Zhu, B. Chen, Y. Bai et al., “Lycopene in protection against obesity and diabetes: a mechanistic review,” *Pharmacological Research*, vol. 159, article 104966, 2020.
- [39] J. J. Mitchell, M. Paiva, and M. B. Heaton, “Vitamin E and β -carotene protect against ethanol combined with ischemia in an embryonic rat hippocampal culture model of fetal alcohol syndrome,” *Neuroscience letters*, vol. 263, no. 2-3, pp. 189–192, 1999.
- [40] V. M. Moia, F. L. Portilho, T. A. Pádua et al., “Lycopene used as anti-inflammatory nanodrug for the treatment of rheumatoid arthritis: animal assay, pharmacokinetics, ABC transporter and tissue deposition,” *Colloids Surf B Biointerfaces*, vol. 188, article 110814, 2020.
- [41] S. Grether-Beck, A. Marini, T. Jaenicke, W. Stahl, and J. Krutmann, “Molecular evidence that oral supplementation with lycopene or lutein protects human skin against ultraviolet radiation: results from a double-blinded, placebo-controlled, crossover study,” *British Journal of Dermatology*, vol. 176, no. 5, pp. 1231–1240, 2017.
- [42] H. Huang, X. Kuang, X. Zhu et al., “Maintaining blood retinal barrier homeostasis to attenuate retinal ischemia-reperfusion injury by targeting the KEAP1/NRF2/ARE pathway with lycopene,” *Cell Signal*, vol. 88, article 110153, 2021.
- [43] W. Stahl and H. Sies, “Lycopene: a biologically important carotenoid for humans?,” *Archives of biochemistry and biophysics*, vol. 336, no. 1, pp. 1–9, 1996.
- [44] K. Wertz, U. Siler, and R. Goralczyk, “Lycopene: modes of action to promote prostate health,” *Archives of biochemistry and biophysics*, vol. 430, no. 1, pp. 127–134, 2004.
- [45] D. Heber and Q. Y. Lu, “Overview of mechanisms of action of lycopene,” *Experimental biology and medicine*, vol. 227, no. 10, pp. 920–923, 2002.
- [46] Z. Wu, X. Zhang, Z. Li, Z. Wen, and Y. Lin, “Activation of autophagy contributes to the protective effects of lycopene against oxidative stress-induced apoptosis in rat chondrocytes,” *Phytotherapy Research*, vol. 35, no. 7, pp. 4032–4045, 2021.
- [47] S. Bi, L. Li, H. Gu et al., “Lycopene upregulates ZO-1 and downregulates claudin-1 through autophagy inhibition in the human cutaneous squamous cell carcinoma cell line COLO-16,” *Journal of Cancer*, vol. 10, no. 2, pp. 510–521, 2019.
- [48] M. Kim, S. H. Kim, J. W. Lim, and H. Kim, “Lycopene induces apoptosis by inhibiting nuclear translocation of beta-catenin in gastric cancer cells,” *Journal of Physiology and Pharmacology*, vol. 70, no. 4, 2019.

- [49] H. G. Kang, S. Lee, P. S. Jeong et al., "Lycopene improves in vitro development of porcine embryos by reducing oxidative stress and apoptosis," *Antioxidants*, vol. 10, no. 2, p. 230, 2021.
- [50] L. M. Mills, H. Wilson, and F. Thies, "Lycopene inhibits lymphocyte proliferation through mechanisms dependent on early cell activation," *Molecular nutrition & food research*, vol. 56, no. 7, pp. 1034–1042, 2012.
- [51] C. M. Yang, I. H. Lu, H. Y. Chen, and M. L. Hu, "Lycopene inhibits the proliferation of androgen-dependent human prostate tumor cells through activation of PPAR γ -LXR α -ABCA1 pathway," *The Journal of nutritional biochemistry*, vol. 23, no. 1, pp. 8–17, 2012.
- [52] X. Y. Dai, X. W. Li, S. Y. Zhu et al., "Lycopene ameliorates Di(2-ethylhexyl) phthalate-induced pyroptosis in spleen via suppression of classic caspase-1/NLRP3 pathway," *Journal of Agricultural and Food Chemistry*, vol. 69, no. 4, pp. 1291–1299, 2021.
- [53] E. Giovannucci, A. Ascherio, E. B. Rimm, M. J. Stampfer, G. A. Colditz, and W. C. Willett, "Intake of carotenoids and retinol in relation to risk of prostate cancer," *JNCI Journal of the National Cancer Institute*, vol. 87, no. 23, pp. 1767–1776, 1995.
- [54] P. R. Gajendragadkar, A. Hubsch, K. M. Mäki-Petäjä, M. Serg, I. B. Wilkinson, and J. Cheriyan, "Effects of oral lycopene supplementation on vascular function in patients with cardiovascular disease and healthy volunteers: a randomised controlled trial," *PLoS One*, vol. 9, no. 6, article e99070, 2014.
- [55] W. Mellert, K. Deckardt, C. Gembardt, S. Schulte, B. V. Ravenzwaay, and R. S. Slesinski, "Thirteen-week oral toxicity study of synthetic lycopene products in rats," *Food and Chemical Toxicology*, vol. 40, no. 11, pp. 1581–1588, 2002.
- [56] P. Reich, H. Shwachman, and J. M. Craig, "Lycopopenia: a variant of carotenemia," *New England Journal of Medicine*, vol. 262, no. 6, pp. 263–269, 1960.
- [57] J. B. Sharma, A. Sharma, A. Bahadur, N. Vimala, A. Satyam, and S. Mittal, "Oxidative stress markers and antioxidant levels in normal pregnancy and pre-eclampsia," *International journal of gynecology & obstetrics*, vol. 94, no. 1, pp. 23–27, 2006.
- [58] D. Choi, K. C. Hwang, K. Y. Lee, and Y. H. Kim, "Ischemic heart diseases: current treatments and future," *Journal of Controlled Release*, vol. 140, no. 3, pp. 194–202, 2009.
- [59] H. Shimokawa and S. Yasuda, "Myocardial ischemia: current concepts and future perspectives," *Journal of cardiology*, vol. 52, no. 2, pp. 67–78, 2008.
- [60] G. Heusch, "Myocardial Ischemia," *Circulation research*, vol. 119, no. 2, pp. 194–196, 2016.
- [61] P. Hafez, S. R. Chowdhury, S. Jose et al., "Development of an in vitro cardiac ischemic model using primary human cardiomyocytes," *Cardiovascular Engineering and Technology*, vol. 9, no. 3, pp. 529–538, 2018.
- [62] G. Morciano, M. Bonora, G. Campo et al., "Mechanistic role of mPTP in ischemia-reperfusion injury," *Mitochondrial Dynamics in Cardiovascular Medicine*, vol. 982, pp. 169–189, 2017.
- [63] D. H. Kazmi, A. Kapoor, A. Sinha et al., "Role of metabolic manipulator trimetazidine in limiting percutaneous coronary intervention-induced myocardial injury," *Indian Heart Journal*, vol. 70, Supplement 3, pp. S365–S371, 2018.
- [64] E. Braunwald and R. A. Kloner, "Myocardial reperfusion: a double-edged sword?," *The Journal of clinical investigation*, vol. 76, no. 5, pp. 1713–1719, 1985.
- [65] D. M. Yellon and D. J. Hausenloy, "Myocardial reperfusion injury," *New England Journal of Medicine*, vol. 357, no. 11, pp. 1121–1135, 2007.
- [66] C. ÇetIn, A. A. Köse, E. Aral et al., "Protective effect of fucoidin (a neutrophil rolling inhibitor) on ischemia reperfusion injury: experimental study in rat epigastric island flaps," *Annals of plastic surgery*, vol. 47, no. 5, pp. 540–546, 2001.
- [67] R. K. Dongworth, U. A. Mukherjee, A. R. Hall et al., "DJ-1 protects against cell death following acute cardiac ischemia-reperfusion injury," *Cell death & disease*, vol. 5, no. 2, article e1082, 2014.
- [68] C. E. Murry, R. B. Jennings, and K. A. Reimer, "Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium," *Circulation*, vol. 74, no. 5, pp. 1124–1136, 1986.
- [69] P. Pagliaro, F. Moro, F. Tullio, M. G. Perrelli, and C. Penna, "Cardioprotective pathways during reperfusion: focus on redox signaling and other modalities of cell signaling," *Antioxidants & redox signaling*, vol. 14, no. 5, pp. 833–850, 2011.
- [70] Z. Q. Zhao, J. S. Corvera, M. E. Halkos et al., "Inhibition of myocardial injury by ischemic postconditioning during reperfusion: comparison with ischemic preconditioning," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 285, no. 2, pp. H579–H588, 2003.
- [71] C. S. Piao, S. Gao, G. H. Lee et al., "Sulforaphane protects ischemic injury of hearts through antioxidant pathway and mitochondrial K_{ATP} channels," *Pharmacological Research*, vol. 61, no. 4, pp. 342–348, 2010.
- [72] M. Mahmoudabady, F. S. Talebian, N. A. Zabihi, S. A. Rezaee, and S. Niazmand, "Teucrium polium L. improved heart function and inhibited myocardial apoptosis in isolated rat heart following ischemia-reperfusion injury," *Journal of Pharmacopuncture*, vol. 21, no. 3, pp. 159–167, 2018.
- [73] H. Liu, C. Wang, Z. Qiao, and Y. Xu, "Protective effect of curcumin against myocardium injury in ischemia reperfusion rats," *Pharmaceutical Biology*, vol. 55, no. 1, pp. 1144–1148, 2017.
- [74] H. Liu, L. Yang, H. J. Wu et al., "Water-soluble acacetin pro-drug confers significant cardioprotection against ischemia/reperfusion injury," *Scientific Reports*, vol. 6, no. 1, p. 36435, 2016.
- [75] F. Wang, Q. Gao, J. Yang et al., "Artemisinin suppresses myocardial ischemia-reperfusion injury via NLRP3 inflammasome mechanism," *Molecular and Cellular Biochemistry*, vol. 474, no. 1-2, pp. 171–180, 2020.
- [76] F. Tullio, C. Angotti, M. G. Perrelli, C. Penna, and P. Pagliaro, "Redox balance and cardioprotection," *Basic Research in Cardiology*, vol. 108, no. 6, p. 392, 2013.
- [77] A. D. Costa and K. D. Garlid, "Intramitochondrial signaling: interactions among mitoKATP, PKC ϵ , ROS, and MPT," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 295, no. 2, pp. H874–H882, 2008.
- [78] P. Pagliaro and C. Penna, "Redox signalling and cardioprotection: translatability and mechanism," *British Journal of Pharmacology*, vol. 172, no. 8, pp. 1974–1995, 2015.
- [79] D. J. Hausenloy, S. B. Ong, and D. M. Yellon, "The mitochondrial permeability transition pore as a target for preconditioning and postconditioning," *Basic research in cardiology*, vol. 104, no. 2, pp. 189–202, 2009.

- [80] A. Skyschally, R. Schulz, P. Gres, H. G. Korth, and G. Heusch, "Attenuation of ischemic preconditioning in pigs by scavenging of free oxyradicals with ascorbic acid," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 284, no. 2, pp. H698–H703, 2003.
- [81] R. A. Forbes, C. Steenbergen, and E. Murphy, "Diazoxide-induced cardioprotection requires signaling through a redox-sensitive mechanism," *Circulation research*, vol. 88, no. 8, pp. 802–809, 2001.
- [82] C. Csonka, K. Kupai, P. Bencsik et al., "Cholesterol-enriched diet inhibits cardioprotection by ATP-sensitive K⁺ channel activators cromakalim and diazoxide," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 306, no. 3, pp. H405–H413, 2014.
- [83] G. F. Kocsis, M. Sárközy, P. Bencsik et al., "Preconditioning protects the heart in a prolonged uremic condition," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 303, no. 10, pp. H1229–H1236, 2012.
- [84] A. Görbe, Z. V. Varga, K. Kupai et al., "Cholesterol diet leads to attenuation of ischemic preconditioning-induced cardiac protection: the role of connexin 43," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 300, no. 5, pp. H1907–H1913, 2011.
- [85] P. Ferdinandy, R. Schulz, and G. F. Baxter, "Interaction of cardiovascular risk factors with myocardial ischemia/reperfusion injury, preconditioning, and postconditioning," *Pharmacological reviews*, vol. 59, no. 4, pp. 418–458, 2007.
- [86] Z. Giricz, A. Görbe, J. Pipis, D. S. Burley, P. Ferdinandy, and G. F. Baxter, "Hyperlipidaemia induced by a high-cholesterol diet leads to the deterioration of guanosine-3',5'-cyclic monophosphate/protein kinase G-dependent cardioprotection in rats," *British journal of pharmacology*, vol. 158, no. 6, pp. 1495–1502, 2009.
- [87] J. Karppi, J. A. Laukkanen, T. H. Makikallio, and S. Kurl, "Low serum lycopene and -carotene increase risk of acute myocardial infarction in men," *The European Journal of Public Health*, vol. 22, no. 6, pp. 835–840, 2012.
- [88] L. Kohlmeier, J. D. Kark, E. Gomez-Gracia et al., "Lycopene and myocardial infarction risk in the EURAMIC study," *American Journal of Epidemiology*, vol. 146, no. 8, pp. 618–626, 1997.
- [89] T. H. Rissanen, S. Voutilainen, K. Nyssönen, R. Salonen, G. A. Kaplan, and J. T. Salonen, "Serum lycopene concentrations and carotid atherosclerosis: the Kuopio ischaemic heart disease risk factor study," *The American journal of clinical nutrition*, vol. 77, no. 1, pp. 133–138, 2003.
- [90] G. Rona, "Catecholamine cardiotoxicity," *Journal of molecular and cellular cardiology*, vol. 17, no. 4, pp. 291–306, 1985.
- [91] K. Harada, Y. Fukata, A. Miwa, S. Kaneta, H. Fukushima, and N. Ogawa, "Effect of KRN2391, a novel vasodilator, on various experimental anginal models in rats," *The Japanese Journal of Pharmacology*, vol. 63, no. 1, pp. 35–39, 1993.
- [92] P. Mladěnka, R. Hrdina, Z. Bobrovová et al., "Cardiac biomarkers in a model of acute catecholamine cardiotoxicity," *Human & experimental toxicology*, vol. 28, no. 10, pp. 631–640, 2009.
- [93] Q. I. He, W. E. Zhou, C. Xiong, G. Tan, and M. Chen, "Lycopene attenuates inflammation and apoptosis in post-myocardial infarction remodeling by inhibiting the nuclear factor- κ B signaling pathway," *Molecular Medicine Reports*, vol. 11, no. 1, pp. 374–378, 2015.
- [94] B. L. B. Pereira, P. P. Reis, F. E. Severino et al., "Tomato (*Lycopersicon esculentum*) or lycopene supplementation attenuates ventricular remodeling after myocardial infarction through different mechanistic pathways," *The Journal of nutritional biochemistry*, vol. 46, pp. 117–124, 2017.
- [95] U. Aman, P. Vaibhav, and R. Balaraman, "Tomato lycopene attenuates myocardial infarction induced by isoproterenol: electrocardiographic, biochemical and anti-apoptotic study," *Asian Pacific journal of tropical biomedicine*, vol. 2, no. 5, pp. 345–351, 2012.
- [96] X. Wang, H. Lv, Y. Gu et al., "Protective effect of lycopene on cardiac function and myocardial fibrosis after acute myocardial infarction in rats via the modulation of p38 and MMP-9," *Journal of molecular histology*, vol. 45, no. 1, pp. 113–120, 2014.
- [97] S. Ojha, S. Goyal, C. Sharma, S. Arora, S. Kumari, and D. S. Arya, "Cardioprotective effect of lycopene against isoproterenol-induced myocardial infarction in rats," *Human & experimental toxicology*, vol. 32, no. 5, pp. 492–503, 2013.
- [98] S. Fan, J. B. Sun, R. Li, X. Song, and J. Li, "Lycopene protects myocardial ischemia injury through anti-apoptosis and antioxidative stress," *European Review for Medical and Pharmacological Sciences*, vol. 23, no. 7, pp. 3096–3104, 2019.
- [99] R. Parvin and N. Akhter, "Protective effect of tomato against adrenaline-induced myocardial infarction in rats," *Bangladesh Medical Research Council Bulletin*, vol. 34, no. 3, pp. 104–108, 2008.
- [100] A. Annapurna, M. A. Ansari, and P. M. Manjunath, "Partial role of multiple pathways in infarct size limiting effect of quercetin and rutin against cerebral ischemia-reperfusion injury in rats," *European Review for Medical and Pharmacological Sciences*, vol. 17, no. 4, pp. 491–500, 2013.
- [101] A. M. Rezk, I. A. H. Ibrahim, M. F. Mahmoud, and A. A. Mahmoud, "Quercetin and lithium chloride potentiate the protective effects of carvedilol against renal ischemia-reperfusion injury in high-fructose, high-fat diet-fed Swiss albino mice independent of renal lipid signaling," *Chemico-Biological Interactions*, vol. 333, article 109307, 2021.
- [102] L. Chen, X. Wu, W. Wang, X. I. A. Wang, and J. Ma, "Quercetin with lycopene modulates enzymic antioxidant genes pathway in isoproterenol cardiotoxicity in rats," *Libyan Journal of Medicine*, vol. 16, no. 1, p. 1943924, 2021.
- [103] H. Liu, L. Zhang, and S. Lu, "Evaluation of antioxidant and immunity activities of quercetin in isoproterenol-treated rats," *Molecules*, vol. 17, no. 4, pp. 4281–4291, 2012.
- [104] S. Asgary, R. Soltani, F. Daraei, L. Salehizadeh, G. Vaseghi, and N. Sarrafzadegan, "The effect of lycopene on serum level of cardiac biomarkers in patients undergoing elective percutaneous coronary intervention: a randomized controlled clinical trial," *ARYA atherosclerosis*, vol. 17, no. 1, pp. 1–7, 2021.
- [105] M. J. Biddle, T. A. Lennie, G. V. Bricker, R. E. Kopec, S. J. Schwartz, and D. K. Moser, "Lycopene dietary intervention: a pilot study in patients with heart failure," *The Journal of cardiovascular nursing*, vol. 30, no. 3, pp. 205–212, 2015.
- [106] K. Jacob, M. J. Periago, V. Böhm, and G. R. Berrueto, "Influence of lycopene and vitamin C from tomato juice on biomarkers of oxidative stress and inflammation," *British Journal of Nutrition*, vol. 99, no. 1, pp. 137–146, 2008.
- [107] C. Tong, C. Peng, L. Wang et al., "Intravenous administration of lycopene, a tomato extract, protects against myocardial ischemia-reperfusion injury," *Nutrients*, vol. 8, no. 3, p. 138, 2016.

- [108] M. Hao, S. Zhu, L. Hu, H. Zhu, X. Wu, and Q. Li, "Myocardial ischemic postconditioning promotes autophagy against ischemia reperfusion injury via the activation of the nNOS/AMPK/mTOR pathway," *International journal of molecular sciences*, vol. 18, no. 3, p. 614, 2017.
- [109] A. Araszkiwicz, M. Grygier, M. Pyda, J. Rajewska, M. Lesiak, and S. Grajek, "Postconditioning attenuates early ventricular arrhythmias in patients with high-risk ST-segment elevation myocardial infarction," *Journal of cardiology*, vol. 65, no. 6, pp. 459–465, 2015.
- [110] S. Cao, Y. Liu, H. Wang et al., "Ischemic postconditioning influences electron transport chain protein turnover in Langendorff-perfused rat hearts," *PeerJ*, vol. 4, article e1706, 2016.
- [111] L. You, L. Li, Q. Xu, J. Ren, and F. Zhang, "Postconditioning reduces infarct size and cardiac myocyte apoptosis via the opioid receptor and JAK-STAT signaling pathway," *Molecular biology reports*, vol. 38, no. 1, pp. 437–443, 2011.
- [112] G. Shinohara, K. Morita, R. Nagahori et al., "Ischemic postconditioning promotes left ventricular functional recovery after cardioplegic arrest in an in vivo piglet model of global ischemia reperfusion injury on cardiopulmonary bypass," *The Journal of Thoracic and Cardiovascular Surgery*, vol. 142, no. 4, pp. 926–932, 2011.
- [113] L. Duan, C. Liang, X. Li et al., "Lycopene restores the effect of ischemic postconditioning on myocardial ischemia-reperfusion injury in hypercholesterolemic rats," *International Journal of Molecular Medicine*, vol. 43, no. 6, pp. 2451–2461, 2019.
- [114] E. R. Gross, A. K. Hsu, and G. J. Gross, "Diabetes abolishes morphine-induced cardioprotection via multiple pathways upstream of glycogen synthase kinase-3beta," *Diabetes*, vol. 56, no. 1, pp. 127–136, 2007.
- [115] J. L. Zhao, Y. J. Yang, S. J. You, C. J. Cui, and R. L. Gao, "Different effects of postconditioning on myocardial no-reflow in the normal and hypercholesterolemic mini-swines," *Microvascular research*, vol. 73, no. 2, pp. 137–142, 2007.
- [116] C. Penna, C. Angotti, and P. Pagliaro, "Protein S-nitrosylation in preconditioning and postconditioning," *Experimental Biology and Medicine*, vol. 239, no. 6, pp. 647–662, 2014.
- [117] Y. Hou, Z. Guo, J. Li, and P. G. Wang, "Seleno Compounds and Glutathione Peroxidase Catalyzed Decomposition of S_2 -Nitrosothiols," *Biochemical and biophysical research communications*, vol. 228, no. 1, pp. 88–93, 1996.
- [118] D. Jour'dheuil, F. S. Laroux, A. M. Miles, D. A. Wink, and M. B. Grisham, "Effect of Superoxide Dismutase on the Stability of S_2 -Nitrosothiols," *Archives of Biochemistry and Biophysics*, vol. 361, no. 2, pp. 323–330, 1999.
- [119] F. Beigi, D. R. Gonzalez, K. M. Minhas et al., "Dynamic denitrosylation via S-nitrosoglutathione reductase regulates cardiovascular function," *Proceedings of the National Academy of Sciences*, vol. 109, no. 11, pp. 4314–4319, 2012.
- [120] X. Li, P. Jia, Z. Huang et al., "Lycopene protects against myocardial ischemia-reperfusion injury by inhibiting mitochondrial permeability transition pore opening," *Drug Design, Development and Therapy*, vol. Volume 13, pp. 2331–2342, 2019.
- [121] R. Yue, X. Xia, J. Jiang et al., "Mitochondrial DNA oxidative damage contributes to cardiomyocyte ischemia/reperfusion-injury in rats: cardioprotective role of lycopene," *Journal of cellular physiology*, vol. 230, no. 9, pp. 2128–2141, 2015.
- [122] P. Bansal, S. K. Gupta, S. K. Ojha et al., "Cardioprotective effect of lycopene in the experimental model of myocardial ischemia-reperfusion injury," *Molecular and cellular biochemistry*, vol. 289, no. 1-2, pp. 1–9, 2006.
- [123] S. Das, H. Otani, N. Maulik, and D. K. Das, "Lycopene, tomatoes, and coronary heart disease," *Free radical research*, vol. 39, no. 4, pp. 449–455, 2005.
- [124] Y. Gao, P. Jia, W. Shu, and D. Jia, "The protective effect of lycopene on hypoxia/reoxygenation-induced endoplasmic reticulum stress in H9C2 cardiomyocytes," *European Journal of Pharmacology*, vol. 774, pp. 71–79, 2016.
- [125] F. Chen, Z. W. Sun, L. F. Ye, G. S. Fu, Y. Mou, and S. J. Hu, "Lycopene protects against apoptosis in hypoxia/reoxygenation-induced H9C2 myocardioblast cells through increased autophagy," *Molecular medicine reports*, vol. 11, no. 2, pp. 1358–1365, 2015.
- [126] J. Xu, H. Hu, B. Chen et al., "Lycopene protects against hypoxia/reoxygenation injury by alleviating ER stress induced apoptosis in neonatal mouse cardiomyocytes," *PLoS One*, vol. 10, no. 8, article e0136443, 2015.
- [127] R. Yue, H. Hu, K. H. Yiu et al., "Lycopene protects against hypoxia/reoxygenation-induced apoptosis by preventing mitochondrial dysfunction in primary neonatal mouse cardiomyocytes," *PLoS One*, vol. 7, no. 11, article e50778, 2012.
- [128] W. Bi, J. Cai, P. Xue et al., "Protective effect of nitronyl nitroxide-amino acid conjugates on liver ischemia-reperfusion induced injury in rats," *Bioorganic & medicinal chemistry letters*, vol. 18, no. 6, pp. 1788–1794, 2008.
- [129] C. Jia, W. Wang, Y. Zhu, and S. Zheng, "Suprahepatic vena cava manipulative bleeding alleviates hepatic ischemia-reperfusion injury in rats," *Digestive and Liver Disease*, vol. 40, no. 4, pp. 285–292, 2008.
- [130] Y. Okatani, A. Wakatsuki, R. J. Reiter, H. Enzan, and Y. Miyahara, "Protective effect of melatonin against mitochondrial injury induced by ischemia and reperfusion of rat liver," *European journal of pharmacology*, vol. 469, no. 1-3, pp. 145–152, 2003.
- [131] H. C. Xing, L. J. Li, K. J. Xu et al., "Intestinal microflora in rats with ischemia/reperfusion liver injury," *Journal of Zhejiang University Science. B*, vol. 6, no. 1, pp. 14–21, 2005.
- [132] H. C. Huang, S. J. Hsu, C. C. Chang et al., "Lycopene treatment improves intrahepatic fibrosis and attenuates pathological angiogenesis in biliary cirrhotic rats," *Journal of the Chinese Medical Association*, vol. 85, no. 4, pp. 414–420, 2022.
- [133] J. Mustra Rakic, C. Liu, S. Veeramachaneni et al., "Dietary lycopene attenuates cigarette smoke-promoted nonalcoholic steatohepatitis by preventing suppression of antioxidant enzymes in ferrets," *The Journal of nutritional biochemistry*, vol. 91, article 108596, 2021.
- [134] A. N. Mekuria, A. K. Tura, B. Hagos et al., "Anti-cancer effects of lycopene in animal models of hepatocellular carcinoma: a systematic review and meta-analysis," *Frontiers in pharmacology*, vol. 11, p. 1306, 2020.
- [135] H. F. Goode, N. R. Webster, P. D. Howdle et al., "Reperfusion injury, antioxidants and hemodynamics during orthotopic liver transplantation," *Hepatology*, vol. 19, no. 2, pp. 354–359, 1994.
- [136] G. Bayramoglu, A. Bayramoglu, Y. Altuner, M. Uyanoglu, and S. Colak, "The effects of lycopene on hepatic ischemia/reperfusion injury in rats," *Cytotechnology*, vol. 67, no. 3, pp. 487–491, 2015.

- [137] R. Xue, J. Qiu, S. Wei et al., "Lycopene alleviates hepatic ischemia reperfusion injury via the Nrf2/HO-1 pathway mediated NLRP3 inflammasome inhibition in Kupffer cells," *Annals of Translational Medicine*, vol. 9, no. 8, p. 631, 2021.
- [138] B. Liu, L. Yan, X. Jiao et al., "Lycopene alleviates hepatic hypoxia/reoxygenation injury through Nrf2/HO-1 pathway in AML12 cell," *Journal of Interferon & Cytokine Research*, vol. 40, no. 8, pp. 406–417, 2020.
- [139] L. Bignotto, J. Rocha, B. Sepodes et al., "Anti-inflammatory effect of lycopene on carrageenan-induced paw oedema and hepatic ischaemia-reperfusion in the rat," *British Journal of Nutrition*, vol. 102, no. 1, pp. 126–133, 2009.
- [140] Z. Tan, Y. Shi, Y. Yan, W. Liu, G. Li, and R. Li, "Impact of endogenous hydrogen sulfide on toll-like receptor pathway in renal ischemia/reperfusion injury in rats," *Ren Fail*, vol. 37, no. 4, pp. 727–733, 2015.
- [141] H. Zhao, A. Alam, A. P. Soo, A. J. George, and D. Ma, "Ischemia-reperfusion injury reduces long term renal graft survival: mechanism and beyond," *EBioMedicine*, vol. 28, pp. 31–42, 2018.
- [142] E. İhtiyar, N. F. Yaşar, N. Erkasap et al., "Effects of doxycycline on renal ischemia reperfusion injury induced by abdominal compartment syndrome," *Journal of Surgical Research*, vol. 167, no. 1, pp. 113–120, 2011.
- [143] A. Kucuk, S. Kabadere, M. Tosun et al., "Protective effects of doxycycline in ischemia/reperfusion injury on kidney," *Journal of physiology and biochemistry*, vol. 65, no. 2, pp. 183–191, 2009.
- [144] J. V. Bonventre and L. Yang, "Cellular pathophysiology of ischemic acute kidney injury," *The Journal of clinical investigation*, vol. 121, no. 11, pp. 4210–4221, 2011.
- [145] Y. A. Hussien, H. Abdalkadim, W. Mahbuba, N. R. Hadi, D. A. Jamil, and H. A. al-Aubaidy, "The nephroprotective effect of lycopene on renal ischemic reperfusion injury: a mouse model," *Indian Journal of Clinical Biochemistry*, vol. 35, no. 4, pp. 474–481, 2020.
- [146] C. Kaya, R. Karabulut, Z. Turkyilmaz et al., "Lycopene has reduced renal damage histopathologically and biochemically in experimental renal ischemia-reperfusion injury," *Ren Fail*, vol. 37, no. 8, pp. 1390–1395, 2015.
- [147] A. Pektaş, H. Gemalmaz, M. Balkaya et al., "The short-term protective effects of lycopene on renal ischemia-reperfusion injury in rats," *Turkish journal of urology*, vol. 40, no. 1, pp. 46–51, 2014.
- [148] Y. Gursoy-Ozdemir, A. Can, and T. Dalkara, "Reperfusion-induced oxidative/nitrative injury to neurovascular unit after focal cerebral ischemia," *Stroke*, vol. 35, no. 6, pp. 1449–1453, 2004.
- [149] J. Aronowski, R. Strong, and J. C. Grotta, "Reperfusion injury: demonstration of brain damage produced by reperfusion after transient focal ischemia in rats," *Journal of Cerebral Blood Flow & Metabolism*, vol. 17, no. 10, pp. 1048–1056, 1997.
- [150] J. M. Hallenbeck and A. J. Dutka, "Background review and current concepts of reperfusion injury," *Archives of Neurology*, vol. 47, no. 11, pp. 1245–1254, 1990.
- [151] S. Kuroda and B. K. Siesjo, "Reperfusion damage following focal ischemia: pathophysiology and therapeutic windows," *Clinical Neuroscience*, vol. 4, no. 4, pp. 199–212, 1997.
- [152] B. C. White, A. Daya, D. J. DeGracia et al., "Fluorescent histochemical localization of lipid peroxidation during brain reperfusion following cardiac arrest," *Acta neuropathologica*, vol. 86, no. 1, pp. 1–9, 1993.
- [153] A. Wu, R. Liu, W. Dai et al., "Lycopene attenuates early brain injury and inflammation following subarachnoid hemorrhage in rats," *International Journal of Clinical and Experimental Medicine*, vol. 8, no. 8, pp. 14316–14322, 2015.
- [154] Z. Xu, C. Liu, R. Wang, X. Gao, C. Hao, and C. Liu, "A combination of lycopene and human amniotic epithelial cells can ameliorate cognitive deficits and suppress neuroinflammatory signaling by choroid plexus in Alzheimer's disease rat," *The Journal of Nutritional Biochemistry*, vol. 88, article 108558, 2021.
- [155] A. Prakash and A. Kumar, "Lycopene protects against memory impairment and mito-oxidative damage induced by colchicine in rats: an evidence of nitric oxide signaling," *European journal of pharmacology*, vol. 721, no. 1–3, pp. 373–381, 2013.
- [156] H. Celik, S. Kucukler, S. Ozdemir et al., "Lycopene protects against central and peripheral neuropathy by inhibiting oxaliplatin-induced ATF-6 pathway, apoptosis, inflammation and oxidative stress in brains and sciatic tissues of rats," *Neurotoxicology*, vol. 80, pp. 29–40, 2020.
- [157] J. Karppi, J. A. Laukkanen, J. Sivenius, K. Ronkainen, and S. Kurl, "Serum lycopene decreases the risk of stroke in men: a population-based follow-up study," *Neurology*, vol. 79, no. 15, pp. 1540–1547, 2012.
- [158] T. Li, Y. Zhang, Y. Qi, and H. Liu, "Lycopene prevents oxygen-glucose deprivation-induced autophagic death in SH-SY5Y cells via inhibition of the oxidative stress-activated AMPK/mTOR pathway," *Molecular Medicine Reports*, vol. 24, no. 2, 2021.
- [159] Y. Zhao, Z. Xin, N. Li et al., "Nano-liposomes of lycopene reduces ischemic brain damage in rodents by regulating iron metabolism," *Free Radical Biology and Medicine*, vol. 124, pp. 1–11, 2018.
- [160] M. H. Jazayeri, H. Amani, A. A. Pourfatollah, A. Avan, G. A. Ferns, and H. Pazoki-Toroudi, "Enhanced detection sensitivity of prostate-specific antigen via PSA- conjugated gold nanoparticles based on localized surface plasmon resonance: GNP-coated anti-PSA/LSPR as a novel approach for the identification of prostate anomalies," *Cancer gene therapy*, vol. 23, no. 10, pp. 365–369, 2016.
- [161] S. Mehranfar, I. Abdi Rad, E. Mostafavi, and A. Akbarzadeh, "The use of stromal vascular fraction (SVF), platelet-rich plasma (PRP) and stem cells in the treatment of osteoarthritis: an overview of clinical trials," *Artificial cells, nanomedicine, and biotechnology*, vol. 47, no. 1, pp. 882–890, 2019.
- [162] Z. Liu, Y. Shen, Y. Wu et al., "An intrinsic therapy of gold nanoparticles in focal cerebral ischemia-reperfusion injury in rats," *Journal of biomedical nanotechnology*, vol. 9, no. 6, pp. 1017–1028, 2013.
- [163] H. Hyun, K. Lee, K. H. Min et al., "Ischemic brain imaging using fluorescent gold nanoprobe sensitive to reactive oxygen species," *Journal of controlled release*, vol. 170, no. 3, pp. 352–357, 2013.
- [164] H. Amani, E. Mostafavi, M. R. Alebouyeh et al., "Would colloidal gold nanocarriers present an effective diagnosis or treatment for ischemic stroke?," *International Journal of Nanomedicine*, vol. 14, pp. 8013–8031, 2019.
- [165] K. Fujita, N. Yoshimoto, T. Kato et al., "Lycopene inhibits ischemia/reperfusion-induced neuronal apoptosis in gerbil

- hippocampal tissue,” *Neurochemical research*, vol. 38, no. 3, pp. 461–469, 2013.
- [166] X. Lei, L. Lei, Z. Zhang, and Y. Cheng, “Neuroprotective effects of lycopene pretreatment on transient global cerebral ischemia-reperfusion in rats: the role of the Nrf2/HO-1 signaling pathway,” *Molecular Medicine Reports*, vol. 13, no. 1, pp. 412–418, 2016.
- [167] Y. Wei, X. Shen, H. Shen, J. Mai, M. Wu, and G. Yao, “Effects of lycopene on cerebral ischemia-reperfusion injury in rats,” *Wei Sheng Yan Jiu*, vol. 39, no. 2, pp. 201–204, 2010.
- [168] G. Hsiao, T. H. Fong, N. H. Tzu, K. H. Lin, D. S. Chou, and J. R. Sheu, “A potent antioxidant, lycopene, affords neuroprotection against microglia activation and focal cerebral ischemia in rats,” *In Vivo*, vol. 18, no. 3, pp. 351–356, 2004.
- [169] S. Pentylala, J. Lee, P. Yalamanchili, S. Vitkun, and S. A. Khan, “Testicular torsion: a review,” *Journal of lower genital tract disease*, vol. 5, no. 1, pp. 38–47, 2001.
- [170] R. C. Williamson, “The continuing conundrum of testicular torsion,” *Journal of British Surgery*, vol. 72, no. 7, pp. 509–510, 1985.
- [171] N. Yildirim, D. Simsek, S. Kose et al., “The protective effect of Ginkgo biloba in a rat model of ovarian ischemia/reperfusion injury: improvement in histological and biochemical parameters,” *Advances in clinical and experimental medicine*, vol. 27, no. 5, pp. 591–597, 2018.
- [172] R. C. Williamson, “Death in the scrotum: testicular torsion,” *New England Journal of Medicine*, vol. 296, no. 6, p. 338, 1977.
- [173] A. Hekimoglu, Z. Kurcer, F. Aral, F. Baba, E. Sahna, and A. Atessahin, “Lycopene, an antioxidant carotenoid, attenuates testicular injury caused by ischemia/reperfusion in rats,” *The Tohoku journal of experimental medicine*, vol. 218, no. 2, pp. 141–147, 2009.
- [174] M. Güzel, M. F. Sönmez, O. Baştuğ et al., “Effectiveness of lycopene on experimental testicular torsion,” *Journal of Pediatric Surgery*, vol. 51, no. 7, pp. 1187–1191, 2016.
- [175] E. P. T. Yilmaz, H. Un, B. Gundogdu et al., “Protective effect of lycopene against reperfusion injury in rats with ovarian torsion: a biochemical and histopathological evaluation,” *Journal of Laboratory Physicians*, vol. 12, no. 1, pp. 32–37, 2020.
- [176] D. A. Kirmizi, E. Baser, A. Okan, M. Kara, E. S. Yalvac, and Z. Doganyigit, “The effect of a natural molecule in ovary ischemia reperfusion damage: does lycopene protect ovary?,” *Experimental Animals*, vol. 70, no. 1, pp. 37–44, 2021.
- [177] M. Z. Schwartz, “Novel therapies for the management of short bowel syndrome in children,” *Pediatric surgery international*, vol. 29, no. 10, pp. 967–974, 2013.
- [178] H. Yasuhara, “Acute mesenteric ischemia: the challenge of gastroenterology,” *Surgery today*, vol. 35, no. 3, pp. 185–195, 2005.
- [179] C. A. Nankervis, P. J. Giannone, and K. M. Reber, “The neonatal intestinal vasculature: contributing factors to necrotizing enterocolitis,” *Seminars in perinatology*, vol. 32, no. 2, pp. 83–91, 2008.
- [180] O. Corcos and A. Nuzzo, “Gastro-intestinal vascular emergencies,” *Best practice & research clinical gastroenterology*, vol. 27, no. 5, pp. 709–725, 2013.
- [181] I. H. Mallick, W. Yang, M. C. Winslet, and A. M. Seifalian, “Ischemia-reperfusion injury of the intestine and protective strategies against injury,” *Digestive diseases and science*, vol. 49, no. 9, pp. 1359–1377, 2004.
- [182] B. Vollmar and M. D. Menger, “Intestinal ischemia/reperfusion: microcirculatory pathology and functional consequences,” *Langenbeck’s archives of surgery*, vol. 396, no. 1, pp. 13–29, 2011.
- [183] J. Grootjans, K. Lenaerts, W. A. Buurman, C. H. C. Dejong, and J. P. M. Derikx, “Life and death at the mucosal-luminal interface: new perspectives on human intestinal ischemia-reperfusion,” *World Journal of Gastroenterology*, vol. 22, no. 9, pp. 2760–2770, 2016.
- [184] Ö. İköz, N. Kahramansoy, H. Erkol, E. Koçoğlu, and T. Fırat, “Effects of lycopene in intestinal ischemia reperfusion injury via intestinal immunoglobulin A,” *Journal of Surgical Research*, vol. 267, pp. 63–70, 2021.
- [185] H. Turula and C. E. Wobus, “The role of the polymeric immunoglobulin receptor and secretory immunoglobulins during mucosal infection and immunity,” *Viruses*, vol. 10, no. 5, p. 237, 2018.
- [186] A. J. Macpherson, K. D. McCoy, F. E. Johansen, and P. Brandtzaeg, “The immune geography of IgA induction and function,” *Mucosal immunology*, vol. 1, no. 1, pp. 11–22, 2008.
- [187] P. L. R. Andrews, W. Cai, J. A. Rudd, and G. J. Sanger, “COVID-19, nausea, and vomiting,” *Journal of gastroenterology and hepatology*, vol. 36, no. 3, pp. 646–656, 2021.
- [188] R. C. Ungaro, T. Sullivan, J. F. Colombel, and G. Patel, “What should gastroenterologists and patients know about COVID-19?,” *Clinical Gastroenterology and Hepatology*, vol. 18, no. 7, pp. 1409–1411, 2020.
- [189] M. Mrugacz, A. Bryl, and K. Zorena, “Retinal vascular endothelial cell dysfunction and neuroretinal degeneration in diabetic patients,” *Journal of clinical medicine*, vol. 10, no. 3, p. 458, 2021.
- [190] M. Tanito, S. Kaidzu, Y. Takai, and A. Ohira, “Association between systemic oxidative stress and visual field damage in open-angle glaucoma,” *Scientific reports*, vol. 6, no. 1, p. 25792, 2016.
- [191] A. Ames, “Energy requirements of CNS cells as related to their function and to their vulnerability to ischemia: a commentary based on studies on retina,” *Canadian journal of physiology and pharmacology*, vol. 70, no. S1, pp. S158–S164, 1992.
- [192] M. Khayat, M. Williams, and N. Lois, “Ischemic retinal vein occlusion: characterizing the more severe spectrum of retinal vein occlusion,” *Survey of ophthalmology*, vol. 63, no. 6, pp. 816–850, 2018.
- [193] H. Noma, K. Yasuda, and M. Shimura, “Cytokines and pathogenesis of central retinal vein occlusion,” *Journal of clinical medicine*, vol. 9, no. 11, p. 3457, 2020.
- [194] F. Ulbrich, T. Lerach, J. Biermann et al., “Argon mediates protection by interleukin-8 suppression via a TLR2/TLR4/STAT3/NF- κ B pathway in a model of apoptosis in neuroblastoma cells in vitro and following ischemia-reperfusion injury in rat retinal vivo,” *Journal of neurochemistry*, vol. 138, no. 6, pp. 859–873, 2016.
- [195] C. R. Kim, J. H. Kim, H. L. Park, and C. K. Park, “Ischemia reperfusion injury triggers TNF α induced-necroptosis in rat retina,” *Current Eye Research*, vol. 42, no. 5, pp. 771–779, 2017.
- [196] T. Wei, Q. Kang, B. Ma, S. Gao, X. Li, and Y. Liu, “Activation of autophagy and paraptosis in retinal ganglion cells after retinal ischemia and reperfusion injury in rats,” *Experimental and therapeutic medicine*, vol. 9, no. 2, pp. 476–482, 2015.

- [197] E. Icel, A. Icel, T. Uçak et al., “The effects of lycopene on alloxan induced diabetic optic neuropathy,” *Cutaneous and ocular toxicology*, vol. 38, no. 1, pp. 88–92, 2019.
- [198] S. K. Gupta, D. Trivedi, S. Srivastava, S. Joshi, N. Halder, and S. D. Verma, “Lycopene attenuates oxidative stress induced experimental cataract development: an in vitro and in vivo study,” *Nutrition*, vol. 19, no. 9, pp. 794–799, 2003.
- [199] J. S. Coselli, S. A. LeMaire, L. D. Conklin, C. Köksoy, and Z. C. Schmittling, “Morbidity and mortality after extent II thoracoabdominal aortic aneurysm repair,” *The Annals of Thoracic Surgery*, vol. 73, no. 4, pp. 1107–1116, 2002.
- [200] M. F. Conrad, J. Y. Ye, T. K. Chung, J. K. Davison, and R. P. Cambria, “Spinal cord complications after thoracic aortic surgery: long-term survival and functional status varies with deficit severity,” *Journal of Vascular Surgery*, vol. 48, no. 1, pp. 47–53, 2008.
- [201] S. L. Drinkwater, A. Goebells, A. Haydar et al., “The incidence of spinal cord ischaemia following thoracic and thoracoabdominal aortic endovascular intervention,” *European Journal of Vascular and Endovascular Surgery*, vol. 40, no. 6, pp. 729–735, 2010.
- [202] S. N. Khan, G. Stansby, and Cochrane Vascular Group, “Cerebrospinal fluid drainage for thoracic and thoracoabdominal aortic aneurysm surgery,” *Cochrane Database of Systematic Reviews*, vol. 10, no. 10, p. CD003635, 2012.
- [203] X. L. Zhu, X. Chen, W. Wang et al., “Electroacupuncture pretreatment attenuates spinal cord ischemia-reperfusion injury via inhibition of high-mobility group box 1 production in a LXA₄ receptor-dependent manner,” *Brain Research*, vol. 1659, pp. 113–120, 2017.
- [204] H. Faheem, A. Mansour, A. Elkordy et al., “Neuroprotective effects of minocycline and progesterone on white matter injury after focal cerebral ischemia,” *Journal of Clinical Neuroscience*, vol. 64, pp. 206–213, 2019.
- [205] X. Jia, R. G. Kowalski, D. M. Sciubba, and R. G. Geocadin, “Critical care of traumatic spinal cord injury,” *Journal of intensive care medicine*, vol. 28, no. 1, pp. 12–23, 2013.
- [206] M. Karimipour, G. H. Farjah, F. Molazadeh, M. Ansari, and B. Pourheidar, “Protective effect of contralateral, ipsilateral, and bilateral remote ischemic preconditioning on spinal cord ischemia reperfusion injury in rats,” *Turkish Neurosurgery*, vol. 29, no. 6, pp. 933–939, 2019.
- [207] C. Hentia, A. Rizzato, E. Camporesi et al., “An overview of protective strategies against ischemia/reperfusion injury: the role of hyperbaric oxygen preconditioning,” *Brain and Behavior*, vol. 8, no. 5, article e00959, 2018.
- [208] Y. Hua, N. Xu, T. Ma, Y. Liu, H. Xu, and Y. Lu, “Anti-inflammatory effect of lycopene on experimental spinal cord ischemia injury via cyclooxygenase-2 suppression,” *Neuroimmunomodulation*, vol. 26, no. 2, pp. 84–92, 2019.
- [209] H. L. Li, Y. C. Chan, D. Z. Cui, and S. W. Cheng, “Acute thrombosis of an infrarenal abdominal aortic aneurysm presenting as bilateral critical lower limb ischemia,” *Vascular and Endovascular Surgery*, vol. 55, no. 2, pp. 186–188, 2021.
- [210] P. Brahmamdam, M. Plummer, J. G. Modrall, S. M. Megison, G. P. Clagett, and R. J. Valentine, “Hand ischemia associated with elbow trauma in children,” *Journal of vascular surgery*, vol. 54, no. 3, pp. 773–778, 2011.
- [211] J. M. Sacks, Y. R. Kuo, E. K. Horibe et al., “An optimized dual-surgeon simultaneous orthotopic hind-limb allotransplantation model in rats,” *Journal of reconstructive microsurgery*, vol. 28, no. 1, pp. 69–75, 2012.
- [212] Y. Schmidt, H. Bannasch, and S. U. Eisenhardt, “Ischemia-reperfusion injury leads to significant tissue damage in free flap surgery,” *Plastic and reconstructive surgery*, vol. 129, no. 1, pp. 174e–175e, 2012.
- [213] M. Utsunomiya, “Endovascular therapy for acute limb ischemia,” *Journal of Atherosclerosis and Thrombosis*, vol. 28, no. 11, pp. 1126–1127, 2021.
- [214] T. Betz, I. Toepel, K. Pfister, M. Steinbauer, and C. Uhl, “Role of aorto(bi)femoral bypass in treatment of patients with critical limb threatening ischemia,” *Surgeon*, vol. 19, no. 2, pp. 103–110, 2021.
- [215] C. Argyriou, G. S. Georgiadis, M. Mantatzis et al., “Hybrid reconstruction of the upper limb in a patient with chronic limb ischemia,” *Vascular*, vol. 23, no. 6, pp. 653–656, 2015.
- [216] D. Barthels and H. Das, “Generation of acute hind limb ischemia in NOD/SCID mice and treatment with nanofiber-expanded CD34(+) hematopoietic stem cells,” *Methods in molecular biology*, vol. 2193, pp. 121–128, 2021.
- [217] F. W. Blaisdell, “The pathophysiology of skeletal muscle ischemia and the reperfusion syndrome: a review,” *Cardiovascular Surgery*, vol. 10, no. 6, pp. 620–630, 2002.
- [218] C. C. Liu, C. C. Huang, W. T. Lin et al., “Lycopene supplementation attenuated xanthine oxidase and myeloperoxidase activities in skeletal muscle tissues of rats after exhaustive exercise,” *British Journal of Nutrition*, vol. 94, no. 4, pp. 595–601, 2005.
- [219] M. Kirişçi, B. Güneri, M. Seyithanoğlu, Ü. Kazancı, A. Doğaner, and H. Güneş, “The protective effects of lycopene on ischemia/reperfusion injury in rat hind limb muscle model,” *Turkish Journal of Trauma and Emergency Surgery*, vol. 26, no. 3, pp. 351–360, 2020.
- [220] N. Karimi, M. Haghani, A. Noorafshan, and S. M. S. Moosavi, “Structural and functional disorders of hippocampus following ischemia/reperfusion in lower limbs and kidneys,” *Neuroscience*, vol. 358, pp. 238–248, 2017.
- [221] D. Garbaisz, Z. Turoczy, P. Aranyi et al., “Attenuation of skeletal muscle and renal injury to the lower limb following ischemia-reperfusion using mPTP inhibitor NIM-811,” *PLoS One*, vol. 9, no. 6, article e101067, 2014.
- [222] M. Kirisci, B. Guneri, M. Seyithanoglu, and U. Kazanci, “Lycopene hampers lung injury due to skeletal muscle ischemia-reperfusion in rat model,” *International Journal for Vitamin and Nutrition Research*, vol. 92, no. 3-4, pp. 240–247, 2020.
- [223] T. Chen, B. Ye, J. Tan, H. Yang, F. He, and R. A. Khalil, “CD146⁺ Mesenchymal stem cells treatment improves vascularization, muscle contraction and VEGF expression, and reduces apoptosis in rat ischemic hind limb,” *Biochemical Pharmacology*, vol. 190, article 114530, 2021.
- [224] J. Y. Cao, B. Wang, T. T. Tang et al., “Exosomal miR-125b-5p deriving from mesenchymal stem cells promotes tubular repair by suppression of p53 in ischemic acute kidney injury,” *Theranostics*, vol. 11, no. 11, pp. 5248–5266, 2021.
- [225] X. Rong, J. Liu, X. Yao, T. Jiang, Y. Wang, and F. Xie, “Human bone marrow mesenchymal stem cells-derived exosomes alleviate liver fibrosis through the Wnt/ β -catenin pathway,” *Stem Cell Research & Therapy*, vol. 10, no. 1, p. 98, 2019.
- [226] A. Monsel, Y. G. Zhu, S. Gennai, Q. Hao, J. Liu, and J. W. Lee, “Cell-based therapy for acute organ injury: preclinical

- evidence and ongoing clinical trials using mesenchymal stem cells,” *Anesthesiology*, vol. 121, no. 5, pp. 1099–1121, 2014.
- [227] Y. Peng, J. L. Zhao, Z. Y. Peng, W. F. Xu, and G. L. Yu, “Exosomal miR-25-3p from mesenchymal stem cells alleviates myocardial infarction by targeting pro-apoptotic proteins and EZH2,” *Cell Death & Disease*, vol. 11, no. 5, p. 317, 2020.
- [228] L. Liu, X. Jin, C. F. Hu, R. Li, Z. Zhou, and C. X. Shen, “Exosomes derived from mesenchymal stem cells rescue myocardial ischaemia/reperfusion injury by inducing cardiomyocyte autophagy via AMPK and Akt pathways,” *Cellular physiology and biochemistry*, vol. 43, no. 1, pp. 52–68, 2017.
- [229] H. Cao, Y. Cheng, H. Gao et al., “In Vivo Tracking of mesenchymal stem cell-derived extracellular vesicles improving mitochondrial function in renal ischemia-reperfusion injury,” *ACS Nano*, vol. 14, no. 4, pp. 4014–4026, 2020.
- [230] A. Sahu, J. Jeon, M. S. Lee, H. S. Yang, and G. Tae, “Nanozyme impregnated mesenchymal stem cells for hepatic ischemia-reperfusion injury alleviation,” *ACS Applied Materials & Interfaces*, vol. 13, no. 22, pp. 25649–25662, 2021.
- [231] R. Li, K. Zhao, Q. Ruan, C. Meng, and F. Yin, “Bone marrow mesenchymal stem cell-derived exosomal microRNA-124-3p attenuates neurological damage in spinal cord ischemia-reperfusion injury by downregulating Ern1 and promoting M2 macrophage polarization,” *Arthritis Research & Therapy*, vol. 22, no. 1, p. 75, 2020.
- [232] X. Liu, M. Zhang, H. Liu et al., “Bone marrow mesenchymal stem cell-derived exosomes attenuate cerebral ischemia-reperfusion injury-induced neuroinflammation and pyroptosis by modulating microglia M1/M2 phenotypes,” *Experimental Neurology*, vol. 341, article 113700, 2021.
- [233] L. H. Silva, M. A. Antunes, C. C. Dos Santos, D. J. Weiss, F. F. Cruz, and P. R. Rocco, “Strategies to improve the therapeutic effects of mesenchymal stromal cells in respiratory diseases,” *Stem Cell Research & Therapy*, vol. 9, no. 1, p. 45, 2018.
- [234] Y. Jin, T. Kato, M. Furu et al., “Mesenchymal stem cells cultured under hypoxia escape from senescence via down-regulation of p16 and extracellular signal regulated kinase,” *Biochemical and Biophysical Research Communications*, vol. 391, no. 3, pp. 1471–1476, 2010.
- [235] S. H. Bhang, S. W. Cho, W. G. La et al., “Angiogenesis in ischemic tissue produced by spheroid grafting of human adipose-derived stromal cells,” *Biomaterials*, vol. 32, no. 11, pp. 2734–2747, 2011.
- [236] J. K. Roh, K. H. Jung, and K. Chu, “Adult stem cell transplantation in stroke: its limitations and prospects,” *Current Stem Cell Research & Therapy*, vol. 3, no. 3, pp. 185–196, 2008.
- [237] Y. Li, F. Xue, S. Z. Xu, X. W. Wang, X. Tong, and X. J. Lin, “Lycopene protects bone marrow mesenchymal stem cells against ischemia-induced apoptosis in vitro,” *European Review for Medical and Pharmacological Sciences*, vol. 18, no. 11, pp. 1625–1631, 2014.
- [238] J. Y. Kim, J. S. Lee, Y. S. Han et al., “Pretreatment with lycopene attenuates oxidative stress-induced apoptosis in human mesenchymal stem cells,” *Biomolecules & Therapeutics*, vol. 23, no. 6, pp. 517–524, 2015.
- [239] Z. Bajory, A. Szabó, I. Király, L. Pajor, and K. Messmer, “Involvement of nitric oxide in microcirculatory reactions after ischemia-reperfusion of the rat urinary bladder,” *European Surgical Research*, vol. 42, no. 1, pp. 28–34, 2009.
- [240] T. Tarcan, K. Rademakers, S. Arlandis, A. von Gontard, G. A. van Koeveeringe, and P. Abrams, “Do the definitions of the underactive bladder and detrusor underactivity help in managing patients: International Consultation on Incontinence Research Society (ICI-RS) Think Tank 2017?,” *Neurourology and Urodynamics*, vol. 37, no. S4, pp. S60–S68, 2018.
- [241] T. Tarcan, K. M. Azadzi, M. B. Siroky, I. Goldstein, and R. J. Krane, “Age-related erectile and voiding dysfunction: the role of arterial insufficiency,” *British journal of urology*, vol. 82, Supplement 1, pp. 26–33, 1998.
- [242] F. Fusco, M. Creta, C. De Nunzio et al., “Progressive bladder remodeling due to bladder outlet obstruction: a systematic review of morphological and molecular evidences in humans,” *BMC urology*, vol. 18, no. 1, p. 15, 2018.
- [243] A. T. Lin and Y. S. Juan, “Ischemia, hypoxia and oxidative stress in bladder outlet obstruction and bladder overdistention injury,” *LUTS: Lower Urinary Tract Symptoms*, vol. 4, Supplement 1, pp. 27–31, 2012.
- [244] Y. S. Juan, W. Y. Lin, C. Kalorin, B. A. Kogan, R. M. Levin, and A. Mannikarottu, “The effect of partial bladder outlet obstruction on carbonyl and nitrotyrosine distribution in rabbit bladder,” *Urology*, vol. 70, no. 6, pp. 1249–1253, 2007.
- [245] A. Mannikarottu, A. D. Lin, C. Whitebeck, R. Leggett, B. Kogan, and R. Levin, “Effect of partial bladder outlet obstruction on nitrotyrosine levels and their correlation with contractile function,” *Neurourol Urodyn*, vol. 25, no. 4, pp. 397–401, 2006.
- [246] G. Bratslavsky, C. Whitbeck, P. Horan, and R. M. Levin, “Effects of in vivo ischemia on contractile responses of rabbit bladder to field stimulation, carbachol, ATP and KCL,” *Pharmacology*, vol. 59, no. 4, pp. 221–226, 1999.
- [247] M. W. Chen, R. M. Levin, P. Horan, and R. B. Buttyan, “Effects of unilateral ischemia on the contractile response of the bladder: protective effect of tadenan (Pygeum africanum extract),” *Molecular Urology*, vol. 3, no. 1, pp. 5–10, 1999.
- [248] M. W. Chen, R. Buttyan, and R. M. Levin, “Genetic and cellular response to unilateral ischemia of the rabbit urinary bladder,” *The Journal of urology*, vol. 155, no. 2, pp. 732–737, 1996.
- [249] F. Pessina, K. Marazova, P. Ninfali, L. Avanzi, S. Manfredini, and G. Sgaragli, “In vitro neuroprotection by novel antioxidants in guinea-pig urinary bladder subjected to anoxia-glucopenia/reperfusion damage,” *Naunyn-Schmiedeberg’s Archives of Pharmacology*, vol. 370, no. 6, pp. 521–528, 2004.