

Author, Year	Assessment Tool	Assessment Periods	Prevalence	Severity Definitions	Severity Prevalence
<b>Giallo, 2014</b>	Kessler-6 (K6)	12 months 2-3 years 4-5 years 6-7 years	12.80% 9.60% 10.10% 11.40%	Reported as either in the symptomatic range (K6 ≥8) or clinical range (>13)	Clinical range: 2.80% 2.20% 2.50% 2.80%
<b>Mayberry, 2007</b>	Edinburgh Postnatal Depression Scale (EPDS)	0-6 months 7-12 months 13-18 months 19-24 months	34.10% 31.20% 28.10% 33.50%	Reported as mild (EPDS 10-12) or moderately-severely (EPDS ≥13) depressed	Mild, mod-severe: 11.00%, 23.10% 15.10%, 16.10% 11.00%, 17.10% 13.10%, 20.40%
<b>Morrissey, 2013</b>	Center for Epidemiologic Studies Depression Scale (CES-D) and Composite International Diagnostic Interview (CIDI)	9 months 2 years 4 years 5.5 years	17.68% 17.64% 19.11% 16.54%	Reported overall proportions of women experiencing moderate (CES-D ≥9) or severe (CES-D >15)	Overall: 17-19% of mothers considered as having mod or severe symptoms, with 5-8% of these being severe
<b>Schultz, 2020</b>	Beck Depression Inventory (BDI-II)	4-5 years	16.70%	Reported as minimal (BDI 0-11), mild (BDI 12-19), moderate (BDI 20-35) or severe (36-63) depression	Min, mild, mod-severe: 52.30%, 14.40%, 16.60%
<b>Tripathy, 2010</b>	Kessler-10 (K10)	2 years 3 years	14.00% 11.00%	Reported as no/mild (K10 10-15), moderate (K10 16-30) or severe (K10 31-50) depression	Mild, mod-severe: 87.00%, 14.00% 90.00%, 11.00%