



Pet dogs' behavior when the owner and an unfamiliar person attend to a faux rival

Emanuela Prato-Previde^{1‡}*, Velia Nicotra^{2,3‡}, Annalisa Pelosi², Paola Valsecchi³

- 1 Dipartimento di Fisiopatologia Medico-Chirurgica e dei Trapianti, Università degli Studi di Milano, Milano, Italy, 2 Dipartimento di Medicina e Chirurgia, Università degli Studi di Parma, Parma, Italy, 3 Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale, Università degli Studi di Parma, Parma, Italy
- ‡ EPP and VN are co-first authors on this work.
- * emanuela.pratoprevide@unimi.it



G OPEN ACCESS

Citation: Prato-Previde E, Nicotra V, Pelosi A, Valsecchi P (2018) Pet dogs' behavior when the owner and an unfamiliar person attend to a faux rival. PLoS ONE 13(4): e0194577. https://doi.org/10.1371/journal.pone.0194577

Editor: Cheryl S. Rosenfeld, University of Missouri Columbia, UNITED STATES

Received: September 14, 2017

Accepted: March 6, 2018

Published: April 18, 2018

Copyright: © 2018 Prato-Previde et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data files are available from Mendeley database: https://data.mendeley.com/library; http://dx.doi.org/10.17632/g8k5t4f645.1.

Funding: This work was partially funded by Velia Nicotra's Doctoral Grant supported by Università degli Studi di Parma.

Competing interests: The authors have declared that no competing interests exist.

Abstract

While dog owners ascribe different emotions to their pets, including jealousy, research on secondary emotions in nonhuman animals is very limited and, so far, only one study has investigated jealousy in dogs (Canis familiaris). This work explores jealousy in dogs one step further. We conducted two studies adapting a procedure devised to assess jealousy in human infants. In each study 36 adult dogs were exposed to a situation in which their owner and a stranger ignored them while directing positive attention towards three different objects: a book, a puppet and a fake dog (Study 1: furry; Study 2: plastic). Overall, the results of both studies do not provide evidence that the behavioral responses of our dogs were triggered by jealousy: we did not find a clear indication that the fake dogs were perceived as real social rivals, neither the furry nor the plastic one. Indeed, dogs exhibited a higher interest (i.e. look at, interact with) towards the fake dogs, but differences in the behavior towards the fake dog and the puppet only emerged in Study 2. In addition, many of the behaviors (protest, stress, attention seeking, aggression) that are considered distinctive features of jealousy were not expressed or were expressed to a limited extent, revealing that dogs did not actively try to regain their owner's attention or interfere with the interaction between the owner and the faux rival. Finally, a differentiated response towards the attachment figure (the owner) and the unfamiliar person (the stranger) did not emerge. Differently from what reported in human infants, dogs' behavior towards the attachment figure and the stranger interacting with the potential competitor (in this case, the fake dog) did not significantly differ: in both studies dogs paid attention to the owner and the stranger manipulating the fake dog to the same extent. In conclusion, we do not exclude that dogs could possess a rudimentary form of jealousy, but we suggest that research on this topic should require the use of a real social interloper (conspecific or human) and more naturalistic procedures.

Introduction

Scientists generally agree that some emotions ("primary" or "basic") have a long evolutionary history and can be found across a wide range of vertebrate species, due to their fundamental



adaptive value [1–6]. Conversely, the presence of "secondary" or "complex" emotions in non-human species is a lively debated topic. Many researchers still assume that these emotions require elaborate cognitive abilities and emerge relatively late in human development. It is also believed that they imply self and interpersonal awareness, which is generally considered a uniquely human characteristic [7]. Nevertheless, the perspective on secondary emotions is changing as evidences deriving from developmental and cross-species studies suggest that at least some secondary emotions, due to their fundamental role in regulating social life, are present in other species. In addition, this view is supported by the presence of specific cortical structures and psychological infrastructures in at least some mammals [3,8–12].

Jealousy is a secondary emotion that appears to have a clear, strong adaptive value in maintaining and protecting social relationships and bonds (i.e. sibling-parent, sexual, friendship), enhancing the individual's fitness [9,13-14]. In the human psychological literature jealousy has been defined as a context-dependent social emotion that requires a social triangle and arises when one individual perceives that an intruder is threatening an important relationship. It is generally expressed by observable negative affective responses (e.g. fear, anger, sadness) and accompanied by overt behaviors directed at restoring the relationship by reducing the threat represented by the interloper and regaining attention and care from a significant social partner [14,15]. It has been proposed that jealousy could also occur outside conscious awareness [9,16], without the need of cortically mediated cognition [13] and in the absence of complex interpretations of the meaning of the social interaction [17]. Thus, at least a primordial form of jealousy could arise in other animals in specific situations where a significant relationship is threatened. The best evidence that jealousy could have a primordial form derives from the literature on human infants, which shows that jealousy can be exhibited within the first two years of life in specific social situations (see [15,18] for reviews). There are several studies indicating that human infants aged from six to twelve months exhibited more protest behaviors, negative vocalizations and proximity seeking (approach and gaze) when their mother was holding an infant-like doll compared with when she held a book and when a stranger held the doll [19-21]. Taken together, these studies suggest that infants are sensitive to the loss of maternal attention, distinguish between social and non-social objects and do not show an undifferentiated general response to any person or object, but react to potential threats to the relationship with the attachment figure. Thus, jealousy could initially have evolved as a behavioral/emotional strategy to protect material and affective resources within the parent-offspring relationship [22]. The fact that jealousy can be recognizable in cognitively immature infants also suggests that this "secondary" emotion could occur with various degrees of complexity across social species, rather than being an all-or-none phenomenon, suddenly appearing in humans during the 2nd year of life or later. In addition, the adaptive function of jealousy in human infants has important theoretical implications since many nonhuman animals, particularly those living in social groups, could face situations that in principle would evoke jealousy in humans [23]. Comparative studies are needed to better understand the evolutionary emergence and the functions of jealousy and its relation with cognitive abilities.

The domestic dog (*Canis lupus familiaris*) has proved to be an interesting and promising animal model to investigate the evolution of social cognition and in the last 20 years researchers showed that dogs have quite sophisticated socio-cognitive abilities that in some cases parallel those reported for human infants [24–31]. Moreover, a growing body of research is shedding light on their emotional life ([12] for a review), suggesting that, besides primary emotions, dogs could also experience, at least to some extent, more complex emotional states, such as empathic-like responses [32,33] and inequity aversion [34–36]. Recently, Harris & Prouvost [37] suggested that dogs could express jealousy when their owner directs his/her attention to a potential rival, providing some support to people's belief that animals, especially dogs, show



jealousy when attention and affection are given to another person or animal [38–40]. Indeed, dogs are suitable subjects for investigating the existence of jealousy as they: 1. form stable groups and differentiate social relationships with conspecifics [41–42]; 2. establish a strong relationship with their owner, characterized by dependency for physical and psychological resources, which is functionally comparable to an infantile attachment [43]; 3. discriminate human emotions [44–46], are sensitive to others' attentional states [47–49] and to unfair treatment [34–36]. Hence, comparing dogs and infants, using the same type of procedures and analyses, may help to unravel proximate and ultimate causes of emotional behaviors.

Adapting a paradigm from human infant studies, Harris and Prouvost [37] tested dogs at their homes in three different conditions in which their owner ignored them while affectionately interacting with different objects. In the "jealousy" condition owners petted and sweetly talked to a realistic-looking stuffed dog that barked and wagged its tail; in the second condition, they directed the same behavior to a jack-o-lantern pail; in the third condition, owners were instructed to read aloud a book which played melodies. At the end of each condition, the owner put the object down within the dog's reach and walked away. It emerged that a significantly higher percentage of dogs showed behaviors that the authors considered indicative of jealousy (pushing/touching the owner and the object, getting between the owner and the object, snapping and whining) when the owner was manipulating the stuffed dog compared to both the jack-o-lantern and the book. These findings represent an interesting starting point to assess jealousy in dogs. However, some methodological issues suggest caution in drawing conclusions: for example, the authors recorded the number of subjects expressing a certain behavior in each condition, but this only provides partial information on the phenomenon, since the robustness of the behavioral response (e.g. duration) is neglected. When they estimated the amount of time (percentage of point samples) spent by dogs attending to the owner (i.e. looking at the owner), no difference emerged between the stuffed dog and the novel object (jack-olantern) conditions. Conversely, dogs directed a significantly higher attention (i.e. looking) to the animated stuffed dog compared to both the jack-o-lantern and the book. These results could be due to different emotional/motivational states rather than jealousy; as many pet dogs are familiar with furry and squeaking toys and are used to play with them, it cannot be excluded that they considered the stuffed dog just a toy to play with. Last but not least, in our opinion, since the function of jealousy is to protect a valuable relationship from an intruder, in order to provide more conclusive evidence it would be important to compare the dogs' behavior towards the owner (the attachment figure) and towards an unfamiliar person giving attention to a potential rival. Human infants differentiate among potential rivals (e.g. a book vs. a realistic doll) and also react in different ways when their mother rather than an unfamiliar adult gives positive attention to a potential rival [18].

The current work explores jealousy in dogs one step further, pursuing two goals: 1. clarify whether dogs can really be deceived by a fake dog, considering it a social rival rather than a toy; 2. evaluate whether the identity of the humans involved plays a role in triggering jealousy behaviors.

The dogs were presented with two types of fake dogs differing in appearance and size (Study 1: furry stuffed dog; Study 2: plastic dog), a novel object (a puppet) that did not appear like a conspecific but had some potentially appealing features for a dog (big eyes and soft fabric), and a picture book (a familiar and non social object). The latter was used as a control stimulus to ascertain to what extent dogs would react to the mere loss of people's attention. The identity of the handler of the potential rivals was controlled introducing an unfamiliar person in the testing situation, as done by Hart and colleagues [19] with human infants. Thus, in our testing situation, both the owner and an unfamiliar person interacted with the three different objects, directing positive attention and affection towards them. Assuming that our



experimental subjects could be deceived by the fake dogs and therefore be driven to jealousy, we expected they would: 1. sniff the ano-genital region of the fake dog, a behavioral pattern that dogs exhibit during social interactions with conspecifics for individual recognition [37]; 2. show some aggression towards the fake dog to prevent the rival to interfere with the relationship with the owner [37,50]; 3. show higher levels of attention/interaction, vocalizations and stress signals in the fake dog condition compared to the puppet condition to gain their owner attention and to disrupt the interaction [37,19]; 4. show attention/interaction, vocalizations and stress signals only when their owner gave affection to the fake dog, since dogs are not affectionately bonded with the stranger [19]; 5. show a limited behavioral response in the book condition, since the book is just a familiar and non-social object [37,19].

Material and methods

Experimental setting and stimuli

Testing took place in two different locations: one at the Canis sapiens Lab of the Università degli Studi di Milano and the other one at the Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale of the Università degli Studi di Parma. Testing rooms were not identical but both measured approximately $4.5 \times 3.5 \, \mathrm{m}$ and were equipped with two chairs (one for the stranger and one for the owner) facing each other at a distance of 120 cm, a soft plastic carpet between them, a water bowl and a small table with a computer positioned on it. The computer was used to guide and standardize the owner's and the stranger's actions throughout the procedure: a PowerPoint presentation provided written instructions at predetermined fixed intervals. An HD video camera was placed on a wall in a corner of the room to record the test.

Three different objects were used during the test: a fake dog, a soft puppet and a picture book. In Study 1 the fake dog was a furry Yorkshire terrier (length 32 cm, height at the withers 18 cm, Fig 1A); in Study 2 the fake dog was a plastic Fox terrier (length 50 cm, height at the withers 30 cm, Fig 1B). In both studies the puppet was a soft hand-made bag with a light grey fleece and two 'eyes' fixed on it (40 x 37 cm, Fig 1C). The objects were all static and did not emit sounds.

Two different female researchers (one in Milano and the other in Parma) played the role of the stranger.

Experimental design

To test our hypotheses, in both studies we set up a 3 (object type: fake dog, puppet, book) x 2 (object handler: owner, stranger) within-subjects design. Thus, each dog had the opportunity to observe and interact with both the owner and the stranger manipulating the three objects.







Fig 1. The three objects used in study 1 and 2. A: Furry fake dog; B: Plastic fake dog; C: Puppet.

https://doi.org/10.1371/journal.pone.0194577.g001



An a priori power analysis (G^* Power 3.1.9.2) for our experimental design (3x2 within-subject) setting alfa = .05, power = .90, an expected medium effect size f = .25, correlation between measures r = .5 and sphericity close to 1 (epsilon = .9), indicated a minimum sample size of 25 subjects.

Ethical statement. All procedures were performed in full accordance with Italian legal regulations (National Directive n. 26/14—Directive 2010/63/UE) and the guidelines for the treatments of animals in behavioral research and teaching of the Association for the Study of Animal Behavior (ASAB). The protocol was approved by the Ethical Committee for the Use of Animals (PROT.N. 105/OPBA/2016) of the Università degli Studi di Parma. A written consent to video-record and use data in an anonymous form was obtained by the owners prior to testing.

Subjects

Study 1. Thirty-six healthy adult dogs (15 males, 21 females; 20 pure breed, 16 mixed breed), ranging in age from 1 to 13 years (mean = 5.24 years, SD = 2.74;), and their owners participated in the study. Dog-owner dyads were recruited both by personal contact and from advertisement distributed within the Università di Milano.

Dogs' inclusion criteria were: being kept exclusively for companionship, living within the human household for at least 1 year, being accustomed to encounter human strangers, being at least 1- year old. Dog-owner dyads were randomly allocated to the different testing sequences and thus we tested 6 dogs for each sequence.

Study 2. Thirty-six healthy adult dogs (17 males, 19 females; 19 pure breed, 17 mixed breed), ranging in age from 1 to 10 years (mean = 4.74 years, SD = 2.5), and their owners participated in the study. Dog-owner dyads were recruited both by personal contact and from advertisement distributed within the Università di Parma.

Inclusion criteria were the same as in Study 1. Dog-owner dyads were randomly allocated to the different testing sequences and thus we tested 6 dogs for each sequence.

Procedure

Researchers met the owner-dog dyads outside the Departments and escorted them to a waiting room were the owner signed a consent form and filled up a questionnaire with information about the dog (age, breed, education and lifestyle). Next, the owner and the dog entered the testing room and the experimenter (a researcher different from the stranger) explained the test procedure to the owner, while the dog was free to explore the room for approximately 10 minutes. The experimenter then left the room. The test started with a 1-min familiarization phase during which the owner initially walked around the room paying attention to some wall posters and then sat on the owner's chair and interacted with his/her dog. The procedure comprised three experimental episodes, each consisting of two 1-min phases during which the owner and the stranger (a researcher unfamiliar to the dog) were both in the room and ignored the dog. In each experimental episode the owner and the stranger alternately handled one of the objects for 1 minute (book, puppet, fake dog) while talking to each other in a friendly manner. For sake of simplicity, the 6 phases will be hereinafter indicated as follows:

FDOW: the fake dog (FD) is handled by the owner (OW);

FDSTR: the fake dog is handled by the stranger (STR);

BOW: the book (B) is handled by the owner;

BSTR: the book is handled by the stranger;



POW: the puppet (P) is handled by the owner;

PSTR: the puppet is handled by the stranger.

These 3 episodes were separated by 1-min intervals identical to the familiarization phase: these intervals aimed at avoiding, or at least reducing, any potential carryover effects from the previous. The order in which the objects were presented during the test and who manipulated them first (i.e. owner or stranger) were partially counterbalanced to avoid order effects on dogs' behavioral responses. Thus, each dog was exposed to a given sequence of phases and the following six combinations were selected:

- 1. BSTR-BOW, FDSTR-FDOW, PSTR-POW;
- 2. BOW-BSTR, FDOW-FDSTR, POW-PSTR;
- 3. FDSTR-FDOW, PSTR-POW, BSTR-BOW;
- 4. FDOW-FDSTR, POW-PSTR, BOW-BSTR;
- 5. PSTR-POW, BTSR-BOW, FDSTR-FDOW;
- 6. POW-PSTR, BOW-BSTR, FDOW-FDSTR.

At the beginning of each episode, the stranger entered the room holding one of the objects. In combinations 1, 3 and 5, upon entering, she sat on the stranger's chair, manipulating the object that was kept on her lap, and engaged in an affectionate conversation about the object with the owner. In sequences 2, 4, and 6 the stranger, upon entering the room, gave the object to the owner, who manipulated it first keeping it on his/her lap. After 1 minute the object was transferred to the other person, who held it on his/her lap and handled it appropriately, while still conversing The dog was ignored by both the owner and the stranger for the two minutes. At the end of each episode, the stranger left the room, taking the object with her.

Data collection and analyses

The test was video recorded and analysed using Solomon Coder beta® 15.01.13 (ELTE TTK, Hungary). Behaviors were recorded continuously in terms of duration and frequency of their occurrence according to the ethogram reported in Table 1. The ethogram was developed considering both the literature on human infants [20, 21] and dogs [37, 43] and was refined after a preliminary analysis of the videos to include other potentially interesting behaviors. The ethogram included behaviors directed towards the owner/stranger, the objects and environment and stress related behaviors. Inter-observer agreement was assessed by means of independent parallel coding of a random sample of 16 dogs out of 72 (8 dogs for each experiment; 22.22% of the total number of dogs). The agreement was assessed by Spearman correlation and it was good for all behaviors (Rho ranging from 0.7 to 0.979; Table 1).

Since there was a small variability in the length of each phase, due to owners' differences in readiness to follow the instructions, durations and frequencies were respectively transformed in percentages of the total time and of total occurrences and used as dependent variables in the statistical analysis.

For both studies, the effects of dog's sex and age, order of the objects presentation (FD-P-B vs. B-FD-P vs. P-B-FD) and order of handlers (OW first vs. STR first) on dogs' behaviors were evaluated by means of preliminary ANOVAs. No significant effects of any variable emerged.

Repeated measures GLM ANOVAs with object (3 levels: book, fake dog, puppet) and handler identity (2 levels: owner, stranger) as factors, and behaviors as dependent variables were carried out using the Greenhouse-Geisser correction when sphericity assumption was violated.



Table 1. Ethogram used in study 1 and 2 and interobserver agreement.

Category	Pattern	Description	Agreement				
^a Object Directed	Chew/bite	The dog chews or bites the object	Rho = 0.809				
Behaviors	Interaction with Object	The dog interacts with the object without ambiguity. This pattern includes behaviors such as touching (with the paws or the muzzle), pushing and sniffing.					
	Look at Object	The dog looks at the object.					
	Social Investigation	The dog clearly smells the ano-genital area, the muzzle, or the ears of the fake dog.	Rho = 0.832				
^a Person Directed	Attention to Owner	The dog is oriented with the head and the body towards the owner and can gaze at him/her face.	Rho = 0.858				
Behaviors	Attention to Stranger	The dog is oriented with the head and the body towards the stranger, and can gaze at her face.					
	Interaction with Owner	The dog actively interacts with the owner. This pattern includes behaviors such as being in contact, touching (with the paws or the muzzle), sniffing and jumping on.					
	Interaction with Stranger	The dog actively interacts with the stranger. This pattern includes behaviors such as being in contact, touching (with the paws or the muzzle), sniffing and jumping on.	Rho = 0.979				
^a Enviroment Directed Behaviors	Explore Room	The dog explores the room. This pattern includes walking around and careful visual/olfactory exploration.	Rho = 0.929				
	Orientation to Door	The dog is oriented with the head and the body towards the door, gazing at it.	Rho = 0.762				
^b Stress Related Behaviors	Stress Signals	Nose-lip licking, shaking, yawning, scratching, stretching, rolling, chewing, raising paw.	Rho = 0.874				
^b Vocal behavior	Vocalizations	This pattern includes whining and barking.	Rho = 0.937				

^a behaviors recorded as duration and transformed into percentage of the total time for purpose of statistical analysis

https://doi.org/10.1371/journal.pone.0194577.t001

Pairwise post-hoc tests with Bonferroni's correction were carried out for significant main and interaction effects. All the statistical analyses were carried out with IBM SPSS Statistics 24.

Results

Data and results of statistical analysis of Studies 1 and 2 are reported in Tables 2-5 and in Figs 2, 3 and 4. In Tables 2 and 4 mean \pm SD of each behavior (duration or frequency) are detailed and in Tables 3 and 5 statistical values of the GLM ANOVAs are reported.

Table 2. (Study 1). Mean percentage± SD of time spent for each behavior across the experimental phases.

Phase	Interaction with Object	Look at Object	Social investigation	Attention to Owner	Attention to Stranger	Interaction with Owner	Interaction with Stranger	Explore Room	Orientation to Door	Stress
BOW	.75±2.06	1.09±2.82	-	18.68±21.99	13.33±17.24	1.399±4.23	1.54±7.03	30.45 ±25.21	3.84±8.84	2.39 ±3.10
FDOW	6.83±7.95	14.80 ±13.95	5.96±7.22	33.06±20.60	5.84±9.53	1.63±3.13	1.39±3.86	10.14 ±17.9	2.10±6.07	2.44 ±2.85
POW	4.81±9.69	4.85±5.77	-	28.89±23.75	4.91±7.57	.75±3.37	1.12±2.71	26.43 ±25.57	3.72±11.59	2.58 ±2.59
BSTR	0.02±.08	.71±1.81	-	19.89±27.07	17.52±21.74	1.39±4.04	.91±3.26	30.55 ±28.42	4.75±12.79	2.11 ±2.15
FDSTR	1.92±3.76	12.88 ±17.77	3.49±5.51	9.52±16.31	31.13±24.02	4.05±15.50	1.53±2.77	16.29±16	1.44±3.47	2.28 ±3.09
PSTR	1.11±2.74	1.49±2.29	-	12.88±19.27	22.46±23.77	4.27±13.79	1.55±3.48	23.18 ±21.37	3.32±8.79	2.39 ±2.72

BOW: the book is handled by the owner; **FDOW**: the fake dog is handled by the owner; **POW**: the puppet is handled by the owner; **BSTR**: the book is handled by the stranger; **FDSTR**: the fake dog is handled by the stranger.

https://doi.org/10.1371/journal.pone.0194577.t002

^b behaviors recorded as frequency and transformed into percentage of total occurrences for purpose of statistical analysis



Table 3. (Study 1) statistical values of the GLM ANOVA for main and interaction effects.

Effects	Interaction with Object	Look at Object	Social Investigation	Attention to Owner	Attention to Stranger	Interaction with Owner	Interaction with Stranger	Explore Room	Orientation to Door	Stress
Object	$F_{[1.6,58.45]} = 9.27$	$F_{[1.13,39.6]} = 35.02$		F _[1.59,55.65] = .22	F _[1.49,52.21] = 1.56	$F_{[1.385,48.491]} = 1.22$	F _[1.557,54.501] = .05	F _[2,70] = 9.37	$F_{[1.59,55.79]} = 1.03$	F _[2,705] = .15
	p = .000	p = .000	-	p = .75	p = .22	p = .219	p = .911	p = .000	p = .36	p = .857
	$\eta_{par}^2 = .21$	$\eta_{par}^2 = .50$		$\eta_{par}^{2} = .006$	$\eta_{par}^2 = .04$	$\eta_{par}^2 = .034$	$\eta_{par}^{2} = .002$	$ \eta_{\text{par}}^2 = $.21	$\eta_{\text{par}}^2 = .03$	$\eta_{par}^{2} = .004$
Handler	$F_{[1,35]} = 15.03$	F _[1,35] = 1.78	$F_{[1,35]} = 2.93$	F _[1,35] = 19.78	F _[1,35] = 32.58	$F_{[1,35]} = 2.266$	$F_{[1,35]} = .003$	F _[1,35] = .016	$F_{[1,35]} = .003$	F[_{1,35]} = .35
	p = .000	p = .19	p = .096	p = .000	p = .000	p = .141	p = .960	p = .69	p = .96	p = .556
	$\eta_{\text{par}}^2 = .30$	$\eta_{par}^2 = .05$	$\eta_{par}^2 = .077$	$\eta_{par}^2 = .36$	$\eta_{par}^2 = .48$	$\eta_{par}^2 = .061$	$\eta_{par}^2 = .00$	$\eta_{par}^2 = .01$	$\eta_{par}^2 = .00$	$\eta_{\text{par}}^2 = .01$
Object*Handler	$F_{[1.74,63.9]} = 3.41$	F _[1.14, 40.15] = .63		F _[2,70] = 11.58	F _[2,70] = 10.36	F _[1.189,41.615] = 1.19	F _[1.505,52.673] = .49	F _[2,70] = 1.87	F _[1.53,53.44] = .31	F[_{2,63.01]} = .014
	p = .046	p = .54	-	p = .000	p = .000	p = .292	p = .561	p = .10	p = .74	p = .986
	$\eta_{\text{par}}^2 = .09$	$\eta_{par}^2 = .02$		$\eta_{par}^2 = .25$	$\eta_{par}^2 = .23$	$\eta_{par}^2 = .033$	$\eta_{par}^2 = .014$	$\eta_{par}^2 = .05$	$\eta_{par}^2 = .01$	$\eta_{\text{par}}^2 = .00$

https://doi.org/10.1371/journal.pone.0194577.t003

Object directed behaviors

Social investigation. In both studies, almost all subjects engaged in Social Investigation (ano-genital region and muzzle) of the fake dogs (Study 1, 30 dogs out of 36 distributed as follow: 12 both in FDOW and FDSTR, 11 only in FDOW and 7 only in FDSTR; Study 2, 28 dogs out of 36 distributed as follow: 16 both in FDOW and FDSTR, 8 only in FDOW and 4 only in FDSTR). Overall, the FD was investigated for about 9.5% of time in Study 1 and 13% of time in Study 2. Time spent investigating the fake dog handled by the owner or the stranger was not significantly different in Study 1 (FDOW vs. FDSTR p = 0.096), while in Study 2 dogs spent a higher amount of time investigating the fake dog in the FDOW than in the FDSTR phase (p = 0.04).

Chew/bite. Aggressive bite was never observed, neither in Study 1 nor in Study 2. Dogs sporadically chewed the fake dog and the puppet. In Study 1, 7 dogs out of 36 (19%) chewed the furry fake dog, 5 of them did it in the FDOW phase and 2 both in the FDOW and FDSTR

Table 4. (Study 2): Mean percentage± SD of time spent for each behavior across the experimental phases.

Phase	Interaction with Object	Look at Object	Social investigation	Attention to Owner	Attention to Stranger	Interaction with Owner	Interaction with Stranger	Explore room	Orientation to door	Stress
BOW	.60±1.89	1.18±1.76	-	5.74±8.53	5.15±6.02	2.44±7.04	4.68±8.84	36.70 ±25.08	6.31±14.32	1.11 ±1.56
FDOW	13.40±13.59	14.91 ±18.57	8.81±11.08	9.78±14.44	1.97±3.77	.91±1.79	2.26±4.26	21.33 ±21.91	3.60±8.72	1.61 ±1.99
POW	6.31±7.08	6.27±8.22	-	5.96±5.77	3.82±4.81	3.63±10.83	3.11±7.47	34.13 ±25.09	6.52±17.76	1.19 ±1.74
BSTR	2.13±6.91	.91±1.85	-	7.55±10.54	6.43±10.86	2.08±6.44	6.43±17.70	40.55 ±31.47	6.10±13.93	1.78 ±2.34
FDSTR	10.04±10.87	11.42 ±16.05	4.56±6.15	9.60±10.57	9.07±9.72	1.93±4.74	3.86±8.36	28.47 ±26.26	2.67±7.62	2.06 ±2.20
PSTR	3.42±5.57	2.68±4.41	-	5.87±7.56	7.42±13.47	3.58±11.53	3.15±9.31	34.79 ±29.24	4.93±8.87	1.5 ±2.16

BOW: the book is handled by the owner; **FDOW**: the fake dog is handled by the owner; **POW**: the puppet is handled by the owner; **BSTR**: the book is handled by the stranger; **FDSTR**: the fake dog is handled by the stranger; **PSTR**: the puppet is handled by the stranger

https://doi.org/10.1371/journal.pone.0194577.t004



Table 5. (Study 2): statistical values of the GLM ANOVA for main and interaction effects.

Effects	Interaction with Object	Look at Object	Social investigation	Attention to Owner	Attention to Stranger	Interaction with Owner	Interaction with Stranger	Explore Room	Orientation to Door	Stress
Object	$F_{[1.45,50.99]} = 24.59$	F _[1.17,41.15] = 17.39		$F_{[1.691,59.172]} = 2.26$	$F_{[2,70]} = .015$	$F_{[1.708,59.763]} = 2.22$	$F_{[1.274,44.588]} = 1.80$	F _[2,70] = 4.11	F _[1.45,50.74] = .99	F _[2,70] = .99
	p = .000	p = .000	-	p = .121	p = .985	p = .125	p = .186	p = .021	p = .36	p = .374
	$\eta_{par}^2 = .41$	$\eta_{par}^2 = .33$		$\eta_{par}^{2} = .061$	$\eta_{par}^2 = .00$	$\eta_{par}^2 = .06$	$\eta_{par}^2 = .049$	$\eta_{par}^{2} = .105$	$\eta_{par}^2 = .03$	$\eta_{par}^{2} = .028$
Handler	$F_{[1,35]} = 1.44$	F _[1,35] = 4.43	$F_{[1,35]} = 4.53$	$F_{[1,35]} = .18$	F _[1,35] = 10.49	$F_{[1,35]} = .041$	$F_{[1,35]} = .85$	F _[1,35] = 2.01	$F_{[1,35]} = .87$	$F_{[1,35]} = 3.85$
	p = .238	p = .046	p = .04	p = .671	p = .003	p = .840	p = .361	p = .166	p = .358	p = .058
	$\eta_{par}^2 = .04$	$\eta_{par}^2 = .11$	$\eta_{par}^{2} = .115$	$\eta_{par}^2 = .005$	$\eta_{par}^{2} = .231$	$\eta_{par}^2 = .001$	$\eta_{par}^2 = .024$	$\eta_{par}^{2} = .054$	$\eta_{par}^2 = .02$	$\eta_{par}^{2} = .099$
Object* Handler	$F_{[2,70]} = 2.76$	F _[1.25,43.85] = .91		$F_{[2,70]} = .42$	$F_{[2,70]} = 3.21$	F _[1.336,46.743] = .26	$F_{[2,70]} = .44$	F _[2,70] = .69	F _[1.53,53.44] = .13	F _[2,70] = .25
	p = .07	p = .409	-	p = .660	p = .046	p = .682	p = .643	p = .504	p = .87	p = .782
	$\eta_{par}^2 = .07$	$\eta_{par}^2 = .02$		$\eta_{par}^2 = .012$	$\eta_{par}^2 = .084$	$\eta_{par}^2 = .007$	$\eta_{par}^2 = .013$	$\eta_{par}^{2} = .019$	$\eta_{par}^2 = .004$	$\eta_{par}^{2} = .007$

https://doi.org/10.1371/journal.pone.0194577.t005

phases. Another dog chewed the puppet only in the POW phase. In Study 2, this behavior was exhibited toward the plastic fake dog and the puppet by only 5 dogs out of 36 (14%): 3 dogs chewed the fake dog and 2 chewed the puppet. No statistical analysis was carried out due to negligible frequency of this behavior.




Fig 2. Mean percentage \pm SE of time spent by dogs interacting with the object in Study 1. Legend: black line, the owner is manipulating the objects; grey line: the stranger is manipulating the objects. Post-hoc values: Fake dog -Owner vs. Book-Owner, p < 0.001; Puppet-Owner vs. Book-Owner, p = 0.035; Fake dog-Stranger vs. Book-Stranger, p = 0.014; Puppet-Stranger vs. Book-Stranger, p = 0.066.

https://doi.org/10.1371/journal.pone.0194577.g002






Fig 3. Mean percentage \pm SE of time spent by dogs paying attention to the owner in Study 1. Legend: black line, the owner is manipulating the objects; grey line: the stranger is manipulating the objects. Post-hoc values: Fake dog-Owner vs. Fake dog-Stranger, p<0.001; Puppet-Owner vs. Puppet-Stranger, p<0.001; Fake dog-Owner vs. Book-Owner, p = 0.017.

https://doi.org/10.1371/journal.pone.0194577.g003

Look at object. In both Study 1 and Study 2 the GLMs revealed a main effect of the object identity: dogs spent significantly more time looking at the fake dog compared to the other objects; the puppet was gazed for a longer time than the book (Study 1: fake dog vs. puppet p<0.001, fake dog vs. book p<0.001, puppet vs. book p=0.002; Study 2: fake dog vs. puppet p=0.006, fake dog vs. book p<0.001, puppet vs. book p=0.001).




Fig 4. Mean percentage \pm SE of time spent by dogs paying attention to the stranger in Study 1(a) and in Study 2(b). Legend: black line, the owner is manipulating the objects; grey line: the stranger is manipulating the objects. Post-hoc values: Fake dog-Stranger vs. Fake dog-Owner, Study 1: p<0.001; Study 2: p<0.001; Puppet-Stranger vs. Puppet-Owner, Study 1: p<0.001; Fake dog-Stranger vs. Book-Stranger, Study 1: p=0.038.

https://doi.org/10.1371/journal.pone.0194577.g004



Interaction with object. In Study 1 (furry fake dog) a significant interaction was found between object and handler identity: overall, dogs interacted significantly longer with the objects when they were manipulated by the owner compared to the stranger (book p = 0.04; fake dog p = 0.001; puppet p = 0.03). As can be seen in Fig 2, dogs interacted longer with the fake dog and with the puppet than with the book, regardless of the identity of the handler (post-hoc values are reported in figure caption).

In Study 2 (plastic fake dog) only the identity of the manipulated object had a significant influence on dogs' behavior: dogs spent significantly more time interacting with the fake dog compared to the puppet (fake dog vs. puppet, p = 0.001) and to the book (fake dog vs. book p < 0.001), and they interacted with the puppet significantly longer than with the book (puppet vs. book p = 0.002).

In sum, in both studies the majority of dogs showed social investigation of the fake dogs, but no aggression towards them, and looked at the fake dog significantly longer than the other two objects. In Study 1, dogs interacted with the furry dog and the puppet to a comparable extent, whereas in Study 2 they interacted with the plastic fake dog significantly longer than with the puppet. Only in Study 1 all objects were more attractive when on the owner's lap.

Person directed behaviors

Attention to owner. In Study 1, the GLM highlighted a significant interaction Object x Handler (Fig 3): in the book condition dogs paid attention to the owner whether or not the book was in his/her hands, whereas in the fake dog and puppet conditions dogs' attention to the owner was driven by the presence of the object. Furthermore, dogs paid attention to their owner significantly longer when he/she was handling the fake dog compared to the book.

In Study 2, even though dogs addressed their attention to the owner mainly in the FDOW and FDSTR phases, the GLM did not highlight any significant effects of object's and handler's identity, nor of the interaction Object x Handler.

Attention to stranger. The GLMs revealed a significant interaction Object x Handler in both studies (Fig 4). In Study 1, both the fake dog and the puppet directed the dogs' attention toward the stranger, and dogs paid attention to the stranger for a significantly higher percentage of time when he/she was handling the furry fake dog compared to the book. No other significant differences emerged. In Study 2, only the fake dog directed the dogs' attention toward the stranger.

Interaction with owner and stranger. The GLM revealed that physical interaction with the owner and the stranger was not differently affected by the objects and handlers' identity.

In sum, in both studies dogs oriented their attention/interaction to the owner and to the stranger manipulating the objects to a similar extent. The fake dogs were the more salient stimuli in eliciting these behaviors.

Vocalizations and stress related behaviors. Very few dogs tried to attract owner's attention using vocalizations (Study 1: 8 dogs out of 36, 22%; Study 2: 5 dogs out of 36, 14%), doing it uniformly across the testing phases. No statistical analysis was carried out due to negligible frequency of this behavior.

The frequency of stress related behaviors during testing was low and no statistical differences across testing phases emerged.

Therefore, these behaviors did not provide any useful insight to interpret dogs' behavioral reactions in terms of jealousy.

Environment directed behaviors

Explore room and orienting to door. In both studies the GLMs showed a main effect of the object's identity on the dogs' exploratory behavior. Exploration significantly decreased



when either the owner or the stranger held the fake dog compared to the other two objects in Study 1, and only compared to the book in Study 2 (Study 1: fake dog vs. book p = 0.002, fake dog vs. puppet p = 0.007, book vs. puppet p = 0.47; Study 2: fake dog vs. book p = 0.016; fake dog vs. puppet p = 0.16; book vs. puppet, p = 1).

As for the time spent being oriented to the door, the GLM did not reveal any difference across the six phases: the dogs spent a similar amount of time oriented to the door whichever objects the owner and the stranger were holding and whoever person was the handler.

Discussion

The existence of jealousy in dogs, as claimed by Harris and Prouvost [37], is appealing, but experimental evidence is preliminary. The current study aimed at further understanding whether dogs react with a set of behavioral patterns that could indicate a 'primordial' form of jealousy when they lose their owner's attention in favour of a potential rival (a fake dog). For this purpose, we carried out two studies using two types of fake dogs (FD), differing in appearance and texture (Study 1: furry FD; Study 2: plastic FD), a novel object (a puppet) and a book. Moreover, as jealousy should be expressed towards a valuable individual, we controlled for the identity of the handler: thus, the objects were manipulated by the owner and by a stranger who had no social bond with the dog.

Although there were some differences in the behavioral pattern showed by dogs tested in Study 1 and Study 2 (e.g. more attention to people in Study 1 than in Study 2; less interaction with the fake dog in Study 1 than in Study 2; see Tables 2 and 4), results of both studies do not provide convincing proof that the behavioral responses of our dogs were triggered by jealousy, since we did not find clear evidence that the fake dogs were perceived as real social rivals. In addition, dogs did not express, or expressed to a limited extent, many of the behaviors that are considered distinctive features of jealousy (aggression, vocalizations, stress related behaviors, attention/interaction with object and owner). Finally, our dogs did not show, as hypothesized, a differentiated response towards the owner and the stranger.

Our first research question was whether dogs would consider the fake dogs as a real conspecific, as suggested by Harris and Prouvost [37], on the basis of the number of dogs that in their study exhibited social investigation of the fake dog (86%) and aggressive behavior toward it (25%). Even though in our studies 86% and 78% of dogs exhibited social investigation respectively toward the furry and the plastic fake dog, this investigation was a momentary action and not a careful inspection (Furry FD mean duration: FDSTR: 1.99 s; FDOW: 3.47 s; Plastic FD: mean FDSTR: 2.48 s; mean FDOW: 4.63). Although social investigation of the plastic FD occurred significantly longer when it was on owner's lap than on stranger's lap, in our opinion, this result can hardly be considered a convincing proof that the FDs were effective in deceiving adult and experienced dogs. It is worth considering that the FDs did not smell like a real dog and that during intraspecific interactions between dogs social investigation allows for individual/sexual recognition through odour cues [51–52]. Olfaction plays such a pivotal role in dogs' social communication that they can discriminate between a familiar and an unfamiliar person using odor cues [53-54] and, remarkably, they show an asymmetric use of nostril when processing odors—emitted by conspecifics/humans—that differ in terms of emotional valence [55–56]. Moreover, dogs form cross-modal representations of humans and conspecifics integrating auditory and visual stimuli [25, 57]. Likely, a handful of seconds was sufficient for our adult dogs to perceive the FDs as faux and this might account for the complete lack of aggressive behavior. Some of the dogs chewed the FDs, but, differently from what reported by Harris and Prouvost [37], snapping and biting were never observed. It is worth noting that their fake dog was a toy for human infants that barked/whined and wagged its tail, while our FDs were



silent and motionless: thus, the aggressive responses they observed could have been motivated by fear of a novel/odd object or by predatory drive rather than by jealousy [58]. It is also possible that the dogs tested at their home by Harris and Prouvost [37] showed aggressive behaviour driven by a protective/territorial motivation [59]. In our opinion a neutral testing environment is more suitable to disentangle aggressive behaviors triggered by jealousy from territorial/protective aggression. However, the choice of the testing location is always critical and presents pros and cons that could be interesting topics for further researches.

If it is reasonable to assume that the FDs are not perceived as real dogs, there are no particular reasons to be jealous: thus, which could be the dogs' underlying motivation to show interest towards them? As in Harris and Prouvost's study [37], our dogs preferentially looked at the furry FD than at the soft puppet, but they did not seem to discriminate between them when interacting (sniffing and touching); this suggests that these objects were both perceived as toys, as they were of similar size and of soft texture, and that the furry FD resembled toys most pet dogs are familiar with. Conversely, in Study 2, the FD and the puppet had a different appearance and a different texture and it is possible that dogs prolonged their interaction with the plastic FD to make sense of it.

We also expected that if dogs were jealous they would have shown signs of distress and protest (vocalizations) when attention and care were directed to the FDs, but not to the other objects [37, 60]. Results show that vocalizations were infrequent and distributed across the test phases without any link to a specific object/person. To a certain extent these results are similar to those of Harris and Prouvost, who found that: 1. vocalization were relatively infrequent; 2. there was no difference across conditions in barking; 3. dogs whined more in the jealousy condition than in the book condition, but not the jack-o-lantern condition [37]. Overall, it seems to us that dogs did not bark/whine to regain their owner's attention and to re-establish the relationship with him/her, but to express frustration for being ignored and for not obtaining full access to a potentially interesting object. Further confirmation derives from the frequency of stress signals, which was low and uniformly distributed across testing phases, indicating an arousal state due to the testing situation. In sum, these findings suggest that the use of the FD as a potential rival in a jealousy evoking paradigm is critical, as the stimulus is unnatural. In the context of temperament assessment, Barnard and colleagues [61] questioned the validity of model devices (a fake dog and a child-like doll): they concluded that adult dogs may perceive these models as social stimuli only at distance and for a few seconds at the beginning of the interaction, since the model does not smell, move or interact like a real conspecific or a real child.

In line with Harris and Provoust's results [37], the book *per se* was largely overlooked, as dogs spent half of the testing time exploring the room when people handled this object. Although to a lesser extent than in the FD conditions, dogs attended to the owner (and the stranger) also in the book condition. This could have at least two different explanations: 1. being sensitive to human inattentiveness [62]; 2. being uncertain about the situation and looking for human cues [30, 63].

Considering that the adaptive function of jealousy is to protect a valuable relationship from an intruder, it should be expected that an unfamiliar person devoting attention and care to a 'conspecific' would not elicit a jealousy response. Thus, to extend Harris and Prouvost's results [37], we introduced a relevant variation in the experimental paradigm, adopting the procedure devised by Hart and colleagues with human infants [19]. A second person, totally unfamiliar to the dogs (but manipulating the same objects) served as control to exclude that dogs' reactions were due to a mere interest towards the more salient/desired object or to the frustration of being deprived of an interesting toy. Results of both studies showed that dogs' attention (being oriented and gazing) to the owner and the stranger was driven by the presence of the



FDs and the puppet in their lap. On the other hand, their attention was not preferentially directed to the attachment figure (the owner), which in the human literature is considered a fundamental requisite to attribute jealousy to an individual. Differently from our results, and in line with an interpretation in terms of jealousy, Hart and colleagues [19] found that children manifested distress and disapproval of the lack of attention and that these reactions were linked with the identity of the person holding the infant-like doll.

Of course our negative outcome does not necessarily mean that dogs do not possess a rudimentary form of jealousy. Indeed, as highlighted above, we believe that our FDs were inappropriate stimuli to trigger jealousy and rivalry in dogs. It is important to underline that comparing adult dogs and human infants, albeit interesting, requires caution: adult dogs and human infants differ in terms of perceptual abilities, as they rely on different sensory modalities in making sense of the world, in their mobility and possibility to interact with the environment. Physical interaction and contact seeking with the attachment figure are pervasive components of the jealousy response in children [14,15] and also occur in situation of mild distress such as the Strange Situation Test [64]. Nevertheless, these behaviors, which are clearly observable in dogs during the paradigm of separation from the attachment figure [43], were scarcely expressed in our testing conditions, providing further support that the dogs did not perceive the presence of the faux rival as a threat to the relationship with their owner.

In sum, current data does not exclude that dogs may have a rudimentary form of jealousy, but shows that available evidence is still inconclusive and that different experimental paradigms and additional rigorous research is required. For example, replications using a more naturalistic procedure and a real social interloper (either conspecific or human), rather than a false social stimulus, could be more suitable in ascertaining whether dogs experience and show a rudimentary form of jealousy, as strongly believed and carefully described by dog owners [38–39]. Our results also suggest that, as for human infants, the investigation of emotions in nonhuman animals is tricky and hampered by limitations, as behavioural expressions may provide ambiguous information about internal emotional states [12, 22].

Supporting information

S1 File. Breed of dogs participating in study1. (DOCX)

S2 File. Breed of dogs participating in study 2. (DOCX)

Acknowledgments

We are grateful to all the dog owners who kindly participated with their dogs in our studies. We thank Chiara Losacco and Irene Zantedeschi for their precious help in data collection and Sofia Fusar Poli for language revision. Velia Nicotra was supported by a doctoral grant from the Università degli Studi di Parma.

Author Contributions

Conceptualization: Emanuela Prato-Previde, Paola Valsecchi.

Data curation: Velia Nicotra, Annalisa Pelosi. Formal analysis: Annalisa Pelosi, Paola Valsecchi.

Investigation: Emanuela Prato-Previde, Velia Nicotra, Paola Valsecchi.



Methodology: Emanuela Prato-Previde, Velia Nicotra.

Project administration: Emanuela Prato-Previde, Velia Nicotra, Paola Valsecchi.

Software: Velia Nicotra.

Supervision: Paola Valsecchi.

Writing – original draft: Emanuela Prato-Previde, Velia Nicotra, Annalisa Pelosi, Paola Valsecchi.

References

- 1. Ekman P. An argument for basic emotions. Cogn Emot. 1992; 6: 169–200.
- Izard CE. Basic emotions, relations among emotions, and emotion-cognition relations. Psychol Rev. 1992; 99: 561–565. PMID: 1502277
- 3. Buck R. The biological affects: a typology. Psychol Rev. 1999; 106(2): 301–336. PMID: 10378015
- 4. Panksepp J. Affective consciousness in animals: perspectives on dimensional and primary process emotion approaches. Proc R Soc B Biol Sci. 2010; 277: 2905–2907.
- Panksepp J. The basic emotional circuits of mammalian brains: do animals have affective lives? Neurosci Biobehav Rev. 2011; 35: 1791–804. https://doi.org/10.1016/j.neubiorev.2011.08.003 PMID: 21872619
- Panksepp J, Watt D. What is basic about basic emotions? Lasting lessons from affective neuroscience. Emot Rev. 2011; 3: 387–396.
- 7. Lewis M. The emergence of human emotions. In: Lewis M, Hayland-Jones JM, Barrett LF, editors. Handbook of emotions. New York: Guilford Press; 2008. pp. 304–319.
- 8. Draghi-Lorenz R, Reddy V, Costall A. Rethinking the development of "nonbasic" emotions: A critical review of existing theories. Dev Rev. 2001; 21: 263–304.
- 9. Harris CR. Evolution of jealousy. Am Sci. 2004; 92: 62–71.
- Panksepp J. Affective neuroscience: the foundations of human and animal emotions. New York: Oxford University Press; 1998.
- de Waal FBM. What is an animal emotion? Ann NY Acad Sci. 2011; 1224: 191–206. https://doi.org/10. 1111/j.1749-6632.2010.05912.x PMID: 21486301
- 12. Kujala MV. Canine emotions as seen through human social cognition. Anim Sentience. 2017; 14(1).
- 13. Panksepp J. The evolutionary sources of jealousy: cross-species approaches to fundamental issues. In: Hart SL, Legerstee M, editors. Handbook of jealousy: theory, research, and multidisciplinary approaches. Malden: Wiley-Blackwell; 2010; pp.101–120.
- Dillon L. Functional aspects of jealousy across the lifespan. Hum Ethol Bull. 2013; 28: 13–26.
- 15. Hart SL. The ontogenesis of jealousy in the first year of life: A theory of jealousy as a biologically-based dimension of temperament. In Hart SL, Legerstee M, editors. Handbook of jealousy: theory, research, and multidisciplinary approaches. Malden: Wiley-Blackwell; 2010. p. 57–82.
- **16.** Massar K, Buunk A, Dechesne M. Jealousy in the blink of an eye: Jealous reactions following subliminal exposure to rival characteristics. Eur J Soc Psychol. 2009; 39:768–79.
- 17. Hobson RP. Is jealousy a complex emotion? In: Hart SL, Legerstee M, editors. Handbook of jealousy: theory, research, and multidisciplinary approaches. Oxford: Wiley-Blackwell; 2010. pp. 293–311.
- 18. Hart SL. Jealousy in infants. Laboratory research on differential treatment. New York: Springer; 2015.
- Hart SL, Field T, del Valle C, Letourneau M. Infants protest their mother's attending to an infant-size doll. Soc Dev. 1998; 7: 54–61.
- 20. Hart SL, Carrington H. Jealousy in 6 month-old infants. Infancy. 2002; 3: 395–402.
- 21. Mize KD, Pineda M, Blau AK, Jones NA. Infant physiological and behavioral responses to a jealousy provoking condition. Infancy. 2014; 19: 338–348.
- Hart SL. Jealousy protest: ontogeny in accord with the 9-month period of human gestation. Evol Psychol. 2016: 1–9.
- 23. Forbes S. Sibling rivalry among the birds and the bees. In: Hart SL, Legerstee M, editors. Handbook of jealousy: theory, research, and multidisciplinary approaches. Oxford: Wiley-Blackwell; 2010. pp 121–143.



- Hare B, Tomasello M. Human-like social skills in dogs? Trends Cogn Sci. 2005; 9: 439–444. https://doi. org/10.1016/j.tics.2005.07.003 PMID: 16061417
- Adachi I, Kuwahata H, Fujita K. Dogs recall their owner's face upon hearing the owner's voice. Anim Cogn. 2007; 10(1): 17–21. https://doi.org/10.1007/s10071-006-0025-8 PMID: 16802145
- Lakatos G, Soproni K, Dóka A, Miklósi A. A comparative approach to dogs' (Canis familiaris) and human infants' comprehension forms of pointing gestures. Anim Cogn. 2009, 12(4): 621–631. https:// doi.org/10.1007/s10071-009-0221-4 PMID: 19343382
- Topál J, Gergely G, Erdohegyi A, Csibra G, Miklósi A. Differential sensitivity to human communication in dogs, wolves, and human infants. Science. 2009; 325: 1269–1272. https://doi.org/10.1126/science. 1176960 PMID: 19729660
- Marshall-Pescini S, Colombo E, Passalacqua C, Merola I, Prato-Previde E. Gaze alternation in dogs and toddlers in an unsolvable task: evidence of an audience effect. Anim Cogn. 2013; 16: 933–943. https://doi.org/10.1007/s10071-013-0627-x PMID: 23543361
- Marshall-Pescini S, Ceretta M, Prato-Previde E. Do domestic dogs understand human actions as goaldirected? PLoS One. 2014; 9(9): e106530. https://doi.org/10.1371/journal.pone.0106530 PMID: 25229452
- Merola I, Prato-Previde E, Marshall-Pescini S. Social referencing in dog-owner dyads? Anim Cogn 2012a; 15: 175–185. https://doi.org/10.1007/s10071-011-0443-0 PMID: 21874515
- Chijiiwa H, Kuroshima H, Hori Y, Anderson JR, Fujita K. Dogs avoid people who behave negatively to their owner: third-party affective evaluation. Animal Behaviour. 2015; 106: 123–127.
- Custance D, Mayer J. Empathetic-like responding by domestic dogs (Canis familiaris) to distress in humans: an exploratory study. Anim Cogn. 2012; 15: 851–859. https://doi.org/10.1007/s10071-012-0510-1 PMID: 22644113
- Quervel-Chaumette M, Faerber V, Faragó T, Marshall-Pescini S, Range F. Investigating empathy-like responding to conspecifics' distress in pet dogs. PLoS One. 2016; 11(4): e0152920. https://doi.org/10. 1371/journal.pone.0152920 PMID: 27124485
- Range F, Horn L, Viranyi Z, Huber L. The absence of reward induces inequity aversion in dogs. Proceedings of the National Academy of Sciences. 2009 Jan 6; 106(1): 340–5.
- Brucks D, Essler JL, Marshall-Pescini S, Range F. Inequity aversion negatively affects tolerance and contact-seeking behaviours towards partner and experimenter. PLoS One. 2016; 11: e0153799. https://doi.org/10.1371/journal.pone.0153799 PMID: 27081852
- Essler J, Marshall-Pescini S, Range F. Domestication does not explain the presence of inequity aversion in dogs. Curr Biol 2017; 27: 1–5. https://doi.org/10.1016/j.cub.2016.10.044
- Harris CR, Prouvost C. Jealousy in dogs. PLoS One. 2014; 9: e94597. https://doi.org/10.1371/journal.pone.0094597 PMID: 25054800
- Morris PH, Doe C, Godsell E. Secondary emotions in non-primate species? Behavioural reports and subjective claims by animal owners. Cogn Emot. 2008; 2: 3–20.
- Morris PH, Knight S, Lesley S. Belief in animal mind: does familiarity with animals influence beliefs about animal emotions? Society & Animals. 2012; 20: 211–224. https://doi.org/10.1163/15685306-12341234
- Martens P, Enders-Slegers MJ, Walker JK. The emotional lives of companion animals: attachment and subjective claims by owners of cats and dogs. Anthrozoös. 2016; 29(1): 73–88. https://doi.org/10.1080/ 08927936.2015.1075299
- 41. Bonanni R, Cafazzo S. The social organization of a population of free-ranging dogs in a suburban area of Rome: a reassessment of the effects of domestication on dogs' behaviour. In: Kaminski J, Marshall-Pescini S, editors. The social dog. Behavior and cognition. Amsterdam: Elsevier; 2014. pp 65–104.
- **42.** Trisko RK, Sandel AA, Smuts B. Affiliation, dominance and friendship among companion dogs. Behaviour. 2016: 153: 693–725.
- **43.** Prato Previde E, Valsecchi P. The immaterial cord: the dog-human attachment bond. In: Kaminski J, Marshall-Pescini S, editors. The social dog. Behavior and cognition. Boston: Academic Press; 2014. pp. 165–189.
- 44. Albuquerque N, Guo K, Wilkinson A, Savalli C, Otta E, Millas D. Dogs recognize dog and human emotions. Biol Lett. 2016; 12: 20150883. https://doi.org/10.1098/rsbl.2015.0883 PMID: 26763220
- Müller CC, Schmitt K, Barber ALA, Huber L. Dogs can discriminate emotional expressions of human faces. Curr Biol. 2015; 25: 601–605. https://doi.org/10.1016/j.cub.2014.12.055 PMID: 25683806
- 46. Merola I, Prato-Previde E, Lazzaroni M, Marshall-Pescini S. Dogs' comprehension of referential emotional expressions: familiar people and familiar emotions are easier. Anim Cogn. 2013; 17:373–85. https://doi.org/10.1007/s10071-013-0668-1 PMID: 23955027



- 47. Gácsi M, Miklósi A, Varga O, Topál J, Csányi V. Are readers of our face readers of our minds? Dogs (Canis familiaris) show situation-dependent recognition of human's attention. Anim Cogn. 2004; 7: 144–153. https://doi.org/10.1007/s10071-003-0205-8 PMID: 14669075
- Schwab C, Huber L. Obey or not obey? Dogs (*Canis familiaris*) behave differently in response to attentional states of their owners. J Comp Psychol. 2006; 120: 169–75. https://doi.org/10.1037/0735-7036. 120.3.169 PMID: 16893253
- **49.** Mongillo P, Bono G, Regolin L, Marinelli L. Selective attention to humans in companion dogs, *Canis familiaris*. Anim Behav. 2010; 80: 1057–1063.
- Volling BL, McElwain NL, Miller AL. Emotion regulation in context: the jealousy complex between young siblings and its relations with child and family characteristics. Child Dev 2002; 73: 581–600 PMID: 11949910
- **51.** Mech LD. The wolf: the ecology and behavior of an endangered species. 1st ed. New York, NY: Natural History Press; 1970.
- 52. Fox M. Behavior of Wolves, Dogs, and Related Canids. Malabar, FL: Krieger Publishing Co; 1972.
- Millot JL, Filiatre JC, Eckerlin A, Gagnon AC, Montagner H. Olfactory and visual cues in the interaction systems between dogs and children. Appl Anim Behav Sci. 1987; 19: 189–195.
- **54.** Filiatre JC, Millot JL, Eckerlin A. Behavioural variability of olfactory exploration of the pet dog in relation to human adults. Appl Anim Behav Sci. 1991; 30(3–4): 341–350.
- 55. Siniscalchi M, Sasso R, Pepe AM, Dimatteo S, Vallortigara G, Quaranta A. Sniffing with the right nostril: lateralization of response to odour stimuli by dogs. Anim Behav. 2011; 8: 399–404.
- Siniscalchi M, d'Ingeo S., Quaranta A. The dog nose 'KNOWS' fear: Asymmetric nostril use during sniffing at canine and human emotional stimuli. Behav Brain Res. 2016; 304: 34–41. https://doi.org/10. 1016/j.bbr.2016.02.011 PMID: 26876141
- Taylor AM, Reby D, McComb K. Cross modal perception of body size in domestic dogs (Canis familiaris). PLoS One 2011; 6(2): e17069. https://doi.org/10.1371/journal.pone.0017069 PMID: 21359228
- Reisner I. Differential diagnosis and management of human directed aggression in dogs. Vet Clin North Am Small Anim Pract. 2003; 33: 303–320. PMID: 12701514
- Blackshaw JK. An overview of types of aggressive behaviour in dogs and methods of treatment. Appl Anim Behav Sci. 1991; 30: 351–361.
- **60.** Szabo N, Dubas JS, van Aken MAG. Jealousy in firstborn toddlers within the context of the primary family triad. Soc Devel. 2014; 23: 325–39.
- **61.** Barnard S, Siracusa C, Reisner I, Valsecchi P, Serpell J. Validity of model devices used to assess canine temperament in behavioral tests. Appl Anim Behav Sci. 2012; 38: 79–87.
- Call J, Bräuer J, Kaminski J, Tomasello M. Domestic dogs (Canis familiaris) are sensitive to the attentional state of humans. J Comp Psychol. 2003; 117(3): 257–63. https://doi.org/10.1037/0735-7036.
 117.3.257 PMID: 14498801
- Merola I, Prato-Previde E, Marshall-Pescini S. Dogs' social referencing towards owners and strangers. PLoS One. 2012b; 7(10): e47653. https://doi.org/10.1371/journal.pone.0047653 PMID: 23071828
- Ainsworth MDF, Blehar MC, Waters E, Wall S. Patterns of attachment: a psychological study of the strange situation. 1978. Hillsdale, NJ: Erlbaum