

The Effects of Smoking on Metabolic Syndrome and Its Components Using Causal Methods in the Iranian Population

Abstract

Background: The aim of this study was to estimate the effect of smoking on metabolic syndrome (MS) and its components applying inverse probability-of-treatment weighting (IPTW) and propensity score (PS) matching. **Methods:** Using data from Tehran Lipid and Glucose Study, 4857 participants aged over 20 years with information on smoking and confounders in the third phase (2005–2008) were included, and the MS was assessed in the fifth phase (2011–2014). IPTW and PS matching were used to adjust for confounders. **Results:** Based on average treatment effect (ATE) estimates, smoking decreased the risk of hypertension (RR: 0.62; 95% CI: 0.43, 0.88), but increased the risk of low HDL cholesterol (1.20; 0.98, 1.48). Similarly, the average treatment effect in the treated (ATT) estimates using IPTW and PS matching suggested that smoking decreased the risk of hypertension (0.63; 0.52, 0.76, and 0.68; 0.54, 0.85), and increased the risk of low HDL cholesterol (1.24; 1.07, 1.43, and 1.28; 1.06, 1.54), respectively. **Conclusions:** Smoking seems to increase the risk of low HDL cholesterol but decreases the risk of hypertension.

Keywords: Metabolic syndrome, propensity score, smoking.

Introduction

Metabolic syndrome (MS) is a set of conditions including central obesity, high blood pressure (BP), increased fasting blood sugar (FBS), increased triglycerides, and decreased high-density lipoprotein cholesterol (HDL-C).^[1] The prevalence of MS is globally increasing and, depending on some background factors, ranges from less than 10% to 84%.^[2] MS patients are at risk of a two-fold increase in type 2 diabetes and a five-fold increase in cardiovascular disease over the next 5 and 10 years, respectively.^[2]

Modifications of life styles such as smoking are useful for chronic diseases prevention.^[3] Smoking is associated with many noncommunicable diseases and contributes to mortality and disability-adjusted life year (DALYs).^[4,5] According to the world health organization, there will be 1.5 billion smokers worldwide by 2050.^[5] The relationship between smoking and MS,^[6,7] BP,^[8,9] abdominal obesity,^[10] FBS,^[11] and triglycerides^[9,12] has been already studied using conventional regression for confounding adjustment. An alternative is propensity score (PS)

methods. PS, defined as the probability of exposure given the set of confounders,^[13] can be used in different procedures for confounding adjustment: matching, inverse probability-of-treatment weighting (IPTW), stratification, and regression adjustment.^[14-17]

PS methods are preferred to outcome regression for inferring causality because i) it is easier to determine whether the exposure models are correctly specified in terms of yielding covariate-balancing PSs using standardized differences, ii) these methods effectively emulate a randomized experiment without any reference to the outcome, and iii) the overlap in the distribution of confounders can be explicitly assessed between two exposure groups: the small number of matched subjects or huge inverse probability-of-treatment weights indicate low overlap.^[18] The aim of this study is to estimate the effect of smoking on MS and its components using IPTW and PS matching.

Methods

Using the third phase data (2005-2008) of Tehran Lipid and Glucose cohort Study (TLGS) as the baseline, 4857 participants aged over 20 years without MS and with information on smoking

Farzad Khodamoradi¹,
Maryam Nazemipour²,
Nasrin Mansournia³,
Kamran Yazdani¹,
Davood Khalili^{4,5},
Mohammad Ali Mansournia¹

¹Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, ²Osteoporosis Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, ³Department of Endocrinology, AJA University of Medical Sciences, Tehran, ⁴Prevention of Metabolic Disorders Research Center, ⁵Department of Biostatistics and Epidemiology, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Address for correspondence: Dr. Mohammad Ali Mansournia, Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, PO Box: 14155-6446, Tehran, Iran. E-mail: mansournia_ma@yahoo.com; mansournia_ma@sina.tums.ac.ir
Dr. Davood Khalili
Prevention of Metabolic Disorders Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, P O Box: 19395-4763, Tehran, Iran.
E-mail: dkhalili@endocrine.ir

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.ir

DOI: 10.4103/ijpvm.ijpvm_45_21

Quick Response Code:



How to cite this article: Khodamoradi F, Nazemipour M, Mansournia N, Yazdani K, Khalili D, Mansournia MA. The effects of smoking on metabolic syndrome and its components using causal methods in the Iranian population. *Int J Prev Med* 2021;12:118.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

and confounders were selected. MS was assessed in the fifth phase (2011–2014). This study was approved by the Research Council of Research Institute for Endocrine Sciences of Shahid Beheshti University of Medical Sciences, and a consent form was obtained from all participants.

The exposure was cigarette smoking status assessed by the question: “is person smoking daily?”. The outcome was MS, defined as having at least three out of the following variables: abdominal obesity (waist circumference ≥95 cm), low HDL-C (<40 in men or <50 mg/dL in women), hypertriglyceridemia (TG ≥150 mg/dL), hypertension (systolic BP ≥130 or diastolic BP ≥85 mmHg) and impaired blood glucose (FBS ≥100 mg/dL).^[19-21]

The confounders were identified using a causal diagram [Figure 1] for the study population.^[22-27] The minimally sufficient set for confounding adjustment, derived based on Pearl’s back-door criterion,^[28] included gender, age, physical activity, marital status, education, and job, measured by a questionnaire, as well as the unmeasured variables income and alcohol. Fractional polynomials were used to identify any nonlinear association between age and smoking in the PS model.^[29]

Statistical methods

IPTW was used to adjust for confounders. We first estimated PS through logistic regression, with smoking as the response variable and confounders as predictors. Then the average treatment effect (ATE) was estimated using weighted risk ratio (RR) between smoking and MS with weights equal to 1/PS for the smokers and 1/(1-PS) for the non-smokers. Moreover, the average treatment effect in the treated (ATT) was estimated with weights equal to 1 for the smokers and PS/(1-PS) for the non-smokers.^[30,31] IPTW produces a pseudo-population in which confounders do not predict the exposure, and the effect in the pseudo-population is the same as that in the population.

Confounders were also adjusted for using PS matching. A PS-matched dataset was created by matching, without replacement, one unexposed person to one exposed based




Figure 1: Causal diagram for the effect of smoking on MS

on the nearest value of PS (±0.05).^[32] Then the ATT effect was estimated using the RR between smoking and MS in the matched sample.^[18]

The 95% confidence intervals (CIs) for the IPTW estimates were derived using robust standard errors.^[33] The 95% CIs for the PS matching was obtained based on nonparametric bootstrapping by 1000 repetitions with 2.5th and 97.5th percentiles as 95% confidence limits.^[34]

The correct specification of the PS model was assessed based on the balance of measured confounders between exposure groups in the matched sample for PS matching, and in the weighted sample for the IPTW. The standardized difference was used to compare the mean and proportion of continuous and binary confounders between the exposed and unexposed, respectively. The standardized difference for continuous confounders is

$$d = \frac{(\bar{x}_{exposed} - \bar{x}_{unexposed})}{\sqrt{\frac{s_{exposed}^2 + s_{unexposed}^2}{2}}}$$

Where $\bar{x}_{exposed}$ and $\bar{x}_{unexposed}$ are the mean estimates, and $s_{exposed}^2$ and $s_{unexposed}^2$ are variance estimates in exposed and unexposed, respectively.

The standardized difference for binary confounders is

$$d = \frac{\hat{p}_{exposed} - \hat{p}_{unexposed}}{\sqrt{\frac{\hat{p}_{exposed}(1 - \hat{p}_{exposed}) + \hat{p}_{unexposed}(1 - \hat{p}_{unexposed})}{2}}}$$

Table 1: Baseline characteristics of the study participants

Characteristic	Smokers, n (%)	Non-smokers, n (%)
Gender (female)	74 (14.5)	2885 (66.4)
Age, years, mean (SD)	40.89 (12.67)	38.89 (13.56)
Marital status		
single	102 (19.9)	918 (21.1)
married	391 (76.4)	3233 (74.4)
divorced	13 (2.5)	56 (1.3)
widowed	6 (1.2)	138 (3.2)
Educational certificate		
elementary	57 (11.1)	592 (13.6)
secondary school	98 (19.1)	641 (14.8)
high school	264 (51.6)	2067 (47.6)
associate degree	31 (6.1)	284 (6.5)
BSc	56 (10.9)	653 (15.0)
MSc or higher degrees	6 (1.2)	108 (2.5)
Occupational status		
employed	378 (73.8)	1606 (37.0)
student	11 (2.1)	291 (6.7)
housewife	51 (10.0)	2039 (46.9)
no work with income	46 (9.0)	325 (7.5)
others	26 (5.1)	84 (1.9)
Physical activity (yes)	310 (60.5)	2812 (64.7)

Where $\hat{p}_{exposed}$ and $\hat{p}_{unexposed}$ are the proportion estimates of the binary confounders in the exposed and unexposed, respectively.

A standardized difference of more than 0.1 was considered as an important difference in mean or proportion of confounders between exposure groups.^[18] All statistical analyses were performed using stata.

Results

Of 4857 participants, 2959 (60.9%) were female, and the mean (SD) of age was 39.10 (13.48) years. There were 512 (10.5%) cigarette smokers at baseline, and 922 (19.0%) developed MS during the study follow-up. We excluded 25 (0.5%) participants due to missing MS data in phase 5. In PS matching, 511 unexposed were matched to 511 exposed. The mean (SD) of weights for ATE and ATT estimates were 2.02 (5.22) and 0.21 (0.30), respectively. The baseline characteristics of participants have been shown

Table 2: Standardized differences for confounders

Confounder	Original data	Weighted data	Matched data
Gender (male)	1.248	0.023	0.006
Age (year)	0.184	0.018	0.004
Marital status			
Single	0.030	0.015	0.005
divorced	0.091	0.017	0.000
widowed	0.138	0.036	0.019
Educational certificate			
elementary	0.076	0.086	0.019
secondary school	0.117	0.014	0.000
associate degree	0.020	0.007	0.008
BSc	0.122	0.006	0.000
MSc or higher degrees	0.098	0.064	0.017
Occupational status			
student	0.223	0.002	0.000
housewife	0.898	0.023	0.006
no work with income	0.055	0.045	0.007
others	0.172	0.010	0.048
Physical activity	0.086	0.020	0.008

in Table 1. Table 2 represents the standardized differences for confounders in original, weighted and matched data. In the original data, eight variables had standardized differences above 0.1, but in both weighted and matched data all variables had standardized differences less than 0.1, a sufficient balance on the confounders. The effects of smoking on MS and its components have been presented in Table 3. Based on ATE estimates, smoking decreased the risk of hypertension, RR: 0.62 (95% CI: 0.43, 0.88), but increased the risk of low HDL-C: 1.20 (95% CI: 0.98, 1.48). Similarly the ATT estimates using IPTW and PS matching suggested that smoking decreased the risk of hypertension, 0.63 (95% CI: 0.52, 0.76) and 0.68 (95% CI: 0.54, 0.85), and increased the risk of low HDL-C, 1.24 (95% CI: 1.07, 1.43) and 1.28 (95% CI: 1.06, 1.54), respectively. There was no strong evidence against no effect of smoking on MS, abdominal obesity, hypertriglyceridemia and impaired blood glucose.

Discussion

We did not find strong evidence against no effect of smoking on MS using causal methods which are consistent with previous studies.^[35,36] However, some studies indicated a positive relationship between smoking and MS.^[37,38] The difference in study population, MS definition, and statistical analysis may justify the controversial results.

Our estimates suggest that smoking increases the risk of low HDL-C which is consistent with the results of previous studies.^[39,40] This finding can be explained by the fact that smoking decreases lipid metabolism through diminished lipoprotein lipase activity.^[40] Also, nicotine can cause lipolysis which in turn may reduce HDL-C.^[6]

Our analyses revealed that smoking lowers BP. According to some studies, smokers have a lower risk of hypertension and a cluster analysis on a national survey in Iran showed that people in a cluster of smokers were less likely to experience elevated BP, though they had a very high level of work-related physical activity.^[41-43] These results may be interpreted as an indirect protective effect of smoking on BP through the body weight.^[44] However,

Table 3: The ATE and ATT estimates of the effect of smoking on MS and its components using IPTW and PS matching

	Risk Ratio (95% CI) (P)			
	Unweighted	ATE using IPTW (n: 4857)	ATT using IPTW (n: 4857)	ATT using PS matching (n: 1022)
MS	1.00 (0.83, 1.21) (0.92)	0.78 (0.55, 1.09) (0.15)	0.83 (0.68, 1.02) (0.08)	0.84 (0.66, 1.07) (0.16)
abdominal obesity	1.10 (0.98, 1.24) (0.07)	0.86 (0.69, 1.08) (0.20)	0.89 (0.78, 1.00) (0.06)	0.86 (0.75, 1.00) (0.06)
hypertriglyceridemia	1.30 (1.15, 1.48) (<0.001)	1.10 (0.87, 1.38) (0.39)	1.08 (0.95, 1.24) (0.20)	1.14 (0.96, 1.36) (0.11)
HDL-C	1.01 (0.89, 1.15) (0.77)	1.20 (0.98, 1.48) (0.06)	1.24 (1.07, 1.43) (0.003)	1.28 (1.06, 1.54) (0.009)
hypertension	0.84 (0.69, 1.09) (0.07)	0.62 (0.43, 0.88) (0.009)	0.63 (0.52, 0.76) (<0.001)	0.68 (0.54, 0.85) (0.001)
impaired blood glucose	1.24 (1.06, 1.44) (0.005)	0.99 (0.74, 1.32) (0.96)	1.04 (0.88, 1.22) (0.61)	0.97 (0.80, 1.19) (0.83)

Abbreviations: MS: metabolic syndrome, HDL-C: HDL cholesterol, ATE: average treatment effect, IPTW: inverse probability-of-treatment weighting, ATT: average treatment effect in the treated, PS: propensity score, CI: confidence interval. Notes: Estimates were adjusted for gender, age, physical activity, marital status, education, and job.

some studies indicated a higher risk of hypertension for the smokers.^[45,46] Moreover, previous studies have shown the positive effect of smoking on cardiovascular disease (2-6 times for >20 vs. <10 cigarette per day)^[47,48] which should always be considered as an important factor.

Our study had some limitations. The confounders alcohol consumption and income, were not available. Also there was measurement bias as smoking was dichotomized and self-reported. Finally some confounders like physical activity had measurement error leading to residual confounding.

Conclusions

In summary, there was no strong evidence against no effect of smoking on MS, abdominal obesity, hypertriglyceridemia and impaired blood glucose. Smoking seems to increase the risk of low HDL-C but decrease the risk of hypertension. More studies are needed to understand better if and how smoking affects MS and its components.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Received: 07 Feb 21 **Accepted:** 27 Feb 21

Published: 29 Sep 21

References

- Huang LN, Wang HJ, Wang ZH, Zhang JG, Jia XF, Zhang B, *et al.* Association of red meat usual intake with serum ferritin and the risk of metabolic syndrome in Chinese adults: A longitudinal study from the China health and nutrition survey. *Biomed Environ Sci* 2020;33:19-29.
- Annani-Akollor ME, Laing EF, Osei H, Mensah E, Owiredu E-W, Afranie BO, *et al.* Prevalence of metabolic syndrome and the comparison of fasting plasma glucose and HbA1c as the glycemic criterion for MetS definition in non-diabetic population in Ghana. *Diabetol Metab Syndr* 2019;11:26.
- Hwang G-Y, Cho Y-J, Chung R-H, Kim S-H. The relationship between smoking level and metabolic syndrome in male health check-up examinees over 40 years of age. *Korean J Fam Med* 2014;35:219.
- Huang C, Chen G, Zhang M, Lu Y, Hua Y, Hu Y, *et al.* Association between environmental tobacco smoke exposure and risk of type 2 diabetes mellitus in Chinese female never smokers: A population-based cohort study. *J Diabetes* 2020;12:339-46.
- Campagna D, Alamo A, Di Pino A, Russo C, Calogero A, Purrello F, *et al.* Smoking and diabetes: dangerous liaisons and confusing relationships. *Diabetol Metab Syndr* 2019;11:1-12.
- Kim BJ, Han JM, Kang JG, Rhee EJ, Kim BS, Kang JH. Relationship of cotinine-verified and self-reported smoking status with metabolic syndrome in 116,094 Korean adults. *J Clin Lipidol* 2017;11:638-45.e2.
- Slagter SN, van Vliet-Ostaptchouk JV, Vonk JM, Boezen HM, Dullaart RP, Kobold ACM, *et al.* Associations between smoking, components of metabolic syndrome and lipoprotein particle size. *BMC Med* 2013;11:195.
- Li G, Wang H, Wang K, Wang W, Dong F, Qian Y, *et al.* The association between smoking and blood pressure in men: A cross-sectional study. *BMC Public Health* 2017;17:797.
- Kim J-Y, Yang Y, Sim Y-J. Effects of smoking and aerobic exercise on male college students' metabolic syndrome risk factors. *J Phys Ther Sci* 2018;30:595-600.
- Kim Y, Jeong SM, Yoo B, Oh B, Kang H-C. Associations of smoking with overall obesity, and central obesity: A cross-sectional study from the Korea National Health and Nutrition Examination Survey (2010-2013). *Epidemiol Health* 2016;38:e2016020.
- Yankey BN, Strasser S, Okosun IS. A cross-sectional analysis of the association between marijuana and cigarette smoking with metabolic syndrome among adults in the United States. *Diabetes Metab Syndr* 2016;10:S89-95.
- Horie M, Noguchi S, Tanaka W, Goto Y, Yoshihara H, Kawakami M, *et al.* Relationships among smoking habits, airflow limitations, and metabolic abnormalities in school workers. *PloS One* 2013;8:e81145.
- Elze MC, Gregson J, Baber U, Williamson E, Sartori S, Mehran R, *et al.* Comparison of propensity score methods and covariate adjustment: Evaluation in 4 cardiovascular studies. *J Am Coll Cardiol* 2017;69:345-57.
- Aryaie M, Sharifi H, Saber A, Nazempour M, Mansournia MA. Longitudinal Causal Effects of Normalized Protein Catabolic Rate on All-Cause Mortality in Patients With End-Stage Renal Disease: Adjusting for Time-Varying Confounders Using the G-Estimation Method. *Am J Epidemiol.* 2021;190:1133-41.
- Mansournia MA, Danaei G, Forouzanfar MH, Mahmoodi M, Jamali M, Mansournia N, *et al.* Effect of physical activity on functional performance and knee pain in patients with osteoarthritis: Analysis with marginal structural models. *Epidemiology* 2012;631-40.
- Mansournia MA, Etminan M, Danaei G, Kaufman JS, Collins G. Handling time varying confounding in observational research. *BMJ* 2017;359:j4587.
- Almasi-Hashiani A, Nedjat S, Ghiasvand R, Safiri S, Nazempour M, Mansournia N, *et al.* The causal effect and impact of reproductive factors on breast cancer using super learner and targeted maximum likelihood estimation: a case-control study in Fars Province, Iran. *BMC Public Health.* 2021;21:1219.
- Austin PC. An introduction to propensity score methods for reducing the effects of confounding in observational studies. *Multivariate Behav Res* 2011;46:399-24.
- Alberti K, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, *et al.* Harmonizing the metabolic syndrome: A joint interim statement of the international diabetes federation task force on epidemiology and prevention; national heart, lung, and blood institute; American heart association; world heart federation; international atherosclerosis society; and international association for the study of obesity. *Circulation* 2009;120:1640-5.
- Delavari A, Forouzanfar MH, Alikhani S, Sharifian A, Kelishadi R. First nationwide study of the prevalence of the metabolic syndrome and optimal cutoff points of waist circumference in the Middle East: The national survey of risk

- factors for noncommunicable diseases of Iran. *Diabetes Care* 2009;32:1092-7.
21. AZIZI F, Hadaegh F, KHALILI D, Esteghamati A, HOSSEIN PF, Delavari A, *et al.* Appropriate definition of metabolic syndrome among Iranian adults: Report of the Iranian National Committee of Obesity. *Arch Iran Med* 2010;13:426-8.
 22. Etminan M, Brophy JM, Collins G, Nazemipour M, Mansournia MA. To Adjust or Not to Adjust: The Role of Different Covariates in Cardiovascular Observational Studies. *Am Heart J*. 2021;237:62-7.
 23. Mansournia MA, Higgins JP, Sterne JA, Hernán MA. Biases in randomized trials: A conversation between trialists and epidemiologists. *Epidemiology (Cambridge, Mass)*. 2017;28:54.
 24. Mansournia MA, Hernán MA, Greenland S. Matched designs and causal diagrams. *Int J Epidemiol* 2013;42:860-9.
 25. Etminan M, Nazemipour M, Sodhi M, Mansournia MA. Potential biases in studies of acid suppressing drugs and COVID-19 infection. *Gastroenterology* 2020. doi: 10.1053/j.gastro.2020.11.053.
 26. Mansournia MA, Nazemipour M, Etminan M. Causal diagrams for immortal time bias. *Int J Epidemiol*. 2021.
 27. Mansournia MA, Collins GS, Nielsen RO, Nazemipour M, Jewell NP, Altman DG, *et al.* A Checklist for statistical assessment of medical papers (the CHAMP statement): Explanation and elaboration. *Br J Sports Med* 2021. doi: 10.1136/bjsports-2020-103652.
 28. Pearl J. *Causality: Models, reasoning and inference* cambridge university press. Cambridge, MA, USA. 2000.
 29. Abdollahpour I, Nedjat S, Mansournia MA, Sahraian MA, Kaufman JS. Estimating the marginal causal effect of fish consumption during adolescence on multiple sclerosis: A population-based incident case-control study. *Neuroepidemiology* 2018;50:111-8.
 30. Mansournia MA, Altman DG. Inverse probability weighting. *BMJ* 2016;352:i189.
 31. Abdollahpour I, Nedjat S, Mansournia MA, Schuster T. Estimation of the marginal effect of regular drug use on multiple sclerosis in the Iranian population. *PloS One* 2018;13:e0196244.
 32. Hernán MA, Robins JM. *Causal inference*. 2016.
 33. Mansournia MA, Nazemipour M, Naimi AI, Collins GS, Campbell MJ. Reflections on modern methods: Demystifying robust standard errors for epidemiologists. *Int J Epidemiol* 2020. doi: 10.1093/ije/dyaa260.
 34. Austin PC, Small DS. The use of bootstrapping when using propensity-score matching without replacement: A simulation study. *Stat Med* 2014;33:4306-19.
 35. Santos A-C, Ebrahim S, Barros H. Alcohol intake, smoking, sleeping hours, physical activity and the metabolic syndrome. *Prev Med* 2007;44:328-34.
 36. Ishizaka N, Ishizaka Y, Toda E-I, Nagai R, Yamakado M. Association between cigarette smoking, white blood cell count, and metabolic syndrome as defined by the Japanese criteria. *Intern Med* 2007;46:1167-70.
 37. Wada T, Urashima M, Fukumoto T. Risk of metabolic syndrome persists twenty years after the cessation of smoking. *Intern Med* 2007;46:1079-82.
 38. Sun K, Liu J, Ning G. Active smoking and risk of metabolic syndrome: A meta-analysis of prospective studies. *PloS One* 2012;7:e47791.
 39. Cheng E, Burrows R, Correa P, Güichapani CG, Blanco E, Gahagan S. Light smoking is associated with metabolic syndrome risk factors in Chilean young adults. *Acta Diabetol* 2019;56:473-9.
 40. Wakabayashi I. Relationship between smoking and metabolic syndrome in men with diabetes mellitus. *Metab Syndr Relat Disord* 2014;12:70-8.
 41. Gharipour M, Kelishadi R, Sarrafzadegan N, Baghaei A, Yazdani M, Anaraki J, *et al.* The association of smoking with components of the metabolic syndrome in non-diabetic patients. *Ann Acad Med Singap* 2008;37:919.
 42. Akbarpour S, Khalili D, Zeraati H, Mansournia MA, Ramezankhani A, Pishkuhi MA, *et al.* Relationship between lifestyle pattern and blood pressure-Iranian national survey. *Sci Rep* 2019;9:1-8.
 43. Akbarpour S, Khalili D, Zeraati H, Mansournia MA, Ramezankhani A, Fotouhi A. Healthy lifestyle behaviors and control of hypertension among adult hypertensive patients. *Sci Rep* 2018;8:1-9.
 44. Leone A. Smoking and hypertension. *J Cardiol Curr Res* 2015;2:00057.
 45. Al-khalifa II, Mohammed SM, Ali ZM. Cigarette smoking as a relative risk factor for metabolic syndrome. *J Endocrinol Metab* 2017;6:178-82.
 46. Bowman TS, Gaziano JM, Buring JE, Sesso HD. A prospective study of cigarette smoking and risk of incident hypertension in women. *J Am Coll Cardiol* 2007;50:2085-92.
 47. Khalili D, Sheikholeslami FH, Bakhtiyari M, Azizi F, Momenan AA, Hadaegh F. The incidence of coronary heart disease and the population attributable fraction of its risk factors in Tehran: A 10-year population-based cohort study. *PloS One* 2014;9:e105804.
 48. Ehteshami-Afshar S, Momenan A, Hajshekholeslami F, Azizi F, Hadaegh F. The impact of smoking status on 9.3 years incidence of cardiovascular and all-cause mortality among Iranian men. *Ann Hum Biol* 2014;41:249-54.