

Invited Mini Review

Harnessing NK cells for cancer immunotherapy:
immune checkpoint receptors and chimeric antigen receptorsNayoung Kim^{1,2,#}, Dong-Hee Lee^{1,2,#}, Woo Seon Choi^{3,4,#}, Eunbi Yi^{3,4}, Hyojeong Kim^{3,4}, Jung Min Kim³, Hyung-Seung Jin^{1,2}
& Hun Sik Kim^{3,4,5,*}¹Department of Convergence Medicine and ²Asan Institute for Life Sciences, Asan Medical Center, University of Ulsan College of Medicine, Seoul 05505, ³Department of Biomedical Sciences, ⁴Stem Cell Immunomodulation Research Center (SCIRC), ⁵Department of Microbiology, Asan Medical Center, University of Ulsan College of Medicine, Seoul 05505, Korea

Natural killer (NK) cells, key antitumor effectors of the innate immune system, are endowed with the unique ability to spontaneously eliminate cells undergoing a neoplastic transformation. Given their broad reactivity against diverse types of cancer and close association with cancer prognosis, NK cells have gained considerable attention as a promising therapeutic target for cancer immunotherapy. NK cell-based therapies have demonstrated favorable clinical efficacies in several hematological malignancies but limited success in solid tumors, thus highlighting the need to develop new therapeutic strategies to restore and optimize anti-tumor activity while preventing tumor immune escape. The current therapeutic modalities yielding encouraging results in clinical trials include the blockade of immune checkpoint receptors to overcome the immune-evasion mechanism used by tumors and the incorporation of tumor-directed chimeric antigen receptors to enhance NK cell anti-tumor specificity and activity. These observations, together with recent advances in the understanding of NK cell activation within the tumor microenvironment, will facilitate the optimal design of NK cell-based therapy against a broad range of cancers and, more desirably, refractory cancers. [BMB Reports 2021; 54(1): 44-58]

INTRODUCTION

Immune checkpoint receptors, such as programmed cell death protein 1 (PD-1) and cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), have recently emerged as molecular targets for

cancer immunotherapy (1). Even before the investigation of the importance of such receptors in T cells (2, 3), the foundation of NK cell immunology was established by determining the quintessential roles of killer cell immunoglobulin-like receptors (KIRs), the inhibitory receptors in human NK cells. In 1986, Kärre *et al.* proposed the “missing-self” hypothesis (4). NK cells kill major histocompatibility complex (MHC) class I-deficient tumor cells but fail to kill MHC class I-expressing tumor cells. The recognition of “missing-self” MHC class I is mediated by KIRs in humans and Ly49s in mice. Later, Kim *et al.* suggested that “licensing” of NK cells is also mediated by the interaction between Ly49s and MHC class I (5). Licensing is a host MHC class I-dependent functional maturation process. Only NK cells that are licensed by self-MHC class I molecules during development are fully functional. Licensing also occurs in human NK cells (6). Thus, immune checkpoint receptors are fundamental for determining NK cell functionality. Nonetheless, NK cells express multiple immune checkpoint receptors, including natural killer group 2A (NKG2A), CTLA-4, PD-1, T cell immunoglobulin mucin 3 (TIM-3), and T cell immunoreceptor with Ig and immunoreceptor tyrosine-based inhibition motif (ITIM) domains (TIGIT), which have been explored as promising therapeutic targets to enhance the specificity and activity of NK cells against a broad range of cancers.

Another promising cancer treatment modality that has raised considerable interest is the incorporation of tumor-directed chimeric antigen receptors (CARs) in immune effector cells. The clinical success of Kymriah[®] and Yescarta[®], two CAR-T cell therapies targeting hematologic malignancies, is sure to promote the growth of CAR-T cell therapies in clinical trials, thereby treating a range of cancers. Nevertheless, the limitations of CAR-T cell therapies, in terms of off-the-shelf utility, safety, and target antigen escape, necessitate alternatives. With an array of innate receptors responding to cellular transformation, NK cells can efficiently kill a range of tumor cells without MHC restriction, thereby complementing MHC-restricted tumor lysis by cytotoxic T cells. With radical differences in tumor cell recognition, cytokine production profile, and *in vivo* persistence, NK-CAR cell therapies are viewed as an attractive alternative or complement to CAR-T cell therapies, as they poten-

*Corresponding author. Tel: +82-2-3010-2207; Fax: +82-2-3010-5307; E-mail: hunkim@amc.seoul.kr

[#]These authors contributed equally to this work.

<https://doi.org/10.5483/BMBRep.2021.54.1.214>

Received 29 September 2020, Revised 23 November 2020,
Accepted 3 December 2020

Keywords: Cancer immunotherapy, Chimeric antigen receptors, Immune checkpoint receptors, Immune escape, Natural killer cells

tially overcome several clinical challenges presented by CAR-T cell therapies. In this review, we summarize recent advances in NK cell-based cancer immunotherapy with a focus on immune checkpoint receptors, some of which are unique to NK cells and CAR-NK cells.

IMMUNE CHECKPOINT RECEPTORS

KIR, LIR, and CD94/NKG2A

Inhibitory KIRs, *i.e.* 2DL1, 2DL2, 2DL3, 2DL5, 3DL1, 3DL2, and 3DL3, have long cytoplasmic tails comprising two ITIMs (7). Inhibitory KIRs recognize human leukocyte antigen (HLA)-A, B, or C (Fig. 1). The main roles of KIRs in NK cells are described above. They have highly polymorphic immunoglobulin (Ig) domains specific to HLA molecules (8). Multiple myeloma (MM) cells upregulate MHC class I molecules, thus blocking inhibitory KIRs could enhance the antitumor effect of NK cells in MM. Treatment with anti-KIR2D antibody (Ab) (Lirilumab; IPH2102) has been proven safe without mediating toxicity and autoimmunity in patients with MM and acute myeloid leukemia (AML) (9, 10). Anti-KIR Ab treatment enhanced *ex vivo* NK cytotoxicity in patients with MM, but the Phase II clinical trial of lirilumab in MM resulted in failure due to lack of efficacy

and presumably a loss of KIR2D expression in NK cells (11). Anti-KIR Abs have been tested alone or in combination with other therapeutics, including lenalidomide, anti-CD20 Ab (rituximab), and immune checkpoint blockades in various hematological disorders, including MM, lymphoma, and myelodysplastic syndromes (12-15). The latest addition to this group is lacutamab (IPH4102), a first-in-class anti-KIR3DL2 Ab. It has been demonstrated to be safe, and 36% of patients with relapsed/refractory cutaneous T cell lymphoma responded to it in a Phase I trial (16). Apart from immune checkpoint blockade, pre-treatment with IL-12/15/18 reduces the expression of KIRs in NK cells and enhances NK cytotoxicity against tumor cells (17), suggesting that *ex vivo* expanded NK cells could be potent anti-tumor therapeutics by themselves or as CAR bearers. Clinical trials using immune checkpoint blockade are summarized in Table 1.

Among leukocyte immunoglobulin-like receptors (LIRs), LIR-1, also known as LIR subfamily B member 1 (LIRB-1), immunoglobulin-like transcript 2 (ILT2), and CD85j, recognizes HLA-G, a non-classical MHC class I molecule. LIR-1 contains ITIM motifs to recruit phosphatases, such as SHP-1 (18). HLA-G is expressed in various tumors and is often associated with reduced NK function or progressive tumors (19). Soluble HLA-G (sHLA-G) also plays a role in mediating regulatory function in some tumors, such as thyroid and colorectal cancers (20, 21). Blocking LIR-1 alone did not enhance the cytotoxicity of NK cells against MM cells (22), but a dual blockade of LIR-1 and NKG2A increased the cytotoxicity of KIR⁻ NK cells against acute leukemic cells *in vitro* (23). NKG2A is well known as an HLA-E receptor but has recently been suggested as an HLA-G receptor (24). However, the action mechanisms for the dual blockade may require further investigation.

CD94/NKG2A is a heterodimeric inhibitory receptor related to C-type lectins, recognizing another non-classical MHC class I molecule, HLA-E. ITIMs are phosphorylated upon receptor engagement and recruit tyrosine phosphatases SHP-1 and SHP-2 (25, 26). SHP-1 mediates dephosphorylation of Vav1 (27). In addition, Crk phosphorylation contributes to the inhibition of NK cells through NKG2A-HLA-E interaction (28). ITIM-based inhibition appears to be dominant over activation in NK cells against normal cells. Recruitment of SHP-1 by MHC-I-specific ITIM-bearing receptors inhibited signaling at a proximal step, such that most downstream signals were prevented (29). HLA-E is overexpressed in human colorectal cancers with poor prognosis (30). Ovarian and cervical cancer cells express HLA-E that limits NKG2A⁺ cytotoxic T cells, thereby resulting in less infiltration of NK cells in HLA-E-expressing gynecological cancer (31). In addition, NKG2A⁻NKG2C⁺KIR⁺CD56^{dim} NK cells are suggested as memory-like NK cells in patients with human cytomegalovirus infection (32). An anti-NKG2A Ab (monalizumab; IPH2201) ameliorates NK cell dysfunction in chronic lymphocytic leukemia (33). Monalizumab is currently under clinical investigation as a single agent in ovarian cancer or in combination with cetuximab (anti-EGFR Ab) and durvalumab




Fig. 1. Interactions between immune checkpoint receptors and their cognate ligands. NK cells express multiple immune checkpoint receptors, which can interact with their cognate ligands on tumor cells as well as other immune cells, in particular, dendritic cells and Tregs. The red circles represent immune checkpoint receptors while the blue circles represent the ligands. SIGLEC7 and SIGLEC9 have common ligands that are sialic acids. The pink squares represent the classical ITIM motif and the light blue squares represent the ISTM motif, which have been implicated in mediating inhibitory signals. The light green squares represent the ITT-like motif. Cytoplasmic domains of other immune checkpoint receptors contain fewer known motifs (not marked as squares). CD73 is a nucleotidase, which does not have conventional inhibitory signaling domains. The black lines indicate receptor-ligand interactions.

Table 1. Current status of clinical trials based on immune checkpoint receptors

Checkpoint receptor	Ab/drug	Combination drugs	Disease	Phase	Clinical trials identifier
KIRs	Anti-KIR (1-7F9, IPH2101)	Single	MM	Phase I	NCT00552396
		Single	MM, SMM	Phase II	NCT01248455
	Lirilumab (IPH2102, BMS-986015)	Lenalidomide	MM	Phase I	NCT01217203
		Nivolumab, Azacitidine	MDS	Phase II	NCT02599649
CD94/NKG2A	Monalizumab (IPH2201)	Single	Gynecologic cancer	Phase I	NCT02459301
		Durvalumab (MEDI4736)	Advanced solid tumors	Phase I/II	NCT02671435
		Cetuximab, Anti-PD-L1	Head and neck carcinoma	Phase I/II	NCT02643550
		Ibrutinib	CLL	Phase I/II	NCT02557516
CTLA-4	Ipilimumab (BMS-734016)	Single	Advanced melanoma	Phase I	NCT00920907
		Nivolumab	Advanced/metastatic melanoma	Phase II	NCT01783938
		Paclitaxel, Cisplatin, Carboplatin	NSCLC	Phase II	NCT01820754
PD-1	Pembrolizumab (MK-3475)	Single	Hepatocellular carcinoma	Phase II	NCT02658019
		Nivolumab	Ipilimumab	Advanced/metastatic melanoma	Phase II
	Durvalumab (MEDI4736)	Tremelimumab	Metastatic pancreatic ductal adenocarcinoma	Phase II	NCT02558894
TIM-3	BGB-A425	Tislelizumab	Advanced or metastatic solid tumors	Phase I/II	NCT03744468
	MBG453	Decitabine, PDR001	AML and high risk MDS	Phase I	NCT03066648
TIGIT	MTIG7192A	Atezolizumab, Carboplatin, Cisplatin, Pemetrexed, Paclitaxel, Etoposide	Advanced/metastatic tumors	Phase I	NCT02794571
LAG-3	LAC-3-Ig (IMP321)	Single	Metastatic breast cancer	Phase I	NCT00349934
CD47	IBI188	Montanide ISA-51	Melanoma	Phase I/II	NCT01308294
		Single	Advanced malignancies	Phase I	NCT03763149
CD73	Oleclumab (MEDI9447)	Paclitaxel, Carboplatin, Durvalumab	Triple negative breast cancer	Phase I/II	NCT03616886
		Ciforadenant (CPI-444)	Pembrolizumab	Advanced cancer	Phase I
CD33 (Siglec 3)	Vadastuximab talirine (SGN-CD33A)	Azacitidine, Decitabine, Placebo	AML	Phase III	NCT02785900

MM, multiple myeloma; SMM, smoldering multiple myeloma; AML, acute myeloid leukemia; MDS, myelodysplastic syndromes; CLL, chronic lymphocytic leukemia; NSCLC, non-small cell lung cancer; MBC, metastatic breast carcinoma.

(anti-PD-L1 Ab) for advanced stage solid cancers (34, 35). Interim results of a Phase II trial of monalizumab and cetuximab in previously treated squamous cell head and neck cancer showed a 31% objective response rate, where monalizumab improved anti-tumor immunity of T and NK cells (36). A combination of monalizumab and durvalumab demonstrated

clinical efficacy and manageable toxicity in a Phase I trial of heavily pretreated metastatic microsatellite colorectal cancer (19). However, NKG2A blockade reportedly works through CD8 T cells rather than NK cells in mouse models that are set to block NKG2A/Qa-1b interaction using HPV16 E6 and E7-expressing tumors (37). Taken together, NKG2A blockade

appears to be a promising immune-oncological therapeutic that promotes T and/or NK cell activation. Notably, NKG2A can recognize HLA-G as well (24), thereby suggesting the previously unexpected benefit of NKG2A blockade in tumor immunity.

CTLA-4 and PD-1

CTLA-4 plays a pivotal role in T cell expansion, whereas PD-1 is a central regulator of T cell effector function. CD80 (B7-1) and CD86 (B7-2) are the common ligands for the costimulatory receptor CD28 as well as the co-inhibitory receptor CTLA-4. However, CTLA-4 binds to ligands with greater affinity than CD28. Despite the absence of inhibitory ITIM, CTLA-4 inhibits the activation of Akt but not PI3K via activating the serine/threonine phosphatase PP2A (38). Engagement of CTLA-4 with CD80 leads to the reduction in IFN- γ production by mouse activated NK cells against mature dendritic cells (39). In head and neck cancer, CTLA-4 is upregulated on Treg cells that suppress NK cell anti-tumor cytotoxicity (40). In melanoma, anti-CTLA-4 treatment leads to Fc receptor-mediated selective depletion of Treg cells (41, 42). Moreover, clinical outcome of CTLA-4 therapy in melanoma is associated with the increased population of mature circulating CD3⁻CD56^{dim}CD16⁺ NK cells (43). Thus, anti-CTLA-4 therapy may enhance anti-tumor cytotoxicity of NK cells in both a direct and indirect manner such as depletion of CTLA-4⁺ Treg cells. Triple immunotherapy with anti-CTLA4 antibodies, monophosphoryl-lipid-A, and indolamine-dioxygenase-1 inhibitor has been reported to enhance NK cell counts and the CD3⁺CD4⁺/Treg and CD3⁺CD8⁺/Treg ratios, in addition to the reduction in tumor mass, in a murine melanoma model (44). Combination therapies could provide additional benefits, although the B7/CTLA-4 axis may not play a key role in NK cell activation (45, 46).

PD-1 has one ITIM and one immunoreceptor tyrosine-based switch motif (ITSM) in its cytoplasmic domain. Specifically, the ITSM tyrosine (Y248) of PD-1 is known to recruit phosphatase SHP-2, which is mandatory for PD-1-mediated inhibition of the PI3K/Akt pathway (47). The cognate ligands for PD-1 are PD-L1 (B7-H1) and PD-L2 (B7-DC). PD-1 expression is found on CD56^{dim}NKG2A⁻KIR⁺CD57⁺ mature NK cells, but not on CD56^{bright} NK cells (48). In ovarian cancer and Kaposi sarcoma, PD-1 expression is elevated on NK cells and associated with impaired NK cell function (49, 50). PD-1⁺ NK cells are considered to be functionally exhausted (32). Blockade of PD-1 enhances cytotoxicity of NK cells against autologous MM cells (51). In Hodgkin lymphoma and diffuse large B-cell lymphoma, PD-1 blocking also reverses the suppression of PD-1⁺ NK cells mediated by tumor-associated macrophage-like monocytes (52). In mice, tumor-infiltrated NK cells express PD-1, which suppresses NK cytotoxicity (53). PD-1/PD-L1 blockade, PD-1/PD-L1 genetic deficiency, or NK cell depletion prevents lung metastasis in a B16 melanoma model and tumor growth in a murine model using CT26 colon tumor cells and a breast cancer orthotopic model using 4T1 cells *in vivo* (53). However, activated human primary NK cells efficiently killed

colorectal cancer cells in organoid culture independently of PD-L1 expression (54), and blockade of PD-L1 failed to increase cytotoxicity of human liver-associated NK cells against hepatocellular carcinoma cell lines *in vitro* (55). In line with these results, PD-1 is expressed only minimally in mouse and human NK cells in various infections and tumor models (56). Nevertheless, PD-1 expression in murine NK cells can be induced in the spleen and liver by glucocorticoid (57), and anti-PD-1 blockade can induce cytokine production, such as IFN- γ , which may boost NK cells indirectly (34).

Four ongoing clinical trials are evaluating the combined effect of infused NK cells and anti-CTLA-4, PD-1, or PD-L1. They are induced pluripotent stem cell (iPSC)-derived NK cells combined with nivolumab or pembrolizumab (NCT03841110), cytokine-induced memory-like NK cells and ipilimumab (NCT 04290546), unmodified allogeneic NK cells and pembrolizumab (NCT03937895), and autologous NK cells combined with avelumab or pembrolizumab (NCT03941262).

TIM-3, CEACAM1, TIGIT, and LAG-3

TIM-3, whose cognate ligands are galectin-9 (Gal-9), phosphatidylserine, high mobility group box 1 (HMGB1), and carcinoembryonic antigen-related cell adhesion molecule 1 (CEACAM-1), does not have a classical signaling motif, but five conserved tyrosine residues (58). In particular, phosphorylated Tyr256 and Tyr263 are required for the Gal-9-mediated BAT3 release from TIM-3 and inhibitory signaling (59). Gal-9 and HMGB1 can be soluble as well as membrane-bound. TIM-3 is regarded as a marker for mature NK cells and TIM-3⁺PD-1⁺ NK cells are considered to be functionally exhausted (60). The expression of TIM-3 is elevated on peripheral NK cells in patients with advanced gastric cancer (61) and lung adenocarcinoma (62). It is also upregulated on the tumor-infiltrated NK cells in over 70% of patients with gastrointestinal stromal tumors (63). Interestingly, PD-1 expression is not found on the TIM-3⁺ tumor-infiltrated NK cells. Anti-TIM3 treatment rescues TIM-3⁺ exhausted NK cells from patients with advanced melanoma (64). Further, TIM-3 expression levels are correlated with the stage of the disease. Several anti-TIM-3 Abs are to be tested in Phase I or II clinical trials: TSR-022 by Tesaro, LY3321367 by Eli Lilly, MGB453 by Novartis, Sym023 by Symphogen, and BGB-A425 by Beigene (65, 66). The antibodies are often applied in combination with anti-PD-1 or anti-LAG-3 Abs in advanced solid tumors or AML. The studies are still recruiting patients, and results will be available in a few years. However, caution is warranted as the blockade of TIM-3 leads to the reduction in NK cell-mediated cytolysis of pancreatic cancer cell lines (67). Moreover, Ab-mediated Gal-9 blocking leads to a decrease in IFN- γ production in NK cells in response to primary AML blasts (68), thereby complicating the outcomes of TIM-3 blocking.

Interestingly, CEACAM1 is also expressed in NK cells and interacts with CEACAM5 (69). Recently, NEO-201, a monoclonal antibody (mAb) specific to the CEACAM family was demonstrated to enhance NK cytotoxicity against various human

tumor cells through CEACAM5 on tumor cells and CEACAM1 on NK cells *in vitro* (70) and *in vivo* (71). CEACAM1-expressing NK cells produce IFN- γ by IL-12/18 in a mouse hepatitis virus infection model and the engagement of CEACAM1 was demonstrated to decrease the production IFN- γ (72). CEACAM1 on activated NK cells inhibits NKG2D-mediated cytolytic function and signaling (73), suggesting that CEACAM1 is an immune checkpoint receptor in NK cells, as well as a cognate ligand to TIM-3. Human CEACAM1 protein has a signaling cytoplasmic domain comprising either a long ITIM-containing domain or a short domain devoid of ITIMs (74). CEACAM1 could have 12 alternatively spliced isoforms that lead to the generation of proteins with potentially different functions.

TIGIT and CD96 are inhibitory receptors that compete with DNAM-1 (CD226), an activating receptor, for CD155 (PVR), and CD112 (Nectin-2). CD155 is highly expressed in many types of tumor cells. TIGIT and CD96 contain the ITIM motif. TIGIT contains an ITT-like motif in addition to an ITIM motif in the cytoplasmic tail, where phosphorylation of ITT-like motif upon ligand binding plays a critical role in inhibitory signaling via the recruitment of SHIP1. Engagement of TIGIT with CD155 induces its phosphorylation through Fyn and Lck and recruits SHIP1 in T cells (58). High TIGIT expression is associated with the exhaustion of tumor-infiltrated NK cells in patients with colorectal cancer (75). The blockade of TIGIT prevents NK cell exhaustion and elicit potent anti-tumor immunity in mice (75). Combined blocking of TIGIT and PD-1 showed significant tumor clearance in mice (76). TIGIT and PD-1 are often co-expressed in tumor-infiltrated NK cells (76), but only TIGIT is associated with NK cell exhaustion (75). As PVR expression is associated with unfavorable prognosis in many solid tumors, such as colon, breast, lung, and pancreatic cancers, the "PVR-TIGIT axis" has been suggested as a novel target in immune checkpoint therapy (77). Notably, tiragolumab, an anti-TIGIT Ab developed by Genentech, is already being evaluated in two independent Phase-III clinical trials for small cell lung cancer and non-small cell lung cancer with atezolizumab, an anti-PD-L1 Ab (66), and chemotherapy. There are two other anti-TIGIT Abs, MTIG7192A and AB154, in Phase I or II trials for various solid tumors. The role of CD96 is relatively less elucidated in NK cells.

Lymphocyte activation gene-3 (LAG-3) is structurally similar to CD4 and binds to MHC class II molecules with a higher affinity than CD4. Fibrinogen-like protein 1 (FGL1) is a recently identified ligand for LAG-3 (78). LAG-3 transduces two independent inhibitory signals through the FXXL motif in the membrane-proximal region and the C-terminal EX repeat (79); the motifs are unique among the known inhibitory receptors. It is expressed on activated NK cells, and chronic stimulation of NKG2C⁺ NK cells can induce high expression of LAG-3 (80). A soluble form of LAG-3-Ig fusion protein, IMP321 induces human NK cells to produce IFN- γ and TNF- α *in vitro* (81). In human trials, IMP321 induced NK cell activation as monothe-

rapy in advanced renal cell carcinoma (82) and combination with paclitaxel in metastatic breast cancer (83). FGL1 blockade also potentiated anti-tumor T cell responses in mice (78), but that of NK cells is not yet known.

CD47, CD73, AND SIGLEC FAMILY PROTEINS

In this section, we introduce some of the emerging immune checkpoint molecules in NK cell biology. CD47 is an integrin-associated protein with a short cytoplasmic domain, interacting with thrombospondin-1 (TSP-1) and signal regulatory protein α (SIRP α), an inhibitory transmembrane protein. CD47 regulates NK cell homeostasis and immune responses to lymphocytic choriomeningitis virus infection (84) and NK cell recruitment and activation in the tumor microenvironment in mice (85). CD47 is quite ubiquitously expressed. Elevated CD47 expression is associated with reduced survival in some cancers. Cord blood cell-derived CD16⁺ NK cells respond well to anti-CD47 Ab-treated T and B-ALL cell lines with an approximately 10% increase in cytotoxicity (86). CD47 blockade with trastuzumab (anti-HER-2 mAb) augmented anti-tumor efficacy, but the effect appears to be due to increased phagocytosis, rather than ADCC (87).

CD73, ecto-5'-nucleotidase, is probably the latest addition to immune checkpoint molecules in NK cells. The expression of CD73 is virtually absent in circulating human and mouse NK cells in healthy individuals, but tumor-infiltrated NK cells express substantial CD73 (88). It defines regulatory NK cells in the tumor environment in patients with breast cancer and sarcoma (89). CD73⁺ NK cells in the tumor microenvironment express LAG-3, VISTA, PD-1, and PD-L1. NK cells transport CD73 upon engagement of 4-1BB on tumor cells, to express IL-10 via STAT3 activation (89). CD73 is suggested as a correlative factor of patient survival and NK cell infiltration in glioblastoma (90) and mediates immunometabolic dysfunction of NK cells under hypoxic conditions in solid tumors (91). Targeting CD73 has also been shown to suppress tumorigenesis. A first-in-class therapeutic anti-CD73 mAb, MEDI9447, is currently being evaluated in Phase I clinical trials in cancer patients (88).

Among sialic acid-binding immunoglobulin-like lectins (Siglecs), Siglec7 and Siglec9 are expressed in NK cells. Sialic acids, cognate ligands of Siglecs are 9-carbon-backbone monosaccharides, which are the glycan residues of glycoproteins and glycolipids. The cytoplasmic domains of Siglec7 and Siglec9 contain an ITIM and an ITIM-like motif (92). Siglec7 and Siglec9 share structural similarity and functionality but have different roles in virus infection and tumors. Siglec7 is expressed on mature or more cytotoxic NK cells and can reduce NK cytotoxicity. Sialic acid-containing glycan has been reported to protect tumor cells from NK cells through Siglec7 (93). Hypersialylated tumor cells can bind to Siglec9, and Siglec9⁺ NK cells express higher levels of KIRs and LIR-1. Siglec7 interacts with gangliosides, while Siglec9 interacts with mucins (92). Importantly, the desi-

ylation of tumor cells by neuraminidases enhanced NK cytotoxicity and cytokine production (94), thereby implying novel therapeutic approaches. Siglec3 (CD33) is just recently identified as an inhibitory receptor on NK cells (95). Siglec3 inhibits cytotoxicity triggered by NKG2D via Vav1 dephosphorylation, but not by Nkp46 (95).

CHIMERIC ANTIGEN RECEPTORS

The potential advantage of CAR-NK cells over CAR-T cells

The majority of CAR-T cell therapies, including Kymriah[®] and Yescarta[®], use autologous T cells collected from cancer patients. However, the use of autologous T cells has well-known disadvantages, including a complex manufacturing process and a low quantity of patient cells (96). Furthermore, T cell dysfunction can occur in patients who have received previous treatment with chemotherapy and/or certain other medications (97). To overcome these limitations, gene-editing tools to knockout T-cell receptors (TCRs) and human leukocyte antigen (HLA) have been employed for the generation of allogeneic CAR-T cells. These tools apply to non-HLA matched patients by reducing the potential risk of graft versus host disease (GVHD) (98-100). However, highly gene-edited cells come with unknown risks.

CAR-NK cell therapies traverse several of the limitations of CAR-T cell therapies. First, NK cells can recognize and kill tumors without HLA matching or prior antigen-sensitization (101). The transfer of allogeneic NK cells has even been shown to mediate graft-versus-tumor (GvT) responses without GvHD (102). Subsequently, NK cells can be obtained from several different sources including peripheral blood, umbilical cord blood, embryonic stem cells, and induced pluripotent stem cells (iPSCs) (103-105).

In addition, NK cell lines such as NK-92 can be utilized as allogeneic off-the-shelf CAR-expressing cell products. In contrast to primary NK cells, CAR-expressing NK-92 cells can be manufactured from a functionally and molecularly characterized single-cell clone under good manufacturing practice-compliant conditions (106). The CRISPR-Cas9 genome editing technology may allow for site-specific integration of the CAR, thereby mitigating the risk of any dysfunction in the NK-92-CAR cells. Additionally, NK-92 cells require irradiation before infusion into patients to avoid potential malignant expansion. Nonetheless, repeated infusion of irradiated CAR-NK92 cells can maintain the efficacy of CAR-NK therapy depending on the dose and frequency.

A major safety concern with CAR T-cell therapy is cytokine release syndrome (CRS). Aberrant activation of CAR T cells can lead to massive production of inflammatory cytokines including IL-6 (107). Several clinical trials have demonstrated that the adoptive transfer therapy of allogeneic CAR-NK cells does not cause severe side effects (108). CAR-NK cells may be potentially safer than CAR-T cells because of their shorter lifespan after infusion (109). There exists little clinical evidence for the comparison of the side effects of CAR-T and CAR-NK

cells (110). However, according to the clinical trial of CAR-NK cell therapy at the University of Texas MD Anderson Cancer Center, the treatment of CAR-NK cells derived from cord blood showed complete remission in 7 of the 11 patients (4 with non-Hodgkin's lymphoma and 3 with chronic lymphocytic leukemia) without CRS, neurotoxicity, or GvHD, which are all potential side effects of CAR-T therapy (111).

Antigen escape is a major obstacle for effective CAR-T therapy. The immune pressure by CAR-T cells results in the outgrowth of antigen loss variants. In hematologic malignancies, CD19 loss after CAR-T therapy drives relapses (112, 113). CAR-NK cells could show effective antitumor activity against target antigen-negative tumors by endogenous activating receptors such as NKG2D, Nkp30, Nkp44, Nkp46, or DNAM-1 that are involved in tumor immune surveillance (114, 115). In addition, cytotoxic response via activating receptors including CD16 (FcγRIII) mediating antibody-dependent cellular cytotoxicity (ADCC) can synergistically enhance the antitumor activity of CAR-NK cells (116).

Enhancing the anti-tumor activity of CAR-NK cells

CARs comprise a signal peptide, a single-chain variable fragment (scFv), hinge region, transmembrane region, and intracellular domains. The composition of conventional CARs is as follows: a CD4, CD8, or IgG hinge; a CD3ξ, CD4, CD8, or CD28 transmembrane domain; a 4-1BB or CD28 costimulatory domain; and a CD3ξ activation domain (117, 118). Herein, we describe the current efforts to optimize the clinical outcome of CAR-NK therapies for each domain.

Antigen recognition domain: The single-chain variable fragment (scFv) is an extracellular component of a CAR and confers antigen specificity. scFv is derived from the variable regions of the heavy chain and light chain of antibodies. Since the main function of scFv is antigen binding, the proper design of scFv against tumor antigens is required for better clinical outcomes. Typically, therapeutic antibodies with high-affinity binding to a specific antigen are preferred for predicted responses with minimal side effects. Although designing an scFv for a CAR is dependent on mAbs specific to the tumor-associated antigen, optimizing CAR scFv sequences for the best clinical outcome is still under discussion. Chmielewski *et al.* generated T cells expressing chimeric TCR pools against human epidermal growth factor receptor 2 (ERBB2, HER2) with a Kd ranging from 3.2×10^{-7} to 1.5×10^{-11} M. Researchers demonstrated that a higher affinity receptor did not correlate with the potency of T cell activation. Moreover, T cells with high-affinity receptors could induce an immune response to normal tissue expressing low antigen levels (119). Morgan *et al.* reported a clinical case study on the administration of CAR-T cells-derived from the FDA-approved therapeutic mAb, trastuzumab (Herceptin), that targets HER2, a tumor-associated antigen used to treat metastatic colon cancer. The patient presented side effects resulting from a dramatic pulmonary infiltration of CAR-T cells by the recognition of low expression levels of HER2 in normal lung

epithelial cells and died 5 days after administration (120). Phase I clinical trial of NK-92/5.28.z (CAR-NK-92 cells targeting HER2 with trastuzumab scFv) is presently ongoing in patients with recurrent HER2-positive glioblastoma (CAR2BRAIN, NCT03383978, clinicaltrials.gov). So far, no remarkable adverse effects have been observed, but more detailed clinical signs should be confirmed in dose-escalating studies (121).

The design of an scFv for CAR-NK therapies is still in its early stages. Because CAR-NK cells are generally more tolerant than CAR-T cells, we would select the one with the strongest affinity. Moreover, NK-92-CAR cells have to be irradiated before infusion, this process can decrease the potency of NK-92-CAR cells. Currently, the sequences for therapeutic mAbs are being adapted for scFv design, but we will consider all possible factors including biochemical (affinity), the transmission of activation signaling in the CAR-NK cells, compatibilities with the cytoplasmic domain of the CAR, and side effects observed in clinical trials, to obtain the best clinical outcome.

Spacer, transmembrane, and costimulatory domains: Since NK cells express CD3z for transmitting activation signaling similarly to T cells (122), the intracellular domain of CD3z is generally used for CAR designs for NK cells. The first generation of CARs generally contained only the CD3z domain for signal transduction. However, due to CD3z's weak potency, costimulatory domains (e.g., CD28 or 4-1BB, or both) were added to the cytoplasmic region of second/third generation CARs (123). A previous study showed that incorporation of DNAX-activation protein 12 (DAP12) instead of the CD3z activation domain allowed the lysis of KIR/HLA-matched prostate stem cell antigen (PSCA)-positive tumor cell lines and was considered suitable for CAR-NK therapy against solid tumors (124). The CD28 and 4-1BB costimulatory domains of third-generation CARs of iPSC-derived NK cells can be replaced with the combination of an NKG2D transmembrane and 2B4 cytoplasmic domain that can induce synergistic activation of NK cells with enhanced antitumor activity of iPSC-CAR NK cells (104). These studies highlighted that the selection of optimal transmembrane and cytoplasmic signaling domains of a CAR unique to NK cells can improve the antitumor activity of CAR-NK cells in a manner different from CAR-T cells.

In addition, distinct hinge domains alter the performance of CAR-T cell therapies. The extracellular spacer has been reported to affect the accessibility of a CAR to approach the target epitope and decides the cell-cell distance depending on the length (125). Moreover, the selection of an optimal hinge region contributes to CAR dimerization and performance. This may be due to structural interactions as well as the flexibility of the CAR (126, 127). Further, the incorporation of 4-1BB costimulatory domains has been demonstrated to ameliorate the exhaustion of T cells caused by CAR-mediated antigen-independent tonic signaling thereby leading to functional differences (128). The difference between CD28/CD3 ξ and 4-1BB/CD3 ξ is associated with kinetics and signal strength (129).

Fourth-generation CARs: Fourth-generation CAR-T cells are engineered to secrete transgenic cytokines. Cytokine-mediated "signal 3" is known to be important for T or NK cell activation and persistence. These CARs, armed with cytokines, were validated to enhance antitumor activities in several studies on T cells. The persistence of CAR-expressing effector cells is associated with improved clinical outcomes (127, 130). Adoptively transferred NK cells have limited persistence *in vivo*, thereby potentially limiting the efficacy of CAR-NK cells. Genetic engineering of CAR-NK cells to express IL-15, which is associated with increased proliferation and survival of NK cells, enhanced the persistence and efficacy of CAR-NK cells (103, 131). *Ex-vivo* NK cells cultured with IL-15 and a pharmacological inhibitor of the glycogen synthase kinase 3 (GSK3) enhanced CD57 upregulation, promoted late-stage maturation, and improved the antitumor activity of the NK cells (132).

Several interesting studies focusing on signal 3 were conducted in CAR-T cells. Kagoya *et al.* developed CARs with truncated IL-2Rbeta and STAT3 binding YXXQ motif in addition to the CD28-CD3z cytoplasmic domain, thereby resulting in enhanced antitumor activity (133). Chmielewski *et al.* developed TRUCKs (T cells redirected for antigen-unrestricted cytokine-initiated killing), an NFAT-based cytokine expression system to increase the cytotoxicity of CAR cells (134, 135). Hurton *et al.* engineered CARs with IL-15 fused to the IL-15Ralpha via a flexible linker (136). However, these mechanistic studies are based on T cells and the conventional CAR structure. The biological mechanisms behind CAR signaling when using different CAR constructs or different cell types such as NK cells are not well understood. It is hypothesized that the exploration of various CARs and activation pathways based on NK cells improves CAR-NK therapy by determining optimal constructs and methodological complications such as cell sources, *ex vivo* culture, infusion dose, and frequency. Next, we will describe current attempts conducted in industries for CAR-NK anti-cancer therapy employing various new technologies.

CAR-NK DEVELOPMENT IN INDUSTRY: CURRENT STATUS

Here, we summarize the current status of CAR-NK developments in the biotech and pharmaceutical industries (Fig. 2). Bellicum Pharmaceuticals (Houston, TX) is developing the novel technology GoCARTM NK cell therapy by introducing both rimiducid-inducible iMC (MyD88-CD40 dimerization) for NK cell activation and proliferation along with rapamycin-inducible iRC9 (Caspase-9) for safety into NK cells. Anti-CD123 and anti-HER2 CAR-NK cells with iMC and autocrine IL-15 showed enhanced persistence and anti-tumor activity in *in vivo* CD123 + AML and HER2 + solid tumor models, respectively (137). Bellicum has announced that GoCARTM NK cells targeting BCMA are in preclinical development for treating multiple myeloma.

Fate Therapeutics Inc. is developing NKCAR-iPSC-NK cells that target the CD19 antigen, FT596, armed with a high-affi-




Fig. 2. Development of CAR-NK cell therapies in the industry. (A) GoCAR-NK utilizes Rimiducid and Rapamycin for inducible activation and suicide signaling, respectively. (B) FT596, combined with Rituximab, can target both CD19 and CD20 with CD16 Fc receptor in B cell malignancy. IL-15 receptor fusion protein with a flexible IL-15 is introduced for further NK cell activation. (C) ONK therapeutics use TRAIL signaling for maximizing tumor cell apoptosis. A high-affinity variant of TRAIL is recognized by the DR5 TRAIL receptor, thereby resulting in activation of FADD-Caspase8 signaling in a target tumor cell. (D) t-haNK is a versatile platform targeting tumors with a combination of CAR and therapeutic antibody. In this figure, we illustrate an example of NSCLC which expresses a high level of PD-L1 and EGFR. ER-retained IL-2 also drives NK cell activation and persistence. (E) NKX-101 uses NKG2D-CAR for tumor antigen recognition, membrane-bound IL-15 for further activation as well. NKG2D ligands (MIC and RAET1 families) are highly expressed in various tumor cells.

nity, non-cleavable CD16 FcR and a novel IL-15 receptor fusion. The high-affinity CD16 receptor allowed the CAR-NK cells to overcome resistance induced by CD19 antigen loss in combination with rituximab (CD20 therapeutic mAb) in a Raji CD19-CD20+ lymphoma model [ASH 2018, 2019]. FT596 is now under Phase I clinical trial for patients with B cell lymphoma and chronic lymphocytic leukemia [NCT04245722].

Avectas and ONK therapeutics are developing a CAR-NK cell therapy by incorporating DR5 TRAIL variants to maximize cytotoxicity in various tumors including CD19 targeting B-cell lymphomas (patent US 10,034,925). The DR5 TRAIL variant showed a maximum of 1,000-fold or greater binding affinity compared to other variants, thereby resulting in TRAIL receptor-mediated apoptosis in target tumor cells.

GEMoAb Monoclonals GmbH is developing a universal CAR

platform (Uni-CAR) which uses a switchable turn-on/off mechanism by binding with cancer-specific targeting modules (TM) (138). Uni-CAR NK-92 cells are redirected and activated by scFv- and IgG4-based TM specific for tumor antigen GD2, thereby resulting in anti-tumor activity in GD2-expressing solid tumor mouse models. *In vivo* pharmacokinetic analysis showed rapid elimination of scFv-based TM with a half-life of 1.6 h.

NantKwest Inc. is developing t-haNK cells that target the PD-L1 in non-small cell lung cancer. The t-haNK cells express an anti-PD-L1 CAR, a high-affinity CD16, and an endoplasmic reticulum retained IL-2 (139). T-haNK is now under Phase I clinical trial for patients with locally advanced or metastatic solid tumors (NCT04050709).

Nkarta Therapeutics Inc. is developing NKX-101: CAR-NK cells consisting of NKG2D receptors in the extracellular domain,

Table 2. Clinical trials with CAR-NK cells

Clinical trials.gov identifier	Title or CAR-NK strategy	Target disease	Source of NK cell	Sponsor	Start date	Status
NCT03692767	Anti-CD22 CAR NK	Relapsed and refractory B cell lymphoma		Allife Medical Science and Technology Co., Ltd.	March 2019	Not yet recruiting
NCT03690310	Anti-CD19 CAR NK	Relapsed and refractory B cell lymphoma		Allife Medical Science and Technology Co., Ltd.	March 2019	Not yet recruiting
NCT03692637	Anti-mesothelin CAR NK	Epithelial ovarian cancer		Allife Medical Science and Technology Co., Ltd.	March 2019	Not yet recruiting
NCT03415100	NKG2D-ligand targeted CAR-NK	Metastatic solid tumors		The Third Affiliated Hospital of Guangzhou Medical University	January 2, 2018	Unknown
NCT04324996	NKG2D-ACE2 CAR-NK	COVID-19	Cord blood	Chongqing Public Health Medical Center	February 21, 2020	Recruiting
NCT03692663	Anti-PSMA CAR NK	Castration-resistant prostate cancer		Allife Medical Science and Technology Co., Ltd.	December 2018	Not yet recruiting
NCT03940820	ROBO1 specific CAR-NK	Solid tumors		Asclepius Technology Company Group (Suzhou) Co., Ltd.	May 2019	Recruiting
NCT03940833	BCMA CAR-NK 92 cells	Relapse/refractory multiple myeloma	NK-92 cell line	Asclepius Technology Company Group (Suzhou) Co., Ltd.	May 2019	Recruiting
NCT03824964	Anti-CD19/CD22 CAR NK	Relapsed and refractory B cell lymphoma		Allife Medical Science and Technology Co., Ltd.	February 1, 2019	Not yet recruiting
NCT02944162	Anti-CD33 CAR-NK	Relapsed/refractory CD33+ AML	NK-92 cell line	PersonGen BioTherapeutics (Suzhou) Co., Ltd.	October 2016	Unknown
NCT02892695	PCAR-119 bridge immunotherapy before stem cell transplant (anti-CD19 CAR-NK)	CD19 positive leukemia and lymphoma	NK-92 cell line	PersonGen BioTherapeutics (Suzhou) Co., Ltd.	September 2016	Unknown
NCT03941457	ROBO1 specific BiCAR-NK	Pancreatic cancer		Asclepius Technology Company Group (Suzhou) Co., Ltd.	May 2019	Recruiting
NCT03931720	ROBO1 Specific BiCAR-NK/T	Malignant tumor		Asclepius Technology Company Group (Suzhou) Co., Ltd.	May 2019	Recruiting
NCT03056339	Umbilical & cord blood (CB) derived CAR-engineered NK; (iC9/CAR.19/IL15-transduced CB-NK)	B lymphoid malignancies	Cord blood	M.D. Anderson Cancer Center	June 21, 2017	Recruiting
NCT04245722	FT596 as a monotherapy and in combination with anti-CD20 monoclonal antibodies	B cell lymphoma, chronic lymphocytic leukemia	iPSC	Fate Therapeutics	March 19, 2020	Recruiting
NCT04050709	QUILT 3.064: PD-L1 t-haNK	Locally advanced or metastatic solid cancers	NK-92 cell line	NantKwest, Inc.	July 18, 2019	Active, not recruiting
NCT03383978	Intracranial injection of NK-92/5.28.z (CAR2BRAIN)	Recurrent HER2-positive glioblastoma	NK-92 cell line	Johann Wolfgang Goethe University Hospital	December 1, 2017	Recruiting

OX40-CD3z in the costimulatory domain, and membrane-bound IL-15. The NKG2D receptor binds to eight NKG2D ligands that are upregulated in a range of leukemic and solid tumors. CAR-NK cells targeting these ligands showed antitumor activity in a murine model of osteosarcoma (140). NKX-101 showed *in vitro* anti-tumor activity and increased cell delivery in an *in vivo* model of colorectal cancer liver metastasis (ASCO 2020).

Takeda, under license from MD Anderson Cancer Center, is developing TAK-007: cord blood-NK cells that are transduced to express CAR targeting CD19 with a CD28 costimulatory domain, IL-15, and an inducible caspase 9 suicide gene. In a recent small-scale clinical trial, TAK-007 showed a response rate of 73% for patients with relapsed or refractory CD19-positive cancers (111). Some additional CAR-NK cell therapies are under development targeting several tumor-associated antigens including MUC1 for solid tumors (PersonGen Biomedicine Co. Ltd), CD38 for AML (Celularity Inc), EGFRvIII for PD-L1 positive solid tumors (PharmAbcine), CD7 for T cell leukemia (Gracell Biotechnology Ltd), GPC-3 for hepatocellular carcinoma (Baylor College of Medicine, Kuur therapeutics), and CSPG4 for triple-negative breast cancer (Baylor College of Medicine). The ongoing clinical trials with NK-CARs are presented in Table 2.

CONCLUSIONS AND PERSPECTIVES

NK cells have the potential to kill a broad spectrum of tumor cells without mutational burden and neoantigen presentation. This property of MHC-unrestricted tumor lysis by NK cells without the risk of GVHD, which is unique among immune effector cells, has positioned NK cells as key components in the arsenal of cancer therapeutics. Recent studies on cancer therapies using NK cells have demonstrated favorable clinical efficacies in hematologic malignancies, but limited success in solid tumors. This may be attributed to the limited capacity of NK cells to infiltrate tumors, persist *in vivo*, or resist in the immunosuppressive tumor microenvironment. Thus, rational strategies for improving NK cell-based therapy have been developed to overcome the existing challenges. These include the blockade of immune checkpoint receptors to rescue dysfunctional NK cells and the incorporation of tumor-directed CAR to augment anti-tumor NK cell specificity and activity.

Since the early studies on KIRs, numerous immune checkpoint receptors revealed to be functionally quintessential in NK cell function. Immune checkpoint blockades that can stimulate NK cells have tremendous potential in cancer therapy. They could not only stimulate NK cells but also T cells in direct and indirect ways, particularly under the situation of blocking immune checkpoint receptors that are expressed on both the cell types. In addition, combination therapies with immune checkpoint blockade and chemotherapy or CAR-T/NK therapy could become a part of standard therapeutic regimens in the near future.

Despite challenges in the genetic manipulation of NK cells,

CAR-NK cells have received increasing attention as next-generation therapeutics against refractory malignancies including solid tumors. In addition to the redirected specificity, they hold promise with “off-the-shelf” clinical utility and low toxicity usually not causing immune-related adverse events. Moreover, further engineering of CAR-NK cells with on-board cytokines (e.g., IL-15) that leads to enhanced *in vivo* persistence and resistance to immunosuppression has paved the way for new therapeutic options to improve clinical efficacy (103, 141). These observations, along with several reliable protocols for GMP-grade large scale expansion of NK cells, will render NK cell-based therapy a viable modality to treat refractory cancers, possibly in rational and optimal combination with other therapies.

ACKNOWLEDGEMENTS

This study was supported by a grant from the National Research Foundation of Korea (2019R1A2C2006475) and an MRC grant (2018R1A5A2020732) funded by the Korean government (MSIT).

CONFLICTS OF INTEREST

The authors have no conflicting interests.

REFERENCES

1. Larkin J, Chiarion-Sileni V, Gonzalez R et al (2015) Combined Nivolumab and Ipilimumab or monotherapy in untreated melanoma. *N Engl J Med* 373, 23-34
2. Ishida Y, Agata Y, Shibahara K and Honjo T (1992) Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J* 11, 3887-3895
3. Selvakumar A, Mohanraj BK, Eddy RL, Shows TB, White PC and Dupont B (1992) Genomic organization and chromosomal location of the human gene encoding the B-lymphocyte activation antigen B7. *Immunogenetics* 36, 175-181
4. Karre K, Ljunggren HG, Piontek G and Kiessling R (1986) Selective rejection of H-2-deficient lymphoma variants suggests alternative immune defence strategy. *Nature* 319, 675-678
5. Kim S, Poursine-Laurent J, Truscott SM et al (2005) Licensing of natural killer cells by host major histocompatibility complex class I molecules. *Nature* 436, 709-713
6. Kim S, Sunwoo JB, Yang L et al (2008) HLA alleles determine differences in human natural killer cell responsiveness and potency. *Proc Natl Acad Sci U S A* 105, 3053-3058
7. Djaoud Z and Parham P (2020) HLAs, TCRs, and KIRs, a triumvirate of human cell-mediated immunity. *Annu Rev Biochem* 89, 717-739
8. Saunders PM, Vivian JP, O'Connor GM et al (2015) A bird's eye view of NK cell receptor interactions with their MHC class I ligands. *Immunol Rev* 267, 148-166

9. Benson DM, Jr., Hofmeister CC, Padmanabhan S et al (2012) A phase 1 trial of the anti-KIR antibody IPH2101 in patients with relapsed/refractory multiple myeloma. *Blood* 120, 4324-4333
10. Vey N, Bourhis JH, Boissel N et al (2012) A phase 1 trial of the anti-inhibitory KIR mAb IPH2101 for AML in complete remission. *Blood* 120, 4317-4323
11. Carlsten M, Korde N, Kotecha R et al (2016) Checkpoint inhibition of KIR2D with the monoclonal antibody IPH2101 induces contraction and hyporesponsiveness of NK cells in patients with myeloma. *Clin Cancer Res* 22, 5211-5222
12. Benson DM Jr, Cohen AD, Jagannath S et al (2015) A phase I trial of the anti-KIR antibody IPH2101 and lenalidomide in patients with relapsed/refractory multiple myeloma. *Clin Cancer Res* 21, 4055-4061
13. Kohrt HE, Thielens A, Marabelle A et al (2014) Anti-KIR antibody enhancement of anti-lymphoma activity of natural killer cells as monotherapy and in combination with anti-CD20 antibodies. *Blood* 123, 678-686
14. Yalniz FF, Daver N, Rezvani K et al (2018) A pilot trial of Lirilumab with or without Azacitidine for patients with myelodysplastic syndrome. *Clin Lymphoma Myeloma Leuk* 18, 658-663 e652
15. Binyamin L, Alpaugh RK, Hughes TL, Lutz CT, Campbell KS and Weiner LM (2008) Blocking NK cell inhibitory self-recognition promotes antibody-dependent cellular cytotoxicity in a model of anti-lymphoma therapy. *J Immunol* 180, 6392-6401
16. Bagot M, Porcu P, Marie-Cardine A et al (2019) IPH4102, a first-in-class anti-KIR3DL2 monoclonal antibody, in patients with relapsed or refractory cutaneous T-cell lymphoma: an international, first-in-human, open-label, phase 1 trial. *Lancet Oncol* 20, 1160-1170
17. Ewen EM, Pahl JHW, Miller M, Watzl C and Cerwenka A (2018) KIR downregulation by IL-12/15/18 unleashes human NK cells from KIR/HLA-I inhibition and enhances killing of tumor cells. *Eur J Immunol* 48, 355-365
18. Shiroishi M, Tsumoto K, Amano K et al (2003) Human inhibitory receptors Ig-like transcript 2 (ILT2) and ILT4 compete with CD8 for MHC class I binding and bind preferentially to HLA-G. *Proc Natl Acad Sci U S A* 100, 8856-8861
19. Khan M, Arooj S and Wang H (2020) NK cell-based immune checkpoint inhibition. *Front Immunol* 11, 167
20. Scarabel L, Garziera M, Fortuna S, Asaro F, Toffoli G and Geremia S (2020) Soluble HLA-G expression levels and HLA-G/irinotecan association in metastatic colorectal cancer treated with irinotecan-based strategy. *Sci Rep* 10, 8773
21. Bertol BC, de Araujo JNG, Sadissou IA et al (2020) Plasma levels of soluble HLA-G and cytokines in papillary thyroid carcinoma before and after thyroidectomy. *Int J Clin Pract* 74, e13585
22. Heidenreich S, Zu Eulenburg C, Hildebrandt Y et al (2012) Impact of the NK cell receptor LIR-1 (ILT-2/CD85j/LILRB1) on cytotoxicity against multiple myeloma. *Clin Dev Immunol* 2012, 652130
23. Godal R, Bachanova V, Gleason M et al (2010) Natural killer cell killing of acute myelogenous leukemia and acute lymphoblastic leukemia blasts by killer cell immunoglobulin-like receptor-negative natural killer cells after NKG2A and LIR-1 blockade. *Biol Blood Marrow Transplant* 16, 612-621
24. Ho GT, Celik AA, Huyton T et al (2020) NKG2A/CD94 is a new immune receptor for HLA-G and distinguishes amino acid differences in the HLA-G heavy chain. *Int J Mol Sci* 21, 4362
25. Burshtyn DN, Scharenberg AM, Wagtmann N et al (1996) Recruitment of tyrosine phosphatase HCP by the killer cell inhibitor receptor. *Immunity* 4, 77-85
26. Olcese L, Lang P, Vely F et al (1996) Human and mouse killer-cell inhibitory receptors recruit PTP1C and PTP1D protein tyrosine phosphatases. *J Immunol* 156, 4531-4534
27. Stebbins CC, Watzl C, Billadeau DD, Leibson PJ, Burshtyn DN and Long EO (2003) Vav1 dephosphorylation by the tyrosine phosphatase SHP-1 as a mechanism for inhibition of cellular cytotoxicity. *Mol Cell Biol* 23, 6291-6299
28. Liu D, Peterson ME and Long EO (2012) The adaptor protein Crk controls activation and inhibition of natural killer cells. *Immunity* 36, 600-611
29. Long EO, Sik Kim H, Liu D, Peterson ME and Rajagopalan S (2013) Controlling natural killer cell responses: integration of signals for activation and inhibition. *Annu Rev Immunol* 31, 227-258
30. Levy EM, Bianchini M, Von Euw EM et al (2008) Human leukocyte antigen-E protein is overexpressed in primary human colorectal cancer. *Int J Oncol* 32, 633-641
31. Gooden M, Lampen M, Jordanova ES et al (2011) HLA-E expression by gynecological cancers restrains tumor-infiltrating CD8(+) T lymphocytes. *Proc Natl Acad Sci U S A* 108, 10656-10661
32. Della Chiesa M, Pesce S, Muccio L et al (2016) Features of memory-like and PD-1(+) human NK cell subsets. *Front Immunol* 7, 351
33. McWilliams EM, Mele JM, Cheney C et al (2016) Therapeutic CD94/NKG2A blockade improves natural killer cell dysfunction in chronic lymphocytic leukemia. *Oncoimmunology* 5, e1226720
34. Chirossone L, Vienne M, Kerdiles YM and Vivier E (2017) Natural killer cell immunotherapies against cancer: checkpoint inhibitors and more. *Semin Immunol* 31, 55-63
35. Burugu S, Dancsok AR and Nielsen TO (2018) Emerging targets in cancer immunotherapy. *Semin Cancer Biol* 52, 39-52
36. Andre P, Denis C, Soulas C et al (2018) Anti-NKG2A mAb is a checkpoint inhibitor that promotes anti-tumor immunity by unleashing both T and NK cells. *Cell* 175, 1731-1743
37. van Montfoort N, Borst L, Korner MJ et al (2018) NKG2A blockade potentiates CD8 T cell immunity induced by cancer vaccines. *Cell* 175, 1744-1755
38. Schildberg FA, Klein SR, Freeman GJ and Sharpe AH (2016) Coinhibitory pathways in the B7-CD28 ligand-receptor family. *Immunity* 44, 955-972
39. Stojanovic A, Fiegler N, Brunner-Weinzierl M and Cerwenka A (2014) CTLA-4 is expressed by activated mouse NK cells and inhibits NK Cell IFN-gamma production in response to mature dendritic cells. *J Immunol* 192, 4184-4191

40. Jie HB, Schuler PJ, Lee SC et al (2015) CTLA-4(+) regulatory T cells increased in Cetuximab-treated head and neck cancer patients suppress NK cell cytotoxicity and correlate with poor prognosis. *Cancer Res* 75, 2200-2210
41. Romano E, Kusio-Kobialka M, Foukas PG et al (2015) Ipilimumab-dependent cell-mediated cytotoxicity of regulatory T cells ex vivo by nonclassical monocytes in melanoma patients. *Proc Natl Acad Sci U S A* 112, 6140-6145
42. Simpson TR, Li F, Montalvo-Ortiz W et al (2013) Fc-dependent depletion of tumor-infiltrating regulatory T cells co-defines the efficacy of anti-CTLA-4 therapy against melanoma. *J Exp Med* 210, 1695-1710
43. Tallero R, Cristiani CM, Staaf E et al (2017) IL-15, TIM-3 and NK cells subsets predict responsiveness to anti-CTLA-4 treatment in melanoma patients. *Oncoimmunology* 6, e1261242
44. Jallad MN, Jurjus AR, Rahal EA and Abdelnoor AM (2020) Triple immunotherapy overcomes immune evasion by tumor in a melanoma mouse model. *Front Oncol* 10, 839
45. Lang S, Vujanovic NL, Wollenberg B and Whiteside TL (1998) Absence of B7.1-CD28/CTLA-4-mediated co-stimulation in human NK cells. *Eur J Immunol* 28, 780-786
46. Cook CH, Chen L, Wen J et al (2009) CD28/B7-mediated co-stimulation is critical for early control of murine cytomegalovirus infection. *Viral Immunol* 22, 91-103
47. Boussiotis VA, Chatterjee P and Li L (2014) Biochemical signaling of PD-1 on T cells and its functional implications. *Cancer J* 20, 265-271
48. Pesce S, Greppi M, Tabellini G et al (2017) Identification of a subset of human natural killer cells expressing high levels of programmed death 1: a phenotypic and functional characterization. *J Allergy Clin Immunol* 139, 335-346 e333
49. Tabellini G, Benassi M, Marcenaro E et al (2014) Primitive neuroectodermal tumor in an ovarian cystic teratoma: natural killer and neuroblastoma cell analysis. *Case Rep Oncol* 7, 70-78
50. Beldi-Ferchiou A, Lambert M, Dogniaux S et al (2016) PD-1 mediates functional exhaustion of activated NK cells in patients with Kaposi sarcoma. *Oncotarget* 7, 72961-72977
51. Benson DM Jr, Bakan CE, Mishra A et al (2010) The PD-1/PD-L1 axis modulates the natural killer cell versus multiple myeloma effect: a therapeutic target for CT-011, a novel monoclonal anti-PD-1 antibody. *Blood* 116, 2286-2294
52. Vari F, Arpon D, Keane C et al (2018) Immune evasion via PD-1/PD-L1 on NK cells and monocyte/macrophages is more prominent in Hodgkin lymphoma than DLBCL. *Blood* 131, 1809-1819
53. Hsu J, Hodgins JJ, Marathe M et al (2018) Contribution of NK cells to immunotherapy mediated by PD-1/PD-L1 blockade. *J Clin Invest* 128, 4654-4668
54. Lanuza PM, Viguera A, Oliván S et al (2018) Activated human primary NK cells efficiently kill colorectal cancer cells in 3D spheroid cultures irrespectively of the level of PD-L1 expression. *Oncoimmunology* 7, e1395123
55. Hwang S, Han J, Baek JS et al (2019) Cytotoxicity of human hepatic intrasinusoidal CD56(bright) natural killer cells against hepatocellular carcinoma cells. *Int J Mol Sci* 20, 1564
56. Judge SJ, Dunai C, Aguilar EG et al (2020) Minimal PD-1 expression in mouse and human NK cells under diverse conditions. *J Clin Invest* 130, 3051-3068
57. Quatrini L, Wieduwild E, Escaliere B et al (2018) Endogenous glucocorticoids control host resistance to viral infection through the tissue-specific regulation of PD-1 expression on NK cells. *Nat Immunol* 19, 954-962
58. Anderson AC, Joller N and Kuchroo VK (2016) Lag-3, Tim-3, and TIGIT: co-inhibitory receptors with specialized functions in immune regulation. *Immunity* 44, 989-1004
59. Rangachari M, Zhu C, Sakuishi K et al (2012) Bat3 promotes T cell responses and autoimmunity by repressing Tim-3-mediated cell death and exhaustion. *Nat Med* 18, 1394-1400
60. Carotta S (2016) Targeting NK cells for anticancer immunotherapy: clinical and preclinical approaches. *Front Immunol* 7, 152
61. Wang Z, Zhu J, Gu H et al (2015) The clinical significance of abnormal Tim-3 expression on NK cells from patients with gastric cancer. *Immunol Invest* 44, 578-589
62. Xu L, Huang Y, Tan L et al (2015) Increased Tim-3 expression in peripheral NK cells predicts a poorer prognosis and Tim-3 blockade improves NK cell-mediated cytotoxicity in human lung adenocarcinoma. *Int Immunopharmacol* 29, 635-641
63. Komita H, Koido S, Hayashi K et al (2015) Expression of immune checkpoint molecules of T cell immunoglobulin and mucin protein 3/galectin-9 for NK cell suppression in human gastrointestinal stromal tumors. *Oncol Rep* 34, 2099-2105
64. Gallois A, Silva I, Osman I and Bhardwaj N (2014) Reversal of natural killer cell exhaustion by TIM-3 blockade. *Oncoimmunology* 3, e946365
65. Zhang C and Liu Y (2020) Targeting NK cell checkpoint receptors or molecules for cancer immunotherapy. *Front Immunol* 11, 1295
66. Chauhan SKS, Koehl U and Kloess S (2020) Harnessing NK cell checkpoint-modulating immunotherapies. *Cancers (Basel)* 12, 1807
67. Van Audenaerde JRM, De Waele J, Marcq E et al (2017) Interleukin-15 stimulates natural killer cell-mediated killing of both human pancreatic cancer and stellate cells. *Oncotarget* 8, 56968-56979
68. Folgiero V, Cifaldi L, Li Pira G, Goffredo BM, Vinti L and Locatelli F (2015) TIM-3/Gal-9 interaction induces IFN gamma-dependent IDO1 expression in acute myeloid leukemia blast cells. *J Hematol Oncol* 8, 36
69. Markel G, Gruda R, Achdout H et al (2004) The critical role of residues 43R and 44Q of carcinoembryonic antigen cell adhesion molecules-1 in the protection from killing by human NK cells. *J Immunol* 173, 3732-3739
70. Fantini M, David JM, Annunziata CM, Morelli MP, Arlen PM and Tsang KY (2020) The monoclonal antibody NEO-201 enhances natural killer cell cytotoxicity against tumor cells through blockade of the inhibitory CEACAM5/

- CEACAM1 immune checkpoint pathway. *Cancer Biother Radiopharm* 35, 190-198
71. Zeligs KP, Morelli MP, David JM et al (2020) Evaluation of the anti-tumor activity of the humanized monoclonal antibody NEO-201 in preclinical models of ovarian cancer. *Front Oncol* 10, 805
 72. Jacques A, Bleau C, Turbide C, Beauchemin N and Lamontagne L (2009) A synergistic interferon-gamma production is induced by mouse hepatitis virus in interleukin-12 (IL-12)/IL-18-activated natural killer cells and modulated by carcinoembryonic antigen-related cell adhesion molecules (CEACAM) 1a receptor. *Immunology* 128, e551-561
 73. Hosomi S, Chen Z, Baker K et al (2013) CEACAM1 on activated NK cells inhibits NKG2D-mediated cytolytic function and signaling. *Eur J Immunol* 43, 2473-2483
 74. Kim WM, Huang YH, Gandhi A and Blumberg RS (2019) CEACAM1 structure and function in immunity and its therapeutic implications. *Semin Immunol* 42, 101296
 75. Zhang Q, Bi J, Zheng X et al (2018) Blockade of the checkpoint receptor TIGIT prevents NK cell exhaustion and elicits potent anti-tumor immunity. *Nat Immunol* 19, 723-732
 76. Johnston RJ, Comps-Agrar L, Hackney J et al (2014) The immunoreceptor TIGIT regulates antitumor and antiviral CD8(+) T cell effector function. *Cancer Cell* 26, 923-937
 77. Gorvel L and Olive D (2020) Targeting the "PVR-TIGIT axis" with immune checkpoint therapies. *F1000Res* 9, 354
 78. Wang J, Sanmamed MF, Datar I et al (2019) Fibrinogen-like protein 1 is a major immune inhibitory ligand of LAG-3. *Cell* 176, 334-347 e312
 79. Maeda TK, Sugiura D, Okazaki IM, Maruhashi T and Okazaki T (2019) Atypical motifs in the cytoplasmic region of the inhibitory immune co-receptor LAG-3 inhibit T cell activation. *J Biol Chem* 294, 6017-6026
 80. Merino A, Zhang B, Dougherty P et al (2019) Chronic stimulation drives human NK cell dysfunction and epigenetic reprogramming. *J Clin Invest* 129, 3770-3785
 81. Brignone C, Grygar C, Marcu M, Schakel K and Triebel F (2007) A soluble form of lymphocyte activation gene-3 (IMP321) induces activation of a large range of human effector cytotoxic cells. *J Immunol* 179, 4202-4211
 82. Brignone C, Escudier B, Grygar C, Marcu M and Triebel F (2009) A phase I pharmacokinetic and biological correlative study of IMP321, a novel MHC class II agonist, in patients with advanced renal cell carcinoma. *Clin Cancer Res* 15, 6225-6231
 83. Brignone C, Gutierrez M, Mefti F et al (2010) First-line chemoimmunotherapy in metastatic breast carcinoma: combination of paclitaxel and IMP321 (LAG-3lg) enhances immune responses and antitumor activity. *J Transl Med* 8, 71
 84. Nath PR, Gangaplara A, Pal-Nath D et al (2018) CD47 expression in natural killer cells regulates homeostasis and modulates immune response to Lymphocytic choriomeningitis virus. *Front Immunol* 9, 2985
 85. Nath PR, Pal-Nath D, Mandal A, Cam MC, Schwartz AL and Roberts DD (2019) Natural killer cell recruitment and activation are regulated by CD47 expression in the tumor microenvironment. *Cancer Immunol Res* 7, 1547-1561
 86. Valipour B, Abedelahi A, Naderali E et al (2020) Cord blood stem cell derived CD16(+) NK cells eradicated acute lymphoblastic leukemia cells using with anti-CD47 antibody. *Life Sci* 242, 117223
 87. Tsao LC, Crosby EJ, Trotter TN et al (2019) CD47 blockade augmentation of trastuzumab antitumor efficacy dependent on antibody-dependent cellular phagocytosis. *JCI Insight* 4, e131882
 88. Allard B, Longhi MS, Robson SC and Stagg J (2017) The ectonucleotidases CD39 and CD73: novel checkpoint inhibitor targets. *Immunol Rev* 276, 121-144
 89. Neo SY, Yang Y, Record J et al (2020) CD73 immune checkpoint defines regulatory NK cells within the tumor microenvironment. *J Clin Invest* 130, 1185-1198
 90. Wang J and Matosevic S (2019) NT5E/CD73 as correlative factor of patient survival and natural killer cell infiltration in glioblastoma. *J Clin Med* 8, 1526
 91. Chambers AM and Matosevic S (2019) Immunometabolic dysfunction of natural killer cells mediated by the Hypoxia-CD73 axis in solid tumors. *Front Mol Biosci* 6, 60
 92. Zheng Y, Ma X, Su D et al (2020) The roles of Siglec7 and Siglec9 on natural killer cells in virus infection and tumour progression. *J Immunol Res* 2020, 6243819
 93. Hudak JE, Canham SM and Bertozzi CR (2014) Glycocalyx engineering reveals a Siglec-based mechanism for NK cell immunoevasion. *Nat Chem Biol* 10, 69-75
 94. Jandus C, Boligan KF, Chijioko O et al (2014) Interactions between Siglec-7/9 receptors and ligands influence NK cell-dependent tumor immunosurveillance. *J Clin Invest* 124, 1810-1820
 95. Hernandez-Caselles T, Miguel RC, Ruiz-Alcaraz AJ and Garcia-Penarrubia P (2019) CD33 (Siglec-3) Inhibitory function: role in the NKG2D/DAP10 activating pathway. *J Immunol Res* 2019, 6032141
 96. Kohl U, Arsenieva S, Holzinger A and Abken H (2018) CAR T cells in trials: recent achievements and challenges that remain in the production of modified T cells for clinical applications. *Hum Gene Ther* 29, 559-568
 97. Maude SL, Laetsch TW, Buechner J et al (2018) Tisagenlecleucel in children and young adults with B-cell lymphoblastic leukemia. *N Engl J Med* 378, 439-448
 98. Ren J, Liu X, Fang C, Jiang S, June CH and Zhao Y (2017) Multiplex genome editing to generate universal CAR T cells resistant to PD1 inhibition. *Clin Cancer Res* 23, 2255-2266
 99. Georgiadis C, Preece R, Nickolay L et al (2018) Long terminal repeat CRISPR-CAR-coupled "universal" T cells mediate potent anti-leukemic effects. *Mol Ther* 26, 1215-1227
 100. Liu X, Zhang Y, Cheng C et al (2017) CRISPR-Cas9-mediated multiplex gene editing in CAR-T cells. *Cell Res* 27, 154-157
 101. Shifrin N, Raulet DH and Ardolino M (2014) NK cell self tolerance, responsiveness and missing self recognition. *Semin Immunol* 26, 138-144
 102. Cheng M, Chen YY, Xiao WH, Sun R and Tian ZG (2013) NK cell-based immunotherapy for malignant

- diseases. *Cell Mol Immunol* 10, 230-252
103. Liu E, Tong Y, Dotti G et al (2018) Cord blood NK cells engineered to express IL-15 and a CD19-targeted CAR show long-term persistence and potent antitumor activity. *Leukemia* 32, 520-531
 104. Li Y, Hermanson DL, Moriarity BS and Kaufman DS (2018) Human iPSC-derived natural killer cells engineered with chimeric antigen receptors enhance anti-tumor activity. *Cell Stem Cell* 23, 181-192 e185
 105. Rezvani K, Rouce R, Liu E and Shpall E (2017) Engineering natural killer cells for cancer immunotherapy. *Mol Ther* 25, 1769-1781
 106. Suck G, Odendahl M, Nowakowska P et al (2016) NK-92: an 'off-the-shelf therapeutic' for adoptive natural killer cell-based cancer immunotherapy. *Cancer Immunol Immunother* 65, 485-492
 107. Karschnia P, Jordan JT, Forst DA et al (2019) Clinical presentation, management, and biomarkers of neurotoxicity after adoptive immunotherapy with CAR T cells. *Blood* 133, 2212-2221
 108. Veluchamy JP, Kok N, van der Vliet HJ, Verheul HMW, de Gruijl TD and Spanholtz J (2017) The rise of allogeneic natural killer cells as a platform for cancer immunotherapy: recent innovations and future developments. *Front Immunol* 8, 631
 109. Moretta A (2002) Natural killer cells and dendritic cells: rendezvous in abused tissues. *Nat Rev Immunol* 2, 957-964
 110. Wang WX, Jiang JT and Wu CP (2020) CAR-NK for tumor immunotherapy: clinical transformation and future prospects. *Cancer Letters* 472, 175-180
 111. Liu E, Marin D, Banerjee P et al (2020) Use of CAR-transduced natural killer cells in CD19-positive lymphoid tumors. *N Engl J Med* 382, 545-553
 112. Ruella M, Barrett DM, Kenderian SS et al (2016) Dual CD19 and CD123 targeting prevents antigen-loss relapses after CD19-directed immunotherapies. *J Clin Invest* 126, 3814-3826
 113. Orlando EJ, Han X, Tribouley C et al (2018) Genetic mechanisms of target antigen loss in CAR19 therapy of acute lymphoblastic leukemia. *Nat Med* 24, 1504-1506
 114. Sayitoglu EC, Georgoudaki AM, Chrobok M et al (2020) Boosting natural killer cell-mediated targeting of sarcoma through DNAM-1 and NKG2D. *Front Immunol* 11, 40
 115. Barrow AD, Martin CJ and Colonna M (2019) The natural cytotoxicity receptors in health and disease. *Front Immunol* 10, 909
 116. Zhang C, Oberoi P, Oelsner S et al (2017) Chimeric antigen receptor-engineered NK-92 cells: an off-the-shelf cellular therapeutic for targeted elimination of cancer cells and induction of protective antitumor immunity. *Front Immunol* 8, 533
 117. Dotti G, Gottschalk S, Savoldo B and Brenner MK (2014) Design and development of therapies using chimeric antigen receptor-expressing T cells. *Immunol Rev* 257, 107-126
 118. Sadelain M, Brentjens R and Riviere I (2013) The basic principles of chimeric antigen receptor design. *Cancer Discov* 3, 388-398
 119. Chmielewski M, Hombach A, Heuser C, Adams GP and Abken H (2004) T cell activation by antibody-like immunoreceptors: Increase in affinity of the single-chain fragment domain above threshold does not increase T cell activation against antigen-positive target cells but decreases selectivity. *J Immunol* 173, 7647-7653
 120. Morgan RA, Yang JC, Kitano M, Dudley ME, Laurencot CM and Rosenberg SA (2010) Case report of a serious adverse event following the administration of T cells transduced with a chimeric antigen receptor recognizing ERBB2. *Mol Ther* 18, 843-851
 121. Burger MC, Zhang CC, Harter PN et al (2019) CAR-engineered NK cells for the treatment of glioblastoma: turning innate effectors into precision tools for cancer immunotherapy. *Front Immunol* 10, 2683
 122. Anderson P, Caligiuri M, Ritz J and Schlossman SF (1989) Cd3-negative natural-killer cells express Zeta-Tcr as part of a novel molecular-complex. *Nature* 341, 159-162
 123. Kofler DM, Chmielewski M, Rapp G et al (2011) CD28 costimulation impairs the efficacy of a redirected T-cell antitumor attack in the presence of regulatory T cells which can be overcome by preventing Lck activation. *Mol Ther* 19, 760-767
 124. Topfer K, Cartellieri M, Michen S et al (2015) DAP12-based activating chimeric antigen receptor for NK cell tumor immunotherapy. *J Immunol* 194, 3201-3212
 125. Chang ZL and Chen YY (2017) CARs: synthetic immunoreceptors for cancer therapy and beyond. *Trends Mol Med* 23, 430-450
 126. Alabanza L, Pegues M, Geldres C et al (2017) Function of novel anti-CD19 chimeric antigen receptors with human variable regions is affected by hinge and transmembrane domains. *Mol Ther* 25, 2452-2465
 127. Brentjens RJ, Riviere I, Park JH et al (2011) Safety and persistence of adoptively transferred autologous CD19-targeted T cells in patients with relapsed or chemotherapy refractory B-cell leukemias. *Blood* 118, 4817-4828
 128. Long AH, Haso WM, Shern JF et al (2015) 4-1BB costimulation ameliorates T cell exhaustion induced by tonic signaling of chimeric antigen receptors. *Nat Med* 21, 581-590
 129. Salter AI, Ivey RG, Kennedy JJ et al (2018) Phosphoproteomic analysis of chimeric antigen receptor signaling reveals kinetic and quantitative differences that affect cell function. *Sci Signal* 11, eaat6753
 130. Kershaw MH, Westwood JA, Parker LL et al (2006) A phase I study on adoptive immunotherapy using gene-modified T cells for ovarian cancer. *Clin Cancer Res* 12, 6106-6115
 131. Sahm C, Schonfeld K and Wels WS (2012) Expression of IL-15 in NK cells results in rapid enrichment and selective cytotoxicity of gene-modified effectors that carry a tumor-specific antigen receptor. *Cancer Immunol Immunother* 61, 1451-1461
 132. Cichocki F, Valamehr B, Bjordahl R et al (2017) GSK3 inhibition drives maturation of NK cells and enhances their antitumor activity. *Cancer Res* 77, 5664-5675
 133. Kagoya Y, Tanaka S, Guo TX et al (2018) A novel chimeric antigen receptor containing a JAK-STAT signaling domain mediates superior antitumor effects. *Nat Med* 24,

- 352-359
134. Chmielewski M and Abken H (2017) CAR T cells releasing IL-18 convert to T-Bet(high) FoxO1(low) effectors that exhibit augmented activity against advanced solid tumors. *Cell Reports* 21, 3205-3219
 135. Chmielewski M and Abken H (2015) TRUCKs: the fourth generation of CARs. *Expert Opin Biol Ther* 15, 1145-1154
 136. Hurton LV, Singh H, Najjar AM *et al* (2016) Tethered IL-15 augments antitumor activity and promotes a stem-cell memory subset in tumor-specific T cells. *Proc Natl Acad Sci U S A* 113, E7788-E7797
 137. Wang XM, Jasinski DL, Medina JL, Spencer DM, Foster AE and Bayle JH (2020) Inducible MyD88/CD40 synergizes with IL-15 to enhance antitumor efficacy of CAR-NK cells. *Blood Adv* 4, 1950-1964
 138. Mitwasi N, Feldmann A, Arndt C *et al* (2020) "UniCAR"-modified off-the-shelf NK-92 cells for targeting of GD2-expressing tumour cells. *Sci Rep* 10, 2141
 139. Fabian KP, Padgett MR, Donahue RN *et al* (2020) PD-L1 targeting high-affinity NK (t-haNK) cells induce direct antitumor effects and target suppressive MDSC populations. *J Immunother Cancer* 8, e000450
 140. Chang YH, Connolly J, Shimasaki N, Mimura K, Kono K and Campana D (2013) A chimeric receptor with NKG2D specificity enhances natural killer cell activation and killing of tumor cells. *Cancer Res* 73, 1777-1786
 141. Imamura M, Shook D, Kamiya T *et al* (2014) Autonomous growth and increased cytotoxicity of natural killer cells expressing membrane-bound interleukin-15. *Blood* 124, 1081-1088