

## Original article

## Immunolocalization of Protein Kinase C Isoenzymes $\alpha$ , $\beta$ I, $\beta$ II and $\gamma$ in Adult and Developing Rat Kidney

Wan–Young Kim, Ph.D.<sup>1</sup>, Gye–Sil Lee, M.D.<sup>1</sup>, Young–Hee Kim, Ph.D.<sup>1</sup>, Eun–Young Park<sup>1</sup>  
Jin–Sun Hwang<sup>1</sup>, Hyang Kim, M.D.<sup>2</sup>, and Jin Kim, M.D.<sup>1</sup>

<sup>1</sup>Department of Anatomy and MRC for Cell Death Disease Research Center,  
The Catholic University of Korea College of Medicine

<sup>2</sup>Department of Internal Medicine, Sungkyunkwan University, Kangbuk Samsung Hospital, Seoul, Korea

Protein kinase C (PKC) plays an important role not only in signal transduction mechanisms in various biological processes, but also in the regulation of growth and differentiation during development. We studied the classical PKC  $\alpha$ ,  $\beta$ I,  $\beta$ II and  $\gamma$ , with regard to their expression in adult and developing rat kidney. PKC $\alpha$  appeared in the ureteric bud at embryonic day (E) 16, and the proximal and distal anlage at E18. After birth, the immunoreactivity of PKC $\alpha$  gradually decreased. In adult, PKC $\alpha$  was expressed intensely in the connecting tubule (CNT), the collecting ducts (CD) and the renal corpuscle, and weakly in the proximal and distal tubules. PKC $\beta$ I appeared in the ureteric bud at E16, and the proximal anlage at E18. After birth, the immunoreactivity of PKC $\beta$ I gradually disappeared from the CD and proximal tubule. In adult, PKC $\beta$ I was expressed in the intercalated cells of the CNT and cortical CD, the proximal straight tubule, and the renal corpuscle. PKC $\beta$ II appeared in distal anlage at E18, and increased markedly after birth. In the CD, PKC $\beta$ II immunoreactivity appeared after birth. In adult, PKC $\beta$ II was expressed in the distal tubule, the CNT and the CD. The immunoreactivity for PKC $\gamma$  appeared only in the proximal anlage at E18, and increased temporally around the time of birth. However, no immunoreactivity for PKC $\gamma$  was observed in adult rat kidney. These results indicate that classical PKC isoforms appear to play a role in the regulation of various renal functions and differentiation within specific functional units of the uriniferous tubule in rat kidney.

**Key Words:** Protein kinase C, Development, Kidney

### Introduction

Protein kinase C (PKC) is a family of protein kinases that specifically phosphorylate serine/threonine residues. The family includes at least 11 isoforms ( $\alpha$ ,  $\beta$ I,  $\beta$ II,  $\gamma$ ,  $\delta$ ,  $\epsilon$ ,  $\zeta$ ,  $\eta$ ,  $\theta$ ,  $\lambda$ , and  $\mu$ ) in mammalian tissue. These isoforms are divided into three subgroups based on their structure and mode of activation. The first group, the classical or conventional PKCs (cPKCs), including the

isoenzymes  $\alpha$ ,  $\beta$ I,  $\beta$ II and  $\gamma$ , are dependent on activation through diacylglycerol (DAG) and  $\text{Ca}^{2+}$ . The second group, the new or novel PKCs (nPKCs), including  $\delta$ ,  $\epsilon$ ,  $\eta$ ,  $\theta$  and  $\mu$ , are activated by DAG. The third group, the atypical PKCs (aPKCs), including  $\zeta$  and  $\lambda$ , are not activated by DAG or  $\text{Ca}^{2+}$  (1–3).

PKC plays a central role in intracellular signal transduction pathways for hormones, neurotransmitters and growth factors, and significantly contributes to the control of various renal functions, including cellular proliferation, differentiation, exocytosis, and ion and water transport (3–5).

In cultured cells, PKC inhibits activation of  $\text{Na}^+$ ,  $\text{K}^+$ –ATPase (4, 6) and activates the  $\text{Na}^+$ / $\text{H}^+$ –exchanger and

Received November 15, 2007. Accepted November 30, 2007.

Corresponding author: Jin Kim, M.D.

Department of Anatomy and Cell Death Disease Research Center, The Catholic University of Korea College of Medicine, 505 Banpo-dong, Seocho-gu, Seoul, 137-701, Korea  
Tel: +82-2-590-1153, Fax: +82-2-590-2992  
E-mail: jinkim@catholic.ac.kr

$\text{Na}^+/\text{HCO}_3^-$  cotransporter<sup>7, 8)</sup>. According to the findings, PKC may be involved in the modulation of intracellular transporters.

There are several studies showing that various PKC isoforms that are expressed in the rat kidney. Kosaka et al.<sup>9)</sup> and Ono et al.<sup>10)</sup> showed the PKC isoenzymes  $\alpha$ ,  $\beta$ , and  $\zeta$ ; Wetsel et al.<sup>11)</sup>,  $\alpha$ ,  $\beta$ II,  $\delta$ ,  $\epsilon$ , and  $\zeta$ ; Caterina et al.<sup>12)</sup> and Aristimuno and Good<sup>13)</sup>,  $\alpha$ ,  $\beta$ ,  $\delta$ ,  $\epsilon$ , and  $\zeta$ ; Ostlund et al.<sup>14)</sup> and Serlachius et al.<sup>15)</sup>,  $\alpha$ ,  $\delta$ ,  $\epsilon$ , and  $\lambda$ ; and Pfaff et al.<sup>16)</sup>,  $\alpha$ ,  $\beta$ I, and  $\beta$ II. Although most of these studies identified the PKC isoenzymes using molecular biologic approaches, little is known about their localization along the nephron.

Hashimoto et al.<sup>17)</sup> and Hirata et al.<sup>18)</sup> detected PKC in brain tissue and Puceat et al.<sup>19)</sup> and Rybin et al.<sup>20)</sup> detected it in heart tissue. Serlachius et al.<sup>15)</sup> suggested a distinct and differential expression and distribution of PKC isoenzymes depending on embryonal development in the kidney. Moreover, they reported that inhibition of PKC activation enhances apoptosis and induces impairment of nephron formation. These findings support that PKC plays a role in growth and differentiation in development<sup>21-25)</sup>.

To identify the function of PKC in the kidney, we studied the differential expression and localization of the PKC isoenzymes  $\alpha$ ,  $\beta$ I,  $\beta$ II and  $\gamma$  in the developing rat kidney using immunohistochemistry.

## Materials and Methods

### 1. Animals and preservation of kidneys

Male Sprague Dawley rats weighing approximately 250 to 300 g were used in all experiments. Prenatal kidneys were obtained from 16-, 18- and 20-day-old fetuses. Postnatal kidneys were obtained from 1-, 3-, 7-, 14- and 21-day-old pups and adult. The animals were anesthetized with an intraperitoneal injection of urethane (16.5%) and perfused with periodate-lysine-paraformaldehyde (PLP) solution for 3-5 minutes through the abdominal aorta. Kidneys were removed, and cut into 2-mm-thick slices, including the renal papilla. Slices were then immersed in PLP solution for 6-12 hours at 4°C.

Tissues were embedded in wax or EPON 812. For immunohistochemistry using a pre-embedding method, PLP-fixed tissues were cut on a vibratome (Lancer Vibratomes Series 10 00; Technical Products International, St. Louis, MO) to a thickness of 50  $\mu\text{m}$ .

## 2. Immunohistochemistry

### 1) Immunostaining of wax sections

The 50- $\mu\text{m}$ -thick wax sections were dewaxed in xylene and hydrated through an ethanol series and washed for 10 minutes. Sections were incubated with 1.4% methanolic  $\text{H}_2\text{O}_2$  for 30 minutes and with 0.5% Triton X-100 (0.01 M PBS, pH 7.4) for 15 minutes. After rinsing three times in PBS, sections were incubated for 1 hour in PBS containing 10% normal goat serum (Vector Laboratories, Burlington, CA, USA). Sections were immunostained with rabbit polyclonal IgGs (Santa Cruz technology, CA, USA) against PKCs  $\alpha$ ,  $\beta$ I,  $\beta$ II and  $\gamma$  as primary antibodies, using a Vectastain ABC kit (Vector Laboratories) according to the manufacturer's instructions. The sections were incubated overnight in PBS solutions containing antibodies diluted 1:2000 for PKC $\alpha$ , 1:2000 for PKC $\beta$ I, 1:1500 for PKC $\beta$ II and 1:5000 for PKC $\gamma$ . Tissue sections were washed three times for 10 minutes and then incubated for 2 hours at room temperature with PBS containing biotin-conjugated goat anti-rabbit IgG (Vector Laboratories), diluted 1:500, as the secondary antibody. Avidin-biotin-peroxidase complex (Vector Laboratories) diluted 1:100 was used as the tertiary reagent. After the sections were washed three times for 10 minutes with 0.05 M Tris-HCl buffer (pH 7.6), the 0.05% diaminobenzidine and 0.0033%  $\text{H}_2\text{O}_2$  were used as chromogen. After the immunostaining, sections were counterstained with hematoxylin.

### 2) Immunostaining of 1-mm-thick EPON sections

To identify the immunoreactivity of PKC in intercalated cells in adult rat kidney, 1 mm semi-thin sections, embedded in EPON-812, were cut into slices

displaying cortex, outer medulla and inner medulla. The EPON was removed using saturated sodium hydroxide. Antibodies against PKC $\alpha$ ,  $\beta$ I,  $\beta$ II and  $\gamma$  were used. H<sup>+</sup>-ATPase (1:2,000) were used on adjacent sections. Antibodies against aquaporin-1 (AQP-1; 1:2,000) were used to differentiate descending thin limbs of Henle from proximal convoluted tubules. Immunostaining was performed with avidin-biotin-peroxidase complex (ABC), and then the sections were examined after staining with the blue-gray-colored Vector SG (Vector Laboratories).

## Results

### 1. Immunohistochemistry

The PKC isoenzymes  $\alpha$ ,  $\beta$ I and  $\beta$ II, but not  $\gamma$ , were

expressed in the adult rat kidney in the tubules, and the distribution in the tubules was variable (Table 1, 2). PKC isoenzymes  $\alpha$ ,  $\beta$ I,  $\beta$ II and  $\gamma$  were expressed in the developing kidney, and distinct and differential expression patterns were shown during development.

### 2. PKC $\alpha$

In the adult kidney, PKC $\alpha$  immunostaining was generally observed in the entire tubule, but was most strongly observed in the connecting tubules and the cortical collecting ducts. On staining with H<sup>+</sup>-ATPase, PKC $\alpha$ -positive cells were strongly evident in the intercalated cells. Type A intercalated cells were stained in the supranuclear portion of the cytoplasm and type B intercalated cells were stained throughout the entire cytoplasm. The connecting tubule cells and principal

**Table 1. Immunoreactivity of Classical Protein Kinase (PKC) Isoforms in Rat Kidney**


	PKC $\alpha$		PKC $\beta$ I		PKC $\beta$ II		PKC $\gamma$	
RC								
PE	+		$\pm\sim+$		-		-	
P	$\pm$		-		-		-	
MC	++		+		-		-	
PT								
PCT	-		-		+		+	
PST	+		$+\sim++$		-		+	
IT								
DTL	-		++		-		-	
ATL	-		-		-		-	
DT								
TAL	$\pm$		$\pm\sim+$		+++		-	
MD	$\pm$		$\pm\sim+$		+++		-	
DCT	$\pm$		$\pm\sim+$		+++		-	
CNT	+	( $++\sim+++$ )	-	( $++\sim+++$ )	++	(-)	-	(-)
CD								
CCD	+	( $++\sim+++$ )	-	( $+\sim+++$ )	++	(-)	-	(-)
OMCD								
OS	$+\sim++$	( $\pm$ )	-	(-)	+	(-)	-	(-)
IS	$++\sim$	( $\pm$ )	-	(-)	+	(-)	-	(-)
IMCD								
i	+++	(-)	-	(-)	+	(-)	-	(-)
t	++		-		++		-	

Abbreviations : RC, renal corpuscle; PE, parietal epithelium; P, podocytes; MC, mesangial cells; PT, proximal tubules; PCT, proximal convoluted tubules; PST, proximal straight tubules; IT, intermediate tubules; DTL, descending thin limb; ATL, ascending thin limb; DT, distal tubules; TAL, thick ascending limb; MD, macular densa; DCT, distal convoluted tubules; CNT, connecting tubules; CD, collecting ducts; CCD, cortical CD; OMCD, outer medullary CD; OS, outer stripe of the OMCD; IS, inner stripe of the OMCD; IMCD, inner medullary CD; i, initial part of the IMCD; t, terminal part of the IMCD. Symbols designate not detectable (-), faint (+), weak (+), moderate (++) , and high (+++) levels of immunoreactivity. Values in parentheses represent the levels of immunore activity in the intercalated cells.

**Table 2. Immunoreactivity of Conventional Protein Kinase C (PKC) Isoforms in the Intercalated Cells of Rat Kidney**

	PKC $\alpha$			PKC $\beta$ I			PKC $\beta$ II			PKC $\gamma$		
	A-IC	B-IC	PC	A-IC	B-IC	PC	A-IC	B-IC	PC	A-IC	B-IC	PC
CNT	+++	++	+	+++	++	-	-	-	++	-	-	-
CCD	+++	++	+	+++	+	-	-	-	++	-	-	-
OMCDo	+	+	+~+++	-	-	-	-	-	+	-	-	-
OMCDi	-~+		+++	-		-	-		+	-		-
IMCDi	-		+++	-		-	-		+	-		-
IMCDt			++			-			++			-

A-IC, type A intercalated cells; B-IC, type B intercalated cells; PC, principal cells; CNT, connecting tubules; CCD, cortical collecting duct; OMCDo, outer stripe part of outer medullary collecting duct; OMCDi, inner stripe part of OMCD; IMCDi, initial part of the inner medullary collecting duct; IMCDt, terminal part of the IMCD. Symbols designate not detectable (-), faint (+), weak (+), moderate (++), and high (+++) levels of immunoreactivity.




**Fig. 1.** Differential interference contrast (DIC) micrographs of wax sections illustrating immunostaining for protein kinase C  $\alpha$  (PKC $\alpha$ ) in the cortical labyrinth (A), medullary ray (B), initial part of the inner medulla (C), and the terminal part of the inner medulla (D) of adult rat kidneys. PKC $\alpha$  immunostaining is observed in the cytoplasm of mesangial cells (arrows) and the parietal epithelium (arrowhead) of renal corpuscles, intercalated cells (open arrows) of the connecting tubule (CNT) and cortical collecting duct (CCD), and on the basolateral plasma membrane of the CNT cells and the principal cells of the collecting duct. Note the PKC $\alpha$ -negative intercalated cells (open arrowheads) in the initial part of the inner medullary collecting duct (IMCDi). IMCDt, terminal part of the IMCD. Magnifications: A-D,  $\times 400$ .

cells were stained weakly at the basolateral plasma membrane. In the medullary collecting ducts, the intercalated cells were PKC $\alpha$  negative and the principal cells


were PKC $\alpha$  positive at the basolateral plasma membrane. The inner stripe of the outer medulla and the initial part of the inner medulla showed strong immu-

noreactivity. In the renal corpuscle, mesangial cells showed moderate immunoreactivity, and we observed weak immunoreactivity in parietal epithelial cells and podocytes. In the proximal tubule, the convoluted part showed weak immunoreactivity on the microvilli and

faint immunoreactivity was observed in the straight portion. The cytoplasm of distal tubule cells showed faint immunoreactivity. We did not observe any PKC $\alpha$  immunoreactivity in the descending or ascending limbs of the Loop of Henle (Fig. 1; Table 1, 2).



**Fig. 2.** DIC micrographs of wax sections illustrating immunostaining for PKC $\alpha$  in 16- (A), 18- (C & F), and 20-day-old (G) fetal kidneys, and 3- (D & H) and 7-day-old (B, E & I) pups. protein kinase C  $\alpha$  (PKC $\alpha$ ) appeared in the ureteric buds (UB) at 16 days of gestation (A) and in the proximal and distal anlage (stars) at 18 days of gestation (C). B. Note that the PKC $\alpha$ -positive tubular profiles (arrows), which are newly formed proximal and distal tubules, are located only in the subcapsular region in 7-day-old pups, whereas PKC $\alpha$  immunoreactivity is decreased in the mature tubules such as the proximal convoluted tubule (PCT) and connecting tubule (CNT) cells located in the inner cortex. C-E, Note the PKC $\alpha$  immunoreactivity in the mesangial cells (arrows) and parietal epithelium (arrowheads) of the developing renal corpuscle. F-G, Immunoreactivity for PKC $\alpha$  in the basolateral plasma membrane of inner medullary collecting duct (IMCD) cells gradually decreased during development. Note the disappearance of apical PKC expression in the intercalated cells (arrows) of the medullary collecting duct (MCD) after birth. Magnifications: A,  $\times 200$ ; B,  $\times 200$ ; C-I,  $\times 528$ .




**Fig. 3.** DIC micrographs of wax sections illustrating protein kinase C  $\beta$  1 (PKC $\beta$ 1) immunostaining in the cortical labyrinth (A), medullary rays (B), outer stripe of the outer medulla (C), inner stripe of the outer medulla (D), and initial part of the inner medulla (E) of the adult rat kidney. PKC $\beta$ 1 immunoreactivity is strong in the intercalated cells (open arrows) of the connecting tubules (CNT) and cortical collecting ducts (CCD), moderate in the proximal straight tubules (PST), and weak in the mesangial cells (arrows) and parietal epithelium of the renal corpuscle. In the inner medulla, PKC $\beta$ 1 immunostaining is observed only in the descending thin limb (stars) of the Loop of Henle. There is no immunoreactivity in the CNT cells, principal cells of the CCD, outer (OMCDo) and inner stripe of the outer medullary collecting duct (OMCDi), and initial inner medullary collecting duct (IMCDi). Note that there is no immunoreactivity in the intercalated cells (open arrowheads) of the OMCD and IMCDi. Magnifications: A–E,  $\times 400$ .

In the developing kidney, PKC $\alpha$  appeared in the ureteric bud at 16 days of gestation, but there was no staining of the renal vesicle and S-shaped body (Fig. 2A). The PKC $\alpha$  immunoreactivity of the collecting tubule gradually decreased during development and showed a mature pattern from 14 days after birth (Fig. 2F–I). PKC $\alpha$  appeared strongly in the mesangial and parietal cells of the developing renal corpuscle in stage III, proximal anlage and distal anlage of the 18-day-old pups, whereas immunoreactivity for PKC $\alpha$  gradually decreased in mature proximal convoluted and distal convoluted tubules (Fig. 2A–E). In the intercalated cells, immunoreactivity was shown in the connecting and

collecting tubules of 18-day-old pups.

### 3. PKC $\beta$ 1

In the adult kidney, there was strong positive PKC $\beta$ 1 staining in the connecting segment and intercalated cells of the cortical collecting duct. Similar to PKC $\alpha$ , type A intercalated cells were positive in the supranuclear area and type B cells were positive throughout the entire cytoplasm. PKC $\beta$ 1 staining was negative in principal cells. In the proximal tubule, the convoluted part was negative and the straight portion was moderately positive. In the renal corpuscle, the mesangial cells were weakly positive, parietal cells faintly positive, and the




**Fig. 4.** DIC micrographs of wax sections illustrating immunostaining for protein kinase C  $\beta$ I (PKC $\beta$ I) in 16- (A), 18- (B) and 20-day-old (E) fetal kidneys, and 3- (F) and 7-day-old (C, D & G) pups. PKC $\beta$ I appeared in the ureteric buds (UB) at 16 days of gestation and in the proximal anlage (open arrows) at 18 days of gestation. PKC $\beta$ I immunostaining appeared in the differentiating proximal tubules (open arrows) in the fetal stage and disappeared from the mature proximal tubule after birth. PKC $\beta$ I-positive proximal anlage are observed in the subcapsular region until 7 days after birth. Note the PKC $\beta$ I immunostaining in the mesangial cells (arrows) and parietal epithelium (arrowheads) of the developing renal corpuscle in D. Immunoreactivity for PKC $\beta$ I in the medullary collecting duct (MCD) cells gradually disappeared after birth (E-G). IMCD, inner medullary collecting duct. Magnifications: A,  $\times 200$ ; B,  $\times 264$ ; C & D-G,  $\times 528$ .

podocytes negative. The thick ascending limb of the Loop of Henle and the distal convoluted tubule were also negative. We did not observe immunoreactivity in the outer and inner medullary collecting tubules. However, we did detect moderately positive staining in the inner medulla, at the apical plasma membrane of the descending thin limb (Fig. 3; Table 1, 2).

In the developing kidney, PKC $\beta$ I immunoreactivity appeared from 16 days of gestation and was strongly positive in the ureteric bud (Fig. 4A). The immunoreactivity of the collecting tubule was strong in the fetus, but decreased markedly after birth, and the principal

cells were negative 3 days after birth (Fig. 4E-G). In the renal corpuscle, mesangial cells, parietal cells, and the proximal anlage were strongly positive in 18-day-old pups (Fig. 4B, C). The renal vesicles and S-shaped bodies were negative (Fig. 4A). During development, the straight portion of the proximal anlage remained moderately immunopositive and the immunoreactivity in the convoluted portion disappeared after birth, being negative from 14 days after birth. In the distal nephron, the distal anlage was negative, with immunoreactivity only becoming positive from 21 days after birth. PKC $\beta$ I immunoreactivity in the intercalated cells showed a



**Fig. 5.** DIC micrographs of 1-mm-thick plastic sections illustrating immunostaining for protein kinase C  $\beta$ II (PKC $\beta$ II) in the cortex (A), medullary rays (B), the border between the outer and inner medulla (C), and the terminal part of the inner medulla (D) of adult rat kidney. Immunoreactivity for PKC $\beta$ II is well localized on the basolateral plasma membrane of distal convoluted tubule (DCT) cells, connecting tubule (CNT) cells, thick ascending limb (TAL) cells, and principal cells throughout the collecting duct. Note that there is no immunoreactivity for PKC $\beta$ II in both the type A intercalated cells (arrow) and type B intercalated cells (arrowhead) in the cortical collecting duct (CCD). There is weak basolateral PKC $\beta$ II labeling in the proximal convoluted tubule (PCT). The open arrow indicates a PKC $\beta$ II-negative intercalated cell in the CNT. ATL, ascending thin limb; G, glomerulus; OMCDi, inner stripe of the outer medullary collecting duct; IMCDi, initial part of the inner medullary collecting duct; IMCDt, terminal part of the IMCD. Magnifications: A,  $\times 368$ ; B,  $\times 640$ ; C & D,  $\times 320$ .

pattern similar to PKC $\alpha$  in appearance and distribution.


#### 4. PKC $\beta$ II

In the adult kidney, immunoreactivity for PKC $\beta$ II was strongly positive in the thick ascending limb of the Loop of Henle, the macula densa, the distal convoluted tubule, and the basolateral membrane of the connecting tubule. In the collecting tubule, the basolateral membrane of the principal cells was moderately positive, but no immunoreactivity was seen in the intercalated cells of the connecting and collecting tubules (Fig. 5B–D).

There was weak basolateral labeling in the proximal convoluted tubule. There was no immunoreactivity in the renal corpuscle or intermediate tubule (Fig. 5A; Table 1, 2).

In the developing kidney, PKC $\beta$ II immunoreactivity appeared in the basolateral membrane of the distal anlage at 18 days of gestation. Immunoreactivity increased markedly in the distal tubule, including the thick ascending limb of the Loop of Henle, the distal convoluted tubule and the connecting tubule from 1 day after birth. PKC $\beta$ II immunoreactivity gradually de-






**Fig. 6.** DIC micrographs of wax sections illustrating immunostaining for protein kinase C  $\beta$ II (PKC $\beta$ II) in 18-day-old fetuses (A), and 1- (B), 3- (C), and 7-day-old (D) pups. A. PKC $\beta$ II immunoreactivity appeared in the distal tubules (asterisks) at 18 days of gestation. B-C. Note a marked increase in PKC $\beta$ II immunostaining in the distal tubules (asterisks) after birth. Insets demonstrating PKC $\beta$ II-negative medullary collecting ducts (MCD) from an 18-day-old fetus (A) and a 1-day-old pup (B). In the thick ascending limb (TAL), PKC $\beta$ II immunoreactivity appeared around the time of birth. P, proximal tubule; S, S-shaped body; III, stage III renal corpuscle; V, renal vesicle. Magnifications: A-D,  $\times 200$ ; insets,  $\times 528$ .


creased from 7 days after birth and had a similar pattern to the adult rat from 21 days. The intercalated cells in the connecting and collecting tubules were negative for PKC  $\beta$ II immunostaining. The principal cells were negative during the initial stages of development, but immunoreactivity gradually increased after birth and showed a similar pattern to adult rats from 14 days after birth (Fig. 6).

#### 5. PKC $\gamma$

There was very weak immunoreactivity for PKC $\gamma$  only in the proximal tubule, whereas no immunoreactivity in other uriniferous tubules (Fig. 7; Table 1, 2). In the developing kidney, PKC $\gamma$  immunoreactivity was strong in the proximal tubule. Immunoreactivity appeared in the proximal anlage at 18 days of gestation. There was strong positive staining in the entire proximal tubule at 1, 3, and



**Fig. 7.** Light micrographs of wax sections illustrating immunostaining for protein kinase C  $\gamma$  (PKC $\gamma$ ) in adult rat kidneys. There is very weak immunostaining for PKC $\gamma$  only in the cytoplasm of proximal convoluted tubules (PCT) and proximal straight tubules (PST). Co, cortex; DCT, distal convoluted tubule; OMCDi, inner stripe part of outer medullary collecting duct; OMCDo, outer stripe part of OMCD; OSOM, outer stripe of outer medulla; ISOM, inner stripe of outer medulla; TAL, thick ascending limb. Magnifications: A–C,  $\times 330$ .




**Fig. 8.** Light micrographs of wax sections illustrating immunostaining for protein kinase C  $\gamma$  (PKC $\gamma$ ) in kidneys from 20-day-old fetuses (A & B), and 1- (C & D), 3- (E), and 21-day-old (F) pups. PKC $\gamma$  immunoreactivity is highly expressed in the proximal tubules (PT) during development and markedly decreased in the PT from 21 days after birth. B and D are higher magnification area indicate by rectangular in A and C, respectively. DT, distal tubule; PCT, proximal convoluted tubule; PST, proximal straight tubule. Magnifications: A, C, E & F,  $\times 83$ ; B & D,  $\times 330$ .

20 days after birth. Subsequently, immunoreactivity decreased and had a similar pattern to adult rats from day 21 after birth (Fig. 8).

### Discussion

PKC plays a central role in intracellular signal

transduction. The various PKC isoforms are expressed in the rat kidney with distinct and differential expression patterns (Fig. 9). As a member of the cPKC group, PKC $\alpha$  expression was predominant in the adult kidney. PKC $\beta$  was localized in the tubules. PKC $\gamma$  is known to be detected in the central nervous system<sup>10</sup>. Our study, using immunohistochemistry and immuno-



**Fig. 9.** Changes in immunoreactivity of classical protein kinase C isoforms in uriniferous tubules during renal development in the rat. RC, renal corpuscle; PT, proximal tubule; DT, distal tubule; CD, collecting duct.

blotting, demonstrates that the expression of PKC $\alpha$ ,  $\beta$ I and  $\beta$ II, but not PKC $\gamma$ , is evident in the tubules of the rat kidney and PKC $\alpha$ ,  $\beta$ I,  $\beta$ II, and  $\gamma$  are expressed in the developing kidney.

Wetsel et al.<sup>11)</sup>, La Porta et al.<sup>12)</sup>, Dong et al.<sup>22)</sup>, Pfaff et al.<sup>26)</sup>, Saxena et al.<sup>27)</sup>, Ostlund et al.<sup>14)</sup>, Aristimuno and Good<sup>13)</sup>, and Serlachius et al.<sup>15)</sup> have reported the expression of PKC $\alpha$  in the kidney. In our study, the expression of PKC $\alpha$  was detected in the cortex, outer stripe of the outer medulla, inner stripe of the outer medulla and, using immunoblotting, in cytosolic and membrane fractions from the inner medulla. Using immunohistochemistry, Dong et al.<sup>28)</sup> and Fukuzaki et al.<sup>29)</sup> reported PKC $\alpha$  expression in the renal corpuscle, proximal straight tubule and collecting duct of the inner medulla of rat and human kidneys. However, our study demonstrates that PKC $\alpha$  staining was diffusely positive, with the exception of the intermediate tubule. Especially, strong positive staining was observed in the connecting tubule, intercalated cells of the cortical collecting tubule, mesangial cells of the renal corpuscle, outer and inner stripes of the outer medulla, and the

principal cells in the collecting duct of the inner medulla.

Wetsel et al.<sup>11)</sup>, Ostlund et al.<sup>14)</sup> and Aristimuno and Good<sup>13)</sup> reported the expression of PKC $\beta$ I in the kidney, but not PKC $\beta$ II. La Porta et al.<sup>12)</sup> identified PKC $\beta$  in renal corpuscles using immunohistochemistry. Our study demonstrates the expression of PKC $\beta$ I and  $\beta$ II. On immunoblotting, PKC $\beta$ I was faintly detected in the cortex and outer stripe of the outer medulla in the cytosolic fractions. Expression was generally weakly detected in the membrane fraction. Using immunohistochemistry, PKC $\beta$ I immunoreactivity was detected in the connecting segment, cortical collecting tubules, proximal straight tubules, mesangial cells of the renal corpuscle and the parietal epithelium. The connecting tubules and intercalated cells of the cortical collecting tubules were strongly positive. The apical plasma membrane of the descending thin limb of Henle was positive. These results were consistent with the immunoblotting findings.

A PKC $\beta$ II band was observed in the membrane fractions. On immunohistochemistry, PKC $\beta$ II was ex-

pressed in the proximal convoluted tubules, distal convoluted tubules, and the basolateral plasma membrane of the connecting and collecting tubules. The distal convoluted tubules, including the thick ascending limb of the Loop of Henle, and the connecting tubules were strongly positive. The intercalated cells in the distal nephron showed distinct and different expression patterns for PKC isoenzymes. In the connecting segment and cortical collecting tubules, the intercalated A cells play a role in  $H^+$  secretion and the intercalated B cells are involved in  $HCO_3^-$  secretion. In the medullary collecting tubules, intercalated cells play a role in  $H^+$  secretion<sup>30</sup>. Our study shows that the type A intercalated cells in the connecting and cortical collecting duct are strongly positive for PKC $\alpha$  and  $\beta$  in their supranuclear cytoplasm. Type B intercalated cells were moderately positive in their cytoplasm and basolateral plasma membranes. Those findings were consistent with the results of mouse kidneys<sup>31</sup> and the location of the  $H^+$ -ATPase<sup>32</sup>, so we suggest that PKC $\alpha$  and  $\beta$  may contribute to secretion of protons. However, PKC $\alpha$ , and  $\beta$  were not expressed in the intercalated cells in the medullary collecting ducts, so we speculate that different control mechanisms exist between intercalated cells in the connecting segment and cortical and medullary collecting tubules.

Several studies have reported that PKC $\gamma$  is not detected in the adult rat kidney<sup>11, 13, 14, 22</sup>. Recently, studies have shown that PKC plays a role in growth and differentiation during development<sup>21-24</sup>. Serlachius et al.<sup>15</sup> suggested that there is distinct and differential expression and distribution of PKC isoenzymes depending on embryonal kidney development. We also observed distinct and differential expression and distribution of PKC isoenzymes depending on kidney development. PKC $\alpha$ ,  $\beta$ I, and  $\beta$ II were expressed and, interestingly, PKC $\gamma$  was detected temporarily in the developing kidney. Positive staining for PKC $\alpha$  and  $\beta$ I appeared in the ureteric bud at 16 days of gestation and was strongly positive before birth, and then gradually decreased after birth. Therefore, we suggest that PKC $\alpha$

and  $\beta$ I play a role in differentiation of the collecting tubule. PKC $\alpha$  expression was positive in the proximal and distal anlage of pups up to 7 days of age. Therefore, we believe that PKC $\alpha$  expression is correlated with differentiation of the proximal and distal tubules. Moreover, PKC $\beta$ I expression is correlated with differentiation of the proximal tubule because it was temporarily expressed in the proximal anlage in early stages of development. PKC $\beta$ II appeared in the basolateral membrane of the distal and connecting tubules at 18 days of gestation, and gradually increased during development. Therefore, we suggest that PKC $\beta$ II expression is not correlated with growth and differentiation. PKC $\gamma$  immunoreactivity appeared and was highly expressed in the proximal anlage of the 18-day-old fetus, then decreased markedly, and disappeared soon after birth. Therefore, PKC $\gamma$  appears to be correlated with differentiation of the proximal tubule.

In summary, our study demonstrates that the classical PKC isoforms, PKC $\alpha$ ,  $\beta$ I and  $\beta$ II, but not PKC $\gamma$ , are expressed in the tubules of adult rat kidneys, and PKC $\alpha$ ,  $\beta$ I,  $\beta$ II, and  $\gamma$  are expressed in the developing kidney, and there are distinct and differential expression patterns for the isoforms according to location and stage of development.

#### Acknowledgements

This work was supported by the Korea Science and Engineering Foundation (KOSEF) through the MRC for Cell Death Disease Research Center at the Catholic University of Korea (R13-2002-005-01001-0).

#### References

- 1) Nishizuka Y: Studies and perspectives of protein kinase C. *Science* **233**:305-312, 1986
- 2) Nishizuka Y: Intracellular signaling by hydrolysis of phospholipids and activation of protein kinase C. *Science* **258**:607-614, 1992
- 3) Nishizuka Y: Protein kinase C and lipid signaling for sustained cellular responses. *FASEB J* **9**:484-496,

- 1995
- 4) Bertorello AM, Aperia A, Walaas SI, Nairn AC, Greengard P: Phosphorylation of the catalytic subunit of Na<sup>+</sup>, K<sup>(+)</sup>-ATPase inhibits the activity of the enzyme. *Proc Natl Acad Sci U S A* **88**:11359–11362, 1991
  - 5) Nowicki S, Kruse MS, Brismar H, Aperia A: Dopamine-induced translocation of protein kinase C isoforms visualized in renal epithelial cells. *Am J Physiol Cell Physiol* **279**:C1812–C1818, 2000
  - 6) Middleton JP, Khan WA, Collinsworth G, Hannun YA, Medford RM: Heterogeneity of protein kinase C-mediated rapid regulation of Na/K-ATPase in kidney epithelial cells. *J Biol Chem* **268**:15958–15964, 1993
  - 7) Horie S, Moe O, Miller RT, Alpern RJ: Long-term activation of protein kinase c causes chronic Na/H antiporter stimulation in cultured proximal tubule cells. *J Clin Invest* **89**:365–372, 1992
  - 8) Ruiz OS, Arruda JA: Regulation of the renal Na<sup>+</sup>-HCO<sub>3</sub><sup>-</sup> cotransporter by cAMP and Ca<sup>2+</sup>-dependent protein kinases. *Am J Physiol* **262**:F560–F565, 1992
  - 9) Kosaka Y, Ogita K, Ase K, Nomura H, Kikkawa U, Nishizuka Y: The heterogeneity of protein kinase C in various rat tissues. *Biochem Biophys Res Commun* **151**:973–981, 1988
  - 10) Ono Y, Fujii T, Ogita K, Kikkawa U, Igarashi K, Nishizuka Y: The structure, expression, and properties of additional members of the protein kinase C family. *J Biol Chem* **263**:6927–6932, 1988
  - 11) Wetsel WC, Khan WA, Merchenthaler I, Rivera H, Halpern AE, Phung HM, Negro-Vilar A, Hannun YA: Tissue and cellular distribution of the extended family of protein kinase C isoenzymes. *J Cell Biol* **117**:121–133, 1992
  - 12) La Porta CA, Comolli R: Biochemical and immunological characterization of calcium-dependent and -independent PKC isoenzymes in renal ischemia. *Biochem Biophys Res Commun* **191**:1124–1130, 1993
  - 13) Aristimo PC, Good DW: PKC isoforms in rat medullary thick ascending limb: selective activation of the delta-isoform by PGE<sub>2</sub>. *Am J Physiol* **272**:F624–F631, 1997
  - 14) Ostlund E, Mendez CF, Jacobsson G, Fryckstedt J, Meister B, Aperia A: Expression of protein kinase C isoforms in renal tissue. *Kidney Int* **47**:766–773, 1995
  - 15) Serlachius E, Svenilsson J, Schalling M, Aperia A: Protein kinase C in the developing kidney: isoform expression and effects of ceramide and PKC inhibitors. *Kidney Int* **52**:901–910, 1997
  - 16) Redling S, Pfaff IL, Leitges M, Vallon V: Immunolocalization of protein kinase C isoenzymes alpha, beta I, beta II, delta, and epsilon in mouse kidney. *Am J Physiol Renal Physiol* **287**:F289–F298, 2004
  - 17) Hashimoto T, Ase K, Sawamura S, Kikkawa U, Saito N, Tanaka C, Nishizuka Y: Postnatal development of a brain-specific subspecies of protein kinase C in rat. *J Neurosci* **8**:1678–1683, 1988
  - 18) Hirata M, Saito N, Kono M, Tanaka C: Differential expression of the beta I- and beta II-PKC subspecies in the postnatal developing rat brain: an immunocytochemical study. *Brain Res Dev Brain Res* **62**:229–238, 1991
  - 19) Puc at M, Hilal-Dandan R, Strulovici B, Brunton LL, Brown JH: Differential regulation of protein kinase C isoforms in isolated neonatal and adult rat cardiomyocytes. *J Biol Chem* **269**:16938–16944, 1994
  - 20) Rybin VO, Steinberg SF: Protein kinase C isoform expression and regulation in the developing rat heart. *Circ Res* **74**:299–309, 1994
  - 21) Housey GM, Johnson MD, Hsiao WL, O'Brian CA, Murphy JP, Kirschmeier P, Weinstein IB: Overproduction of protein kinase C causes disordered growth control in rat fibroblasts. *Cell* **52**:343–354, 1988
  - 22) Dong L, Stevens JL, Fabbro D, Jaken S: Regulation of protein kinase C isozymes in kidney regeneration. *Cancer Res* **53**:4542–4549, 1993
  - 23) Mischak H, Goodnight JA, Kolch W, Martiny-Baron G, Schaechtle C, Kazanietz MG, Blumberg PM, Pierce JH, Mushinski JF: Overexpression of protein kinase C-delta and -epsilon in NIH 3T3 cells induces opposite effects on growth, morphology, anchorage dependence, and tumorigenicity. *J Biol Chem* **268**:6090–6096, 1993
  - 24) Berra E, Diaz-Meco MT, Dominguez I, Municio MM, Sanz L, Lozano J, Chapkin RS, Moscat J: Protein kinase C zeta isoform is critical for mitogenic signal transduction. *Cell* **74**:555–563, 1993
  - 25) Szallasi Z, Kosa K, Smith CB, Dlugosz AA, Williams EK, Yuspa SH, Blumberg PM: Differential regulation by anti-tumor-promoting 12-deoxyphorbol-13-phenylacetate reveals distinct roles of the classical and novel protein kinase C isozymes in biological responses of primary mouse keratinocytes. *Mol Pharmacol* **47**:258–265, 1995
  - 26) Pfaff IL, Wagner HJ, Vallon V: Immunolocalization of protein kinase C isoenzymes alpha, beta I and beta II in rat kidney. *J Am Soc Nephrol* **10**:1861–1873, 1999
  - 27) Saxena R, Saksa BA, Hawkins KS, Ganz MB: Protein kinase C beta I and beta II are differentially expressed in the developing glomerulus. *FASEB J* **8**:646–653, 1994
  - 28) Dong LQ, Stevens JL, Jaken S: Biochemical and immunological characterization of renal protein kinase C. *Am J Physiol* **261**:F679–F687, 1991
  - 29) Fukuzaki A, Kaneto H, Ikeda S, Orikasa S: Characterization of protein kinase C expression in human kidney. *Tohoku J Exp Med* **178**:263–269, 1996
  - 30) Madsen KM, Tisher CC: Structural-functional relationships along the distal nephron. *Am J Physiol* **250**:F1–F15, 1986
  - 31) Kim WY, Jung JH, Park EY, Yang CW, Kim H, Nielsen

S, Madsen KM, Kim J: Expression of protein kinase C isoenzymes alpha, beta, and delta in subtypes of intercalated cells of mouse kidney. *Am J Physiol Renal Physiol* **291**:F1052–F1060, 2006

32) Madsen KM, Verlander JW, Kim J, Tisher CC: Morphological adaptation of the collecting duct to acid–base disturbances. *Kidney Int* **40**(Suppl 33): S57–S63, 1991