

POSTER PRESENTATION

Open Access

# What is the importance of solitary focal bony FDG-uptake on 18F-FDG PET-CT of known cancer patients?

LI Sonoda\*, A Lakhani, S Ghosh-Ray

From International Cancer Imaging Society (ICIS) 14th Annual Teaching Course  
Heidelberg, Germany. 9-11 October 2014

## Aim

18F-FDG-PET-CT plays an important role in oncology staging. While the presence of multiple FDG-avid lesions on PET-CT in the context of known malignancy is generally considered metastases, the exact significance of solitary FDG-avid-lesions remains unknown. This study was undertaken to evaluate the significance of solitary bony lesions on PET-CT of oncology patients.

## Methods

Retrospective review of 15,645 PET-CT studies was performed. Further evaluation of solitary bony FDG-avid lesions was carried out by conventional imaging, follow-up and biopsy studies. Spontaneous resolution on subsequent PET-CT without a change in therapy was considered benign while progression was considered malignant.

## Results

361 (3%) cases were found to have single FDG-avid skeletal lesions, of which 16 were due to uptake at the primary bony malignancy, and 42 were not further-investigated/passed away, hence excluded. Of the remaining 303 lesions 276 (91%) were confirmed as metastases, 27 (9%) proven benign (10 by imaging, 5 by biopsy and 12 by follow-up).

Of 276 metastases ( $\text{SUV}_{\text{max}} 9.6+/-6.6$ ); 191 were lytic, 45 sclerotic, 21 mixed and 19 normal on CT. Of 27 benign ( $\text{SUV}_{\text{max}} 3.8+/-2.8$ ); 2 were lytic, 7 sclerotic, 2 mixed and 16 normal on CT. PPV of PET-CT on lytic, sclerotic, mixed and normal lesions on CT are 99%, 87%, 91% and 54% respectively. There was significant difference in  $\text{SUV}_{\text{max}}$  between malignant/benign lesions ( $P<0.001$ ).

PET-CT correctly upstaged in 83/303 (27%) cases, but incorrectly upstaged or suggested further investigation in 18/303 (6%) cases.

## Conclusion

Solitary skeletal FDG-uptake on 18F-FDG-PET-CT in patients with known malignancy is just as significant as multiple skeletal FDG-uptake, carrying high risk of metastases.

Published: 9 October 2014

doi:10.1186/1470-7330-14-S1-P6

Cite this article as: Sonoda et al.: What is the importance of solitary focal bony FDG-uptake on 18F-FDG PET-CT of known cancer patients? *Cancer Imaging* 2014 14(Suppl 1):P6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: luke@sonoda.co.uk  
Paul Strickland Scanner Centre, Mount Vernon Hospital, London, UK

© 2014 Sonoda et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.