

Original Article



Shellfish/crustacean oral allergy syndrome among national service pre-enlistees in Singapore

Bernard Yu-Hor Thong^{1*}, Shalini Arulanandam², Sze-Chin Tan¹, Teck-Choon Tan¹, Grace Yin-Lai Chan¹, Justina Wei-Lyn Tan¹, Mark Chong-Wei Yeow², Chwee-Ying Tang¹, Jinfeng Hou¹, and Khai-Pang Leong¹

¹Department of Rheumatology, Allergy and Immunology, Tan Tock Seng Hospital, Singapore 308433

²Medical Classification Centre, Ministry of Defence, Singapore 109680



Received: Jan 21, 2018

Accepted: Apr 8, 2018

*Correspondence to

Bernard Yu-Hor Thong

Department of Rheumatology, Allergy and Immunology, Tan Tock Seng Hospital, 11 Jalan Tan Tock Seng, Singapore 308433.

Tel: +65-63577822

Fax: +65-63572686

E-mail: bernard_thong@ttsh.com.sg

Copyright © 2018. Asia Pacific Association of Allergy, Asthma and Clinical Immunology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Bernard Yu-Hor Thong

<https://orcid.org/0000-0002-6338-8482>

Author Contributions

Conceptualization: Bernard Yu-Hor Thong. Data curation: Bernard Yu-Hor Thong, Shalini Arulanandam, Sze-Chin Tan, Teck-Choon Tan, Grace Yin-Lai Chan, Justina Wei-Lyn Tan, Mark Chong-Wei Yeow, Chwee-Ying Tang, Jinfeng Hou, Khai-Pang Leong. Formal analysis: Bernard Yu-Hor Thong. Investigation: Bernard Yu-Hor Thong. Project administration: Bernard Yu-Hor Thong. Resources: Bernard Yu-Hor Thong. Validation: Bernard Yu-Hor Thong, Shalini Arulanandam, Sze-Chin Tan, Teck-Choon Tan, Grace Yin-Lai Chan, Justina Wei-Lyn Tan, Mark Chong-Wei Yeow, Chwee-Ying

ABSTRACT

Background: All Singaporean males undergo medical screening prior to compulsory military service. A history of possible food allergy may require referral to a specialist Allergy clinic to ensure that special dietary needs can be taken into account during field training and deployment.

Objective: To study the pattern of food allergy among pre-enlistees who were referred to a specialist allergy clinic to work up suspected food allergy.

Methods: Retrospective study of all pre-enlistees registered in the Clinical Immunology/Allergy New Case Registry referred to the Allergy Clinic from 1 August 2015 to 31 May 2016 for suspected food allergy.

Results: One hundred twenty pre-enlistees reporting food allergy symptoms other than rash alone were referred to the Allergy Clinic during the study period. Of these, 77 (64.2%) had food allergy. Among those with food allergy, mean age was 19.1 ± 1.5 years. They comprised predominantly Chinese (66.2%) and Malays (20.8%). The most commonly reported foods were shellfish/crustaceans (78%), peanut (15.6%), and egg (6.5%). Self-limiting oral allergy syndrome, OAS (itchy lips and throat with/without lip angioedema) was the most common manifestation ($n = 33$, 42.9%) followed by anaphylaxis ($n = 23$, 29.9%). Majority of OAS was from shellfish/crustacean (90.6%); of which shrimp (30.3%), crab (15.2%), and lobster (3.0%) were the most common. Mild childhood asthma (69.7%), allergic rhinitis (6.3%), and eczema (6.1%) were the most common atopic conditions among individuals with shellfish/crustacean OAS. This pattern was similar for shellfish/crustacean anaphylaxis. Skin prick tests were most commonly positive for shrimp (OAS 87.1% vs. anaphylaxis 100%), crab (OAS 95.8% vs. 90.9%), and lobster (OAS 91.7% vs. 63.6%).

Conclusion: OAS to shellfish/crustaceans was more common than anaphylaxis among this study population of young males referred for food allergy symptoms other than rash alone.

Keywords: Anaphylaxis; Hypersensitivity; Shellfish

INTRODUCTION

Prior to compulsory military service after the age of 18 years, Singaporean males are screened in the military and referred to our Allergy Clinic for confirmation of food allergy if they declare a history of suspected food allergy, with symptoms other than rash alone.

Tang, Jinfeng Hou, Khai-Pang Leong. Writing - original draft: Bernard Yu-Hor Thong, Shalini Arulanandam. Writing - review & editing: Bernard Yu-Hor Thong, Shalini Arulanandam, Teck-Choon Tan, Sze-Chin Tan, Grace Yin-Lai Chan, Justina Wei-Lyn Tan, Mark Chong-Wei Yeow, Chwee-Ying Tang, Jinfeng Hou, Khai-Pang Leong.

Allergic diseases are highly prevalent among children and adolescents in Singapore. This includes asthma, allergic rhinitis, atopic dermatitis, anaphylaxis, and food allergy. Screening pre-enlistees for food allergy allows special dietary needs to be catered for in military camps and during outfield training. The aim of this study was to describe the pattern of food allergy among pre-enlistees who reported a history of suspected food allergy.

MATERIALS AND METHODS

All pre-enlistees registered in the Clinical Immunology/Allergy New Case Registry who were referred to the Allergy Clinic from 1 August 2015 to 31 May 2016 for suspected food allergy were retrospectively studied. Their demographic profile, suspected food allergy/allergies, concomitant atopy, and outcomes of skin prick tests (SPTs), blood tests for allergen specific IgE and/or oral food challenges (OFCs) were studied. Patients who had a definite diagnosis of food allergy in childhood based on clinical history and SPT and/or specific IgE measurements did not have the tests repeated. OFC were offered to patients where childhood history was equivocal, and SPT/specific IgE were negative, or to demonstrate tolerance following dietary elimination over time. This study was part of the Clinical Immunology/Allergy New Case Registry's approved domain specific review board reference number TTSH/2005-0016 for which additional informed consent was not required as this was a retrospective registry review with minimal risks to research subjects.

RESULTS

There were 120 male pre-enlistees comprising 26% of all referrals to the Allergy Clinic during the study period, of whom 77 (64.2%) had food allergy. Among those with food allergy, mean age was 19.1 ± 1.5 years. They comprised predominantly Chinese (66.2%) and Malays (20.8%) followed by 6.5% Indians and other ethnicities respectively. The most commonly reported foods were shellfish/crustaceans (78%), peanut (15.6%), and egg (6.5%).

Self-limiting oral allergy syndrome, OAS (itchy lips and throat with/without lip angioedema) was the most common manifestation ($n = 33$, 42.9%) followed by anaphylaxis ($n = 23$, 29.9%). The remaining 27.2% had urticarial and/or periorbital angioedema or food sensitive eczema. Among those with anaphylaxis, the most common manifestations were dyspnoea/wheeze (73.9%), angioedema (65.2%) most commonly of the lips, urticaria (60.9%), and diarrhoea (34.8%).

Majority of OAS was from shellfish/crustacean (90.6%); of which shrimp (30.3%), crab (15.2%), and lobster (3.0%) were the most common. Mild childhood asthma (69.7%), allergic rhinitis (6.3%), and eczema (6.1%) were the most common atopic condition among individuals with shellfish/crustacean OAS. This pattern was similar for shellfish/crustacean anaphylaxis.

SPTs were most commonly positive for shrimp (OAS 87.1% vs. anaphylaxis 100%), crab (OAS 95.8% vs. 90.9%) and lobster (OAS 91.7% vs. 63.6%). Two pre-enlistees had food sensitive eczema to cow's milk and peanut respectively. There were no cases of food-dependent exercise-induced anaphylaxis. The above information is summarized in **Table 1**.

Table 1. Most common foods and concomitant atopy among patients with food induced oral allergy syndrome and anaphylaxis

Variable	Oral allergy syndrome (n = 33)	Anaphylaxis (n = 23)
Culprit food		
Prawn	10 (30.3)	8 (34.8)
Crab	5 (15.2)	4 (17.4)
Squid	2 (6.1)	0 (0)
Lobster	1 (3.0)	0 (0)
Other foods		
Peanut	0 (0)	5 (21.7)
Egg	0 (0)	1 (4.3)
Abalone	0 (0)	1 (4.3)
Limpet	0 (0)	1 (4.3)
Bird nest	0 (0)	1 (4.3)
Buckwheat	0 (0)	1 (4.3)
Atopy		
Asthma	23 (69.7)	11 (47.8)
Allergic rhinitis	10 (30.3)	7 (30.4)
Eczema	2 (6.1)	1 (4.3)
Positive skin prick tests		
Shrimp	27/31 (87.1)	11/11 (100)
Crab	23/24 (95.8)	10/11 (90.9)
Lobster	22/24 (91.7)	7/11 (63.6)
Squid	11/20 (55.0)	1/8 (12.5)
Clam	9/20 (45.0)	2/9 (22.2)
Oyster	6/19 (31.6)	3/4 (75.0)
Scallop	6/20 (30.0)	1/9 (11.1)
Mussel	0 (0)	4/10 (40.0)

Values are presented as number (%).

Skin prick tests denominators vary with each respective food allergen as not all patients had all the skin tests done to the same panel of food allergens.

DISCUSSION

Allergic diseases are common in Singapore among children and adolescents (defined as up to 21 years old in the Singapore content), especially asthma [1], allergic rhinitis [2], and atopic dermatitis [3]. House dust mite is the most important inhalant allergen driving allergic sensitization in these 3 conditions in Singapore and Southeast Asia [4, 5], more so than grass and tree pollen. Anaphylaxis from food allergy (especially seafood/shellfish) and drug allergy/hypersensitivity (especially nonsteroidal anti-inflammatory drugs) are also common [6, 7].

In the international study of asthma and allergies in childhood (ISAAC) published in 1996 [8] and 2004 [9], the Singapore prevalence of doctor-diagnosed asthma was 20%, parent-reported rhinitis 44%, and chronic rashes 12% in 1996. In 2009, the Growing Up in Singapore Towards healthy Outcomes (GUSTO) study was initiated [10]. This is a large birth cohort study comprising 1,163 pregnant mothers and their children, with an integrated Allergy domain describing the epidemiology of allergic manifestations, phenotype, and the association between the development of metabolic diseases and allergy. Some of the key findings from the GUSTO cohort to date are as follows. Firstly, early childhood rhinitis up to the age of 18 months was associated with a history of parental atopy; atopic comorbidities of eczema and wheeze, but not allergen sensitization [11]. Secondly, early onset (before 18 months of age) eczema and wheeze are risk factors for later allergen sensitization as demonstrated by positive SPT to inhalant allergens (house dust mites: *Dermatophagoides pteronyssinus*, *Dermatophagoides farinae*, and *Blomia tropicalis*) and food allergens (egg, peanut, and cow's milk) [12]. Thirdly, there is a possible influence of maternal diet during pregnancy

on the development of allergic outcomes (eczema, rhinitis, and wheeze) in offspring up to the first 36 months of life [13]. Fourthly, delayed introduction of allergenic foods (egg, peanut, and shellfish) after 10 months in 50% or more of infants did not increase the prevalence of food allergy between the ages of 12–18 months: egg allergy 0.35%–1.8%, peanut allergy 0.1%–0.3%, and shellfish allergy 0.2%–0.9% [14]. This contrasts with the recent recommendation of early introduction of peanuts to prevent the onset of peanut allergy in high risk infants based on the “Learning Early About Peanut” study [15].

Food allergy in Singapore is relatively common among children and adolescents where shellfish allergy is overall more common than egg, cow's milk, and peanut allergy [16, 17]. Some unusual food allergens have also been reported locally, namely bird's nest [18], galactooligosaccharide – a prebiotic found in milk formulas [19], dust-mite contaminated flour [20], and tropical fish (threadfin, Indian anchovy, pomfret, and tengirri) [21]. Case series of food-dependent exercise induced anaphylaxis from shellfish [22] and wheat [23] have also been reported. A novel OAS from crustacean-shellfish allergy presenting with predominantly oral symptoms (tingling, lip swelling) has also been described in Singapore [24], with tropomyosin in house dust mites believed to be a possible cross-reactive sensitizing inhalant allergen [25]. This is consistent with the findings in our study.

There is paucity of literature on allergic disorders experienced by military servicemen or pre-enlistees in the Asia-Pacific region. Most of the literature in military medicine describe case reports/series on the epidemiology of insect venom hypersensitivity [26–28], asthma [29]/ exercise-induced bronchoconstriction [30], anaphylaxis [31, 32], and skin disorders [33, 34]. The literature on therapeutics in allergic diseases in the military covers mostly venom immunotherapy (VIT) and the use of epinephrine autoinjectors in anaphylaxis. VIT is the definitive treatment for insect venom anaphylaxis, enabling active duty servicemen to remain medically fit and ready for deployment around the world [35, 36]. Epinephrine autoinjector use in military patient populations [37, 38] has also been described. From the Asia-Pacific region, there has only been a single study describing the prevalence of allergic disease and wheezing among Korean military personnel [39] using a modified ISAAC questionnaire. The prevalence of current wheeze (10.5%), allergic rhinitis (14.0%), and eczema (9.7%) confirmed a significant prevalence of allergic disease in young adults in the Korean military.

Food allergy can be a limiting factor in the deployability of servicemen, as providing for a range of allergen-free rations in the field can be challenging; and there is an ever-present risk of a severe allergic reactions occurring in training areas with limited access to tertiary medical care.

Self-declaration of food allergy at pre-enlistment was implemented in 2015 as there appeared to be an increasing prevalence of food allergy among pre-enlistees. Servicemen with food allergies have their food allergy documented in their electronic personnel records, such that they can be supplied with suitable allergen-free diets in their camp cookhouses.

OAS to shellfish/crustaceans among pre-enlistees is consistent with the phenotype described in recent studies on shrimp allergy in Singapore [24]. The risk of accidental shrimp ingestion in a shrimp OAS patient triggering off anaphylaxis is unknown, and there are no studies presently to identify and risk-prognosticate such patients. Shrimp allergic individuals react to several high-molecular weight allergens, making molecular diagnosis with component resolved diagnostics difficult [40]. Tropomyosin and sarcoplasmic-calcium binding protein sensitization has been

shown to be associated with clinical reactivity to shrimp, arginine kinase and hemocyanin appear to be cross-reacting allergens between shrimp and arthropods [41]. In contrast in pollen-fruit OAS, the primary sensitizing and cross-reactive allergens are more clearly defined, and reactions are generally mild with little risk for anaphylaxis [42, 43]. For servicemen with shellfish anaphylaxis, the risk of accidental ingestion when outfield vis-à-vis difficulty in obtaining early access to emergency medical services continues to pose a risk to them being field-deployable [44]. However, for the majority of shellfish/crustacean-allergic pre-enlistees who have mild oropharyngeal symptoms, the avoidance of seafood ingredients in field rations should be sufficient to allow them to be field-deployable, as the risk of anaphylaxis in this group is likely to be low.

Oral immunotherapy (OIT) for food allergy may in future become a therapeutic option in achieving long-term tolerance (rather than just desensitization) with daily ingestion of the culprit food allergen [45]. For now, OIT remains a research tool, limited mainly to cow's milk, egg, and peanut allergy even though they have been mentioned in recently written clinical practice guidelines from Spain [46] and Japan [47]. More well designed randomized clinical trials are needed before OIT can become standard of care (like VIT). Although house dust mite sublingual immunotherapy has been reported to improve symptoms of shrimp allergy [48], this alone is unlikely to be effective where nontropomyosin allergens are involved. T-cell epitope peptide immunotherapy [49] and addition of other adjuvants may be needed to improve the efficacy and safety of immunotherapy.

A limitation of this study was that all pre-enlistees with shellfish/crustacean OAS were not contemporaneously offered repeat SPT (regardless how long before they had last been done), followed by confirmatory OFC to determine their threshold of reactivity, as in a previous local study [24]. The variability of the threshold of reactivity between individuals may pose a risk of systemic reactions following accidental ingestion. OFCs may play a role in determining the anaphylaxis-risk and thus field-deployability of an individual with OAS.

In conclusion, allergy to shellfish/crustaceans was the most common food allergy among pre-enlistees, and OAS to shellfish/crustaceans was more common than anaphylaxis.

REFERENCES

1. Soh JY, Ng B, Tan Z, Xu S, Hing WC, Wu TS, Chan YH, Lee BW. Ten-year prescription trends of asthma medications in the management of childhood wheeze. *Allergy Asthma Proc* 2014;35:e1-8.
[PUBMED](#) | [CROSSREF](#)
2. Chiang WC, Chen YM, Tan HK, Balakrishnan A, Liew WK, Lim HH, Goh SH, Loh WY, Wong P, Teoh OH, Goh A, Chay OM. Allergic rhinitis and non-allergic rhinitis in children in the tropics: prevalence and risk associations. *Pediatr Pulmonol* 2012;47:1026-33.
[PUBMED](#) | [CROSSREF](#)
3. Lee BW, Detzel PR. Treatment of childhood atopic dermatitis and economic burden of illness in Asia Pacific countries. *Ann Nutr Metab* 2015;66 Suppl 1:18-24.
[PUBMED](#) | [CROSSREF](#)
4. Tham EH, Lee AJ, Bever HV. Aeroallergen sensitization and allergic disease phenotypes in Asia. *Asian Pac J Allergy Immunol* 2016;34:181-9.
[PUBMED](#) | [CROSSREF](#)
5. Llanora GV, Ming LJ, Wei LM, van Bever HP. House dust mite sensitization in toddlers predict persistent wheeze in children between eight to fourteen years old. *Asia Pac Allergy* 2012;2:181-6.
[PUBMED](#) | [CROSSREF](#)

6. Ganapathy S, Lwin Z, Ting DH, Goh LS, Chong SL. Anaphylaxis in children: experience of 485 episodes in 1,272,482 patient attendances at a tertiary Paediatric Emergency Department from 2007 to 2014. *Ann Acad Med Singapore* 2016;45:542-8.
[PUBMED](#)
7. Liew WK, Chiang WC, Goh AE, Lim HH, Chay OM, Chang S, Tan JH, Shih E, Kidon M. Paediatric anaphylaxis in a Singaporean children cohort: changing food allergy triggers over time. *Asia Pac Allergy* 2013;3:29-34.
[PUBMED](#) | [CROSSREF](#)
8. Goh DY, Chew FT, Quek SC, Lee BW. Prevalence and severity of asthma, rhinitis, and eczema in Singapore schoolchildren. *Arch Dis Child* 1996;74:131-5.
[PUBMED](#) | [CROSSREF](#)
9. Wang XS, Shek LP, Ma S, Soh SE, Lee BW, Goh DY. Time trends of co-existing atopic conditions in Singapore school children: prevalence and related factors. *Pediatr Allergy Immunol* 2010;21(1 Pt 2):e137-41.
[PUBMED](#) | [CROSSREF](#)
10. Soh SE, Lee SS, Hoon SW, Tan MY, Goh A, Lee BW, Shek LP, Teoh OH, Kwek K, Saw SM, Godfrey K, Chong YS, Gluckman P, van Bever HP. The methodology of the GUSTO cohort study: a novel approach in studying pediatric allergy. *Asia Pac Allergy* 2012;2:144-8.
[PUBMED](#) | [CROSSREF](#)
11. Hardjojo A, Goh A, Shek LP, Van Bever HP, Teoh OH, Soh JY, Thomas B, Tan BH, Chan YH, Ramamurthy MB, Goh DY, Soh SE, Saw SM, Kwek K, Chong YS, Godfrey KM, Gluckman PD, Lee BW. Rhinitis in the first 18 months of life: exploring the role of respiratory viruses. *Pediatr Allergy Immunol* 2015;26:25-33.
[PUBMED](#) | [CROSSREF](#)
12. Loo EX, Sim JZ, Goh A, Teoh OH, Chan YH, Saw SM, Kwek K, Gluckman PD, Godfrey KM, Van Bever H, Chong YS, Lee BW, Kramer MS, Shek LP. Predictors of allergen sensitization in Singapore children from birth to 3 years. *Allergy Asthma Clin Immunol* 2016;12:56.
[PUBMED](#) | [CROSSREF](#)
13. Loo EXL, Ong L, Goh A, Chia AR, Teoh OH, Colega MT, Chan YH, Saw SM, Kwek K, Gluckman PD, Godfrey KM, Van Bever H, Lee BW, Chong YS, Chong MF, Shek LP. Effect of maternal dietary patterns during pregnancy on self-reported allergic diseases in the first 3 years of life: results from the GUSTO Study. *Int Arch Allergy Immunol* 2017;173:105-13.
[PUBMED](#) | [CROSSREF](#)
14. Tham EH, Lee BW, Chan YH, Loo EXL, Toh JY, Goh A, Teoh OH, Yap F, Tan KH, Godfrey KM, Chong MFF, Van Bever HPS, Chong YS, Shek LP. Low food allergy prevalence despite delayed introduction of allergenic foods-data from the GUSTO cohort. *J Allergy Clin Immunol Pract* 2017 Jul 19 [Epub]. pii: S2213-2198(17)30423-3. <https://doi.org/10.1016/j.jaip.2017.06.001>.
15. Togias A, Cooper SF, Acebal ML, Assa'ad A, Baker JR Jr, Beck LA, Block J, Byrd-Bredbenner C, Chan ES, Eichenfield LF, Fleischer DM, Fuchs GJ 3rd, Furuta GT, Greenhawt MJ, Gupta RS, Habich M, Jones SM, Keaton K, Muraro A, Plaut M, Rosenwasser LJ, Rotrosen D, Sampson HA, Schneider LC, Sicherer SH, Sidbury R, Spergel J, Stukus DR, Venter C, Boyce JA. Addendum guidelines for the prevention of peanut allergy in the United States: Report of the National Institute of Allergy and Infectious Diseases-sponsored expert panel. *J Allergy Clin Immunol* 2017;139:29-44.
[PUBMED](#) | [CROSSREF](#)
16. Lee AJ, Shek LP. Food allergy in Singapore: opening a new chapter. *Singapore Med J* 2014;55:244-7.
[PUBMED](#) | [CROSSREF](#)
17. Lee AJ, Gerez I, Shek LP, Lee BW. Shellfish allergy: an Asia-Pacific perspective. *Asian Pac J Allergy Immunol* 2012;30:3-10.
[PUBMED](#)
18. Goh DL, Chew FT, Chua KY, Chay OM, Lee BW. Edible "bird's nest"-induced anaphylaxis: an under-recognized entity? *J Pediatr* 2000;137:277-9.
[PUBMED](#) | [CROSSREF](#)
19. Tay SY, Tham E, Yeo CT, Yi FC, Chen JY, Cheong N, Chua KY, Lee BW. Anaphylaxis following the ingestion of flour contaminated by house dust mites: a report of two cases from Singapore. *Asian Pac J Allergy Immunol* 2008;26:165-70.
[PUBMED](#)
20. Soh JY, Chiang WC, Huang CH, Woo CK, Ibrahim I, Heng K, Pramanick A, Lee BW. An unusual cause of food-induced anaphylaxis in mothers. *World Allergy Organ J* 2017;10:3.
[PUBMED](#) | [CROSSREF](#)
21. Lim DL, Neo KH, Yi FC, Chua KY, Goh DL, Shek LP, Giam YC, Van Bever HP, Lee BW. Parvalbumin: the major tropical fish allergen. *Pediatr Allergy Immunol* 2008;19:399-407.
[PUBMED](#) | [CROSSREF](#)

22. Teo SL, Gerez IF, Ang EY, Shek LP. Food-dependent exercise-induced anaphylaxis: a review of 5 cases. *Ann Acad Med Singapore* 2009;38:905-9.
[PUBMED](#)
23. Thalayasingam M, Allameen NA, Soh JY, Bigliardi P, Van Bever H, Shek LP. Wheat-dependent exercise-induced anaphylaxis: a retrospective case review from a tertiary hospital. *Postgrad Med J* 2014;90:488-92.
[PUBMED](#) | [CROSSREF](#)
24. Thalayasingam M, Gerez IF, Yap GC, Llanora GV, Chia IP, Chua L, Lee CJ, Ta LD, Cheng YK, Thong BY, Tang CY, Van Bever HP, Shek LP, Curotto de Lafaille MA, Lee BW. Clinical and immunochemical profiles of food challenge proven or anaphylactic shrimp allergy in tropical Singapore. *Clin Exp Allergy* 2015;45:687-97.
[PUBMED](#) | [CROSSREF](#)
25. Wong L, Huang CH, Lee BW. Shellfish and house dust mite allergies: is the link tropomyosin? *Allergy Asthma Immunol Res* 2016;8:101-6.
[PUBMED](#) | [CROSSREF](#)
26. Thong BY, Leong KP, Chng HH. Insect venom hypersensitivity: experience in a clinical immunology/allergy service in Singapore. *Singapore Med J* 2005;46:535-9.
[PUBMED](#)
27. Turbyville JC, Dunford JC, Nelson MR. Hymenoptera of Afghanistan and the central command area of operations: assessing the threat to deployed U.S. service members with insect venom hypersensitivity. *Allergy Asthma Proc* 2013;34:179-84.
[PUBMED](#) | [CROSSREF](#)
28. Goldberg A, Confino-Cohen R, Mekori YA. Deliberate hymenoptera sting challenge as a diagnostic tool in highly selected venom-allergic patients. *Ann Allergy Asthma Immunol* 1995;75:30-2.
[PUBMED](#)
29. Brooks SM. Occupational medicine model and asthma military recruitment. *Mil Med* 2015;180:1140-6.
[PUBMED](#) | [CROSSREF](#)
30. Stocks J, Tripp M, Lin T. Methacholine challenge is insufficient to exclude bronchial hyper-responsiveness in a symptomatic military population. *J Asthma* 2014;51:886-90.
[PUBMED](#) | [CROSSREF](#)
31. Zeindler PR, Gervais A. Anaphylaxis: assessment of a disease-based military medical standard. *Mil Med* 2011;176:889-95.
[PUBMED](#) | [CROSSREF](#)
32. Handfield KS, Dolan CK, Kaplan M. Cholinergic urticaria with anaphylaxis: hazardous duty of a deployed US marine. *Cutis* 2015;95:241-3.
[PUBMED](#)
33. Gan WH, Low R, Koh D. Dermatological conditions in military conscripts. *Occup Med (Lond)* 2013;63:435-8.
[PUBMED](#) | [CROSSREF](#)
34. Chong WS. Dermatology in the military field: What physicians should know? *World J Clin Cases* 2013;1:208-11.
[PUBMED](#) | [CROSSREF](#)
35. Davis KL, Kolisnyk JT, Klote MM, Yacovone MA, Martin BL, Nelson MR. Implications of venom hypersensitivity for a deploying soldier. *Mil Med* 2007;172:544-7.
[PUBMED](#) | [CROSSREF](#)
36. Mikals K, Beakes D, Banks TA. Stinging the conscience: a case of severe hymenoptera anaphylaxis and the need for provider awareness. *Mil Med* 2016;181:e1400-3.
[PUBMED](#) | [CROSSREF](#)
37. Baker TW, Stolfi A, Johnson TL. Use of epinephrine autoinjectors for treatment of anaphylaxis: which commercially available autoinjector do patients prefer? *Ann Allergy Asthma Immunol* 2009;103:356-8.
[PUBMED](#) | [CROSSREF](#)
38. Haymore BR, Carr WW, Frank WT. Anaphylaxis and epinephrine prescribing patterns in a military hospital: underutilization of the intramuscular route. *Allergy Asthma Proc* 2005;26:361-5.
[PUBMED](#)
39. Lee SM, Ahn JS, Noh CS, Lee SW. Prevalence of allergic diseases and risk factors of wheezing in Korean military personnel. *J Korean Med Sci* 2011;26:201-6.
[PUBMED](#) | [CROSSREF](#)
40. Matricardi PM, Kleine-Tebbe J, Hoffmann HJ, Valenta R, Hilger C, Hofmaier S, Aalberse RC, Agache I, Asero R, Ballmer-Weber B, Barber D, Beyer K, Biedermann T, Bilò MB, Blank S, Bohle B, Bosshard PP, Breiteneder H, Brough HA, Caraballo L, Caubet JC, Cramer R, Davies JM, Douladiris N, Ebisawa

- M, Elgenmann PA, Fernandez-Rivas M, Ferreira F, Gadermaier G, Glatz M, Hamilton RG, Hawranek T, Hellings P, Hoffmann-Sommergruber K, Jakob T, Jappe U, Jutel M, Kamath SD, Knol EF, Korosec P, Kuehn A, Lack G, Lopata AL, Mäkelä M, Morisset M, Niederberger V, Nowak-Węgrzyn AH, Papadopoulos NG, Pastorello EA, Pauli G, Platts-Mills T, Posa D, Poulsen LK, Raulf M, Sastre J, Scala E, Schmid JM, Schmid-Grendelmeier P, van Hage M, van Ree R, Vieths S, Weber R, Wickman M, Muraro A, Ollert M. EAACI Molecular Allergology User's Guide. *Pediatr Allergy Immunol* 2016;27 Suppl 23:1-250.
[PUBMED](#) | [CROSSREF](#)
41. Pascal M, Grishina G, Yang AC, Sánchez-García S, Lin J, Towle D, Ibañez MD, Sastre J, Sampson HA, Ayuso R. Molecular diagnosis of shrimp allergy: efficiency of several allergens to predict clinical reactivity. *J Allergy Clin Immunol Pract* 2015;3:521-9.e10.
[PUBMED](#) | [CROSSREF](#)
42. Fernández-Rivas M. Fruit and vegetable allergy. *Chem Immunol Allergy* 2015;101:162-70.
[PUBMED](#) | [CROSSREF](#)
43. Popescu FD. Cross-reactivity between aeroallergens and food allergens. *World J Methodol* 2015;5:31-50.
[PUBMED](#) | [CROSSREF](#)
44. Stokes S, Hudson S. Managing anaphylaxis in a jungle environment. *Wilderness Environ Med* 2012;23:51-5.
[PUBMED](#) | [CROSSREF](#)
45. Gernez Y, Nowak-Węgrzyn A. Immunotherapy for food allergy: are we there yet? *J Allergy Clin Immunol Pract* 2017;5:250-72.
[PUBMED](#) | [CROSSREF](#)
46. Martorell A, Alonso E, Echeverría L, Escudero C, García-Rodríguez R, Blasco C, Bone J, Borja-Segade J, Bracamonte T, Claver A, Corzo JL, De la Hoz B, Del Olmo R, Domínguez O, Fuentes-Aparicio V, Guallar I, Larramona H, Martín-Muñoz F, Matheu V, Michavila A, Ojeda I, Ojeda P, Piquer M, Poza P, Reche M, Rodríguez Del Río P, Rodríguez M, Ruano F, Sánchez-García S, Terrados S, Valdesoiro L, Vazquez-Ortiz M. Expert panel selected from members of the Spanish Society of Pediatric Allergology, Asthma and Clinical Immunology (SEICAP) and the Spanish Society of Allergology and Clinical Immunology (SEAIC). Oral Immunotherapy for Food Allergy: A Spanish Guideline. *Immunotherapy Egg and Milk Spanish Guide (ITEMS Guide)*. Part I: cow milk and egg oral immunotherapy: introduction, methodology, rationale, current state, indications, contraindications, and oral immunotherapy build-up phase. *J Investig Allergol Clin Immunol* 2017;27:225-37.
[PUBMED](#) | [CROSSREF](#)
47. Ebisawa M, Ito K, Fujisawa T Committee for Japanese Pediatric Guideline for Food Allergy, The Japanese Society of Pediatric Allergy and Clinical Immunology, The Japanese Society of Allergology. Japanese guidelines for food allergy 2017. *Allergol Int* 2017;66:248-64.
[PUBMED](#) | [CROSSREF](#)
48. Cortellini G, Spadolini I, Santucci A, Cova V, Conti C, Corvetta A, Passalacqua G. Improvement of shrimp allergy after sublingual immunotherapy for house dust mites: a case report. *Eur Ann Allergy Clin Immunol* 2011;43:162-4.
[PUBMED](#)
49. Wai CY, Leung NY, Leung PS, Chu KH. T cell epitope immunotherapy ameliorates allergic responses in a murine model of shrimp allergy. *Clin Exp Allergy* 2016;46:491-503.
[PUBMED](#) | [CROSSREF](#)