

Racial Differences in Tuberculosis Infection in United States Communities: The Coronary Artery Risk Development in Young Adults Study

Payam Nahid,¹ David J. Horne,² Leah G. Jarlsberg,¹ Alexander P. Reiner,³ Dennis Osmond,¹ Philip C. Hopewell,¹ and Kirsten Bibbins-Domingo⁴

¹Curry International Tuberculosis Center, Division of Pulmonary and Critical Care Medicine, San Francisco General Hospital, University of California, San Francisco, California; ²Division of Pulmonary and Critical Care Medicine, University of Washington, Seattle, Washington; ³Department of Epidemiology, School of Public Health, University of Washington, Seattle, Washington; and ⁴Division of General Internal Medicine, San Francisco General Hospital, University of California, San Francisco, California

Previously reported associations between race/ethnicity and tuberculosis infection have lacked sufficient adjustment for socioeconomic factors. We analyzed race/ethnicity and self-reported tuberculosis infection data from the Coronary Artery Risk Development in Young Adults (CARDIA) study, a well-characterized cohort of 5115 black and white participants, and found that after adjusting for sociodemographic and clinical factors, black participants were more likely to report tuberculosis infection and/or disease (odds ratio, 2.0; 95% confidence interval, 1.5–2.9).

Tuberculosis (TB) is among the leading conditions for which disparities in disease rates between blacks and white exist [1]. According to the Centers for Disease Control and Prevention (CDC), of the 4378 cases of TB reported in United States (US)-born patients in 2010, 40% were among non-Hispanic blacks, a rate 8 times higher than that among non-Hispanic whites [2]. Similarly, the prevalence of latent TB infection (LTBI) in US-born individuals is 5.7% among blacks, significantly higher than

the 2.5% prevalence in Mexican-Americans and 1.1% in whites [3]. The proposed causes of this disparity include environmental and host factors [4], as well as differences in comorbid conditions that affect risk of developing TB once infected, such as human immunodeficiency virus (HIV) infection [5]. In some studies, low socioeconomic status is reported to account for a significant proportion of the increased risk for TB seen in blacks in the US [6, 7]; however, other studies indicate that genetic susceptibility to infection may explain some of the racial differences in disease rates [8, 9]. Overall, racial disparities in TB are substantial in the US [10], and whether these differences are accounted for by socioeconomic status continues to be debated.

We used the Coronary Artery Risk Development in Young Adults (CARDIA) study, a population-based longitudinal study of risk factors for cardiovascular disease with a cohort of 5115 black and white participants, to determine whether black race is associated with TB infection after adjusting for socioeconomic status and other traditional TB risk factors. The thorough characterization of individual-level data on socioeconomic indicators of CARDIA participants provided an opportunity to control for this potential confounder.

METHODS

Study Population and Setting

Participants were unrelated men and women who participated in the CARDIA study, an ongoing epidemiologic study of cardiovascular risk development in young adults [11]. To be eligible for the CARDIA study, participants must have identified themselves as non-Hispanic white or black and as having a permanent address in 1 of 4 US urban communities (Birmingham, Alabama; Chicago, Illinois; Minneapolis, Minnesota; or Oakland, California). The CARDIA cohort was designed to be balanced by race, sex, education (less than high school vs high school graduation), and age. A total of 5115 individuals were recruited and examined at the baseline examination in 1985–1986 when participants were 18–30 years of age. Follow-up examinations occurred after 2, 5, 7, 10, 15, and 20 years. The CARDIA study has had a high retention rate at year 20, with 87.5% of the original cohort completing the annual telephone interview for outcome ascertainment and 71.8% completing the in-person examination [12]. The current analysis of risk factors for TB infection was conducted on 3112 participants for whom there were complete clinical and demographic data at baseline and follow-up examinations. The study was approved by the Committee on Human Research at the University of California,

Received 22 February 2011; accepted 29 April 2011.

Correspondence: Payam Nahid, MD, MPH, Curry International Tuberculosis Center, Division of Pulmonary and Critical Care Medicine, San Francisco General Hospital, University of California, San Francisco, 1001 Potrero Ave, 5K1, San Francisco, CA 94110 (pnahid@ucsf.edu).

Clinical Infectious Diseases 2011;53(3):291–294

© The Author 2011. Published by Oxford University Press on behalf of the Infectious Diseases Society of America. All rights reserved. For Permissions, please email: journals.permissions@oup.com. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/2.5/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

1058-4838/2011/533-0013\$14.00

DOI: 10.1093/cid/cir378

San Francisco, and all participating CARDIA centers. Written informed consent was obtained from participants at each examination.

Categorization of Tuberculosis Infection

At years 10, 15, and 20, all available CARDIA participants were asked: "Has a doctor or nurse ever said that you have tuberculosis or a positive skin test for tuberculosis?" Participants who answered yes to this question at any of these exams were categorized as being TB infected, whereas participants who reported consistent negative response to the question at exam years 10, 15, and 20 were categorized as TB uninfected.

In addition to age, sex, and race, predictors for TB infection analyzed included socioeconomic measures (household income, education, housing status, history of incarceration, and crowding [as measured by the reported number of household members]), as well as comorbid characteristics associated with risk for TB infection and disease including diabetes, smoking, hematologic cancers, kidney disease, and self-reported HIV status. All covariates and risk factors were assessed prior to reported TB infection so as to assure correct chronology in terms of risk and exposure and timing of TB infection.

Statistical Analysis

The definitions and primary objectives of this study were formulated prior to data analysis. All statistical analysis was done using SAS software, version 9.2. Associations with TB infection were adjusted using multivariate logistic regression models. Predictors with $P < .2$ in bivariate analysis were included in the multivariate model. Because of differential dropout of participants with characteristics of importance to the study, we performed analyses using stabilized inverse probability-of-censoring weights to reduce potential bias. Weights were developed using all the characteristics associated with dropout, which included black race, male sex, site of enrollment, incarceration history, self-report of ever smoking, and having less than a high school education at baseline [13].

RESULTS

Of the 5115 participants initially enrolled in CARDIA, 1410 black participants (53.5% of all black participants) and 1702 white participants (68.7% of all white participants) had complete demographic and clinical data for this analysis and were included in this study. Of these, 185 (5.9%) reported a history of TB infection or disease, which corresponded to 8.5% of black participants and 3.8% of white participants (unadjusted odds ratio [OR], 2.3; 95% confidence interval [CI], 1.7–3.2). The strongest associations with TB infection, other than race, included reported HIV infection, smoking, household income, home ownership, and household size as measured by the reported number of household members (Table 1).

From the multivariate logistic regression model for characteristics associated with TB infection shown in Table 1, black race and HIV infection were each independently and positively associated with TB infection ($P < .05$).

Black participants, and participants with the characteristics of male gender, site of enrollment in Minnesota, incarceration history, self-report of ever smoking, and having less than a high school education at baseline, were more likely to be lost to follow-up in the CARDIA study. To address potential bias resulting from differential dropout, we performed our analysis using stabilized inverse probability weights; our independent predictors of TB infection, including black race (OR, 2.0; 95% CI, 1.4–3.0), remained unchanged (analysis not shown).

DISCUSSION

In this study, we identified racial differences in the reported rate of TB infection in participants enrolled in the CARDIA study, an ongoing epidemiologic study of coronary artery risk development in young black and white adults. The CARDIA study collects individual-level data on a variety of socioeconomic indicators, including level of education, housing status, household size, and income level. In our analyses, the increased risk of TB infection in black participants persisted after adjusting for these socioeconomic indicators. The increased risk for TB infection in black participants also persisted after adjusting for other known risk factors for TB such as HIV infection, diabetes, renal disease, hematologic cancers, intravenous drug use, and incarceration, with black participants having a 2-fold increased odds of self-reported TB infection compared with white participants ($P < .001$). Our findings suggest that factors other than comorbid conditions and socioeconomic status may contribute to the racial disparity seen in TB rates between blacks and whites in the US [1]. Whether this disparity is due to unmeasured factors related to race as a social construct or due to biologic susceptibility factors is still subject to debate. An analysis of the influence of race/ethnicity on rates of active TB in the United States found that adjusting for 6 socioeconomic indicators accounted for approximately half the increased risk of TB previously ascribed to race/ethnicity among US-born blacks, Hispanics, and Native Americans [6]. However, this study used ecologic-level data for socioeconomic indicators, specifically US Census data and zip code-specific demographic substratum values, whereas our study used individual-level data. Taken together, the numerous publications that show a link between race and increased risk of TB in US-born populations raise the question of whether targeted testing and treatment for LTBI should be recommended by race. Whether such a strategy would be cost-effective and whether it would have the potential to address the persistent disparity seen in TB between blacks and whites in the United States deserves further study.

Table 1. Predictors of Tuberculosis Infection in 3112 Participants Enrolled in CARDIA

	Tuberculosis infected (<i>n</i> = 185)	Unadjusted OR (95% CI), <i>P</i> value	Adjusted OR (95% CI), <i>P</i> value
Race			
Black	8.5%	2.3 (1.7–3.2), <.001	2.0 (1.5–2.9), <.001
White	3.8%		
Sex			
Male	5.6%	0.9 (0.6–1.2), .42	
Female	6.2%		
Enrollment site			
Oakland, CA	5.4%	1.0	
Minneapolis, MN	4.8%	0.9 (0.6–1.4), .61	
Birmingham, AL	6.5%	1.2 (0.8–1.9), .34	
Chicago, IL	7.4%	1.4 (0.9–2.1), .11	1.3 (0.9–1.8), .11
Mean BMI at year 10 ± SD (odds per 5-unit increase)	28 ± 7	1.1 (1.0–1.2), .05	1.0 (0.9–1.2), .52
Ever self-reported kidney disease^a			
Yes	6.9%	1.2 (0.8–1.9), .44	
No	5.8%		
Fasting glucose >125 mg/dL at year 10			
Yes	1.8%	0.3 (0.0–2.0), .21	
No	6.0%		
Ever self-reported diabetes mellitus^a			
Yes	6.0%	1.0 (0.5–2.0), .96	
No	5.9%		
Ever self-reported hematologic cancer^b			
Yes	6.8%	1.1 (0.4–3.2), .78	
No	5.9%		
Ever self-reported HIV infection^c			
Yes	20.0%	4.0 (1.3–12.2), .01	3.4 (1.1–10.9), .04
No	5.9%		
Ever self-reported injection drug use^b			
Yes	9.1%	1.6 (0.8–3.3), .18	1.6 (0.8–3.3), .22
No	5.8%		
Ever self-reported incarceration history^d			
Yes	6.9%	1.2 (0.6–2.2), .61	
No	5.9%		
Ever self-reported smoking^a			
Yes	6.5%	1.2 (0.9–1.6), .20	
No	5.5%		
Education at year 10			
7–12 years (High school or less)	6.5%	1.1 (0.7–1.7), .68	
13–16 years (Any college)	5.7%	1.0 (0.7–1.4), .84	
>16 years (Graduate school)	5.9%	1.0	
Household income at year 10^e			
<2 times FPL	8.1%	1.8 (1.2–2.6), .001	1.0 (0.7–1.6), .91
2–4 times FPL	7.1%	1.6 (1.1–2.2), .01	1.2 (0.8–1.7), .32
>4 times FPL	4.6%	1.0	1.0
Mean number of people in household ± SD at year 10 (odds per 1-person increase)	3.3 ± 1.7	1.1 (1.0–1.2), .02	1.1 (1.0–1.2), .09
Homeowner at year 10			
Yes	4.9%	0.6 (0.5–0.8), .002	0.7 (0.5–1.0), .08
No	7.5%		

NOTE. BMI, body mass index; CARDIA, Coronary Artery Risk Development in Young Adults study; CI, confidence interval; FPL, federal poverty line; HIV, human immunodeficiency virus; OR, odds ratio. Only variables with *P* value <.2 were included in the multivariate (adjusted) model.

^a Responses are cumulative from years 0, 2, 5, 7, and 10 and precede assessment for tuberculosis.

^b Question was not added to the survey until year 5. Responses are from years 5, 7, and 10.

^c Question regarding HIV history was not added to the survey until year 15. Responses are from years 15 and 20.

^d Question regarding incarceration history was only on the survey during years 0 and 2.

^e FPL calculation is based on poverty thresholds adjusted for household size, United States Census Bureau, 1995 (year 10 of the CARDIA study).

Our study has limitations. First, CARDIA was focused on cardiovascular disease and consequently did not pursue diagnostic testing to confirm self-reports of HIV and TB. To address this, we limited our analysis to participants who provided consistent responses to the TB questions. Second, we could not clearly distinguish between patients with LTBI and those who may have had active TB. Consequently, we cannot comment on susceptibility to infection versus progression to disease. Third, although guidelines do not list race/ethnicity as indications for LTBI testing [14], it is possible that there was differential testing for TB based on race, as has been reported for other primary care procedures [15]. However, the increased risk for LTBI in non-Hispanic blacks reported in an analysis of National Health and Nutrition Examination Survey data on 7386 participants with tuberculin skin test results suggests that the association between race and TB is not an artifact of increased testing [3]. We attempted to address this issue by controlling for characteristics that disproportionately affect blacks in the United States and may be indications for LTBI testing, including incarceration history and housing status, and race retained its association with self-report of TB. Nonetheless, differential testing as well as residual confounding may still exist.

In summary, we have shown that within a prospective, ongoing epidemiologic study of coronary artery disease in young black and white adults in 4 US communities, black study participants were 2 times more likely to report a history of TB infection compared with white participants after adjusting for a variety of socioeconomic factors, HIV status, intravenous drug use, diabetes, renal disease, and other traditional risk factors for TB. Our study suggests that factors other than socioeconomic status also contribute to the racial disparity in tuberculosis noted in US-born blacks.

Acknowledgments

The authors acknowledge and are grateful for the editorial assistance of Amy Markowitz, who was supported by National Institutes of Health/National Center for Research Resources University of California San Francisco-Clinical Translational Science Institute grant (UL1 RR024131).

Financial support. Work on this manuscript was supported by the National Institutes of Health through National Heart, Lung, and Blood Institute grants (K23HL092629 and F32HL094031) and contracts (University of Alabama at Birmingham, Coordinating Center, N01-HC-95095; University of Alabama at Birmingham, Field Center, N01-HC-48047; University of Minnesota, Field Center and Diet Reading Center [year 20 Exam], N01-HC-48048; Northwestern University, Field Center, N01-HC-48049; Kaiser Foundation Research Institute, N01-HC-48050; University of California, Irvine, Echocardiography Reading Center [years 5 and 10], N01-HC-45134;

Harbor-UCLA Research Education Institute, Computed Tomography Reading Center [year 15 Exam], N01-HC-05187; Wake Forest University [year 20 Exam], N01-HC-45205; New England Medical Center [year 20 Exam], N01-HC-45204), as well as a National Institute of Allergy and Infectious Diseases grant (AI034238).

Potential conflicts of interest. P. N. has received grant support to his institution from the National Institutes of Health (NIH) and the Centers for Disease Control and Prevention (CDC). P. H. has received grant support to his institution from NIH, CDC, and the World Health Organization. All other authors: No reported conflicts.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed in the Acknowledgments

References

1. Keppel KG, Percy JN, Wagener DK. Trends in racial and ethnic-specific rates for the health status indicators: United States, 1990-98. *Healthy People 2000 Stat Notes* **2002**; 23:1-16.
2. Trends in tuberculosis—United States, 2010. *MMWR Morb Mortal Wkly Rep* **2011**; 60:333-7.
3. Bennett DE, Courval JM, Onorato I, et al. Prevalence of tuberculosis infection in the United States population: the National Health and Nutrition Examination Survey, 1999-2000. *Am J Respir Crit Care Med* **2008**; 177:348-55.
4. Holmes CB, Hausler H, Nunn P. A review of sex differences in the epidemiology of tuberculosis. *Int J Tuberc Lung Dis* **1998**; 2:96-104.
5. Racial disparities in tuberculosis—selected southeastern states, 1991-2002. *MMWR Morb Mortal Wkly Rep* **2004**; 53:556-9.
6. Cantwell MF, McKenna MT, McCray E, Onorato IM. Tuberculosis and race/ethnicity in the United States: impact of socioeconomic status. *Am J Respir Crit Care Med* **1998**; 157(Pt 1):1016-20.
7. McKenna MT, McCray E, Jones JL, Onorato IM, Castro KG. The fall after the rise: tuberculosis in the United States, 1991 through 1994. *Am J Public Health* **1998**; 88:1059-63.
8. Stead WW, Senner JW, Reddick WT, Lofgren JP. Racial differences in susceptibility to infection by *Mycobacterium tuberculosis*. *N Engl J Med* **1990**; 322:422-7.
9. Crowle AJ, Elkins N. Relative permissiveness of macrophages from black and white people for virulent tubercle bacilli. *Infect Immun* **1990**; 58:632-8.
10. O'Donnell MR, Chamblee S, von Reyn CF, et al. Racial disparities in primary and reactivation tuberculosis in a rural community in the southeastern United States. *Int J Tuberc Lung Dis* **2010**; 14:733-40.
11. Friedman GD, Cutter GR, Donahue RP, et al. CARDIA: study design, recruitment, and some characteristics of the examined subjects. *J Clin Epidemiol* **1988**; 41:1105-16.
12. Bibbins-Domingo K, Pletcher MJ, Lin F, et al. Racial differences in incident heart failure among young adults. *N Engl J Med* **2009**; 360:1179-90.
13. Alonso A, Segui-Gomez M, de Irala J, Sanchez-Villegas A, Beunza JJ, Martinez-Gonzalez MA. Predictors of follow-up and assessment of selection bias from dropouts using inverse probability weighting in a cohort of university graduates. *Eur J Epidemiol* **2006**; 21:351-8.
14. American Thoracic Society. Targeted tuberculin testing and treatment of latent tuberculosis infection. *MMWR Recomm Rep* **2000**; 49:1-51.
15. Franks P, Fiscella K, Meldrum S. Racial disparities in the content of primary care office visits. *J Gen Intern Med* **2005**; 20:599-603.