

RESEARCH

Open Access

Mild hypothermia alone or in combination with anesthetic post-conditioning reduces expression of inflammatory cytokines in the cerebral cortex of pigs after cardiopulmonary resuscitation

Patrick Meybohm^{1*}, Matthias Gruenewald¹, Kai D Zacharowski², Martin Albrecht¹, Ralph Lucius³, Nikola Fösel¹, Johannes Hensler¹, Karina Zitta¹, Berthold Bein¹

Abstract

Introduction: Hypothermia improves survival and neurological recovery after cardiac arrest. Pro-inflammatory cytokines have been implicated in focal cerebral ischemia/reperfusion injury. It is unknown whether cardiac arrest also triggers the release of cerebral inflammatory molecules, and whether therapeutic hypothermia alters this inflammatory response. This study sought to examine whether hypothermia or the combination of hypothermia with anesthetic post-conditioning with sevoflurane affect cerebral inflammatory response after cardiopulmonary resuscitation.

Methods: Thirty pigs (28 to 34 kg) were subjected to cardiac arrest following temporary coronary artery occlusion. After seven minutes of ventricular fibrillation and two minutes of basic life support, advanced cardiac life support was started according to the current American Heart Association guidelines. Return of spontaneous circulation was achieved in 21 animals who were randomized to either normothermia at 38°C, hypothermia at 33°C or hypothermia at 33°C combined with sevoflurane (each group: n = 7) for 24 hours. The effects of hypothermia and the combination of hypothermia with sevoflurane on cerebral inflammatory response after cardiopulmonary resuscitation were studied using tissue samples from the cerebral cortex of pigs euthanized after 24 hours and employing quantitative RT-PCR and ELISA techniques.

Results: Global cerebral ischemia following resuscitation resulted in significant upregulation of cerebral tissue inflammatory cytokine mRNA expression (mean \pm SD; interleukin (IL)-1 β 8.7 \pm 4.0, IL-6 4.3 \pm 2.6, IL-10 2.5 \pm 1.6, tumor necrosis factor (TNF) α 2.8 \pm 1.8, intercellular adhesion molecule-1 (ICAM-1) 4.0 \pm 1.9-fold compared with sham control) and IL-1 β protein concentration (1.9 \pm 0.6-fold compared with sham control). Hypothermia was associated with a significant ($P < 0.05$ versus normothermia) reduction in cerebral inflammatory cytokine mRNA expression (IL-1 β 1.7 \pm 1.0, IL-6 2.2 \pm 1.1, IL-10 0.8 \pm 0.4, TNF α 1.1 \pm 0.6, ICAM-1 1.9 \pm 0.7-fold compared with sham control). These results were also confirmed for IL-1 β on protein level. Experimental settings employing hypothermia in combination with sevoflurane showed that the volatile anesthetic did not confer additional anti-inflammatory effects compared with hypothermia alone.

Conclusions: Mild therapeutic hypothermia resulted in decreased expression of typical cerebral inflammatory mediators after cardiopulmonary resuscitation. This may confer, at least in part, neuroprotection following global cerebral ischemia and resuscitation.

* Correspondence: meybohm@anaesthesie.uni-kiel.de

¹Department of Anaesthesiology and Intensive Care Medicine, University Hospital Schleswig-Holstein, Campus Kiel, Schwannenweg 21, Kiel, 24105, Germany

Introduction

Although initial return of spontaneous circulation (ROSC) from cardiac arrest is achieved in about 30 to 40% of cases, only 10 to 30% of the patients admitted to the hospital will be discharged with good outcome [1]. One third of those who survive, suffer persistent neurological impairments [2]. Mild therapeutic hypothermia has emerged as the most effective strategy to reduce neurological impairment after successful cardiopulmonary resuscitation (CPR) [3]. The precise mechanisms by which mild hypothermia protects brain cells remain to be elucidated, but it is very likely that hypothermia acts upon multiple pathways including reduction in cerebral metabolism and oxygen consumption, attenuation of neuronal damage, and inhibition of excitatory neurotransmitter release [4].

There is growing evidence on the damaging nature of the inflammatory response following brain ischemia. Various inflammatory cytokines have been implicated as important mediators of ischemia/reperfusion injury following both focal and global cerebral ischemia [5]. Most of the previous experimental studies induced global cerebral ischemia by bilateral carotid artery occlusion as a correlate of cardiac arrest, but inflammatory response mechanisms following carotid artery occlusion and anti-inflammatory mechanisms of hypothermia may be different from those observed after cardiac arrest and manual CPR. Thus, it is unknown whether cardiac arrest also triggers the release of cerebral inflammatory molecules, and whether therapeutic hypothermia alters this inflammatory response.

Neuronal injury may also result in necrotic and apoptotic cell death. In contrast to necrosis (cell death by acute injury), apoptosis is a well-regulated physiological process. Cells undergoing apoptosis are characterized by cytoplasmic shrinkage, nuclear condensation, and formation of membrane-bound vesicles. Key elements of the apoptotic pathway include changes in the gene expression of the pro-apoptotic protein Bax and the apoptosis-suppressing protein Bcl-2. The extent to which hypothermia affects cerebral apoptosis-related proteins after successful CPR is not clear [4].

The mismatch between early survival and final outcome after CPR emphasizes the importance of further research on potential adjuvants in addition to mild hypothermia. Specifically, pharmacological post-conditioning may offer an attractive opportunity to further ameliorate damage to the brain in the post-resuscitation period. While the volatile anesthetic sevoflurane has emerged as a pre-conditioning-like agent with significant neuroprotective effects in models of focal and global cerebral ischemia [6], its potential neuroprotective and anti-inflammatory properties have not yet been investigated in the context of post-resuscitation care. Thus, a

combination of hypothermia and anesthetic post-conditioning with sevoflurane may extend neuroprotection, as it has recently been shown for the noble anesthetic gas xenon combined with hypothermia after neonatal hypoxia-ischemia [7].

We hypothesized that hypothermia attenuates cerebral inflammatory response in a pig model of global cerebral ischemia following cardiac arrest. We further hypothesized that the volatile anesthetic sevoflurane, when administered during reperfusion after successful CPR, confers additional anti-inflammatory effects.

Materials and methods

The project was approved by the Animal Investigation Committee of the University Schleswig-Holstein, Campus Kiel, Germany, and animals were managed in accordance with the guidelines of the University Schleswig-Holstein, Campus Kiel, Germany, and the Utstein-style guidelines [8]. All animals received human care in compliance with the *Guide for the Care and Use of Laboratory Animals* published by the National Institute of Health (NIH Publication No. 88.23, revised 1996).

Animals

This is an experimental study on 40 healthy pigs (cardiac arrest: n = 30; sham control: n = 5; excluded from study n = 5) aged three to four months of both gender, weighing 28 to 34 kg. Anesthesia was initiated by intramuscular injection of 8 mg/kg azaperone and 0.05 mg/kg atropine, and completed by intravenous injection of 1 to 2 mg/kg propofol and 0.3 µg/kg sufentanil. After endotracheal intubation, pigs were ventilated with a volume-controlled ventilator (Draeger, Sulla 808V, Luebeck, Germany) and the following setting: a FiO₂ of 0.3 at 20 breaths/minute, a tidal volume of 8 mL/kg to maintain normocapnia, and a positive end-expiratory pressure of 5 mm Hg. Ventilation was monitored using an inspired/expired gas analyzer that measured oxygen and end-tidal carbon dioxide (suction rate, 200 mL/min; M-PRESTN; Datex-Ohmeda Inc., Helsinki, Finland). Total intravenous anesthesia (TIVA) was maintained by continuous infusion of 4 to 8 mg/kg/h propofol and 0.3 µg/kg/h sufentanil; muscle relaxation was achieved by continuous infusion of 0.2 mg/kg/h pancuronium. Depth of anesthesia was judged according to blood pressure, heart rate and Bispectral Index (BISXP, Aspect Medical Systems, Natick, MA, USA) [9]. In order to assure an appropriate depth of anesthesia we performed also indirect measures such as tail clamping, monitoring of the corneal reflex and lacrimation, as well as changes in hemodynamics and heart rate. If assessment suggested inadequate level of anesthesia, additional sufentanil and propofol was injected. Ringer's solution was

administered continuously throughout the preparation phase to replace fluid loss during instrumentation. Standard leads II and V₅ electrocardiogram were used to monitor cardiac rhythm.


A 7F saline-filled central venous catheter was inserted in the right internal jugular vein for drug administration. A 4F thermistor-tipped catheter for arterial thermodilution (Pulsion Medical Systems, Munich, Germany) was inserted percutaneously into the right femoral artery. The arterial catheter was connected to the PiCCO system (PiCCO plus, Software Version 6.0, Pulsion Systems, Munich, Germany), and the resulting signal processed to determine mean arterial blood pressure, heart rate, and blood temperature. In addition, the arterial catheter allowed discontinuous measurement of transpulmonary cardiac output by injecting 10 mL ice cold saline into the proximal port of the central venous catheter. The mean of three consecutive measurements randomly assigned to the respiratory cycle was used for determination of cardiac output. Cardiac index was calculated as the ratio of cardiac output/body surface area (body surface area = 0.0734*(body weight in kg)^{0.656} [10]). Intravascular catheters were attached to pressure transducers (Smiths Medical, Kirchseeon, Germany) that were aligned at the level of the right atrium.

Experimental protocol

The experimental time line is presented in Figure 1. Because the majority of patients experience cardiac arrest due to myocardial ischemia [11], and because this scenario has only been considered in few animal experiments, our study is based on an experimental porcine model of cardiac arrest following acute coronary artery ischemia reflecting a realistic clinical setting. Five healthy animals served as sham controls, which were anesthetized with TIVA until the end of the experiment. Thirty-five pigs underwent left anterior descending (LAD) coronary artery occlusion for 60 minutes according to the technique previously described [12]. Five pigs fibrillated spontaneously following left anterior descending coronary artery occlusion, which were excluded from further analysis. Thirty pigs were then subjected to cardiac arrest 20 minutes after LAD occlusion. Ventricular fibrillation was electrically-induced by an alternating current of 5 to 10 V in a standardized manner, and mechanical ventilation was discontinued. After a seven-minute non-intervention interval of untreated ventricular fibrillation, basic life support CPR was simulated for two minutes applying external manual closed chest compressions at a rate of 100 per minute, and a compression-to-ventilation ratio of 30:2. Subsequently, advanced cardiac life support was started with 100 J biphasic defibrillation attempt (M-Series Defibrillators, Zoll Medical Corporation, Chelmsford, Massachusetts,

USA), all subsequent attempts were performed with 150 J every two minutes. Ventilations were performed with 100% oxygen at 20 breaths/minute. All pigs received 45 µg/kg epinephrine and 0.4 U/kg vasopressin alternating as suggested by the American Heart Association guidelines [13]. ROSC was defined as maintenance of an unassisted pulse and a systolic aortic blood pressure of ≥60 mm Hg lasting for 10 consecutive minutes according to the Utstein-style guidelines [8]. Since neurological recovery is very unlikely after 30 minutes of normothermic cardiac arrest, CPR was terminated, when resuscitation remained unsuccessful after 23 minutes of CPR. After ROSC, animals were randomized either to normothermia (38°C) plus TIVA (NT), hypothermia (33°C) plus TIVA (HT), or hypothermia (33°C) combined with 2.0 Vol% end-tidal sevoflurane and 0.3 µg/kg/h sufentanil (HT+SEV). Since hypothermia was shown to increase blood concentrations of propofol by about 30% [14], we reduced continuous infusion of propofol during hypothermia targeting bispectral index values below 60. Body core temperature was monitored continuously by the arterial catheter, and normothermic body temperature was maintained at 38.0°C with a heating blanket, since the physiological rectal temperature of pigs is supposed to be about 38°C [15]. Hypothermia was induced by 1,000 mL saline (4°C) and maintained by a cooling device (Icy catheter and CoolGard 3000; Alsius Corp, Irvine, CA, USA) that was introduced into the femoral vein. According to the landmark study by Bernard et al. [16] we used a target body temperature of 33°C for 12 hours. Thereafter, re-warming was initiated (0.5°C per hour). One hour after ROSC, FiO₂ was reduced to 0.4. During the post-resuscitation period, animals received crystalloid infusions to keep central venous pressure above 8 mm Hg and mean arterial blood pressure above 50 mm Hg. If this first step failed, additional norepinephrine was administered to keep mean arterial blood pressure above 50 mm Hg. We further aimed at serum glucose levels less than 150 mg/dL by intermittent insulin bolus administration. Animals were killed by an overdose of sufentanil, propofol and potassium chloride 24 hours after ROSC. Tissue samples of the cerebral cortex were collected within 15 seconds following euthanasia *via* a craniotomy that was established before euthanasia, and then immediately snap-frozen in liquid nitrogen (stored at -80°C) to minimize time-dependent effects of cerebral ischemia following euthanasia on cytokine expression. Autopsy was routinely performed for documentation of potential injuries to the thoracic and abdominal cavity during CPR.

Hemodynamic data, including mean arterial blood pressure, heart rate, end-tidal carbon dioxide, and cardiac index were determined at baseline (BL), following ROSC, and 7 and 24 hours after ROSC, respectively.



Quantitative real-time RT-PCR

Transcript levels of interleukin (IL)-1 β , IL-6, IL-10, tumor necrosis factor (TNF) α , intercellular adhesion molecule (ICAM)-1, and the apoptosis-associated proteins Bcl-2 and Bax were investigated in the cerebral cortex tissue of all surviving animals and compared with tissue of sham control animals. Tissue samples were analyzed by a person blinded to treatment assignment. Fully detailed description of quantitative real-time RT-PCR is presented in the Additional File 1 and Table S1 [17-20].

Enzyme-linked immunosorbent assay (ELISA)

Protein concentrations of IL-1 β were determined by a swine specific ELISA (BioSource International, Inc. Camarillo, CA, USA) in homogenates of frozen tissues according to the manufacturer's protocol. All ELISA assays were carried out in duplicates.

Statistical analysis

Statistics were performed using commercially available statistics software (GraphPad Prism version 5.02 for Windows, GraphPad Software, San Diego, CA, USA). Survival rates were compared using Fisher's exact test. Statistical analysis was performed with a one-way analysis of variance (ANOVA) followed by a Bonferroni post hoc test to correct for multiple measurements. RT-PCR data analysis was performed according to a relative standard curve method using an Excel spreadsheet, and statistical significance was tested using two-sided Pair-wise fixed Reallocation Randomisation Test, as provided in the REST2005 program [20]. The Mann-Whitney test was used for analysis of protein concentrations of IL-1 β where normal distribution was not expected. Variables

are expressed as mean \pm SD unless otherwise specified. Statistical significance was considered at a two-sided *P* value of ≤ 0.05 .

Results

Cardio-pulmonary resuscitation

Twenty-one animals were successfully resuscitated. Detailed resuscitation data are presented in Table 1. In the NT group, five out of seven animals survived for 24 hours compared to all animals in the HT and HT+SEV group (*P* = 0.46 vs. NT). Two animals of the NT group died due to hemodynamic instability during the post-resuscitation period.

Post-resuscitation hemodynamics

Post-resuscitation systemic hemodynamic variables are presented in Table 2. Heart rate, mean arterial blood pressure and cardiac index did not significantly differ between groups. Cumulative crystalloid fluid load and cumulative norepinephrine doses were not significantly different between groups 24 hours after ROSC (volume load (*P* = 0.540), norepinephrine doses (*P* = 0.812); NT: 4241 \pm 1244 mL, 4.4 \pm 1.6 mg; HT: 3987 \pm 932 mL, 4.9 \pm 2.1 mg; HT+SEV: 4627 \pm 1056 mL, 5.1 \pm 1.8 mg).

Cerebral inflammatory response

Global cerebral ischemia following resuscitation resulted in a significant upregulation of cerebral tissue inflammatory cytokine mRNA expression (NT: IL-1 β 8.7 \pm 4.0, IL-6 4.3 \pm 2.6, IL-10 2.5 \pm 1.6, TNF α 2.8 \pm 1.8, ICAM-1 4.0 \pm 1.9-fold compared with sham control) and IL-1 β protein concentration (1.9 \pm 0.6-fold compared with sham control). Hypothermia was associated with significantly (*P* < 0.05 versus

Table 1 Cardiopulmonary resuscitation data

	NT	HT	HT+SEV	P values
ROSC rate [n]	7/10	7/10	7/10	-
CPR time to successful resuscitation [min]	9.7 ± 2.8	10.3 ± 3.4	10.5 ± 3.1	0.939
Cumulative epinephrine dose [µg/kg]	100 ± 44	101 ± 47	93 ± 33	0.828
Cumulative vasopressin dose [IU/kg]	0.8 ± 0.2	0.8 ± 0.3	0.8 ± 0.3	0.897
Cumulative defibrillation energy [J]	755 ± 420	703 ± 413	795 ± 199	0.854
CorPP 10 [mm Hg]	31 ± 9	26 ± 11	28 ± 6	0.559
CorPP 15 [mm Hg]	39 ± 28	40 ± 25	38 ± 24	0.890
Time to target temperature 33°C [min]	-	47 ± 10	45 ± 15	-

ROSC, return of spontaneous circulation; CPR, cardiopulmonary resuscitation. Time to successful resuscitation, cumulative epinephrine and vasopressin dose, cumulative defibrillation energy, coronary perfusion pressure (CorPP) 10 and 15 minutes after induction of ventricular fibrillation, and induction time to target temperature of 33°C. NT indicates normothermia; HT, hypothermia; HT+SEV, hypothermia combined with sevoflurane. Data are mean ± SD.

normothermia) less upregulation of mRNA expression (IL-1β 1.7 ± 1.0, IL-6 2.2 ± 1.1, IL-10 0.8 ± 0.4, TNFα 1.1 ± 0.6, ICAM-1 1.9 ± 0.7-fold compared with sham control) and IL-1β protein concentration (1.3 ± 0.4-fold compared with sham control). Sevoflurane did not confer statistically significant (versus hypothermia) additional protective effects neither on mRNA (IL-1β 1.2 ± 0.6, IL-6 2.0 ± 0.9, IL-10 0.7 ± 0.3, TNFα 0.9 ± 0.4, ICAM-1 1.8 ± 0.6-fold compared with sham control) nor on protein levels (1.1 ± 0.2-fold compared with sham control; Figures 2 and 3).

Table 2 Hemodynamic data

	NT	HT	HT+SEV	P value
Baseline				
HR, beats/minute	107 ± 21	105 ± 14	96 ± 15	0.342
MAP, mm Hg	65 ± 13	70 ± 11	71 ± 13	0.314
ETCO ₂ , mm Hg	40 ± 5	36 ± 4	37 ± 5	0.180
CI, L/min/m ²	7.4 ± 1.8	6.8 ± 1.3	7.4 ± 1.7	0.580
ROSC				
HR, beats/minute	94 ± 18	99 ± 33	96 ± 21	0.988
MAP, mm Hg	55 ± 6	65 ± 21	60 ± 8	0.225
ETCO ₂ , mm Hg	39 ± 9	42 ± 4	39 ± 7	0.363
CI, L/min/m ²	4.4 ± 0.5	4.5 ± 1.9	4.6 ± 1.0	0.982
Seven hours ROSC				
HR, beats/minute	131 ± 17	140 ± 22	127 ± 20	0.821
MAP, mm Hg	58 ± 4	59 ± 12	61 ± 8	0.820
ETCO ₂ , mm Hg	37 ± 7	35 ± 4	35 ± 2	0.731
CI, L/min/m ²	6.2 ± 0.4	5.5 ± 1.1	7.1 ± 1.2	0.071
24 hours ROSC				
HR, beats/minute	154 ± 25	143 ± 19	124 ± 24	0.297
MAP, mm Hg	46 ± 6	57 ± 12	54 ± 4	0.249
ETCO ₂ , mm Hg	37 ± 4	41 ± 1	37 ± 2	0.180
CI, L/min/m ²	5.6 ± 0.2	7.1 ± 1.6	8.7 ± 1.8	0.139

Hemodynamic data were determined at baseline, following return of spontaneous circulation (ROSC), and 7 and 24 hours after ROSC. HR indicates heart rate; MAP, mean arterial blood pressure; ETCO₂, end-tidal carbon dioxide, CI, cardiac index; NT, normothermia; HT, hypothermia; HT+SEV, hypothermia combined with sevoflurane. Data are mean ± SD.

Bax and Bcl-2 mRNA expression

We found a significant ($P < 0.01$) upregulation of both Bcl-2 mRNA and Bax expression after global cerebral ischemia (NT: Bcl-2 3.2 ± 1.8-fold, Bax 2.3 ± 1.3-fold compared with sham control). Hypothermia was associated with significantly ($P < 0.05$) less upregulation of mRNA expression (Bcl-2 1.2 ± 0.5-fold, Bax 1.2 ± 0.6-fold compared with sham control). Sevoflurane did not confer additional effects (Bcl-2 1.1 ± 0.4-fold, Bax 1.1 ± 0.4-fold compared with sham control; Figure 4).

Discussion

Neurological dysfunction resulting from cardiac arrest largely contributes to morbidity and mortality after initially successful CPR [21]. Employing a pig model we showed that (i) global cerebral ischemia following cardiac arrest and CPR results in upregulation of pro-inflammatory cytokine expression in the cerebral tissue, (ii) mild hypothermia significantly reduces cerebral tissue inflammatory response, and (iii) pharmacological post-conditioning with sevoflurane does not confer additional anti-inflammatory effects on cerebral tissue.

Cerebral inflammatory response following resuscitation

Mechanisms of brain injury following cerebral ischemia are complex with multiple modulators, signaling pathways, proteins and enzymes being involved that may facilitate cell death or survival [22]. Post-ischemic inflammation has been shown to play a critical role in cerebral ischemia/reperfusion injury [23]. Specifically, there is strong evidence suggesting that a disproportionate and persistent production of cytokines can significantly increase the risk and extent of brain injury [5,24]. In terms of a systemic inflammatory response, increased serum levels of different cytokines and chemokines have recently been presented in a rat model of cardiac arrest [25], and in patients successfully resuscitated from out-of-hospital cardiac arrest [26,27]. The role of the cerebral inflammatory response after cardiac arrest, however,




Figure 2 Cerebral cytokine mRNA expression. Transcript levels of the cerebral cytokines interleukin (IL)-1β, IL-6, IL-10, tumor necrosis factor (TNF)α and intercellular adhesion molecule (ICAM)-1 were determined by quantitative RT-PCR. NT, normothermia; HT, hypothermia; HT+SEV, hypothermia combined with sevoflurane. Data are expressed as mean ± SD (x-fold upregulation compared with Sham control). * $P < 0.05$, † $P < 0.01$ vs. Sham. § $P < 0.05$, # $P < 0.01$ vs. NT. RT-PCR data analysis was performed using two-sided *Pair-wise fixed Reallocation Randomisation Test*.




Figure 3 Protein concentration of interleukin-1β. Protein concentration of interleukin (IL)-1β was determined by a swine specific enzyme-linked-immunosorbent assay. NT, normothermia; HT, hypothermia; HT+SEV, hypothermia combined with sevoflurane. Data are expressed as mean ± SD (x-fold upregulation compared with Sham control). * $P < 0.05$ vs. Sham. § $P < 0.05$, # $P < 0.01$ vs. NT (using Mann-Whitney test).

has poorly been investigated. Most of the previous experimental studies induced global cerebral ischemia by bilateral carotid artery occlusion as a surrogate of cardiac arrest, but inflammatory response mechanisms following carotid artery occlusion and anti-inflammatory mechanisms of hypothermia are different from the ones observed after cardiac arrest and resuscitation [27]. Youngquist et al. [28] have recently shown increased TNFα and IL-6 protein concentration in the

cerebrospinal fluid following cardiac arrest. Since the presence of a lesion pattern of cortical involvement, termed as *extensive cortical lesion pattern* in MR imaging, has very recently been shown to be a very good predictor of poor neurologic prognosis after cardiac arrest [29], we focused on neuroinflammation in the cerebral cortex tissue. In our study, global cerebral ischemia following cardiac arrest resulted in a significant upregulation of mRNA expression of several cytokines




Figure 4 Cerebral Bcl-2 and Bax mRNA expression. Transcript levels of the cerebral apoptosis-associated proteins Bcl-2 and Bax were determined by quantitative RT-PCR. NT, normothermia; HT, hypothermia; HT+SEV, hypothermia combined with sevoflurane. Data are expressed as mean \pm SD (x-fold upregulation compared with Sham control). * $P < 0.05$ vs. Sham. § $P < 0.05$ vs. NT. RT-PCR data analysis was performed using two-sided *Pair-wise fixed Reallocation Randomisation Test*.

in the cerebral cortex tissue. In addition, we observed a significant rise in IL-1 β protein concentration in the cerebral cortex tissue that may be most probably due to local synthesis primarily by microglial cells, astrocytes and/or endothelial cells [30] rather than transport across the blood-brain barrier. This is emphasized by the data of Mizushima and colleagues who demonstrated that the integrity of the blood-spinal cord and blood-brain barriers to radiolabelled TNF α remains intact following resuscitation in a mouse model of cardiac arrest [31].

Effects of hypothermia on cerebral inflammatory response

Several mechanisms by which hypothermia exerts its protective effects have been characterized, including reduction in excitotoxin accumulation and inhibition of molecular pathways such as apoptosis and necrosis [4]. The role of inflammation in global cerebral ischemia induced by bilateral carotid artery occlusion and focal cerebral ischemia has extensively been studied, but effects of hypothermia on global cerebral ischemia/reperfusion injury following cardiac arrest has been investigated to a much lesser extent. Webster et al. have previously found that mild hypothermia attenuated microglial activation and nuclear translocation of NF κ B, and thereby reduced activation of the downstream inflammatory pathway [32]. Considering the relatively

late onset of the inflammatory response and the prolonged destructive process following cerebral ischemia/reperfusion, there appears to be a reasonable therapeutic time window using mild hypothermia to favourably affect the inflammatory pathway [33]. To date, the majority of publications may suggest that hypothermia simply blocks any ischemia-induced damaging cascade. However, contrary to this popular belief, the expression of certain beneficial genes is actually upregulated by mild hypothermia [4]. Hicks and colleagues [34] further demonstrated that prolonged hypothermia during later reperfusion improved neurological outcome after experimental global ischemia and was associated with selective changes in the pattern of stress-induced protein expression. From our data we conclude that mild hypothermia initiated after successful resuscitation from cardiac arrest reduces pro-inflammatory cytokine, IL-10, and ICAM-1 mRNA expression compared to normothermia. Inhibition of adhesion molecule expression and microglial activation has also been confirmed by Deng and colleagues in rat models of both focal cerebral ischemia and brain inflammation [35]. Thus, the beneficial effects of hypothermia on neuroprotection are considered to be due, in part, to suppression of post-injury pro-inflammatory factors by microglia. However, the role of hypothermia in modulating anti-inflammatory cytokines, for example, IL-10, remains controversial. While mild

hypothermia has been shown to increase plasma IL-10 concentration in endotoxemic rats, thus potentially mediating the anti-inflammatory effects of hypothermia [36,37], Matsui et al. [38] and Russwurm et al. [39] have previously demonstrated that mild hypothermia inhibits IL-10 production in peripheral blood mononuclear cells. Interestingly, in lipopolysaccharide-activated cultured microglia cells isolated from rats, hypothermia has also been found to reduce production of IL-6, IL-10, and nitric oxide, suggesting that the neuroprotective effects of hypothermia might involve not only the inhibition of pro-inflammatory factors, but also the inhibition of anti-inflammatory factor(s) [40]. Comparably, we found less upregulation of IL-10 mRNA expression in the cerebral tissue in the hypothermia group compared to normothermia after successful CPR.

Since IL-1 β was one of the cytokines that was strongly up-regulated on mRNA level in our study, we decided to evaluate IL-1 β expression also on the protein level. Analysis of cerebral cortex tissue using a swine specific ELISA system revealed significantly increased IL-1 β protein concentration compared with the sham control group after cardiac arrest and normothermia but not after hypothermia. Interestingly, Callaway et al. have recently demonstrated that hypothermia after cardiac arrest did not alter serum inflammatory markers, suggesting that circulating cytokines may not play a specific role regarding the neuroprotective effect of hypothermia [25]. In contrast, it is well conceivable, that local cerebral cytokines released by brain cells will affect more extensively various cerebral ischemia/reperfusion injury cascades and will have a much broader effect on brain damage than systemically elevated levels of cytokines.

Concerning reliable biochemical markers of brain tissue damage, increased serum levels of the low molecular weight protein S100B have been reported after cardiac arrest correlating with neurological complications. However, mild therapeutic hypothermia did not affect S100B serum levels in survivors of cardiac arrest in several clinical studies [41,42]. In addition, Xiao and colleagues have previously shown that cardiac arrest significantly increased brain myeloperoxidase activity, but again, mild hypothermia had no effect. Thus, the hypothermia-elicited neuroprotection seemed not to be neutrophil-dependent, at least in that rat model of asphyxial cardiac arrest [43].

Effects of pharmacological post-conditioning on cerebral inflammatory response

Volatile anesthetic agents have emerged as pre-conditioning-like agents with significant neuroprotective effects and the ability to reduce excitotoxic induced cell death, to decrease cerebral metabolic rate, to activate inducible nitrous oxide synthase and p38 mitogen-

activated protein kinases, and to improve neurological deficits in models of both focal and global cerebral ischemia [6,44,45]. Most experimental studies have documented improved functional performance when neuroprotective agents were given before the insult. In patients with cardiac arrest, however, pretreatment is virtually impossible because of the unpredictable onset of ischemia. Therefore, as in our study, potential protective interventions should be initiated during or after experimental ischemia to affect reperfusion injury. In this context, pharmacological postconditioning with volatile anesthetics in addition to mild hypothermia may offer an attractive opportunity to further ameliorate brain damage and inflammation in the post-resuscitation period. The effects of volatile agents on the inflammatory response after cardiac arrest have not yet been elucidated. In endotoxemic rats, inhalation of sevoflurane significantly attenuated plasma levels of TNF α and IL-1 β [46]. In addition, sevoflurane post-conditioning showed anti-inflammatory and anti-necrotic effects in cultured kidney proximal tubule cells [47], and sevoflurane attenuated the inflammatory response upon stimulation of alveolar macrophages with endotoxin in vitro [48]. In our study, however, sevoflurane administered instead of propofol during reperfusion after successful CPR did not further attenuate local cerebral inflammatory response. These observations are comparable to those obtained in a study by Fries et al. where the volatile anesthetic isoflurane did not reduce neurological dysfunction and histopathological alterations induced by cardiac arrest [49]. However, it is conceivable that hypothermia alone has such potent anti-inflammatory properties compared to normothermia, that an additional effect of sevoflurane could not be revealed in the present study. Moreover, potential protective effects of volatile anesthetics depend on energy-dependent signal transduction, for example, protein synthesis and phosphorylation [50], that may be affected by hypothermia-induced decrease of metabolic rate as well as suppression of protein synthesis.

Cerebral apoptosis-related mRNA expression

In the cerebral cortex tissue, we found a significant upregulation of both Bcl-2 and Bax mRNA expression after global cerebral ischemia. Comparably, Mishra and colleagues have recently reported increased apoptosis in a pig model of cerebral hypoxia for 60 minutes, indicated by an increased ratio of Bax/Bcl-2 protein concentration, activation of caspase-9, lipid peroxidation, and DNA fragmentation in mitochondria of the cerebral cortex [51].

Besides the regulation of inflammatory molecules, mild therapeutic hypothermia significantly attenuated the mRNA expression of the apoptosis-regulating proteins Bax and Bcl-2 in our study. These results are

partly comparable to the findings of Eberspächer et al. [52,53], where hypothermia prevented an ischemia-induced increase of the pro-apoptotic protein Bax, but did not change or even increase expression of the anti-apoptotic protein Bcl-2. Potential discrepancies between the work presented here and those in the literature could be due to the type of species and duration of ischemia. In our pig model seven minutes of cardiac arrest were followed by resuscitation compared with the latter studies investigating a rat model of common carotid artery occlusion plus hemorrhagic hypotension [52,53]. In a similar rat model of cerebral ischemia, Pape et al investigated the effects of sevoflurane on neuronal damage and expression of apoptic factors. Sevoflurane was administered before, during and after cerebral ischemia, and has been found to modulate the balance between pro- and anti-apoptotic key proteins towards a reduction of active programmed cell death by increasing the hippocampal concentration of the anti-apoptotic proteins Bcl-2, and by inhibiting the ischemia-induced upregulation of the pro-apoptotic protein Bax [54].

In our pig model of cardiac arrest, however, sevoflurane post-conditioning combined with mild hypothermia did not confer additional effects in terms of apoptotic-related mRNA expression. Again, it is conceivable that hypothermia alone has such potent anti-apoptotic effects, that an additional effect of sevoflurane could not be revealed in the present study.

Limitations

Although we used a porcine model of cardiac arrest following myocardial ischemia reflecting a common clinical scenario, there are several points that need to be addressed in future studies: (i) both long-term survival and neurological outcome were not evaluated because of limitations posed by governmental regulations; therefore, we did not assess the relationship between the upregulation of cytokines and post-resuscitation cerebral dysfunction. (ii) Blinding the investigator was not possible throughout the experiment due to the cooling technique, but tissue samples were analyzed by a person blinded to treatment assignment.

Conclusions

In conclusion, (i) global cerebral ischemia following cardiac arrest results in up-regulation of pro-inflammatory cytokines; (ii) hypothermia after cardiac arrest reduces up-regulation of various cytokines in the cerebral tissue. This may promote, at least in part, neuroprotection. (iii) The volatile anesthetic sevoflurane, when administered during reperfusion after successful CPR, did not confer statistically significant additional anti-inflammatory effects in the above setting.

Key messages

- Global cerebral ischemia following cardiac arrest results in up-regulation of local pro-inflammatory cytokines expression.
- Mild hypothermia after cardiac arrest attenuates cerebral inflammatory response.
- Sevoflurane does not confer additional anti-inflammatory effects.
- Further studies on the relationship between cerebral inflammatory response and post-resuscitation cerebral dysfunction are warranted.

Additional file 1: Extended Method section - Quantitative real-time RT-PCR. Detailed description of quantitative real-time RT-PCR, primer sequences and TaqMan probes.
Click here for file
[<http://www.biomedcentral.com/content/supplementary/cc8879-S1.doc>]

Abbreviations

BL: baseline; CPR: cardiopulmonary resuscitation; ELISA: enzyme-linked immunosorbent assay; HT: hypothermia; ICAM-1: intercellular adhesion molecule-1; IL: interleukin; LAD: left anterior descending (coronary artery); NT: normothermia; ROSC: return of spontaneous circulation; RT-PCR: reverse transcriptase polymerase chain reaction; SEV: sevoflurane; TIVA: total intravenous anesthesia; TNF α : tumor necrosis factor α ; VF: ventricular fibrillation.

Acknowledgements

This work has been supported by the German Interdisciplinary Association of Critical Care Medicine (PM) and by the German Research Foundation (BB). The founders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The authors are indebted to H. Fiedler, B. Zastrow, B. Kuhr, and V. Haensel-Bringmann for technical assistance. We thank C. Rodde, S. Piontek, M. Koelln, G. Jopp, Prof. I. Cascorbi and M. Ufer for laboratory analysis. The manuscript was presented in part at the Annual Meeting of the Society of Neurosurgical Anesthesia and Critical Care, Orlando, FL, USA, 17th October 2008, and at the 3rd International Hypothermia Symposium, Lund, Sweden, 5th September 2009.

Author details

¹Department of Anaesthesiology and Intensive Care Medicine, University Hospital Schleswig-Holstein, Campus Kiel, Schwannenweg 21, Kiel, 24105, Germany. ²Clinic of Anaesthesiology, Intensive Care Medicine and Pain Therapy, University Hospital Frankfurt, Theodor-Stern-Kai 7, Frankfurt am Main, 60590, Germany. ³Institute of Anatomy, Christian-Albrechts-University of Kiel, Otto-Hahn-Platz 8, Kiel, 24118, Germany.

Authors' contributions

PM, KDZ and BB conceived and designed the experiments. PM, MG, KDZ and MA performed the experiments. MG, MA, RL, NF, JH and KZ analyzed the data. PM, KDZ, MA and BB wrote the paper. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 30 September 2009 Revised: 24 December 2009

Accepted: 16 February 2010 Published: 16 February 2010

References

1. de Vreede-Swagemakers JJ, Gorgels AP, Dubois-Arbouw WI, van Ree JW, Daemen MJ, Houben LG, Wellens HJ: **Out-of-hospital cardiac arrest in the 1990s: a population-based study in the Maastricht area on incidence, characteristics and survival.** *J Am Coll Cardiol* 1997, **30**:1500-1505.

2. Longstreth WT Jr, Inui TS, Cobb LA, Copass MK: **Neurologic recovery after out-of-hospital cardiac arrest.** *Ann Intern Med* 1983, **98**:588-592.
3. Holzer M, Bernard SA, Hachimi-Idrissi S, Roine RO, Sterz F, Mullner M: **Hypothermia for neuroprotection after cardiac arrest: systematic review and individual patient data meta-analysis.** *Crit Care Med* 2005, **33**:414-418.
4. Zhao H, Steinberg GK, Sapolsky RM: **General versus specific actions of mild-moderate hypothermia in attenuating cerebral ischemic damage.** *J Cereb Blood Flow Metab* 2007, **27**:1879-1894.
5. Dirnagl U, Simon RP, Hallenbeck JM: **Ischemic tolerance and endogenous neuroprotection.** *Trends Neurosci* 2003, **26**:248-254.
6. Sanders RD, Ma D, Maze M: **Anaesthesia induced neuroprotection.** *Best Pract Res Clin Anaesthesiol* 2005, **29**:461-474.
7. Thoresen M, Hobbs CE, Wood T, Chakkarapani E, Dingley J: **Cooling combined with immediate or delayed xenon inhalation provides equivalent long-term neuroprotection after neonatal hypoxia-ischemia.** *J Cereb Blood Flow Metab* 2009, **29**:707-714.
8. Idris AH, Becker LB, Ornato JP, Hedges JR, Bircher NG, Chandra NC, Cummins RO, Dick W, Ebmeyer U, Halperin HR, Hazinski MF, Kerber RE, Kern KB, Safar P, Steen PA, Swindle MM, Tisitlik JE, von Planta I, von Planta M, Wears RL, Weil MH: **Utstein-style guidelines for uniform reporting of laboratory CPR research. A statement for healthcare professionals from a task force of the American Heart Association, the American College of Emergency Physicians, the American College of Cardiology, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, the Institute of Critical Care Medicine, the Safar Center for Resuscitation Research, and the Society for Academic Emergency Medicine. Writing Group.** *Circulation* 1996, **94**:2324-2336.
9. Johnson KB, Egan TD, Kern SE, McJames SW, Cluff ML, Pace NL: **Influence of hemorrhagic shock followed by crystalloid resuscitation on propofol: a pharmacokinetic and pharmacodynamic analysis.** *Anesthesiology* 2004, **101**:647-659.
10. Kelley KW, Curtis SE, Marzan GT, Karara HM, Anderson CR: **Body surface area of female swine.** *J Anim Sci* 1973, **36**:927-930.
11. Neumar RW, Nolan JP, Adrie C, Aibiki M, Berg RA, Bottiger BW, Callaway C, Clark RS, Geocadin RG, Jauch EC, Kern KB, Laurent I, Longstreth WT, Merchant RM, Morley P, Morrison LJ, Nadkarni V, Peberdy MA, Rivers EP, Rodriguez-Nunez A, Sellke FW, Spaulding C, Sunde K, Hoek Vanden T, International Liaison Committee on Resuscitation; Emergency Cardiovascular Care Committee, American Heart Association; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiopulmonary, Perioperative, and Critical Care; Council on Clinical Cardiology; Council on Stroke: **Post-cardiac Arrest Syndrome: epidemiology, pathophysiology, treatment, and prognostication. A consensus statement from the International Liaison Committee on Resuscitation (American Heart Association, Australian and New Zealand Council on Resuscitation, European Resuscitation Council, Heart and Stroke Foundation of Canada, interamerican Heart Foundation, Resuscitation Council of Asia, and the Resuscitation Council of Southern Africa); the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; and the Stroke Council.** *Circulation* 2008, **118**:2452-2483.
12. Roesner JP, Petzelbauer P, Koch A, Mersmann J, Zacharowski PA, Boehm O, Reingruber S, Pasterner W, Mascher D, Wolzt M, Barthuber C, Nöldge-Schomburg GE, Scheeren TW, Zacharowski K: **The fibrin-derived peptide Bbeta15-42 is cardioprotective in a pig model of myocardial ischemia-reperfusion injury.** *Crit Care Med* 2007, **35**:1730-1735.
13. **2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care.** *Circulation* 2005, **112**: IV1-203.
14. Tortorici MA, Kochanek PM, Poloyac SM: **Effects of hypothermia on drug disposition, metabolism, and response: A focus of hypothermia-mediated alterations on the cytochrome P450 enzyme system.** *Crit Care Med* 2007, **35**:2196-2204.
15. Hannon JP, Bossone CA, Wade CE: **Normal physiological values for conscious pigs used in biomedical research.** *Lab Anim Sci* 1990, **40**:293-298.
16. Bernard SA, Gray TW, Buist MD, Jones BM, Silvester W, Gutteridge G, Smith K: **Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia.** *N Engl J Med* 2002, **346**:557-563.
17. Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP: **Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper-Excel-based tool using pair-wise correlations.** *Biotechnol Lett* 2004, **26**:509-515.
18. Rozen S, Skaletsky H: **Primer3 on the WWW for general users and for biologist programmers.** *Methods Mol Biol* 2000, **132**:365-386.
19. Pfaffl MW: **A new mathematical model for relative quantification in real-time RT-PCR.** *Nucleic Acids Res* 2001, **29**:e45.
20. Pfaffl MW, Horgan GW, Dempfle L: **Relative expression software tool (REST) for group-wise comparison and statistical analysis of relative expression results in real-time PCR.** *Nucleic Acids Res* 2002, **30**:e36.
21. Oddo M, Ribordy V, Feihl F, Rossetti AO, Schaller MD, Chiolerio R, Liaudet L: **Early predictors of outcome in comatose survivors of ventricular fibrillation and non-ventricular fibrillation cardiac arrest treated with hypothermia: a prospective study.** *Crit Care Med* 2008, **36**:2296-2301.
22. Harukuni I, Bhardwaj A: **Mechanisms of brain injury after global cerebral ischemia.** *Neurol Clin* 2006, **24**:1-21.
23. Zheng Z, Yenari MA: **Post-ischemic inflammation: molecular mechanisms and therapeutic implications.** *Neurol Res* 2004, **26**:884-892.
24. Zhu Y, Saito K, Murakami Y, Asano M, Iwakura Y, Seishima M: **Early increase in mRNA levels of pro-inflammatory cytokines and their interactions in the mouse hippocampus after transient global ischemia.** *Neurosci Lett* 2006, **393**:122-126.
25. Callaway CW, Rittenberger JC, Logue ES, McMichael MJ: **Hypothermia after cardiac arrest does not alter serum inflammatory markers.** *Crit Care Med* 2008, **36**:2607-2612.
26. Fries M, Kunz D, Gressner AM, Rossaint R, Kuhlen R: **Procalcitonin serum levels after out-of-hospital cardiac arrest.** *Resuscitation* 2003, **59**:105-109.
27. Adrie C, Adib-Conquy M, Laurent I, Monchi M, Vinsonneau C, Fitting C, Fraisse F, Dinh-Xuan AT, Carli P, Spaulding C, Dhainaut JF, Cavaillon JM: **Successful cardiopulmonary resuscitation after cardiac arrest as a "sepsis-like" syndrome.** *Circulation* 2002, **106**:562-568.
28. Youngquist ST, Niemann JT, Heyming TW, Rosborough JP: **The central nervous system cytokine response to global ischemia following resuscitation from ventricular fibrillation in a porcine model.** *Resuscitation* 2009, **80**:249-252.
29. Topcuoglu MA, Oguz KK, Buyukserbetci G, Bulut E: **Prognostic value of magnetic resonance imaging in post-resuscitation encephalopathy.** *Intern Med* 2009, **48**:1635-1645.
30. Szelenyi J: **Cytokines and the central nervous system.** *Brain Res Bull* 2001, **54**:329-338.
31. Mizushima H, Banks WA, Dohi K, Shioda S, Matsumoto K: **Effect of cardiac arrest on brain weight and the permeability of the blood-brain and blood-spinal cord barrier to albumin and tumor necrosis factor-alpha.** *Life Sci* 1999, **65**:2127-2134.
32. Webster CM, Kelly S, Koike MA, Chock VY, Giffard RG, Yenari MA: **Inflammation and NFkappaB activation is decreased by hypothermia following global cerebral ischemia.** *Neurobiol Dis* 2009, **33**:301-312.
33. Polderman KH: **Mechanisms of action, physiological effects, and complications of hypothermia.** *Crit Care Med* 2009, **37**:S186-202.
34. Hicks SD, DeFranco DB, Callaway CW: **Hypothermia during reperfusion after asphyxial cardiac arrest improves functional recovery and selectively alters stress-induced protein expression.** *J Cereb Blood Flow Metab* 2000, **20**:520-530.
35. Deng H, Han HS, Cheng D, Sun GH, Yenari MA: **Mild hypothermia inhibits inflammation after experimental stroke and brain inflammation.** *Stroke* 2003, **34**:2495-2501.
36. Fujimoto K, Fujita M, Tsuruta R, Tanaka R, Shinagawa H, Izumi T, Kasaoka S, Maekawa T: **Early induction of moderate hypothermia suppresses systemic inflammatory cytokines and intracellular adhesion molecule-1 in rats with caerulein-induced pancreatitis and endotoxemia.** *Pancreas* 2008, **37**:176-181.
37. Sarcia PJ, Scumpia PO, Moldawer LL, DeMarco VG, Skimming JW: **Hypothermia induces interleukin-10 and attenuates injury in the lungs of endotoxemic rats.** *Shock* 2003, **20**:41-45.
38. Matsui T, Ishikawa T, Takeuchi H, Tsukahara M, Maekawa T: **Mild hypothermia inhibits IL-10 production in peripheral blood mononuclear cells.** *Acta Anaesthesiol Scand* 2004, **48**:205-210.
39. Russwurm S, Stonans I, Schwerter K, Stonane E, Meissner W, Reinhart K: **Direct influence of mild hypothermia on cytokine expression and**

- release in cultures of human peripheral blood mononuclear cells. *J Interferon Cytokine Res* 2002, **22**:215-221.
40. Matsui T, Kakeda T: IL-10 production is reduced by hypothermia but augmented by hyperthermia in rat microglia. *J Neurotrauma* 2008, **25**:709-715.
 41. Tiainen M, Roine RO, Pettila V, Takkunen O: Serum neuron-specific enolase and S-100B protein in cardiac arrest patients treated with hypothermia. *Stroke* 2003, **34**:2881-2886.
 42. Derwall M, Stoppe C, Brucken D, Rossaint R, Fries M: Changes in S-100 protein serum levels in survivors of out-of-hospital cardiac arrest treated with mild therapeutic hypothermia: a prospective, observational study. *Crit Care* 2009, **13**:R58.
 43. Xiao F, Zhang S, Arnold TC, Alexander JS, Huang J, Carden DL, Conrad SA: Mild hypothermia induced before cardiac arrest reduces brain edema formation in rats. *Acad Emerg Med* 2002, **9**:105-114.
 44. Blanck TJ, Haile M, Xu F, Zhang J, Heerd P, Veselis RA, Beckman J, Kang R, Adamo A, Hemmings H: Isoflurane pretreatment ameliorates postischemic neurologic dysfunction and preserves hippocampal Ca²⁺/calmodulin-dependent protein kinase in a canine cardiac arrest model. *Anesthesiology* 2000, **93**:1285-1293.
 45. Eberspächer E, Eckel B, Engelhard K, Muller K, Hoffman WE, Blobner M, Werner C: Effects of sevoflurane on cognitive deficit, motor function, and histopathology after cerebral ischemia in rats. *Acta Anaesthesiol Scand* 2009, **53**:774-782.
 46. Hofstetter C, Boost KA, Flondor M, Basagan-Mogol E, Betz C, Homann M, Muhl H, Pfeilschifter J, Zwissler B: Anti-inflammatory effects of sevoflurane and mild hypothermia in endotoxemic rats. *Acta Anaesthesiol Scand* 2007, **51**:893-899.
 47. Lee HT, Kim M, Jan M, Emala CW: Anti-inflammatory and antinecrotic effects of the volatile anesthetic sevoflurane in kidney proximal tubule cells. *Am J Physiol Renal Physiol* 2006, **291**:F67-78.
 48. Steurer M, Schlapfer M, Steurer M, Z'Graggen BR, Booy C, Reyes L, Spahn DR, Beck-Schimmer B: The volatile anaesthetic sevoflurane attenuates lipopolysaccharide-induced injury in alveolar macrophages. *Clin Exp Immunol* 2009, **155**:224-230.
 49. Fries M, Coburn M, Nolte KW, Timper A, Kottmann K, Kuru TH, Weis J, Rossaint R: Early administration of xenon or isoflurane may not improve functional outcome and cerebral alterations in a porcine model of cardiac arrest. *Resuscitation* 2009, **80**:584-590.
 50. Weber NC, Schlack W: The concept of anaesthetic-induced cardioprotection: mechanisms of action. *Best Pract Res Clin Anaesthesiol* 2005, **19**:429-443.
 51. Mishra OP, Randis T, Ashraf QM, Delivoria-Papadopoulos M: Hypoxia-induced Bax and Bcl-2 protein expression, caspase-9 activation, DNA fragmentation, and lipid peroxidation in mitochondria of the cerebral cortex of newborn piglets: the role of nitric oxide. *Neuroscience* 2006, **141**:1339-1349.
 52. Eberspächer E, Werner C, Engelhard K, Pape M, Gelb A, Hutzler P, Henke J, Kochs E: The effect of hypothermia on the expression of the apoptosis-regulating protein Bax after incomplete cerebral ischemia and reperfusion in rats. *J Neurosurg Anesthesiol* 2003, **15**:200-208.
 53. Eberspächer E, Werner C, Engelhard K, Pape M, Laacke L, Winner D, Hollweck R, Hutzler P, Kochs E: Long-term effects of hypothermia on neuronal cell death and the concentration of apoptotic proteins after incomplete cerebral ischemia and reperfusion in rats. *Acta Anaesthesiol Scand* 2005, **49**:477-487.
 54. Pape M, Engelhard K, Eberspächer E, Hollweck R, Kellermann K, Zintner S, Hutzler P, Werner C: The long-term effect of sevoflurane on neuronal cell damage and expression of apoptotic factors after cerebral ischemia and reperfusion in rats. *Anesth Analg* 2006, **103**:173-179.

doi:10.1186/cc8879

Cite this article as: Meybohm et al.: Mild hypothermia alone or in combination with anesthetic post-conditioning reduces expression of inflammatory cytokines in the cerebral cortex of pigs after cardiopulmonary resuscitation. *Critical Care* 2010 **14**:R21.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

