



Cardiac progenitors instruct second heart field fate through Wnts

Matthew Miyamoto^a, Suraj Kannan^a, Matthew J. Anderson^b , Xihe Liu^a , David Suh^a , Myo Htet^a , Biyi Li^a , Tejasvi Kakani^a, Sean Murphy^a , Emmanouil Tampakakis^a, Mark Lewandoski^b , Peter Andersen^a , Hideki Uosaki^{a,c} , and Chulan Kwon^{a,1}

Edited by Roel Nusse, Stanford University School of Medicine, Stanford, CA; received October 19, 2022; accepted December 22, 2022

The heart develops in a synchronized sequence of proliferation and differentiation of cardiac progenitor cells (CPCs) from two anatomically distinct pools of cells, the first heart field (FHF) and second heart field (SHF). Congenital heart defects arise upon dysregulation of these processes, many of which are restricted to derivatives of the FHF or SHF. Of the conserved set of signaling pathways that regulate development, the Wnt signaling pathway has long been known for its importance in SHF development. The source of such Wnts has remained elusive, though it has been postulated that these Wnts are secreted from ectodermal or endodermal sources. The central question remains unanswered: Where do these Wnts come from? Here, we show that CPCs autoregulate SHF development via Wnt through genetic manipulation of a key Wnt export protein (Wls), scRNA-seq analysis of CPCs, and use of our precardiac organoid system. Through this, we identify dysregulated developmental trajectories of anterior SHF cell fate, leading to a striking single ventricle phenotype in knockout embryos. We then applied our findings to our precardiac organoid model and found that Wnt2 is sufficient to restore SHF cell fate in our model of disrupted endogenous Wnt signaling. In this study, we provide a basis for SHF cell fate decision—proliferation vs. differentiation—autoregulated by CPCs through Wnt.

heart development | second heart field | Wnt | Wls | organoid

Wnt signaling is required for second heart field (SHF) development (1). Some have thought that the source of these Wnts is from ectodermal or endodermal sources (2, 3). However, we found that *Wnts* are enriched in first heart field (FHF) progenitors while Wnt activity is higher in SHF progenitors (4). This led us to test whether the FHF is influencing SHF development. While others have perturbed Wnt signaling in Wnt-receiving cells through manipulation of β -catenin levels (1), the obligatory transcriptional effector of Wnt signaling, we instead focused on eliminating the ability of cells to secrete Wnts. To suppress Wnt secretion, we used a mouse strain harboring loxp sites flanking the gene encoding Wls (5), a protein required for export of Wnt ligands (6, 7). This allowed us to study which populations of cells that secrete Wnt affect heart development (Fig. 1A).

As there are no validated Cre drivers that induce recombination with high efficiency in the early FHF, we first deleted Wls in precardiac mesoderm using *Mesp1-Cre* (8). Interestingly, upon knockout of Wls, we found a heart defect starting from embryonic day (e) 8.5 (Fig. 1B). Control embryos formed the primitive outflow tract/right ventricle (OT/RV) and left ventricles (LVs), expected at this developmental stage. However, a structure reminiscent of the cardiac crescent—which is transiently present from e7.75 to e8.25—and early heart tube persisted in knockout embryos at e8.5. The heart defect became more pronounced by e9.5, where, in comparison to expected LV, RV, and OT formation in control embryos, knockout embryos exhibit a single ventricle-like structure (Fig. 1C).

The presence of a cardiac crescent-like/early heart tube structure (FHF) in knockout embryos suggests that the single ventricle phenotype may be due to dysregulation of SHF development. To test this, we utilized an *Hcn4-Green Fluorescent Protein (GFP)* reporter mouse strain to visualize FHF progenitors. As expected, GFP⁺ cells were localized in the LV of control embryos. However, knockout embryos showed an expansion of the GFP-expressing domain across the single ventricle (Fig. 1D). Additionally, immunostaining showed that the entire ventricle was positive for the LV marker *Tbx5* in knockout embryos, while only the LV was positive for *Tbx5* in control embryos (Fig. 1E), consistent with a decrease in SHF derivatives. These results suggest that mesodermal Wnts are required for SHF development.

To determine the source of mesodermal Wnts more specifically, we leveraged multiple precardiac Cre drivers with shared expression domains to conditionally knockout Wls. We first used *Islet1-Cre* mice, as their Cre expression begins shortly after gastrulation in FHF progenitors before marking the SHF (4, 9, 10). We chose this particular Cre driver as it is activated in both heart fields and expressed early in heart development (11), which

Author affiliations: ^aDivision of Cardiology, Department of Medicine, Johns Hopkins University, Baltimore, MD, 21205; ^bGenetics of Vertebrate Development Section, Cancer and Developmental Biology Laboratory, National Cancer Institute, NIH, Frederick, MD, 21702; and ^cDivision of Regenerative Medicine, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke, Tochigi 329-0498, Japan

Author contributions: M.M., S.K., P.A., H.U., and C.K. designed research; M.M., S.K., M.J.A., X.L., D.S., M.H., B.L., T.K., S.M., E.T., M.L., P.A., and H.U. performed research; M.M., S.K., M.J.A., X.L., D.S., M.H., B.L., S.M., E.T., M.L., P.A., H.U., and C.K. analyzed data; and M.M. and C.K. wrote the paper.

The authors declare no competing interest.

Copyright © 2023 the Author(s). Published by PNAS. This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

¹To whom correspondence may be addressed. Email: ckwon13@jhmi.edu.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2217687120/-/DCSupplemental>.

Published January 17, 2023.

we confirmed by observing Red Fluorescent Protein (RFP) expression in both LV and RV in Cre-recombined control embryos (Fig. 1*F*). Interestingly, we observed a similar single ventricle phenotype in *Islet1-Cre; Wls*-knockout embryos (Fig. 1*F*), albeit with variable penetrance, perhaps due to differing domains of recombination previously observed in this particular *Islet1-Cre* driver. These results suggest that the source of Wnts is from a common population of cells derived from both *Mesp1*⁺ and *Islet1*⁺ cells—the derivatives of the FHF and SHF. To test which heart field is responsible for the phenotype, we blocked Wnt secretion in the *Tbx1*⁺ domain of the anterior SHF (aSHF). No heart defect was observed in *Tbx1-Cre; Wls*-knockout embryos (Fig. 1*G*). This indicates that Wnts from the *Tbx1* lineage are dispensable for SHF development. Together, these results support that cardiac progenitor cells (CPCs), likely the FHF, provide Wnts for SHF development.

To understand the observed defect at the transcriptional level, we isolated *Mesp1*⁺ cells from *Mesp1-Cre; Ai9; Wls*^{f/f} and *Wls*^{f/f} mice at three developmental time points (e8.0, e8.5, and e9.5) for scRNA-seq. We performed fluorescence-activated cell sorting to isolate RFP⁺ cells and generated sequencing libraries. We performed dimensionality reduction through UMAP and annotated clusters through known marker genes (Fig. 2*A*). Our full cluster identification and quality control strategy is detailed in *SI Appendix, Online Methods*. We then analyzed Wnts in each population of mesoderm. Based on levels of expression of relevant Wnts, and our *Wls* loss of function studies, we focused on three populations as putative sources of instructive Wnts—posterior SHF (pSHF), FHF, and OT cardiomyocytes—though the OT cardiomyocytes are known to derive from *Tbx1*⁺ cells. The FHF cluster highly expressed *Wnt2*, while the pSHF cluster expressed low/undetectable levels of Wnts (Fig. 2*B*). These data further support our hypothesis that the FHF may be a source of instructive Wnts.

We then asked how *Wls* deletion affected mesodermal cells outside a generalized response due to secondary effects of having a defective heart. To do this, we explored genes that were differentially expressed in unique clusters of cells. Notably, this pool of genes was enriched for Gene Ontology terms related to cell cycle, Wnt signaling, and mesodermal organ tissue and development. This suggested that,

outside the nonspecific stress response in all tissues, *Wls* deficiency leads to perturbation of tissue development in specific mesodermal processes. The majority of these differentially expressed genes were identified in aSHF, pSHF, and sclerotome (Fig. 2*C*), suggesting that these tissues are most perturbed by loss of Wnt signaling.

Given the perturbation of aSHF cells, we performed trajectory reconstruction of control and knockout cells in the aSHF. We reconstructed a branched trajectory in which control and knockout cells are indistinguishable early at e8.0, before reaching disparate states at later time points (Fig. 2*D*). We were interested in genes differentially expressed across the control and knockout paths at the branch point, focusing on genes with differential expression at the end states. We classified four clusters of genes based on expression dynamics (Fig. 2*D*). In general, GO terms relevant to heart development were enriched in upregulated gene clusters while GO terms related to cell cycle were enriched in downregulated clusters (Fig. 2*D*), pointing to potential impaired proliferation. Given that proper SHF development requires maintenance of proliferation and delayed differentiation, the upregulation of aSHF genes and downregulation of proliferation genes suggests dysregulation of cell identity after premature differentiation. We verified this *in vivo* by examining proliferation of *Mesp1*⁺/*Tbx5*⁺ cells to quantify proliferation of the aSHF and found consistent decreased proliferation of SHF progenitors based on this staining (Fig. 2*E*). Taken together, these results point to a decreased cell proliferation in the aSHF as a key mechanism for the knockout heart phenotype.

We then performed a basic ligand–receptor analysis (Fig. 2*F*) to identify potential candidate signaling pairs affected by *Wls* knockout. Interestingly, we found that genes do not change significantly in the FHF as a result of *Wls* knockout; thus, there are no receptors or ligands whose expression is changing in the FHF. We instead focused on receptors whose expression is changing in the aSHF (as in Fig. 2*E*), with the corresponding ligand expressed in at least 25% of cells in the FHF. As expected, many receptors are increased in expression level in the knockout, potentially indicative of a compensatory response to loss of Wnt signaling. In particular, the Wnt receptor *Fzd4*, which has been shown to interact with *Wnt2*, increases in level in the knockout.




Fig. 1. Knockout of *Wntless* in CPCs leads to FHF-derived single chamber phenotype. (A) Schematic detailing conditional knockout of the gene *Wls*. (B) e8.5 *Mesp1-Cre; Wls*^{+/+} (control), and *Mesp1-Cre; Wls*^{f/f} (knockout) embryos. (C) e9.5 control and knockout embryos. (D) Fluorescent overlay images of *Hcn4-GFP* reporter in control and knockout embryos. (E) Immunostaining of control and knockout sections of embryo hearts for cTnt (green), a cardiomyocyte marker, and *Tbx5* (red), an LV marker with emphases on the right (*) and left (**) sides of the resulting chambers. (F) Front images of *Islet1-Cre* control and knockout embryos. Fifty percent shows significant heart defects. (G) Front images of *Tbx1-Cre* control and knockout embryos (OT, outflow tract; IFT, inflow tract; RV, right ventricle; LV, left ventricle; V, ventricle). White scale bars indicate 500 μ m.




Fig. 2. SHF development is dysregulated through decreased proliferation without *Mesp1*⁺ Wnts. (A) UMAP clustering based on cell type. (B) Expression of Wnts across cell populations. (C) Quantification of differentially expressed genes (DEGs) across time points and cell populations. (D) aSHF developmental trajectories from control and knockout embryos and heatmaps identifying differential gene trends with expression dynamics of cell cycle and cardiomyogenesis genes. (E) Quantification of percentage of *Mesp1*⁺/*Tbx5*⁻ cells undergoing proliferation. White boxes highlight the *Tbx5*-positive regions which were excluded from quantification. Control average = 2.77%. Knockout average = 1.71%. **P* = 0.039. (F) Differentially expressed ligand-receptor pairs in knockout embryos in the aSHF. (G) Frontal (Top) and lateral (Bottom) views of whole-mount in situ hybridization of candidate Wnts with *Tbx5* cohybridization in 4ss embryos (HF, head fold; CC, cardiac crescent; A, anterior; P, posterior). The white scale bar indicates 500 μ m. Auto fluorescent debris is noted in *Wnt5a/5b/11* embryos. (H) Flow cytometry quantifications of *Hcn4*-GFP; *Tbx1*-RFP-positive cells at day 7 of differentiation after IWP-2 and IWP-2+Wnt2 treatment. IWP-2 %GFP⁺ average = 9%; %RFP⁺ average = 20.44%. IWP-2+Wnt2 %GFP⁺ average = 8.61%; %RFP⁺ average = 27.43%. GFP⁺ cells *P* = 0.86 (NS = not significant); RFP⁺ cells **P* = 0.016.

To determine the specificity of Wnt expression to the FHF, we performed whole-mount in situ hybridization (12) of candidate Wnts with cohybridization of *Tbx5*, and found that, among our candidate genes identified from scRNA-seq, only *Wnt2* was localized to the FHF (Fig. 2G). Based on this, we used our precardiac organoid system, which recapitulates FHF/SHF development at both transcriptomic and functional levels, to determine whether *Wnt2* affects formation of the SHF in the setting of blocked Wnt secretion (Fig. 2H). We blocked Wnt secretion at day 4 of differentiation using the small molecule IWP-2 as previously described (4) and treated organoids with recombinant *Wnt2* protein. Remarkably, upon cotreatment with *Wnt2* protein, we observed an increase in SHF formation in IWP-2 treated cells, with no observable impact on the FHF. This demonstrates that *Wnt2*, which is expressed by the FHF, is sufficient to promote SHF development.

In summary, we show that *Wls*-deficient CPCs dysregulate developmental trajectories of aSHF cells, leading to impaired proliferation. These results demonstrate a critical role of precardiac (likely FHF) Wnts in SHF fate decisions, highlighting the importance of coordinated Wnt signals in heart field development and chamber formation. Further research into the instructive role of Wnt signaling in human models will help elucidate these complex cell-cell morphogenic signaling relationships and better understand the etiology of congenital heart disease.

Data, Materials, and Software Availability. scRNA-seq data have been deposited in Synapse; GitHub (<https://www.synapse.org/#!Synapse:syn24200678/files/>; <https://github.com/skannan4/wls>).

ACKNOWLEDGMENTS. This work was supported by the NIH, the Saving tiny Hearts Society, the American Heart Association, and the Department of Defense.

1. A. Klaus, Y. Saga, M. M. Taketo, E. Tzahor, W. Birchmeier, Distinct roles of Wnt/ β -catenin and Bmp signaling during early cardiogenesis. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 18531-18536 (2007), 10.1073/pnas.0703113104.
2. C. Kemp, E. Willems, S. Abdo, L. Lambiv, L. Leyns, Expression of all Wnt genes and their secreted antagonists during mouse blastocyst and postimplantation development. *Dev. Dyn.* **233**, 1064-1075 (2005).
3. E. Tzahor, A. B. Lassar, Wnt signals from the neural tube block ectopic cardiogenesis. *Genes Dev.* **15**, 255-260 (2001).
4. P. Andersen et al., Preadiac organoids form two heart fields via Bmp/Wnt signaling. *Nat. Commun.* **9**, 1-13 (2018).
5. A. C. Carpenter, S. Rao, J. M. Wells, K. Campbell, R. A. Lang, Generation of mice with a conditional null allele for *Wntless*. *Genesis* **48**, 554-558 (2010).
6. K. Bartscherer, N. Pelte, D. Ingelfinger, M. Boutros, Secretion of Wnt ligands requires Evi, a conserved transmembrane protein. *Cell* **125**, 523-533 (2006).
7. C. Bänziger et al., *Wntless*, a conserved membrane protein dedicated to the secretion of Wnt proteins from signaling cells. *Cell* **125**, 509-522 (2006).
8. Y. Saga et al., *MesP1* is expressed in the heart precursor cells and required for the formation of a single heart tube. *Development* **126**, 3437-3447 (1999).
9. Q. Ma, B. Zhou, W. T. Pu, Reassessment of *Isl1* and *Nkx2-5* cardiac fate maps using a *Gata4*-based reporter of Cre activity. *Dev. Biol.* **323**, 98-104 (2008).
10. Y. Sun et al., *Isl1* is expressed in distinct cardiovascular lineages, including pacemaker and coronary vascular cells. *Dev. Biol.* **304**, 286-296 (2007).
11. L. Yang et al., *Isl1* Cre reveals a common Bmp pathway in heart and limb development. *Development* **133**, 1575-1585 (2006).
12. M. J. Anderson, V. Magidson, R. Kageyama, M. Lewandoski, *Fgf4* maintains *Hes7* levels critical for normal somite segmentation clock function. *Elife* **9**, 1-22 (2020).