



## Early View

### Correspondence

## **Factors limiting the utility of bronchoalveolar lavage in the diagnosis of Covid-19**

Deepak Aggarwal, Varinder Saini

Please cite this article as: Aggarwal D, Saini V. Factors limiting the utility of bronchoalveolar lavage in the diagnosis of Covid-19. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.03116-2020>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©ERS 2020. This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.

## Title Page

Title: Factors limiting the utility of bronchoalveolar lavage in the diagnosis of Covid-19

Authors (in order)

1. Deepak Aggarwal, MD. Associate Professor, Dept. of Pulmonary Medicine,  
Government Medical College & Hospital, Sector-32,  
Chandigarh. Pin: 160030, India
2. Varinder Saini, MD. Professor & Head, Dept. of Pulmonary Medicine,  
Government Medical College & Hospital, Sector-32,  
Chandigarh. Pin: 160030, India

Corresponding Author: Deepak Aggarwal, Associate Professor, Dept. Of Pulmonary Medicine, Block-D, Level-5, Government Medical College & Hospital, Sector-32, Chandigarh.

Pin-160030, India. E mail: [drdeepak@hotmail.com](mailto:drdeepak@hotmail.com). Mobile +91-9646121584

Word Count: 290

Key words: Covid-19, bronchoalveolar lavage, diagnosis

Conflict of interest: No conflict of Interest

Take home message: Bronchoalveolar lavage has a specific place in the diagnosis of Covid-19. The clinical performance of BAL rRt-PCR is dynamic in nature and depends on different clinical and demographic factors.

## Main Text

We read with interest study by Geri et al<sup>1</sup> in which the authors have demonstrated a strong agreement between negative nasopharyngeal (NPs)/nasal (Ns) swab and bronchoalveolar lavage (BAL) real-time reverse transcriptase-polymerase chain reaction (rRT-PCR) in the diagnosis of Covid-19 among hospitalized patients. The study findings contradicted earlier

report<sup>2</sup> and suggested a limited utility of BAL. However, the results need to be interpreted comprehensively before drawing any conclusion.

In the present study, BAL was negative for SARS-CoV-2 by rRT-PCR in majority of cases that included 38 patients (48%) with strong clinical and radiological suspicion for Covid-19. This finding implies either a high false negative rate of BAL rRT-PCR or an alternate diagnosis. Hence, authors should give a detail account of the final diagnoses and treatment outcomes of the patients and correlate these with the rRT-PCR results. This will give a better picture of the clinical performance of rRT-PCR in both BAL and upper respiratory samples.

Clinical test performance of rRT-PCR (in BAL/Ns/NPs) is a dynamic parameter that depends not only on its analytic sensitivity but also on the pretest probability. The pretest probability may in turn depend on the SARS-CoV-2 exposure history, disease symptoms and local disease prevalence.<sup>3</sup> The unexpected low positivity seen with BAL rRT-PCR in the study despite high pretest probability might be due to delayed time of sampling and/or disease stage. All these factors should have been considered while comparing the diagnostic yield in the study

Bronchoalveolar lavage has a specific place in the diagnostic algorithm of Covid-19 and is usually performed in a patient with lower respiratory tract involvement and high clinical suspicion but negative Ns/NPs result. However, negative BAL results should be interpreted comprehensively in light of different clinical and demographic factors on a case-to-case basis.

## References

1. Geri P, Salton F, Zuccatosta L, et al. Limited role for bronchoalveolar lavage to exclude Covid-19 after negative upper respiratory tract swabs: a multicenter study. *Eur Respir J* 2020 Aug 6;2001733. doi: 10.1183/13993003.01733-2020.
2. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA* 2020; 323: 1843–1844.
3. Woloshin S, Patel N, Kesselheim AS. False Negative Tests for SARS-CoV-2 Infection - Challenges and Implications. *N Engl J Med* 2020;383(6):e38. doi: 10.1056/NEJMp2015897.