

RESEARCH ARTICLE


# Rate dependent influence of arterial desaturation on self-selected exercise intensity during cycling

Saro D. Farra<sup>1</sup>, Stephen S. Cheung<sup>1,2</sup>, Scott G. Thomas<sup>1</sup>, Ira Jacobs<sup>1\*</sup>

<sup>1</sup> Faculty of Kinesiology & Physical Education, University of Toronto, Toronto, Ontario, Canada,

<sup>2</sup> Department of Kinesiology, Brock University, St. Catharines, Ontario, Canada

\* [ira.jacobs@utoronto.ca](mailto:ira.jacobs@utoronto.ca)



## Abstract

The purpose of this study was to clarify if Ratings of Perceived Exertion (RPE) and self-selected exercise intensity are sensitive not only to alterations in the absolute level of arterial saturation ( $S_{pO_2}$ ) but also the rate of change in  $S_{pO_2}$ . Twelve healthy participants ( $31.6 \pm 3.9$  y,  $175.5 \pm 7.7$  cm,  $73.3 \pm 10.3$  kg,  $51 \pm 7$  mL·kg<sup>-1</sup>·min<sup>-1</sup>  $\dot{V}O_{2peak}$ ) exercised four times on a cycle ergometer, freely adjusting power output (PO) to maintain RPE at 5 on Borg's 10-point scale with no external feedback to indicate their exercise intensity. The fraction of inspired oxygen ( $F_{iO_2}$ ) was reduced during three of those trials such that  $S_{pO_2}$  decreased during exercise from starting values (>98%) to 70%. These trials were differentiated by the time over which the desaturation occurred:  $3.9 \pm 1.4$  min,  $-8.7 \pm 4.2\% \cdot \text{min}^{-1}$  (FAST),  $11.0 \pm 3.7$  min,  $-2.8 \pm 1.3\% \cdot \text{min}^{-1}$  (MED), and  $19.5 \pm 5.8$  min,  $-1.5 \pm 0.8\% \cdot \text{min}^{-1}$  (SLOW) ( $P < 0.001$ ). Compared to stable PO throughout the control condition (no  $S_{pO_2}$  manipulation), PO significantly decreased across the experimental conditions (FAST =  $2.8 \pm 2.1$  W·%  $S_{pO_2}^{-1}$ ; MED =  $2.5 \pm 1.8$  W·%  $S_{pO_2}^{-1}$ ; SLOW =  $1.8 \pm 1.6$  W·%  $S_{pO_2}^{-1}$ ;  $P < 0.001$ ). The rates of decline in PO during FAST and MED were similar, with both greater than SLOW. Our results confirm that decreases in absolute  $S_{pO_2}$  impair exercise performance and that a faster rate of oxygen desaturation magnifies that impairment.

## OPEN ACCESS

**Citation:** Farra SD, Cheung SS, Thomas SG, Jacobs I (2017) Rate dependent influence of arterial desaturation on self-selected exercise intensity during cycling. PLoS ONE 12(3): e0171119. doi:10.1371/journal.pone.0171119

**Editor:** Alejandro Lucia, Universidad Europea de Madrid, SPAIN

**Received:** August 17, 2016

**Accepted:** January 15, 2017

**Published:** March 3, 2017

**Copyright:** © 2017 Farra et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper.

**Funding:** This work was supported by the Province of Ontario's Research Program in Applied Sport Science. S. S. Cheung was supported by a Canada Research Chair.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

There are a myriad of mediators that have been associated with the development of fatigue during voluntary exercise such as increasing core body temperature ( $T_C$ ) [1, 2], accumulating concentrations of muscle and blood metabolites [3, 4], energy availability [5–7], and insufficient oxygen ( $O_2$ ) availability and/or transport [8, 9]. The research conducted to date in the area of exercise-induced fatigue demonstrates how the magnitude of changes in various ambient stressors, or the absolute magnitude of physiological strain associated with those stressors, augment the development of exercise-induced fatigue. Much less is known about how the rate of change of these same stressors and strains influence the homeostatic control systems that are thought to regulate the neuromuscular system during exercise. Should the rate of change in physiological strain be shown to influence exercise performance, it would support the

proposition that the neuromuscular system is sensitive to feed-forward homeostatic control during exercise [10, 11], because feed-forward systems can be characterized as being able to differentiate between the instantaneous physiological state and the instantaneous rate of change within the system [12, 13].

It has been suggested that the feed-forward regulation of voluntary exercise performance is based on one's ratings of perceived exertion (RPE) [14], which is defined as the conscious manifestation of the degree of strain experienced during physical work [15]. Although several theories exist regarding the physiological and psychological determinants of RPE [16], research suggests that it is an integration of multiple sources of information including corollary discharge of the efferent output from the motor cortex [17–20], afferent feedback from the periphery [21–26], as well as psychological factors such as previous experience and knowledge of exercise duration [27], motivation [16], positive and negative affect [28] and awareness [29]. Although isolating the feed-forward contribution to voluntary exercise performance can be challenging, Tucker [11] proposed a conceptual model, known as 'constant RPE exercise', that may be useful to further this line of research. In this model, the investigator dictates a target RPE (known as the RPE template), and the subject then maintains that RPE by freely adjusting exercise intensity. According to this model, the control of exercise intensity is the result of error correction feedback, as oscillations in exercise performance occur as the central nervous system (CNS) attempts to minimize the difference between the conscious RPE and the 'RPE template'. When participants exercise at a moderate intensity, they reliably reproduced the exercise intensity and the associated physiological steady state throughout treadmill [30] and cycling [26, 31] exercise in the absence of any knowledge about the exercise intensity or their physiological responses [26, 30, 31]. Other forms of self-paced exercise, such as a time trial does not allow for this consistency, as exercise intensity and the associated physiological responses are constantly changing. Such a consistent and reproducible steady state over the duration of an exercise trial lends itself to experimental manipulation seeking to separate the impact of the absolute magnitude of physiological strain from the rate of physiological strain development on exercise performance as the manipulation can be applied irrespective of time.

It is well established that exercise performance is impaired in environments where oxygen ( $O_2$ ) supply is reduced. Exposure to hypoxia is associated with centrally mediated reductions in muscle activation [32, 33], and the mechanisms responsible for this central impairment shift with increasing hypoxia severity. While exercising in mild hypoxia ( $F_1O_2 = 0.15$  to  $0.16$ ) [32, 34] increasing levels of afferent feedback from fatiguing muscles have been proposed to limit central motor drive (CMD) [3, 35] so that the development of peripheral muscle fatigue does not surpass a critical threshold [35, 36]. In this theoretical framework, hypoxia exacerbates the accumulation of metabolites that stimulate group III & IV afferents [37], triggering inhibitory sensory feedback to the central nervous system (CNS). While exercising in severe hypoxia ( $F_1O_2 = 0.10$ ), the hypoxic stress diminishes the pressure gradient for gas exchange between the lungs and the active tissues [38], which may interfere with neurotransmitter turnover [39] or impair cerebral aerobic metabolism [40–42]. However, the supporting research, for both mild and severe hypoxia, reports how absolute changes in hypoxic stress impaired exercise performance and not whether the rate of change in either hypoxic stress and/or strain may modulate self-selected exercise intensity. Some research speculates that self-selected exercise intensity is sensitive to the rate of physiological strain development [43–45]. With respect to traditional models of exercise-induced fatigue accompanying arterial hypoxemia, it is unclear how and why the rate of physiological strain development would influence exercise performance [38, 46–48]. However, newer theories suggest that the CNS uses both the absolute magnitude and the rate of change in physiological strain as independent modulators of self-

selected exercise intensity [11]. The mechanism of this proposed form of regulation remains currently unknown.

Therefore, the aim of this study was to investigate how different rates of arterial desaturation impact self-selected exercise intensity. As previous research suggests that the rate of physiological strain development influences exercise performance, this study was designed to test the hypothesis that faster rates of arterial deoxygenation will augment the declines in muscle activity and self-selected power output (PO) while cycling at a fixed RPE, as well as the peak power output (PPO) generated during a 5 s maximal effort sprint.

## Materials and methods

### Participants

Twelve participants (8 male and 4 female) volunteered to participate in this study. Sample size calculations were planned to detect a 15 W difference in our main outcome variable, PO at an RPE of 5 on Borg’s 10-point scale. Based on a within-subject standard deviation of 11 W (determined during pilot testing), this sample size was required to detect a treatment difference at the 0.05 level, with a power of 0.80. Participants were healthy, competitive cyclists. The mean physical and physiological characteristics are listed in Table 1. Prior to their commencement, participants were informed of the risks associated with the study and a written informed consent was obtained. Participants were asked to avoid ingesting any food or drink (besides water) for 2 h before testing, to refrain from consuming caffeinated food or beverages for 12 h before testing, and not to exercise for 24 h before any testing session. The Health Sciences Research Ethics Board of the University of Toronto approved the protocol for this study. All experiments conformed to the standards set by the Declaration of Helsinki.

### Protocol

Participants reported to the laboratory on six occasions, with sessions being separated by at least 48 h. Throughout the study, all exercise was preceded by a 15 min warm up on the cycle ergometer at a self-chosen intensity corresponding to a RPE of 5 on Borg’s 10-point scale. During the first visit, basic physiological and physical characteristics were measured, including height, weight, and peak aerobic power ( $\dot{V}O_{2peak}$ ) during cycle ergometry. Participants began the incremental test at 90 W. The workload increased in a step-wise fashion by 30 W every 3 min until 1) the investigators terminated the test because participants were not able to maintain their cadence within 20 RPM of their self-selected target for more than 30 s, or 2) the participant reached volitional fatigue. During the test, participants were connected to a metabolic cart that measured respiratory parameters on a breath-by-breath basis.  $\dot{V}O_{2peak}$  ( $ml \cdot kg^{-1} \cdot min^{-1}$ ) was defined as the highest 30 s average  $O_2$  consumption that was recorded during the test.

**Table 1. Physical and physiological subject characteristics.**

	Age (y)	Height (cm)	Weight (kg)	$P_{max}$ (W)	$\dot{V}O_{2Peak}$ ( $ml \cdot kg^{-1} \cdot min^{-1}$ )	$HR_{max}$ (BPM)	$\dot{V}_{Emax}$ ( $L \cdot min^{-1}$ )
Male	31.3 ± 3.7	179.9 ± 3.4	79.1 ± 6.2	330 ± 32	53 ± 5	190 ± 6	161 ± 24
Female	30.2 ± 5.2	166.7 ± 5.7	61.8 ± 6.1	215 ± 17	46 ± 4	181 ± 11	133 ± 27
Group	30.9 ± 4.1	175.5 ± 7.7	73.3 ± 10.4	293 ± 63	51 ± 6	187 ± 8	154 ± 27

$P_{max}$ , Highest exercise intensity achieved during the incremental step test to exhaustion

doi:10.1371/journal.pone.0171119.t001

During the second visit, the relationship between the partial pressure of inspired O<sub>2</sub> (P<sub>I</sub>O<sub>2</sub>) and arterial saturation (S<sub>p</sub>O<sub>2</sub>) was modeled under isocapnic conditions.

During the four remaining visits, the study utilized a single-blind, crossover design where experimental and control trials were completed in random order. Participants cycled on an isokinetic cycle ergometer at an intensity they felt corresponded to a RPE of 5 on Borg's 10-point scale [25] for 30 min, with the final stage of the  $\dot{V}O_{2peak}$  test as their anchor for what would constitute a "10"; effort was adjusted with fatigue, such that subjective sensation of whole body exertion remained steady at target level of 5 [45]. On three trials, the fraction of inspired O<sub>2</sub> (F<sub>I</sub>O<sub>2</sub>) was reduced to desaturate arterial blood from starting values (>98%) to 70% while maintaining isocapnia. This desaturation was designed to occur approximately linearly over 3 different time periods: 5 min, 15 min, and 25 min for the FAST, MED, and SLOW conditions respectively. When S<sub>p</sub>O<sub>2</sub> reached 70%, F<sub>I</sub>O<sub>2</sub> was surreptitiously switched back to 21% for the remainder of the 30 min trial. During the control trial (CON), F<sub>I</sub>O<sub>2</sub> remained at 21%. Participants were blinded to P<sub>I</sub>O<sub>2</sub> and were kept naive as to the hypotheses of the study. More specifically, participants were made aware of the absolute level of arterial desaturation they were to undergo, but were not informed that different desaturation rates would be used during each of the trials.

Participants also performed several 5 s maximal effort sprints to measure PPO at the following intervals: two sprints prior to commencing the constant RPE trial (PRE), one sprint at a S<sub>p</sub>O<sub>2</sub> of 70% (POST), and two sprints 2 min after POST when S<sub>p</sub>O<sub>2</sub> recovered back to initial values (REC). During PRE and REC, each sprint was separated by 3 min and the trial with the highest PPO was selected for analysis. For some participants, the constant RPE trials were terminated prior to achieving a S<sub>p</sub>O<sub>2</sub> of 70%, when the subject indicated that they wished to stop the trial, or when the investigator observed the participant displaying symptoms that indicated severe hypoxic strain. The participants that were able to achieve a S<sub>p</sub>O<sub>2</sub> of 70% did so repeatedly across conditions. However, when a participant did not achieve a S<sub>p</sub>O<sub>2</sub> of 70% during the first hypoxic trial, the investigator terminated all subsequent constant RPE exercise trials so that the final S<sub>p</sub>O<sub>2</sub>, and the S<sub>p</sub>O<sub>2</sub> during the 5 s sprints, would be similar across conditions. All participants were made aware of the termination criteria.

## Hypoxia administration

**Isocapnic hypoxia system.** The isocapnic hypoxia system (IHS), developed in our laboratory [49], was used to decrease S<sub>p</sub>O<sub>2</sub> while clamping the partial pressure of end-tidal carbon dioxide (PET<sub>CO2</sub>) within 2 mmHg of starting values throughout and in between all exercise trials. Isocapnia was maintained utilizing the principles of sequential gas delivery [50–52]. Briefly, the IHS included a 3-way breathing manifold connected to an inspiratory and expiratory reservoir. The configuration of the manifold was such that it first delivered fresh gas from the inspiratory reservoir, and when this source of gas was depleted, it sequentially delivered previously expired gas from the expiratory reservoir. The flow of fresh gas into the inspiratory reservoir, as well as its composition, were controlled by custom computer software (LabView, National Instruments, Austin, TX) regulating two independent mass flow controllers (Model # 32907–77, Cole-Parmer, Montreal), with each mass flow controller adjusting the flow of one source gas (Tank 1–21% O<sub>2</sub>, balance nitrogen [N<sub>2</sub>]; Tank 2–5% O<sub>2</sub>, balance N<sub>2</sub>). Throughout the trial, the investigator adjusted the gas flow rate manually and the software automatically regulated changes in gas composition according to user-defined parameters as indicated below. At the start of each trial, the investigators monitored PET<sub>CO2</sub> and adjusted the rate of fresh gas delivery as required to clamp PET<sub>CO2</sub> during the exercise trials.

**Modeling the  $P_1O_2$ - $S_pO_2$  relationship.** Participants were connected to the IHS and cycled at an intensity that was associated with their first ventilatory breakpoint, determined during the  $\dot{V}O_{2peak}$  test [53, 54]. Participants exercised at this intensity under isocapnic conditions, while  $F_I O_2$  levels were decreased every 3 min (21%, 18%, and 14%).  $S_pO_2$  was recorded at the end of each stage.  $F_I O_2$  was multiplied by barometric pressure to calculate  $P_1O_2$  (mmHg). The relationship between  $P_1O_2$  and  $S_pO_2$  was individually modelled for each subject with a second order polynomial function. Eq 1 exemplifies this function for one subject:

$$P_1O_2 = 0.12x^2 - 16.5x + 679.5 \quad (1)$$

where  $x = S_pO_2$  (%).

**Progressive arterial desaturation.** During the first 5 min of the four constant RPE trials, the IHS was supplied with 21%  $O_2$  so that participants could establish a level of voluntary effort that corresponded to the target RPE. At the start of the of the desaturation procedure, the  $F_I O_2$  of the inspire was continuously reduced over time according to a second order polynomial function (Eq 2). For each experimental condition, a different function was developed to decrease  $S_pO_2$  linearly over the target duration of the trial (Eq 2), according to the  $P_1O_2$ - $S_pO_2$  relationship (Eq 1) previously established. Eqs 2A–2C exemplify these functions for one subject:

$$P_1O_2 = 4.1x^2 - 36.2x + 157.2 \quad (2A, \text{FAST})$$

$$P_1O_2 = 0.5x^2 - 12.1x + 157.2 \quad (2B, \text{MED})$$

$$P_1O_2 = 0.2x^2 - 7.2x + 157.2 \quad (2C, \text{SLOW})$$

where  $x = \text{Time (min)}$ .

**Data acquisition.** All data were synchronously recorded (Power lab 16/35, ADInstruments, Australia), digitized and stored on a laptop computer for later analysis.

**$S_pO_2$  and Heart Rate (HR).**  $S_pO_2$  and HR were sampled at 1 Hz using a pulse oximeter (Model # 7500, Nonin Medical, USA). The left earlobe was prepared by applying hyperemic cream (Finalgon, Boehringer-Ingelheim, France) for 8 min. The probe was attached to the earlobe after cleaning the area with isopropyl alcohol.

**Cycle ergometry.** All cycling exercise was performed on an electronically braked cycle ergometer (Excalibur Sport, Lode, Groningen, Netherlands), which sampled PO at 5 Hz. This ergometer can be used for exercising at a specific absolute PO within a broad range of pedaling frequencies (hyperbolic mode), or it can be used to regulate the rate of pedaling (RPM) at a target cadence and register the power that is being generated (isokinetic mode). During the  $\dot{V}O_{2peak}$  test the ergometer was set to its hyperbolic mode. During all constant RPE exercise trials and 5 s sprints, the ergometer operated in its isokinetic mode where participants self-selected their preferred RPM. The power that they chose to generate was recorded continuously and participants were required to complete all subsequent cycling exercise at that cadence. Because the ergometer regulated cadence at a set rate, participants could only alter PO by increasing or decreasing their force on the pedals to maintain RPE at a 5 of Borg's 10-point. No visual or verbal feedback was provided except for the occasional reminder to sustain a consistent perceived effort throughout the trial. The participants used the same footwear and pedals for every session. Each subject's set-up was recorded in the first session and was subsequently reproduced for the remaining tests.

**Surface electromyography.** Surface electromyography (sEMG) signals from the vastus medialis (VM) and vastus lateralis (VL) were measured via bipolar Ag-Ag-Cl surface

electrodes (inter-electrode distance 2 cm). Signals were amplified with an isolated differential amplifier (Dual Bioamp FE 135, ADInstruments, Australia), band pass filtered (10–500Hz), and digitized with a sampling frequency of 2000 Hz. Electrodes were positioned over the muscle belly along the assumed angle of pennation after shaving and cleaning the area with isopropyl alcohol. Electrodes and wires were first anchored using adhesive tape (Ref # 71443–02, Hypafix, Germany), which was then covered with a mesh sleeve (Cat # GL-705, Surgilast, Canada) to reduce movement artifact. sEMG signals for both muscles were quantified by Root-Mean-Square (sEMG<sub>RMS</sub>). During constant RPE exercise, sEMG<sub>RMS</sub> values were normalized to the initial 30 s average value when S<sub>p</sub>O<sub>2</sub> was roughly 100%. During the 5 s sprint sEMG<sub>RMS</sub> were normalized to PRE values.

**Near infrared spectroscopy.** Indicators of cerebral and muscle oxygenation were measured with near infra-red spectroscopy (NIRS) (Niro 300, Hamamatsu, Japan) with a sampling rate of 1 Hz. NIRS measured tissue oxygenation index (TOI) and oxy-hemoglobin (O<sub>2</sub>Hb) in response to arterial desaturation. The cerebral probe was positioned over the left pre-frontal cortex (PFC) 1 cm above the eyebrow and 1 cm to the left of the skull centre [55]. The muscle probe was positioned over the right VL along the vertical axis of the thigh, approximately 10–14 cm from the knee joint [56]. The probes were fixed using a dense rubber vinyl holder and held in place with adhesive tape (Ref # 71443–02, Hypafix, Germany) and covered with a mesh sleeve (Cat # GL-705, Surgilast, Canada). Measures are expressed as relative changes from the start of the trial.

**Minute ventilation.**  $\dot{V}_E$  was determined on a breath-by-breath basis using a heated pneumotach (Model # 3813, Hans Rudolph, USA) positioned on the expiratory limb of the sequential gas delivery breathing manifold. Airflow was sampled at 100 Hz and signals were low pass filtered (1 Hz).  $\dot{V}_E$  was calculated as:

$$\dot{V}_E = \frac{V_T}{(T_B/60)} \quad (3)$$

where,

$V_T$  = Tidal Volume (L) = integral of flow – time signal and,

$T_B$  = Breathing Time (s) = time between the start of two successive expirations.

**Partial pressure of O<sub>2</sub> and CO<sub>2</sub>:** The partial pressure of O<sub>2</sub> and CO<sub>2</sub> were continuously measured at 100 Hz using a gas analyzer (O2CapB, Oxigraf, USA) sampling near the mouth. Inspired and end-tidal values for O<sub>2</sub> and CO<sub>2</sub> were determined on a breath-by-breath basis using the built in “cyclic measurements” functions of our data acquisition software (LabChart V8.0 Pro, ADInstruments, Australia).

## Data processing

To analyze decrements in self-selected PO relative to reductions in S<sub>p</sub>O<sub>2</sub>, 15 s averages were calculated for PO and S<sub>p</sub>O<sub>2</sub>. To estimate the time delay between the start of arterial desaturation and the subsequent decrease in PO [57], the point at which PO in each hypoxic trial began to deviate relative to CON was determined by visual inspection. To determine the slopes of the S<sub>p</sub>O<sub>2</sub> vs. time, PO vs. S<sub>p</sub>O<sub>2</sub> and sEMG<sub>RMS</sub> vs. S<sub>p</sub>O<sub>2</sub> relationships using linear regression, 5 s averages were calculated for S<sub>p</sub>O<sub>2</sub>, PO, and sEMG<sub>RMS</sub>. To illustrate the change in HR,  $\dot{V}_E$ , and NIRS responses over the course of the constant RPE exercise bout, 30 s averages were calculated at defined stages during the trial. INITIAL represents the average of the first 30 s, while FINAL denotes the average of the last 30 s. For the three experimental conditions, MIDDLE



represent the 30 s average at a  $S_{pO_2}$  of 85% (midpoint with respect to  $S_{pO_2}$ ), while for CON it denotes the midpoint with respect to time. During the 5 s sprints, PPO was calculated as the average PO over the 5 s effort, while the difference in  $sEMG_{RMS}$  between POST and PRE as well as REC and PRE were calculated.

### Statistical analysis

Results are presented as mean values  $\pm$  standard deviation (SD). Differences were considered significant when  $P < 0.05$ . A one-way repeated measures analysis of variance (RM-ANOVA) was performed to evaluate the differences in the slope of the  $S_{pO_2}$  vs time, PO vs  $S_{pO_2}$ , and  $sEMG_{RMS}$  vs.  $S_{pO_2}$  between the experimental conditions. All other responses during the constant RPE trials were analyzed using a two-way RM-ANOVA (Condition: FAST, MED, SLOW, CON X Stage: INITIAL, MIDDLE, FINAL). Four participants voluntarily stopped some exercise trials prior to achieving a  $S_{pO_2}$  of 70%. Therefore PO at 70%  $S_{pO_2}$  was not included in that analysis. Responses during the 5 s effort were analyzed using a two-way RM-ANOVA (Condition: FAST, MED, SLOW, CON X  $S_{pO_2}$ : POST, REC). When the assumption of sphericity was violated, a Greenhouse-Geisser correction was used. Significant findings were followed up with *post hoc* pairwise analysis using a Bonferroni adjustment. All statistics were calculated using SPSS software (Version 22 for Mac, IBM).

## Results

### Arterial saturation

$S_{pO_2}$  was at or above 98% at the start of each exercise trial (FAST =  $98 \pm 1\%$ ; MED =  $98 \pm 1\%$ ; SLOW =  $98 \pm 1\%$ ; CON =  $100 \pm 1\%$ ).  $S_{pO_2}$  remained above 98% throughout CON, which lasted  $22.7 \pm 3.7$  min. Technical equipment difficulties caused early termination of exercise during CON for some subjects before completing the full trial.  $S_{pO_2}$  decreased from starting values to approximately 70% (or the lowest tolerable  $S_{pO_2}$ ; three participants reached 75%; one participant achieved 80%), in  $3.9 \pm 1.4$  min,  $11.0 \pm 3.7$  min,  $19.5 \pm 5.8$  min for the FAST, MED, and SLOW conditions respectively, with the rate of arterial desaturation significantly different between the conditions (Table 2) ( $P < 0.001$ ). Representative data from one subject, illustrating the  $S_{pO_2}$  vs. time response, are shown in Fig 1 (top).

**Table 2. Rates of change during constant RPE exercise trials for FAST, MED, and SLOW.**  $S_{pO_2}$  remained at or above 98% throughout CON.

	FAST	MED	SLOW
$S_{pO_2}$ (%•min <sup>-1</sup> )	$-8.7 \pm 4.2^{1, 2, 3}$	$-2.8 \pm 1.3^{1, 2}$	$-1.5 \pm 0.8^1$
PO (W•% $S_{pO_2}^{-1}$ )	$2.8 \pm 2.1^2$	$2.5 \pm 1.8^2$	$1.8 \pm 1.6$
VM $sEMG_{RMS}$ (%•% $S_{pO_2}^{-1}$ )	$1.3 \pm 0.6^2$	$1.1 \pm 0.5^2$	$0.7 \pm 0.7$
VL $sEMG_{RMS}$ (%•% $S_{pO_2}^{-1}$ )	$1.2 \pm 0.6^2$	$0.9 \pm 0.6$	$0.6 \pm 0.5$
VL TOI (%•% $S_{pO_2}^{-1}$ )	$0.3 \pm 0.1$	$0.3 \pm 0.1$	$0.3 \pm 0.1$
VL O <sub>2</sub> Hb (%•% $S_{pO_2}^{-1}$ )	$0.2 \pm 0.1$	$0.2 \pm 0.1$	$0.2 \pm 0.1$
PFC TOI (%•% $S_{pO_2}^{-1}$ )	$0.4 \pm 0.2$	$0.4 \pm 0.2$	$0.4 \pm 0.1$
PFC O <sub>2</sub> Hb (%•% $S_{pO_2}^{-1}$ )	$0.3 \pm 0.1$	$0.4 \pm 0.1$	$0.3 \pm 0.1$


Values are means  $\pm$  SD; n = 12

<sup>1</sup> Significantly different from CON ( $P < 0.05$ )

<sup>2</sup> Significantly different from SLOW ( $P < 0.05$ )

<sup>3</sup> Significantly different from MED ( $P < 0.05$ )

doi:10.1371/journal.pone.0171119.t002



**Fig 1. Representative data from one subject illustrating changes in  $S_{pO_2}$  (%) over time (min) (top) and PO (W) over time (min) (bottom).** Constant RPE exercise was performed for 30 min. The subject was breathing normoxic gas for the first 5 min during the exercise period so that they could establish a level of voluntary effort that corresponded to a RPE of 5. The desaturation procedure was initiated at 0 min, which was the official start of the experimental trials.

doi:10.1371/journal.pone.0171119.g001

### Power output

The average cadence selected by the participants was  $91 \pm 6$  RPM. There was a significant interaction between  $S_{pO_2}$  and condition on PO during the constant RPE trials ( $P = 0.002$ ) (Table 3). The PO voluntarily chosen and maintained during the first 15 s of exercise was similar on all trials. With reductions in  $S_{pO_2}$ , PO progressively decreased across conditions at significantly different rates ( $P < 0.001$ ). The rate at which PO decreased in FAST and MED was similar, and both had a greater rate of decrease when compared to SLOW (Table 2). During CON, there was no significant change in PO during the trial (INITIAL =  $146 \pm 66$  W; MIDDLE =  $149 \pm 64$  W; FINAL =  $146 \pm 58$  W). Representative data from one subject, illustrating the PO vs. time response, are shown in Fig 1 (bottom).

Upon the onset of a decrease in  $S_{pO_2}$ , the approximate time delay for beginning the decrease in PO was significantly different between each condition (FAST =  $0.1 \pm 0.2$  min;



**Table 3. Power output (W) at different absolute  $S_{pO_2}$  during constant RPE exercise trials for FAST, MED, and SLOW.** The initial power output chosen by participants during CON was  $146 \pm 66$  and remained at this level through out the trial.

	~100%	90%	80%
	$S_{pO_2}$	$S_{pO_2}$	$S_{pO_2}$
FAST	$149 \pm 68$	$125 \pm 54$	$94 \pm 40^2$
MED	$149 \pm 67$	$128 \pm 62$	$99 \pm 46^2$
SLOW	$146 \pm 66$	$140 \pm 66$	$115 \pm 55$


Values are means  $\pm$  SD; n = 12

<sup>2</sup> Significantly different from SLOW ( $P < 0.05$ )

doi:10.1371/journal.pone.0171119.t003

MED =  $1.8 \pm 1.0$  min; SLOW =  $5.6 \pm 2.3$  min;  $P < 0.05$ ). Representative data from one subject, illustrating the time delay in decreasing PO, are shown in Fig 2.

There was a significant interaction between condition and  $S_{pO_2}$  on PPO ( $P < 0.001$ ). Post hoc testing revealed that PRE PPO was similar between all of the conditions. During CON, there was a significant decrease in PPO from PRE to POST, however PPO remained at a similar level between POST and REC. When  $S_{pO_2}$  was reduced to 70%, FAST, MED, and SLOW POST PPO significantly decreased to similar levels and all were less than CON. When  $S_{pO_2}$  recovered back to ~ 100% after the constant RPE trials, FAST, MED, and SLOW REC PPO significantly increased from POST and were all similar to CON, but did not recover back to PRE values (Table 4).



**Fig 2. Representative data from one subject illustrating the time delay between the start of arterial desaturation and the subsequent decrease in PO.** Dashed lines indicate when FAST, MED, and SLOW began to deviate from CON. The shaded area represents the lag time between the start of arterial desaturation and the ensuing decrease in PO.

doi:10.1371/journal.pone.0171119.g002

**Table 4. Power output and sEMG<sub>RMS</sub> responses during the 5 s sprints.**

	Stage	FAST	MED	SLOW	CON
PPO (W)	PRE	737 ± 265	733 ± 263	742 ± 268	744 ± 271
	POST	551 ± 204 <sup>1, a</sup>	532 ± 188 <sup>1, a</sup>	570 ± 214 <sup>1, a</sup>	662 ± 248 <sup>a</sup>
	REC	685 ± 244 <sup>a, b</sup>	661 ± 250 <sup>a, b</sup>	669 ± 253 <sup>a, b</sup>	668 ± 235 <sup>a</sup>
VM sEMG <sub>RMS</sub> (Δ PRE)	POST	-28 ± 15	-32 ± 11	-23 ± 15	-18 ± 13
	REC	-8 ± 17 <sup>b</sup>	-16 ± 11 <sup>b</sup>	-14 ± 13 <sup>b</sup>	-15 ± 17
VL sEMG <sub>RMS</sub> (Δ PRE)	POST	-22 ± 13	-22 ± 11	-19 ± 11	-16 ± 9
	REC	-10 ± 7 <sup>b</sup>	-12 ± 11 <sup>b</sup>	-13 ± 12 <sup>b</sup>	-18 ± 7

Values are means ± SD; PPO n = 12; sEMG<sub>RMS</sub> n = 10

<sup>1</sup> Significantly different from CON (P < 0.05)

<sup>a</sup> Significantly different from PRE (P < 0.05)

<sup>b</sup> Significantly different from POST (P < 0.05)

doi:10.1371/journal.pone.0171119.t004

### Surface electromyography

With decreases in S<sub>p</sub>O<sub>2</sub>, sEMG<sub>RMS</sub> decreased at significantly different rates from the beginning of each constant RPE trial in VM (P < 0.001). The FAST and MED were similar, but each had a greater rate of decrease when compared to SLOW. sEMG<sub>RMS</sub> also decreased at significantly different rates in the VL (P < 0.001). Post hoc testing revealed that the rate of decrease was greater in FAST when compared to SLOW (Table 2). During CON, there was no significant change in sEMG<sub>RMS</sub> from initial values for either VM (INITIAL = 100 ± 0%; MIDDLE = 103 ± 10%; FINAL = 100 ± 16%) or VL (INITIAL = 100 ± 0%; MIDDLE = 103 ± 11%; FINAL = 99 ± 11%) during the trial.

There was a significant interaction between condition and S<sub>p</sub>O<sub>2</sub> on sEMG<sub>RMS</sub> for the VM (P = 0.017) and VL (P = 0.002) during the 5 s sprint. Follow up tests revealed that sEMG<sub>RMS</sub> values increased from POST to REC during FAST, MED, and SLOW. However, sEMG<sub>RMS</sub> during CON remained at similar values between POST and REC. Both the VM and VL exhibited these trends (Table 4).

### Tissue oxygenation index

There was a significant interaction between condition and stage on TOI of the VL (P < 0.001) and PFC (P < 0.001). Follow up tests revealed that INITIAL TOI values in the VL were similar across all conditions. With reductions in S<sub>p</sub>O<sub>2</sub>, the decrease in TOI of the VL during FAST, MED, and SLOW was similar and all were less than CON by the end of the trial. Follow up tests also revealed that starting TOI values in the PFC were similar across all conditions. With reductions in S<sub>p</sub>O<sub>2</sub>, the decrease in TOI of the PFC during FAST, MED, and SLOW was similar and all were less than CON by the end of the trial (Table 5).

### Oxyhemoglobin

There was a significant interaction between condition and stage on O<sub>2</sub>Hb of the VL (P < 0.001) and PFC (P < 0.001). Follow up tests revealed that starting O<sub>2</sub>Hb values in the VL were similar across all conditions. With reductions in S<sub>p</sub>O<sub>2</sub>, the decrease in O<sub>2</sub>Hb of the VL during FAST, MED, and SLOW was similar and all were less than CON. Follow up tests also revealed that starting O<sub>2</sub>Hb values in the PFC were similar across all conditions. With reductions in S<sub>p</sub>O<sub>2</sub>, the decrease in O<sub>2</sub>Hb of the PFC during FAST, MED, and SLOW was similar. However O<sub>2</sub>Hb during CON increased throughout the trial (Table 5).

**Table 5. NIRS responses at different trial stages during constant RPE exercise.**

	Stage	FAST	MED	SLOW	CON
<u>Vastus Lateralis</u>					
TOI (%)	Initial	56 ± 8	58 ± 6	56 ± 9	56 ± 6
	Middle	52 ± 8	53 ± 7	51 ± 9	55 ± 6
	Final	49 ± 8 <sup>1</sup>	50 ± 7 <sup>1</sup>	48 ± 8 <sup>1</sup>	54 ± 7
O <sub>2</sub> Hb (% change from initial)	Initial	-0.2 ± 0.8	0.1 ± 0.9	0.0 ± 0.8	-0.1 ± 0.6
	Middle	-2.7 ± 1.6 <sup>1</sup>	-3.2 ± 1.4 <sup>1</sup>	-2.9 ± 1.9 <sup>1</sup>	0.0 ± 2.2
	Final	-4.7 ± 2.1 <sup>1</sup>	-5.1 ± 2.6 <sup>1</sup>	-5.4 ± 3.2 <sup>1</sup>	0.3 ± 2.8
<u>Pre-Frontal Cortex</u>					
TOI (%)	Initial	66 ± 5	66 ± 4	64 ± 6	64 ± 6
	Middle	60 ± 5	61 ± 5	58 ± 7 <sup>1</sup>	64 ± 6
	Final	57 ± 6 <sup>1</sup>	56 ± 7 <sup>1</sup>	53 ± 6 <sup>1</sup>	63 ± 7
O <sub>2</sub> Hb (% change from initial)	Initial	-0.1 ± 1.1	0.4 ± 1.4	-0.4 ± 1.2	0.4 ± 0.8
	Middle	-3.8 ± 2.4 <sup>1</sup>	-4.2 ± 2.4 <sup>1</sup>	-4.2 ± 2.4 <sup>1</sup>	2.6 ± 1.9
	Final	-7.5 ± 3.0 <sup>1</sup>	-8.7 ± 4.3 <sup>1</sup>	-8.1 ± 3.6 <sup>1</sup>	3.1 ± 2.9

Values are means ± SD; n = 12

<sup>1</sup> Significantly different from CON (P < 0.05)

doi:10.1371/journal.pone.0171119.t005

### Heart rate

There was a significant interaction between condition and stage on HR (P = 0.001). HR increased with reductions in S<sub>p</sub>O<sub>2</sub> from 100 to 85% and then remained at a similar value as S<sub>p</sub>O<sub>2</sub> continued to decrease to 70% for FAST, MED, and SLOW. During CON, HR continuously increased throughout the trial. However, further analysis revealed that starting HR values and the HR response throughout the trials were similar across the four experimental conditions (Table 6).

### Minute ventilation

There was a significant interaction between condition and stage on  $\dot{V}_E$  (P < 0.001). Follow up tests revealed that starting  $\dot{V}_E$  values were similar across all conditions. Although  $\dot{V}_E$  increased throughout the trial for each condition, the change in  $\dot{V}_E$  was similar for FAST, MED, and SLOW with decreasing S<sub>p</sub>O<sub>2</sub> and all were greater than CON (Table 6).

**Table 6. Cardiorespiratory responses at different trial stages during constant RPE exercise.**

	Stage	FAST	MED	SLOW	CON
HR (BPM)	Initial	141 ± 18	144 ± 19	137 ± 21	143 ± 17
	Middle	146 ± 18	151 ± 17	150 ± 19	147 ± 19
	Final	146 ± 16	153 ± 18	152 ± 20	153 ± 18
$\dot{V}_E$ (L•min <sup>-1</sup> )	Initial	64 ± 21	63 ± 18	60 ± 20	60 ± 17
	Middle	75 ± 26 <sup>1</sup>	77 ± 25 <sup>1</sup>	75 ± 25 <sup>1</sup>	63 ± 20
	Final	78 ± 29 <sup>1</sup>	84 ± 24 <sup>1</sup>	84 ± 28 <sup>1</sup>	65 ± 21
PET <sub>CO2</sub> (mmHg)	Initial	42 ± 4	42 ± 3	41 ± 3	40 ± 4
	Middle	40 ± 4	42 ± 3	41 ± 3	38 ± 4
	Final	41 ± 4	42 ± 3	41 ± 3	37 ± 4

Values are means ± SD; n = 12

<sup>1</sup> Significantly different from CON (P < 0.05)

doi:10.1371/journal.pone.0171119.t006

## Partial pressure of end-tidal CO<sub>2</sub>

There was a significant effect of S<sub>p</sub>O<sub>2</sub> on fresh gas delivery ( $P < 0.001$ ). Follow up tests revealed that the initial rates of fresh gas delivery at the commencement of the trial and the decrease over the course of the exercise bout were similar for FAST, MED, and SLOW. With decreases in fresh gas delivery, PET<sub>CO<sub>2</sub></sub> values remained within 2 mmHg throughout each trial and in between FAST, MED, and SLOW (Table 6).

## Discussion

This study is the first report of the influence of different rates of change in S<sub>p</sub>O<sub>2</sub> on self-selected exercise intensity during constant RPE exercise. Despite exercising for less time, reaching a similar absolute change in S<sub>p</sub>O<sub>2</sub> and achieving a comparable estimated oxygenation status in cerebral and muscle tissues, the relationship between self-selected exercise intensity and muscle activation with S<sub>p</sub>O<sub>2</sub> was altered, such that a faster arterial deoxygenation rate was associated with a greater decline in submaximal self-selected work rate. These results suggest that the rate of arterial deoxygenation has a central depressant effect on submaximal self-selected exercise intensity, which is independent from the absolute level of S<sub>p</sub>O<sub>2</sub>. In contrast, the decline in PPO, and the associated sEMG, during the 5 s sprint was not affected by the rate of change of S<sub>p</sub>O<sub>2</sub>, suggesting that the magnitude of impairment to the overall capacity of the neuromuscular system was similar.

This investigation corroborates previous research illustrating that hypoxia impairs continuous submaximal exercise performance. Using various exercise modalities, previous research has illustrated that hypoxia increases the time to traverse a given distance [58, 59], and decreases the tolerance time at a given absolute work rate [60, 61]. Several theories of hypoxia-induced fatigue have been proposed that could explain some of the fatigue observed between CON and the experimental conditions. First decreasing brain oxygenation [40] has been reported to affect the ability to sustain a given exercise intensity when S<sub>p</sub>O<sub>2</sub> is below a critical level of approximately 70% [62]. However, as four of our participants chose to terminate the trial prior to achieving a S<sub>p</sub>O<sub>2</sub> of 70%, differences in exercise performance across conditions were analyzed when S<sub>p</sub>O<sub>2</sub> was as high as 80% (Table 3), well above the level that cerebral deoxygenation has been suggested to influence exercise. Second, respiratory muscle demand during continuous high intensity exercise exacerbates the increase in sympathetic vasoconstrictor tone in the exercising limb and contributes to locomotor muscle fatigue by decreasing limb vascular conductance and blood flow [32, 34]. However,  $\dot{V}_E$ , and presumably the work of breathing, were similar across the experimental conditions (Table 5) suggesting that their influence on sympathetic vasoconstriction was similar. Furthermore, the work of breathing associated with submaximal exercise intensities similar to this study, did not contribute to the vasoconstrictor activity present in working locomotor muscles [63]. Third, direct recordings from group III and IV afferents from the triceps surae muscle of artificially ventilated, barbiturate-anesthetized cats found that hypoxia ( $P_aO_2 = 23$  mmHg) increased the baseline discharge frequency of these afferents at rest as well as potentiated the activity within these afferents during electrically-induced static muscle contraction when compared to normoxia ( $P_aO_2 = 108$ ) [64]. The increase in baseline firing frequency has been suggested to be independent of metabolic changes within the muscle, as the metabolites that stimulate these afferents, such as lactic acid and H<sup>+</sup>, were unaffected when acute hypoxic stimuli were given at rest [64, 65]. Given their prolific innervation of muscle tissue, small changes in their individual discharge rates, would substantially influence their collective input into the CNS [66]. Fourth, a decrease in O<sub>2</sub> availability decreases  $\dot{V}O_{2max}$  in a dose dependent manner [59, 67], while RPE is reflective of relative physical strain [15]. It stands to reason that with a decrease in S<sub>p</sub>O<sub>2</sub>,  $\dot{V}O_{2max}$  would

decrease, thus increasing relative exercise intensity for a given absolute workload. To maintain the same RPE during progressive hypoxia, participants would have to continuously decrease PO to keep their perceived exertion on target.

Although this investigation confirms that a decrease in the absolute magnitude of O<sub>2</sub> availability impairs submaximal exercise performance, the novel finding of this study is that the impairment to self-selected exercise intensity was magnified when arterial desaturation occurred at a faster rate, at a time when O<sub>2</sub> availability was similar across conditions (i.e. S<sub>p</sub>O<sub>2</sub> = 80%). As this study was the first to employ an experimental design to investigate the independent effect of different arterial desaturation rates on self-selected exercise intensity, it is difficult to compare our results with those of previous reports. However, it is unlikely that the alterations to  $\dot{V}O_{2max}$ , as well as to the bioenergetic processes within the peripheral tissues, resulting from our hypoxic manipulations had a major differential influence on self-selected exercise intensity and sEMG between FAST, MED, and SLOW when S<sub>p</sub>O<sub>2</sub> was at 80%. While O<sub>2</sub> delivery was compromised, O<sub>2</sub> availability was similar, which should lead to a comparable influence on  $\dot{V}O_{2max}$ , tissue aerobic metabolism and subsequently self-selected exercise intensity. As NIRS has been used to directly compare the metabolic state of tissues in various environmental conditions [40], our data illustrate comparable metabolic environments within the respective tissues (Table 4). With respect to the brain, a similar oxygenation status should have similar effects on cerebral metabolism and subsequently CMD generation. With regard to muscle, a comparable metabolic milieu should lead to comparable levels of inhibitory afferent feedback as well as similar effects on muscle metabolism. As the magnitude of tissue deoxygenation was similar between conditions, its direct impact on aerobic metabolism was also likely similar between conditions: i.e. it likely did not have a differential influence on self-selected exercise intensity.

Our data suggest that the generation of self-selected exercise intensity at a constant RPE is sensitive to both the rate of change and absolute magnitude of arterial deoxygenation. The mechanism(s) by which the rate of arterial desaturation modulates the integration of various factors that influence RPE and subsequently the relationship between RPE, self-selected exercise intensity and sEMG remains unknown. However, given the striking similarities in the neuroanatomy involved, the receptors that initiate signals regulating cardiopulmonary function during exercise in hypoxia, may also regulate CMD and muscle activation. Central to this paradigm is the model proposed by Mateika and Duffin [68] highlighting the role of central chemoreceptors, peripheral chemoreceptors, and group III & IV afferents in maintaining a respiratory steady state during exercise. Central chemoreceptors respond to increases in interstitial or CSF H<sup>+</sup> concentration [69] arising from increasing P<sub>a</sub>CO<sub>2</sub> levels. As PET<sub>CO2</sub>, and presumably P<sub>a</sub>CO<sub>2</sub>, was maintained within 2 mmHg within each trial and across conditions, central chemoreceptor output should have been controlled, likely not contributing to the modulation of exercise intensity. Based on the available evidence, we can speculate that the metabolic perturbations associated with exercise, combined with the additional stresses of hypoxia, stimulate feedback from either peripheral chemoreceptors and/or group III & IV afferents that project onto and alter the output of the solitary tract nucleus (NTS) [68]. The output from the NTS has several targets including the insular cortex [70]. Williamson *et al.* [71] describe the insular cortex as a likely area involved in the generation of perceived exertion, while Tanaka and Watanabe [72] describe the insular cortex as an area that projects inhibitory signals onto the motor cortex. The hypoxia-induced decrements in self-selected exercise intensity and muscle activation were amplified during constant RPE exercise but not during the 5 s sprint, suggesting that the rate of arterial deoxygenation activates a central adjustment modifying the relationship between RPE and PO rather than alterations in neuromuscular capacity.

Although, the proposition that RPE acts as a feed-forward controller has some experimental support [73], its role in feed-forward control remains unclear. However, based on the findings of this investigation, several processes may be postulated that could modulate the physiological and psychological processes involved in the generation of self-selected exercise intensity, which would require further research to confirm.

## Hypothesized mechanisms of homeostatic control

**Central information processing.** As indices of gross respiratory strain,  $S_pO_2$ , HR,  $\dot{V}_E$ , and NIRS parameters were similar between FAST, MED, and SLOW at the end of the trial, the magnitude of the associated afferent signaling was presumably also similar. If confirmed, it would suggest that the CNS independently utilizes the rate of change and absolute magnitude of physiological strain development as concomitant modulators of RPE, CMD, and self-selected exercise intensity. Using computational modeling, Puccini *et al* [74] demonstrated that cortical networks effectively anticipate incoming synaptic inputs by analyzing the magnitude of the stimulus, in combination with its rate of change. Such anticipation supports a proposition that the feed-forward homeostatic control of the neuromuscular system is a function of supraspinal processes.

**Sensory adaptation.** The sensitivity of sensory receptors change in response to the stimulus over time [75]. After reaching a maximum discharge rate, some reports have demonstrated a progressive and significant decline in peripheral chemoreceptor firing activity over time [76] while others have not [77], despite a constant level of hypoxia. Further work is required to determine if sensory adaptation decreases the sensitivity of human peripheral receptors with progressive changes in stimuli over time. Should this theory be confirmed then the attenuated fatigue response during SLOW when compared to FAST and MED could be explained by a decreased level of feedback projecting onto supraspinal centres that alter the generation of RPE and CMD as well as possibly a reduced level of inhibitory feedback projecting onto the MN at the spinal level. In other words, the longer the time required to reach an absolute  $S_pO_2$ , less feedback would arise to modulate self-selected exercise intensity.

**Rate sensitive output from peripheral receptors.** There are several reviews articles summarizing the stimuli that stimulate central and peripheral chemoreceptors as well as group III & IV afferents [68, 78]. The studies cited in those reviews examined the impact of absolute changes in metabolite concentration on the output of receptors. We are not aware of any investigations that have determined if peripheral receptors are also sensitive to the rate of change in their stimuli. Should this theory be confirmed then the magnified fatigue response during FAST and MED when compared to SLOW could be explained by a greater level of feedback projecting onto supraspinal centres that alter the generation of RPE and CMD as well as possibly a greater level of inhibitory feedback projecting onto the MN at the spinal level. In other words, the faster the rate of physiological strain development, the greater the output from the sensor for a given  $S_pO_2$ .

**Spinal modulation.** According to Marcora [19], RPE is centrally generated by corollary discharge from motor to sensory areas of the cerebral cortex and not related to afferent feedback from the periphery. This theory has received experimental support in investigations where curare was used to partially block NMJ transmission. With the curare block, the increase in CMD required to exercise at the same workload elicited a greater perceived effort [18]. Accordingly, the decreases in muscle activation during constant RPE exercise could be the result of inhibitory spinal modulation, as CMD would have remained at a similar level. Should this theory be confirmed then the magnified fatigue response during FAST and MED when compared to SLOW could be explained by either sensory adaptation or a rate sensitive output from the peripheral receptors.



## Limitations

The normal function of the human body during exercise is reliant on the effective integration of a multitude of physiological systems, which we have yet to fully comprehend and appreciate. Although our measures indicated similar amounts of gross respiratory strain across conditions, we did not measure the regional biochemical alterations within the active tissues. For example, the synthesis and release of excitatory neurotransmitters such as acetylcholine [79, 80], aromatic monoamines [79, 81], as well as glutamate and aspartate [82, 83] are reduced during acute hypoxia, while the synthesis and release of the inhibitory neurotransmitters and neuromodulators such as  $\gamma$ -aminobutyric acid (GABA) [83–85], adenosine [86], beta-alanine [82, 83], taurine [87], endogenous opioids [88, 89], and lactate [82, 90] are increased. The shift in the balance between excitatory and inhibitory neuro-effectors would hyperpolarize neuronal membranes, possibly impairing motor function by limiting the generation of CMD. However, given the time required for the body to achieve a cellular steady state during acute hypoxia, the likelihood of neuronal hyperpolarization increases with longer hypoxic exposure, minimizing its influence on the shorter trials. Therefore, if there were differences in the biochemical environment between conditions, the difference in the performance decrement would have likely been underestimated in this study, strengthening the conclusions regarding the independent depressant influence of a faster rate of arterial deoxygenation.

Furthermore, differences in sEMG during constant RPE trials between conditions suggest that central factors contributed to the rate dependent relationship between  $S_pO_2$  and self-selected exercise intensity. The underlying assumption is that for a given  $S_pO_2$ , all else in the  $O_2$  transport pathway remains similar between conditions. Although the NIRS parameters used in this study suggest that a similar level of cerebral and muscle oxygenation was achieved across conditions, further work is required to confirm this assumption. For example, future work could explore the influence of different deoxygenation rates on the regulation of peripheral factors such as muscle blood flow, blood pressure, and cardiac output.

## Conclusions

When controlling for absolute decreases in  $S_pO_2$ , a faster rate of arterial deoxygenation was associated with a greater decrement in self-selected exercise intensity than a slower rate when exercising at a RPE of 5 on Borg's 10-point scale but not on PPO during a 5 s maximal effort sprint. This decrement in performance was accompanied by a great reduction in muscle activation, suggesting that central processes were involved. Whether the rate sensitive component within the proposed framework is a function of central information processing leading to changes in CMD or the modulation of CMD at the spinal level, does not conceptually change the feed-forward model as each possibility allow the organism to differentiate between the instantaneous state and the instantaneous rate of change [13]. Further research should be undertaken to clarify if peripheral factors contributed to these observations, if the regulation of exercise intensity is similarly sensitive to the rate of change of other physiological stimuli, and to elucidate the mechanisms responsible for mediating this effect.

## Author Contributions

**Conceptualization:** SF.

**Data curation:** SF IJ.

**Formal analysis:** SF IJ.

**Funding acquisition:** SF IJ.

**Investigation:** SF.

**Methodology:** SF SC ST IJ.

**Project administration:** SF IJ.

**Resources:** SC SC IJ.

**Supervision:** SF SC ST IJ.

**Validation:** SF IJ.

**Visualization:** SF IJ.

**Writing – original draft:** SF.

**Writing – review & editing:** SF SC ST IJ.

## References

1. González-Alonso J, Teller C, Andersen SL, Jensen FB, Hyldig T, Nielsen B. Influence of body temperature on the development of fatigue during prolonged exercise in the heat. *J Appl Physiol.* 1999; 86(3):1032–9. PMID: [10066720](#)
2. Nybo L, Nielsen B. Hyperthermia and central fatigue during prolonged exercise in humans. *J Appl Physiol.* 2001; 91(3):1055–60. PMID: [11509498](#)
3. Amann M, Blain GM, Proctor LT, Sebranek JJ, Pegelow DF, Dempsey JA. Implications of group III and IV muscle afferents for high-intensity endurance exercise performance in humans. *J Physiol.* 2011; 589(Pt 21):5299–309. doi: [10.1113/jphysiol.2011.213769](#) PMID: [21878520](#)
4. Gandevia SC, Allen GM, Butler JE, Taylor JL. Supraspinal factors in human muscle fatigue: Evidence for suboptimal output from the motor cortex. *J Physiol.* 1996; 490(Pt 2):529–36.
5. Bergström J, Hermansen L, Hultman E, Saltin B. Diet, muscle glycogen and physical performance. *Acta Physiol Scand.* 1967; 71(2):140–50. doi: [10.1111/j.1748-1716.1967.tb03720.x](#) PMID: [5584523](#)
6. Hermansen L, Hultman E, Saltin B. Muscle glycogen during prolonged severe exercise. *Acta Physiol Scand.* 1967; 71(2):129–39. doi: [10.1111/j.1748-1716.1967.tb03719.x](#) PMID: [5584522](#)
7. Holloszy JO, Kohrt WM. Regulation of carbohydrate and fat metabolism during and after exercise. *Annu Rev Nutr.* 1996; 16:121–38. doi: [10.1146/annurev.nu.16.070196.001005](#) PMID: [8839922](#)
8. Garner SH, Sutton JR, Burse RL, McComas AJ, Cymerman A, Houston CS. Operation everest II: Neuromuscular performance under conditions of extreme simulated altitude. *J Appl Physiol.* 1990; 68(3):1167–72. PMID: [2341341](#)
9. Pugh LG, Gill MB, Lahiri S, Milledge JS, Ward MP, West JB. Muscular exercise at great altitudes. *J Appl Physiol.* 1964; 194:31–40.
10. Noakes TD, Marino FE. Arterial oxygenation, central motor output and exercise performance in humans. *J Physiol.* 2007; 585(Pt 3):919–21. doi: [10.1113/jphysiol.2007.145110](#) PMID: [17962324](#)
11. Tucker R. The anticipatory regulation of performance: The physiological basis for pacing strategies and the development of a perception-based model for exercise performance. *Br J Sports Med.* 2009; 43(6):392–400. doi: [10.1136/bjism.2008.050799](#) PMID: [19224911](#)
12. Allgood GO. Mapping function and structure for an anticipatory system: What impact will it have and is it computationally feasible, today? *Systems, Man, and Cybernetics, 2000 IEEE International Conference on.* 2000; 3:2198–2203.
13. Rosen R. *Anticipatory systems. Philosophical, mathematical and methodological foundations.* New York: Pergamon; 1985.
14. Tucker R, Noakes TD. The physiological regulation of pacing strategy during exercise: A critical review. *Br J Sports Med.* 2009; 43(6):e1. doi: [10.1136/bjism.2009.057562](#) PMID: [19224909](#)
15. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc.* 1982; 14(5):377–81.
16. Abbiss CR, Laursen PB. Models to explain fatigue during prolonged endurance cycling. *Sports Med.* 2005; 35(10):865–98. PMID: [16180946](#)
17. de Morree HM, Klein C, Marcora SM. Perception of effort reflects central motor command during movement execution. *Psychophysiology.* 2012; 49(9):1242–53. doi: [10.1111/j.1469-8986.2012.01399.x](#) PMID: [22725828](#)

18. Gallagher KM, Fadel PJ, Strømstad M, Ide K, Smith SA, Query RG, et al. Effects of partial neuromuscular blockade on carotid baroreflex function during exercise in humans. *J Physiol.* 2001; 533(Pt 3):861–70. doi: [10.1111/j.1469-7793.2001.t01-1-00861.x](https://doi.org/10.1111/j.1469-7793.2001.t01-1-00861.x) PMID: [11410641](https://pubmed.ncbi.nlm.nih.gov/11410641/)
19. Marcora SM. Perception of effort during exercise is independent of afferent feedback from skeletal muscles, heart, and lungs. *J Appl Physiol.* 2009; 106(6):2060–2. doi: [10.1152/jappphysiol.90378.2008](https://doi.org/10.1152/jappphysiol.90378.2008) PMID: [18483166](https://pubmed.ncbi.nlm.nih.gov/18483166/)
20. Smith SA, Query RG, Fadel PJ, Gallagher KM, Strømstad M, Ide K, et al. Partial blockade of skeletal muscle somatosensory afferents attenuates baroreflex resetting during exercise in humans. *J Physiol.* 2003; 551(Pt 3):1013–21. doi: [10.1113/jphysiol.2003.044925](https://doi.org/10.1113/jphysiol.2003.044925) PMID: [12819303](https://pubmed.ncbi.nlm.nih.gov/12819303/)
21. Borg GA. Perceived exertion: A note on "history" and methods. *Med Sci Sports.* 1973; 5(2):90–3. PMID: [4721012](https://pubmed.ncbi.nlm.nih.gov/4721012/)
22. Morgan WP, Pollock ML. Psychological characterization of the elite distance runner. *Ann N Y Acad Sci.* 1977; 301:382–403. PMID: [270929](https://pubmed.ncbi.nlm.nih.gov/270929/)
23. Noble BJ, Metz KF, Pandolf KB, Bell CW, Cafarelli E, Sime WE. Perceived exertion during walking and running. II. *Med Sci Sports.* 1973; 5(2):116–20. PMID: [4721005](https://pubmed.ncbi.nlm.nih.gov/4721005/)
24. Noble BJ, Metz KF, Pandolf KB, Cafarelli E. Perceptual responses to exercise: A multiple regression study. *Med Sci Sports.* 1973; 5(2):104–9. PMID: [4721003](https://pubmed.ncbi.nlm.nih.gov/4721003/)
25. Noble BJ, Borg GA, Jacobs I, Ceci R, Kaiser P. A category-ratio perceived exertion scale: Relationship to blood and muscle lactates and heart rate. *Med Sci Sports Exerc.* 1983; 15(6):523–8. PMID: [6656563](https://pubmed.ncbi.nlm.nih.gov/6656563/)
26. Skinner JS, Hutsler R, Bergsteinová V, Buskirk ER. The validity and reliability of a rating scale of perceived exertion. *Med Sci Sports.* 1973; 5(2):94–6. PMID: [4721013](https://pubmed.ncbi.nlm.nih.gov/4721013/)
27. Faulkner J, Eston RG. Perceived exertion research in the 21st century: Developments, reflections, and questions for the future. *J Exerc Sci Fit.* 2008; 6(1):1–14.
28. Renfree A, West J, Corbett M, Rhoden C, St Clair Gibson A. Complex interplay between determinants of pacing and performance during 20-km cycle time trials. *Int J Sports Physiol Perform.* 2012; 7(2):121–9. PMID: [22173069](https://pubmed.ncbi.nlm.nih.gov/22173069/)
29. St Clair Gibson A, Lambert EV, Rauch LHG, Tucker R, Baden DA, Foster C, Noakes TD. The role of information processing between the brain and peripheral physiological systems in pacing and perception of effort. *Sports Med.* 2006; 36(8):705–22. PMID: [16869711](https://pubmed.ncbi.nlm.nih.gov/16869711/)
30. Eston RG, Davies BL, Williams JG. Use of perceived effort ratings to control exercise intensity in young healthy adults. *Eur J Appl Physiol Occup Physiol.* 1987; 56(2):222–224. PMID: [3569229](https://pubmed.ncbi.nlm.nih.gov/3569229/)
31. Eston RG, Williams JG. Reliability of ratings of perceived effort regulation of exercise intensity. *Br J Sports Med.* 1988; 22(4):153–155. PMID: [3228684](https://pubmed.ncbi.nlm.nih.gov/3228684/)
32. Amann M, Romer LM, Subudhi AW, Pegelow DF, Dempsey JA. Severity of arterial hypoxaemia affects the relative contributions of peripheral muscle fatigue to exercise performance in healthy humans. *J Physiol.* 2007; 581(Pt 1):389–403. doi: [10.1113/jphysiol.2007.129700](https://doi.org/10.1113/jphysiol.2007.129700) PMID: [17317739](https://pubmed.ncbi.nlm.nih.gov/17317739/)
33. Kayser B. Exercise starts and ends in the brain. *Eur J Appl Physiol.* 2003; 90(3–4):411–9. doi: [10.1007/s00421-003-0902-7](https://doi.org/10.1007/s00421-003-0902-7) PMID: [12883902](https://pubmed.ncbi.nlm.nih.gov/12883902/)
34. Goodall S, Ross EZ, Romer LM. Effect of graded hypoxia on supraspinal contributions to fatigue with unilateral knee-extensor contractions. *J Appl Physiol.* 2010; 109(6):1842–51. doi: [10.1152/jappphysiol.00458.2010](https://doi.org/10.1152/jappphysiol.00458.2010) PMID: [20813979](https://pubmed.ncbi.nlm.nih.gov/20813979/)
35. Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev.* 2001; 81(4):1725–89. PMID: [11581501](https://pubmed.ncbi.nlm.nih.gov/11581501/)
36. Amann M, Dempsey JA. Locomotor muscle fatigue modifies central motor drive in healthy humans and imposes a limitation to exercise performance. *J Physiol.* 2008; 586(1):161–73. doi: [10.1113/jphysiol.2007.141838](https://doi.org/10.1113/jphysiol.2007.141838) PMID: [17962334](https://pubmed.ncbi.nlm.nih.gov/17962334/)
37. Kaufman MP, Rybicki KJ, Waldrop TG, Ordway GA. Effect of ischemia on responses of group III and IV afferents to contraction. *J Appl Physiol Respir Environ Exerc Physiol.* 1984; 57(3):644–50. PMID: [6092310](https://pubmed.ncbi.nlm.nih.gov/6092310/)
38. Squires RW, Buskirk ER. Aerobic capacity during acute exposure to simulated altitude, 914 to 2286 meters. *Med Sci Sports Exerc.* 1982; 14(1):36–40. PMID: [7070255](https://pubmed.ncbi.nlm.nih.gov/7070255/)
39. Amann M, Kayser B. Nervous system function during exercise in hypoxia. *High Alt Med Biol.* 2009; 10(2):149–64. doi: [10.1089/ham.2008.1105](https://doi.org/10.1089/ham.2008.1105) PMID: [19555297](https://pubmed.ncbi.nlm.nih.gov/19555297/)
40. Millet GY, Muthalib M, Jubeau M, Laursen PB, Nosaka K. Severe hypoxia affects exercise performance independently of afferent feedback and peripheral fatigue. *J Appl Physiol.* 2012; 112(8):1335–44. doi: [10.1152/jappphysiol.00804.2011](https://doi.org/10.1152/jappphysiol.00804.2011) PMID: [22323647](https://pubmed.ncbi.nlm.nih.gov/22323647/)

41. Rasmussen P, Dawson EA, Nybo L, van Lieshout JJ, Secher NH, Gjedde A. Capillary-oxygenation-level-dependent near-infrared spectrometry in frontal lobe of humans. *J Cereb Blood Flow Metab.* 2007; 27(5):1082–93. doi: [10.1038/sj.jcbfm.9600416](https://doi.org/10.1038/sj.jcbfm.9600416) PMID: [17077816](https://pubmed.ncbi.nlm.nih.gov/17077816/)
42. Rasmussen P, Nielsen J, Overgaard M, Krogh-Madsen R, Gjedde A, Secher NH, Petersen NC. Reduced muscle activation during exercise related to brain oxygenation and metabolism in humans. *J Physiol.* 2010; 588(Pt 11):1985–95. doi: [10.1113/jphysiol.2009.186767](https://doi.org/10.1113/jphysiol.2009.186767) PMID: [20403976](https://pubmed.ncbi.nlm.nih.gov/20403976/)
43. Marino FE, Lambert MI, Noakes TD. Superior performance of african runners in warm humid but not in cool environmental conditions. *J Appl Physiol.* 2004; 96(1):124–30. doi: [10.1152/jappphysiol.00582.2003](https://doi.org/10.1152/jappphysiol.00582.2003) PMID: [12949014](https://pubmed.ncbi.nlm.nih.gov/12949014/)
44. Tucker R, Rauch L, Harley YXR, Noakes TD. Impaired exercise performance in the heat is associated with an anticipatory reduction in skeletal muscle recruitment. *Pflugers Arch.* 2004; 448(4):422–30. doi: [10.1007/s00424-004-1267-4](https://doi.org/10.1007/s00424-004-1267-4) PMID: [15138825](https://pubmed.ncbi.nlm.nih.gov/15138825/)
45. Tucker R, Marle T, Lambert EV, Noakes TD. The rate of heat storage mediates an anticipatory reduction in exercise intensity during cycling at a fixed rating of perceived exertion. *J Physiol.* 2006; 574(Pt 3):905–15. doi: [10.1113/jphysiol.2005.101733](https://doi.org/10.1113/jphysiol.2005.101733) PMID: [16497719](https://pubmed.ncbi.nlm.nih.gov/16497719/)
46. Balsom PD, Gaitanos GC, Ekblom B, Sjödín B. Reduced oxygen availability during high intensity intermittent exercise impairs performance. *Acta Physiol Scand.* 1994; 152(3):279–85. doi: [10.1111/j.1748-1716.1994.tb09807.x](https://doi.org/10.1111/j.1748-1716.1994.tb09807.x) PMID: [7872005](https://pubmed.ncbi.nlm.nih.gov/7872005/)
47. Ekblom B, Huot R, Stein EM, Thorstensson AT. Effect of changes in arterial oxygen content on circulation and physical performance. *J Appl Physiol.* 1975; 39(1):71–5. PMID: [1150596](https://pubmed.ncbi.nlm.nih.gov/1150596/)
48. Romer LM, Haverkamp HC, Lovering AT, Pegelow DF, Dempsey JA. Effect of exercise-induced arterial hypoxemia on quadriceps muscle fatigue in healthy humans. *Am J Physiol Regul Integr Comp Physiol.* 2006; 290(2):R365–75. doi: [10.1152/ajpregu.00332.2005](https://doi.org/10.1152/ajpregu.00332.2005) PMID: [16166208](https://pubmed.ncbi.nlm.nih.gov/16166208/)
49. Farra SD, Kessler C, Duffin J, Wells GD, Jacobs I. Clamping end-tidal carbon dioxide during graded exercise with control of inspired oxygen. *Respir Physiol Neurobiol.* 2016; 231:28–36. doi: [10.1016/j.resp.2016.05.013](https://doi.org/10.1016/j.resp.2016.05.013) PMID: [27236039](https://pubmed.ncbi.nlm.nih.gov/27236039/)
50. Slessarev M, Han J, Mardimae A, Prisman E, Preiss D, Volgyesi G, et al. Prospective targeting and control of end-tidal CO<sub>2</sub> and O<sub>2</sub> concentrations. *J Physiol.* 2007; 581(3):1207–1219.
51. Sommer LZ, Iscoe S, Robicsek A, Kruger J, Silverman J, Rucker J, et al. A simple breathing circuit minimizing changes in alveolar ventilation during hyperpnoea. *Eur Respir J.* 1998; 12(3):698–701. PMID: [9762802](https://pubmed.ncbi.nlm.nih.gov/9762802/)
52. Somogyi RB, Vesely AE, Preiss D, Prisman E, Volgyesi G, Azami T, et al. Precise control of end-tidal carbon dioxide levels using sequential rebreathing circuits. *Anaesth Intensive Care.* 2005; 33(6):726–32. PMID: [16398376](https://pubmed.ncbi.nlm.nih.gov/16398376/)
53. McLellan TM, Skinner JS. Blood lactate removal during active recovery related to the aerobic threshold. *Int J Sports Med.* 1982; 3(4):224–229.
54. McLellan TM. Ventilatory and plasma lactate response with different exercise protocols: A comparison of methods. *Int J Sports Med.* 1985; 6(1):30–5. doi: [10.1055/s-2008-1025809](https://doi.org/10.1055/s-2008-1025809) PMID: [3988412](https://pubmed.ncbi.nlm.nih.gov/3988412/)
55. Patrick Neary J, Roberts ADW, Leavins N, Harrison MF, Croll JC, Sexsmith JR. Prefrontal cortex oxygenation during incremental exercise in chronic fatigue syndrome. *Clin Physiol Funct Imaging.* 2008; 28(6):364–72. doi: [10.1111/j.1475-097X.2008.00822.x](https://doi.org/10.1111/j.1475-097X.2008.00822.x) PMID: [18671793](https://pubmed.ncbi.nlm.nih.gov/18671793/)
56. Bhambhani Y, Maikala R, Esmail S. Oxygenation trends in vastus lateralis muscle during incremental and intense anaerobic cycle exercise in young men and women. *Eur J Appl Physiol.* 2001; 84(6):547–558. doi: [10.1007/s004210000372](https://doi.org/10.1007/s004210000372) PMID: [11482550](https://pubmed.ncbi.nlm.nih.gov/11482550/)
57. Johnson BD, Joseph T, Wright G, Battista RA, Dodge C, Balweg A, et al. Rapidity of responding to a hypoxic challenge during exercise. *Eur J Appl Physiol.* 2009; 106(4):493–9. doi: [10.1007/s00421-009-1036-3](https://doi.org/10.1007/s00421-009-1036-3) PMID: [19326141](https://pubmed.ncbi.nlm.nih.gov/19326141/)
58. Amann M, Eldridge MW, Lovering AT, Stickland MK, Pegelow DF, Dempsey JA. Arterial oxygenation influences central motor output and exercise performance via effects on peripheral locomotor muscle fatigue in humans. *J Physiol.* 2006; 575(Pt 3):937–52. doi: [10.1113/jphysiol.2006.113936](https://doi.org/10.1113/jphysiol.2006.113936) PMID: [16793898](https://pubmed.ncbi.nlm.nih.gov/16793898/)
59. Clark SA, Bourdon PC, Schmidt W, Singh B, Cable G, Onus KJ, et al. The effect of acute simulated moderate altitude on power, performance and pacing strategies in well-trained cyclists. *Eur J Appl Physiol.* 2007; 102(1):45–55. doi: [10.1007/s00421-007-0554-0](https://doi.org/10.1007/s00421-007-0554-0) PMID: [17882451](https://pubmed.ncbi.nlm.nih.gov/17882451/)
60. Amann M, Romer LM, Subudhi AW, Pegelow DF, Dempsey JA. Severity of arterial hypoxaemia affects the relative contributions of peripheral muscle fatigue to exercise performance in healthy humans. *J Physiol.* 2007; 581(Pt 1):389–403. doi: [10.1113/jphysiol.2007.129700](https://doi.org/10.1113/jphysiol.2007.129700) PMID: [17317739](https://pubmed.ncbi.nlm.nih.gov/17317739/)
61. Goodall S, González-Alonso J, Ali L, Ross EZ, Romer LM. Supraspinal fatigue after normoxic and hypoxic exercise in humans. *J Physiol.* 2012; 590(Pt 11):2767–82.

62. Verges S, Rupp T, Jubeau M, Wuyam B, Esteve F, Levy P, et al. Cerebral perturbations during exercise in hypoxia. *Am J Physiol Regul Integr Comp Physiol*. 2012; 302(8):R903–16. doi: [10.1152/ajpregu.00555.2011](https://doi.org/10.1152/ajpregu.00555.2011) PMID: [22319046](https://pubmed.ncbi.nlm.nih.gov/22319046/)
63. Wetter TJ, Harms CA, Nelson WB, Pegelow DF, Dempsey JA. Influence of respiratory muscle work on VO<sub>2</sub> and leg blood flow during submaximal exercise. *J Appl Physiol*. 1999; 87(2):643–51. PMID: [10444624](https://pubmed.ncbi.nlm.nih.gov/10444624/)
64. Hill JM, Pickar JG, Parrish MD, Kaufman MP. Effects of hypoxia on the discharge of group III and IV muscle afferents in cats. *J Appl Physiol*. 1992; 73(6):2524–9. PMID: [1490966](https://pubmed.ncbi.nlm.nih.gov/1490966/)
65. Sahlin K, Katz A. Hypoxaemia increases the accumulation of inosine monophosphate (IMP) in human skeletal muscle during submaximal exercise. *Acta Physiol Scand*. 1989; 136(2):199–203. doi: [10.1111/j.1748-1716.1989.tb08653.x](https://doi.org/10.1111/j.1748-1716.1989.tb08653.x) PMID: [2782093](https://pubmed.ncbi.nlm.nih.gov/2782093/)
66. Gandevia SC. Neural control in human muscle fatigue: Changes in muscle afferents, motoneurons and motor cortical drive [corrected]. *Acta Physiol Scand*. 1998; 162(3):275–83. doi: [10.1046/j.1365-201X.1998.0299f.x](https://doi.org/10.1046/j.1365-201X.1998.0299f.x) PMID: [9578373](https://pubmed.ncbi.nlm.nih.gov/9578373/)
67. Wehrlin JP, Hallén J. Linear decrease in VO<sub>2</sub>max and performance with increasing altitude in endurance athletes. *Eur J Appl Physiol*. 2006; 96(4):404–12. doi: [10.1007/s00421-005-0081-9](https://doi.org/10.1007/s00421-005-0081-9) PMID: [16311764](https://pubmed.ncbi.nlm.nih.gov/16311764/)
68. Mateika JH, Duffin J. A review of the control of breathing during exercise. *Eur J Appl Physiol Occup Physiol*. 1995; 71(1):1–27. PMID: [7556128](https://pubmed.ncbi.nlm.nih.gov/7556128/)
69. West JA. *Respiratory physiology: The essentials*. 6 ed. Baltimore, Maryland: Lippincott Williams & Wilkins; 2000.
70. Craig AD. Interoception: The sense of the physiological condition of the body. *Curr Opin Neurobiol*. 2003; 13(4):500–5. PMID: [12965300](https://pubmed.ncbi.nlm.nih.gov/12965300/)
71. Williamson JW, Fadel PJ, Mitchell JH. New insights into central cardiovascular control during exercise in humans: A central command update. *Exp Physiol*. 2006; 91(1):51–8. doi: [10.1113/expphysiol.2005.032037](https://doi.org/10.1113/expphysiol.2005.032037) PMID: [16239250](https://pubmed.ncbi.nlm.nih.gov/16239250/)
72. Tanaka M, Watanabe Y. Supraspinal regulation of physical fatigue. *Neurosci Biobehav Rev*. 2012; 36(1):727–34. doi: [10.1016/j.neubiorev.2011.10.004](https://doi.org/10.1016/j.neubiorev.2011.10.004) PMID: [22040772](https://pubmed.ncbi.nlm.nih.gov/22040772/)
73. Eston R, Faulkner J, St Clair Gibson A, Noakes T, Parfitt G. The effect of antecedent fatiguing activity on the relationship between perceived exertion and physiological activity during a constant load exercise task. *Psychophysiology*. 2007; 44(5):779–86. doi: [10.1111/j.1469-8986.2007.00558.x](https://doi.org/10.1111/j.1469-8986.2007.00558.x) PMID: [17617170](https://pubmed.ncbi.nlm.nih.gov/17617170/)
74. Puccini GD, Sanchez-Vives MV, Compte A. Integrated mechanisms of anticipation and rate-of-change computations in cortical circuits. *PLoS Comput Biol*. 2007; 3(5):e82. doi: [10.1371/journal.pcbi.0030082](https://doi.org/10.1371/journal.pcbi.0030082) PMID: [17500584](https://pubmed.ncbi.nlm.nih.gov/17500584/)
75. Webster MA. Evolving concepts of sensory adaptation. *F1000 Biol Rep*. 2012; 421.
76. Li K, Ponte J, Sadler CL. Carotid body chemoreceptor response to prolonged hypoxia in the rabbit: Effects of domperidone and propranolol. *J Physiol*. 1990; 430:1–11. PMID: [2128334](https://pubmed.ncbi.nlm.nih.gov/2128334/)
77. Barnard P, Andronikou S, Pokorski M, Smatresk N, Mokashi A, Lahiri S. Time-dependent effect of hypoxia on carotid body chemosensory function. *J Appl Physiol*. 1987; 63(2):685–91. PMID: [3654428](https://pubmed.ncbi.nlm.nih.gov/3654428/)
78. Kaufman MP, Forster HV. Reflexes controlling circulatory, ventilatory and airway responses to exercise. *Comprehensive Physiology*. 1996
79. Davis JN, Carlsson A. The effect of hypoxia on monoamine synthesis, levels and metabolism in rat brain. *J Neurochem*. 1973; 21(4):783–90. PMID: [4148239](https://pubmed.ncbi.nlm.nih.gov/4148239/)
80. Gibson GE, Shimada M, Blass JP. Alterations in acetylcholine synthesis and cyclic nucleotides in mild cerebral hypoxia. *J Neurochem*. 1978; 31(4):757–60. PMID: [212531](https://pubmed.ncbi.nlm.nih.gov/212531/)
81. Brown RM, Snider SR, Carlsson A. Changes in biogenic amine synthesis and turnover induced by hypoxia and/or foot shock stress. II. The central nervous system. *J Neural Transm*. 1974; 35(4):293–305. PMID: [4548084](https://pubmed.ncbi.nlm.nih.gov/4548084/)
82. Duffy TE, Nelson SR, Lowry OH. Cerebral carbohydrate metabolism during acute hypoxia and recovery. *J Neurochem*. 1972; 19(4):959–77. PMID: [5019592](https://pubmed.ncbi.nlm.nih.gov/5019592/)
83. Erecińska M, Nelson D, Wilson DF, Silver IA. Neurotransmitter amino acids in the CNS. I. Regional changes in amino acid levels in rat brain during ischemia and reperfusion. *Brain Res*. 1984; 304(1):9–22. PMID: [6146383](https://pubmed.ncbi.nlm.nih.gov/6146383/)
84. Iversen K, Hedner T, Lundborg P. GABA concentrations and turnover in neonatal rat brain during asphyxia and recovery. *Acta Physiol Scand*. 1983; 118(1):91–4. doi: [10.1111/j.1748-1716.1983.tb07247.x](https://doi.org/10.1111/j.1748-1716.1983.tb07247.x) PMID: [6688698](https://pubmed.ncbi.nlm.nih.gov/6688698/)

85. Wood JD, Watson WJ, Ducker AJ. The effect of hypoxia on brain gamma-aminobutyric acid levels. *J Neurochem.* 1968; 15(7):603–8. PMID: [5692041](#)
86. Winn HR, Rubio R, Berne RM. Brain adenosine concentration during hypoxia in rats. *Am J Physiol.* 1981; 241(2):H235–42. PMID: [6791511](#)
87. Hagberg H, Lehmann A, Sandberg M, Nyström B, Jacobson I, Hamberger A. Ischemia-induced shift of inhibitory and excitatory amino acids from intra- to extracellular compartments. *J Cereb Blood Flow Metab.* 1985; 5(3):413–9. doi: [10.1038/jcbfm.1985.56](#) PMID: [4030918](#)
88. Chernick V, Madansky DL, Lawson EE. Naloxone decreases the duration of primary apnea with neonatal asphyxia. *Pediatr Res.* 1980; 14(4 Pt 1):357–9.
89. Chernick V, Craig RJ. Naloxone reverses neonatal depression caused by fetal asphyxia. *Science.* 1982; 216(4551):1252–3. PMID: [7200636](#)
90. Neubauer JA, Simone A, Edelman NH. Role of brain lactic acidosis in hypoxic depression of respiration. *J Appl Physiol.* 1988; 65(3):1324–31. PMID: [3182502](#)