

Aberrant expression of long non-coding RNAs (lncRNAs) is involved in brain glioma development

Yi Ding¹, Xinfu Wang¹, Junchen Pan², Minjun Ji³, Zhengxiang Luo⁴, Penglai Zhao⁴, Yansong Zhang⁴, Gang Wang¹

¹Department of Neurosurgery, Children's Hospital of Nanjing Medical University, Nanjing, China

²Department of Neurosurgery, Nanjing BenQ hospital, Nanjing, China

³Department of Neurosurgery, Nanjing Medical University, Nanjing, China

⁴Department of Neurosurgery, Nanjing Brian Hospital Affiliated to Nanjing Medical University, Nanjing, China

Submitted: 27 March 2017

Accepted: 1 August 2017

Arch Med Sci 2020; 16 (1): 177–188

DOI: <https://doi.org/10.5114/aoms.2020.91290>

Copyright © 2019 Termedia & Banach

Corresponding authors:

Yansong Zhang
Department
of Neurosurgery
Nanjing Brian Hospital
Affiliated to Nanjing
Medical University
025 Nanjing, China
Phone: +86 25 86862793
E-mail: yan_song_
zhang@126.com

Gang Wang
Department
of Neurosurgery
Nanjing Children's Hospital
Nanjing, China
E-mail: wang509111@163.com

Abstract

Introduction: Aberrant expression of long non-coding RNAs (lncRNAs) has been implicated in various diseases, including cancer. However, little is known about lncRNAs in human brain gliomas.

Material and methods: We examined lncRNA profiles from three glioma specimens using lncRNA expression profiling microarrays. Quantitative real-time RT-PCR was used to analyze the differential expression of raw intensities of lncRNA expression in glioma and peritumoral tissues.

Results: We found 4858 lncRNAs to be differentially expressed between tumor tissue and peritumoral tissue. Of these, 2845 lncRNAs were up-regulated (fold change > 3.0) and 2013 were down-regulated (fold change < 1/3). A total of 4084 messenger RNAs were also differentially expressed, including 2280 up-regulated transcripts (fold change > 3.0) and 1804 that were down-regulated (fold change < 1/3). Consistent with the microarray data, qPCR confirmed differential expression of these 6 lncRNAs (ak125809, ak098473, uc002ehu.1, bc043564, NR_027322, and uc003qmb.2) between tumor and peritumoral tissue. We next established co-expression networks of differentially expressed lncRNAs and mRNAs. Many mRNAs, such as LOC729991, NUDCD1, SHC3, PDGFA, and MDM2, and lncRNAs, such as ENST00000425922, ENST00000455568, uc002ukz.1, ENST00000502715, and NR_027873, have been shown to play important roles in glioma development. Consistent with this, pathway analysis revealed that "GLIOMA" (KEGG Pathway ID: hsa05214) was significantly enriched in tumor tissue.

Conclusions: Our data suggest that altered expression of lncRNAs may be a critical determinant of tumorigenesis in glioma patients.

Key words: lncRNAs, glioma, microarrays.

Introduction

Gliomas are the most common and most severe form of primary brain cancer, accounting for approximately 52% of all primary brain tumors and 20% of intracranial tumors [1]. Gliomas arise from malignant transformation of glial cells and can be subdivided into astrocytomas, oligodendrogliomas, ependymomas, and mixed gliomas according to their clinical and pathological features. Malignant glioma ranks second (behind only stroke) among all nerve-associated diseases in mortality rate [1].

Even with current surgical and chemotherapy options, the median survival time of patients with malignant glioma or glioblastoma is only 14 months [2]. While great strides in glioma research have been made over the last two decades, the exact pathological mechanism underlying the disease has not been defined [3]. Thus, additional research is required to not only better understand the disease itself but also to improve prognosis and therapeutic options for patients.

Long non-coding RNAs (lncRNAs) are regulatory RNA transcripts typically over 200 nucleotides in length [4]. In mammals, lncRNAs are 6–8 times more abundant than mRNAs and are encoded by 4–9% of the entire genome [5]. While originally considered the “noise” of transcription, lncRNAs are now known to be vital in multiple molecular processes, including chromatin modification [5, 6], transcriptional activation [7], and post-transcriptional regulation [8]. lncRNAs have also been shown to be critical in a number of physiological and pathological processes [9–13].

Aberrant expression of lncRNAs has been linked to tumorigenesis [14, 15]. For example, one study used a hepatocellular carcinoma-specific cDNA microarray to identify increased expression of the lncRNA HULC in tumor tissue [15]. Additionally, lncRNA-HULC was also detectable in the plasma of hepatocellular carcinoma patients, thus making it a potential diagnostic biomarker [15]. lncRNA-HULC (hepatocellular carcinoma up-regulated long non-coding RNA) was also shown to regulate the expression of multiple genes in hepatocellular carcinoma cells, strengthening the link between this lncRNA and tumorigenesis [12]. Overexpression of the lncRNA maternally expressed gene 3 (MEG3) inhibited proliferation and induced apoptosis of both U87 MG and U251 human glioma cell lines [16]. The authors showed that this effect was due to the interaction among MEG3, p53, MDM2, and GDF15 [17]. Another study examined the expression profiles of both lncRNAs and mRNAs in glioblastoma; the authors found two lncRNAs, ASLNC22381 and ASLNC2081, that were related to the disease via their regulation of insulin-like growth factor 1 (IGF-1) [18]. Zhang *et al.* used a combination of publicly available data and their own microarray analysis to assess lncRNA expression in astrocytic and oligodendroglial glioma [19]. Interestingly, they found a significant correlation between the expression of specific lncRNAs and both malignancy grade and histological differentiation [19]. The follow-up to these findings has been limited due to insufficient quantity of samples and to limited lncRNAs included on microarray platforms. Despite this, much of this earlier work laid the foundation for a role for lncRNAs in glioma development – a topic that deserves more extensive study.

Here, we used lncRNA expression profiling microarray to identify differentially expressed lncRNAs in clinical glioma samples compared to normal tissue. Six lncRNAs (ak125809, ak098473, uc002ehu.1, bc043564, NR_027322, and uc003qmb.2) were selected and validated by quantitative real-time RT-PCR (qPCR). We established an mRNA/lncRNA co-expression network to identify key regulators of glioma development. The data we present here may provide a better understanding of the disease and new therapeutic strategies to combat it in patients.

Material and methods

Sample collection

Forty cases of glioma and peritumoral tissue were obtained from the Neurosurgery Department of Nanjing Brain Hospital Affiliated with Nanjing Medical University. Peritumoral tissue was obtained from both a surgical approach and peripheral edema tissue. Pathological categories were evaluated by the hospital’s pathologist (ID: NJNKYY2012LS027). Written informed consent was obtained from all patients. This study was approved by the Ethics Committee of the University. Following collection, specimens were shipped and preserved in liquid nitrogen. Details are shown in supplement material 1. The protocol for the tissue collection was as follows: (1) Record the information of tumor tissue on the tissue cryopreserved tube; (2) wash the tumor tissue with saline, use ophthalmic scissors, ophthalmic tweezers to remove blood vessels, envelopes and necrotic tissue; (3) cut the tumor tissue into 2 mm thick slices; (4) flatten tissue on sterile gauze, and remove the surface liquid of tissue sections; (5) quickly immerse the tissue in liquid nitrogen, not less than 5 minutes; (6) quickly move the tissue into the tissue cryopreservation tube, tighten the tube and move into the liquid nitrogen tank for preservation.

RNA extraction

Total RNA was isolated with TRIzol reagent (Invitrogen, CA, USA) following the single-step method [20]: 1) Glioma and adjacent tissues were pulverized with a mortar and pestle in liquid nitrogen. Tissue samples were homogenized in 1 ml of Trizol per 50 mg of tissue until liquid nitrogen evaporated. 2) The homogenate was pipetted up and down several times and transferred to an Eppendorf tube. 3) Homogenized samples were incubated for 5 min at room temperature and then centrifuged at 12,000 rpm for 5 min according to the protocol. Next, 200 μ l of chloroform was added per 1 ml of TRIzol. Samples were incubated for 15 min at room temperature and centrifuged at 12,000 rpm for

15 min at 4°C. The upper soluble layer was isolated and mixed with 0.5 ml of isopropanol per 1 ml of TRIzol. Samples were incubated for 10 min at room temperature and centrifuged at 12,000 rpm for 10 min at 4°C. The supernatant was removed and washed with 1 ml of 75% ethanol. Samples were then centrifuged at 8,000 rpm for 5 min at 4°C. This was followed by a brief incubation at room temperature to allow the samples to dry. Finally, RNA samples were dissolved in 50 µl of DEPC water and stored at -80°C until use. The concentration of each sample was determined using the NanoDrop ND-1000 Spectrophotometer (NanoDrop Technologies, Inc., USA). RNA integrity was evaluated by electrophoresis on a denaturing agarose gel.

lncRNA and mRNA microarrays

Total RNA from both glioma (0301, 0605, and 114982) and peritumoral tissue (0601) was isolated and converted into double-stranded complementary DNA (cDNA). The synthesized cDNAs were labeled and hybridized to Arraystar Human lncRNA Microarray V2.0 (Arraystar, Rockville, MD) containing probes for 33,045 lncRNAs and 30,215 mRNAs identified from both publications and authoritative databases, such as RefSeq, UCSC Knowngenes, and Ensembl. Each transcript was represented by a specific exon or splice junction probe, which could accurately identify individual transcripts. Positive control probes and negative control probes for housekeeping genes were also printed onto the array [21].

Microarray hybridization and data analysis

Arraystar lncRNA Array Protocol: step 1 – Prepare the RNA sample using the following reagents and kits: TRIzol Reagent (Invitrogen Life Technologies), Biopulverizer (Biospec), and Mini-Bead-Beater-16 (Biospec); step 2 – Total RNA clean-up and quality control using the following reagents and instruments: RNeasy Mini Kit (Qiagen, p/n 74104), NanoDrop ND-1000, and Baseline-ZERO DNase (EPICENTRE, Cat. Nos. DB0711K); step 3 – Prepare

labeling reaction with the following: Quick Amp Labeling Kit and One-Color (Agilent p/n 5190-0442); step 4 – Purify the labeled RNA using the RNeasy Mini Kit (Qiagen, p/n 74104) and NanoDrop ND-1000; step 5 – Hybridization using the Agilent Gene Expression Hybridization Kit (Agilent, p/n 5188-5242), stainless hybridization chamber (Agilent, p/n G2534A), Hybridization Chamber gasket slides (Agilent, p/n G2534-60003), Hybridization oven (Agilent, p/n G2545A), and Hybridization oven rotator for Agilent Microarray Hybridization Chambers (Agilent, p/n G2530-60029); step 6 – Microarray Wash using Gene Expression Wash Buffer 1 (Agilent, p/n 5188-5325), Gene Expression Wash Buffer 2 (Agilent, p/n 5188-5326), Magnetic stir bar (Corning, p/n 401435), Magnetic stir plate (Corning, p/n 6795-410), and Slide-staining dish with slide rack (Thermo Shandon, p/n 121); step 7 – Scanning performed with Agilent Microarray Scanner (Agilent p/n G2565BA); step 8 – Extract data using Agilent Feature Extraction software. Following scanning on the Agilent Scanner G2505B, acquired images were analyzed with Agilent Feature Extraction software (version 10.7.3.1). Data analysis and normalization were performed using the GeneSpring GX v11.5.1 software package (Agilent Technologies). All microarray work and data processing were carried out by KangChen Bio-tech (Shanghai, P.R. China) [22].

Real-time quantitative polymerase chain reaction (RT-qPCR)

Total RNA was extracted using TRIzol (Invitrogen, CA, USA). Reverse transcription was performed using the PrimeScript RT reagent Kit with gDNA Eraser (Perfect Real Time) (TaKaRa, Dalian, China) according to the manufacturer's protocol. Expression of lncRNAs (ak125809, ak098473, uc002ehu.1, bc043564, NR_027322, and uc003qmb.2) was determined using the SYBRGreen detection method (TaKaRa, Dalian, China). Samples were run on an ABI7500 System (Applied Biosystems, Foster City, CA). Primer sequences for lncRNA detection are listed in Table I. β -actin was

Table I. Primers for qPCR

lncRNA	Forward 5'-3'	Reverse 5'-3'
ak125809	GGGAGGTTTTGCTACTTCTGTG	TGGTCTTTGGGAGTGTTTGT
ak098473	GGAATGTCTCCCTCACCTC	AGCTTGATCCCTCGTCTTTG
uc002ehu.1	TGTAGGATGCCACTGGAAATG	CTTCTGCGTGACAACTGAGGAT
bc043564	CTTTTTCGATCGGAACCGCC	GAGGACGCTCTTCTCCTTG
NR_027322	ACCTACCCGTACTIONATCCC	TGGTGTTCTCGGCTTCTGTC
uc003qmb.2	TGTGGTATTGTTCTCTGGATTG	ACTTCGGAGGCAGTAGTAAATCA
β -actin	TCCTCTCCCAAGTCCACACA	GCACGAAGGCTCATCATTCA

used as a normalization control. The qPCR reaction was carried out in a total volume of 20 μ l. The reaction consisted of 10 μ l of 2 \times SYBR Green Master mix, 1 μ l of PCR Forward Primer (5 μ mol/ μ l), 1 μ l of PCR Reverse Primer (5 μ mol/ μ l), 1 μ l cDNA (10 ng), and 7 μ l of RNase-free water. The RT-qPCR reaction was performed as follows: initial denaturation at 95°C for 10 min, 40 cycles of denaturation at 95°C for 15 s, annealing at 60°C for 30 s, and extension at 72°C for 30s, and a final extension at 72°C for 5 min. Each sample was analyzed in triplicate. The threshold cycle (CT) value was calculated by the mean value. Quantitative differential expression was calculated according to the $2^{-\Delta\Delta Ct}$ method [23]. Statistical analysis was conducted using Student's *t* test and SPSS software (Version 16.0 SPSS Inc.). Statistical significance was set at $p < 0.05$.

Pathway analysis

Differentially expressed mRNAs were examined by pathway analysis using the latest KEGG (Kyoto Encyclopedia of Genes and Genomes) database. A number of specific biological pathways were identified by enrichment of the differentially expressed mRNAs. For statistical significance, the *p*-value cut-off was set at 0.05.

Establishing co-expression networks

Co-expression networks of lncRNAs and mRNAs were constructed by differential gene expression. Different networks were drawn based on calculations of different types of expression data. For example, the co-expression network in the control group was calculated by differential gene expression in control samples, while that in the experimental group was calculated by differential gene expression in experimental samples. We calculated the Pearson correlation coefficient for each pair of genes. Regulatory relationships were identified when the Pearson coefficients achieved statistical significance; subsequently, significant correlation pairs were chosen to construct a network. An array *M* was constructed based on the regulatory relationship between genes and mapped into the network diagram. Within the network diagram, each node represents a single gene, and the edges between two nodes represent an interaction between genes, i.e. the correlation between lncRNA-mRNA, lncRNA-lncRNA, and mRNA-mRNA.

The degree of connectivity was used to assess the importance of the gene in the network. Degrees were used to quantify interactions between target genes. For any given gene, the higher its degree, the more regulatory relationships it is involved in with other genes in the network. Thus, the higher the degree, the more central the gene

is in the network. Clustering coefficients were used to calculate the density of interactions among neighboring genes. The higher the clustering coefficient, the greater was the possibility that the gene was located in a core region and that neighboring genes were part of the core network as well. Core genes, which had high degree values, were defined as genes having extensive connections to other genes in the network. Comparisons between different networks were used to discover different gene connections in different networks.

Statistical analysis

Each experiment was performed in triplicate, and all data were represented as average values. Differences between average values were analyzed using Student's *t* test. All statistical analyses were performed with SPSS software (Version 13.0; SPSS, Inc., Chicago, IL, USA).

Important biological pathways were identified with Fisher's test and the χ^2 test. Both *p*-value and FDR (the *p*-value cut-off is 0.05) were used as selection criteria. Enrichment was calculated based on the following formula: $Re = (nf/n)/(Nf/N)$ ($Re = ENRICHMENT$), where *nf* is the number of differentially expressed genes in a particular gene category, *n* is the total number of genes in that category, *Nf* is the number of differentially expressed genes in the entire microarray, and *N* is the total number of genes included in the microarray [24–26].

Results

Aberrantly expressed lncRNAs in glioma

We performed microarray analyses to identify lncRNAs and mRNAs that were differentially expressed between glioma tumor tissue (O301, O605, and 114982) and peritumoral tissue (O601) (Figure 1 A). Differences in expression are represented as fold change of tumor/normal (T/N). A fold change ≥ 3 was used to filter the data (a negative fold change value indicated down-regulation and a positive value indicated up-regulation). In some cases, log₂ fold changes are listed. *P*-value and fold change were both calculated by normalized expression data. Using these parameters, we identified thousands of lncRNAs that were differentially expressed between glioma and normal tissue. Importantly, these lncRNAs have been described in several publications and are also included in a number of databases, including UCSC_knowngene, Ensembl, RefSeq_NR, H-invDB, Fantom, Fantom_stringent, RNAdb, NRED, misc_lncRNA, and UCR.

We next compared lncRNA expression between tumor tissue and matched peritumoral tissue. While more than 23,000 lncRNAs were detected in all tissues, only a few thousand were found to be significantly differentially expressed. An average




Figure 1. Microarray analysis was applied to detect the lncRNAs and mRNAs in glioma compared to normal peritumoral tissue. **A** – Differentially expressed lncRNAs were detected in gliomas. **B** – Differentially expressed mRNAs were detected in gliomas. **C** – Clustering data of lncRNAs in gliomas were analyzed. **D** – Clustering data of mRNAs in gliomas were analyzed

of 4,858 (4330–5153) lncRNAs were differentially expressed in glioma compared to peritumoral tissue. These consisted of 2,845 (2,610–3,228) up-regulated (fold change > 3.0) lncRNAs and 2,013 (1,720–2,395) down-regulated (fold change < 1/3) lncRNAs. In tissues obtained from the three glioma patients, there were 1,290 lncRNAs that were up-regulated or down-regulated more than 3-fold; additionally, 494 were changed (either up or down) more than 5-fold, and 145 lncRNAs altered by greater than 10-fold. Of all differentially expressed lncRNAs, uc002ukq.2 (log₂ fold change T/N = 8.50108) was the most significantly up-regulated and AK002107 (log₂ fold change T/N = –10.64121) was the most significantly down-regulated.

Of all differentially expressed lncRNAs, a majority were up-regulated in glioma; in contrast, a smaller percentage of lncRNAs were down-regulated in tumor tissue (Figure 1 B). Differentially expressed lncRNAs and mRNAs are provided in Table II. In general, the greater the fold change value, the more likely that a particular lncRNA is involved in tumor development. To narrow down our list, we chose 6 lncRNAs (ak125809, ak098473, uc002ehu.1, bc043564, NR_027322, and uc003qmb.2) showing the greatest fold changes and the highest raw intensity values for further investigation.

Aberrantly expressed mRNAs in glioma

We identified 18,315 mRNA transcripts that were differentially expressed between tumor

Table II. Summary of data from microarray for three pairs of glioma and adjacent normal tissues

Relative long noncoding RNA expression between three glioma and adjacent normal tissues						
Sample		Long non-coding RNA				
ID		Fold change > 3	Fold change > 5	Fold change > 10	Total	Changed lncRNA
Couple1	Up-regulation	2029	945	254	3228	5153
(0301 vs. 0601)*	Down-regulation	1051	567	307	1925	
Couple2	Up-regulation	1668	741	201	2610	4330
(0605 vs. 0601)	Down-regulation	1084	403	233	1720	
Couple3	Up-regulation	1538	794	464	2696	5091
(114982 vs. 0601)	Down-regulation	1330	680	385	2395	
Relative messenger RNA expression between three glioma and adjacent normal tissues						
Sample		Messenger RNA				
		Fold change > 3	Fold change > 5	Fold change > 10	Total	Changed mRNA
Couple1	Up-regulation	1363	673	244	2280	4143
(0301 vs. 0601)	Down-regulation	964	445	454	1863	
Couple2	Up-regulation	1106	558	272	1936	3548
(0605 vs. 0601)	Down-regulation	916	398	298	1612	
Couple3	Up-regulation	1353	856	417	2626	4564
(114982 vs. 0601)	Down-regulation	903	536	499	1938	
Glioma tissues: 0301, 0605, 114982. adjacent normal tissue: 0601						

tissue and normal tissue. An average of 2,280 (1,936–2,626) mRNAs were up-regulated more than 3-fold in tumor tissue and an average of 1,804 (1,612–1,938) mRNAs were down-regulated (fold change < 1/3) in tumors. In the three glioma samples, 1,735 mRNAs were differentially expressed greater than 3-fold, 762 mRNAs greater than 5-fold, and 272 mRNAs greater than 10-fold (Table II).

Pathway analysis

Pathway analysis was performed to identify the major signaling pathways activated in tumor cells. We identified 19 pathways that were associated with aberrantly up-regulated transcripts; the most highly enriched pathway was Cell cycle – Homo sapiens (human) consisting of 34 differentially expressed target genes. In contrast, 39 pathways were associated with down-regulated transcripts; the most highly enriched pathway in this group was Salivary secretion – Homo sapiens (human) consisting of 21 target genes (recommended *p*-value cut-off = 0.05). Based on these analyses, we can also conclude that the “GLIOMA” signaling pathway (KEGG Pathway ID: hsa05214) was clearly involved in glioma development (Figure 2).

Establishing gene co-expression networks

We established lncRNA-mRNA co-expression networks to further investigate the connection between these two types of molecules. Only lncRNAs and mRNAs that were at least 5-fold differentially expressed between tumor and normal tissue were selected for construction of the co-expression network (Figure 3). Co-expression networks were constructed based on the expression values of differentially expressed lncRNAs and mRNAs in each sample. Since gene expression in tumors reflected the regulatory relationships among genes, the co-expression networks of differentially expressed lncRNAs and mRNAs were constructed based on tumor-specific conditions.

The network property of each node was analyzed based on the co-expression network. The degree of each gene in the network was used to predict mRNA/lncRNA interaction. A list of mRNAs and lncRNAs with the highest degrees is provided in Table III.

Using the mRNA/lncRNA network, we identified 119 lncRNAs and 335 mRNAs showing some degree of interaction in glioma (correlation = 0.99, *p* = 0.007). Of these, LOC729991 and NUDCD1 mRNAs had the highest degree value. ENST00000425922 and ENST00000455568 lncRNAs were the most




Figure 2. Pathway analysis of aberrantly expressed transcripts in gliomas. **A** – Key genes belonging to the “GLIOMA” signaling pathway (KEGG: hsa05214), including PDGF, SHC, MDM2, and EGFR, also exhibit close relationships with lncRNAs

extensively connected with other mRNAs and lncRNAs in the network. Additionally, key genes of the “GLIOMA” signaling pathway, including SHC3, PDGFA, and MDM2, all had high degree scores.

RT-qPCR verification

lncRNAs that were differentially expressed in all three tumor groups and had fold changes > 10 were arranged in descending order based on normalized intensity. The top six lncRNAs in this group (ak125809, ak098473, uc002ehu.1, bc043564, NR_027322, and uc003qmb.2) were validated by qPCR. We found that lncRNAs ak125809, ak098473, and uc002ehu.1 were up-regulated in over 90% of the 40 glioma cases. In contrast, lncRNAs bc043564, NR_027322, and uc003qmb.2 were down-regulated in tumor tissue compared to normal. These data are consistent with the microarray results ($p = 0.03$) (Figure 4).

Discussion

The genetic basis of glioma development has been extensively studied, and the molecular path-

ways that become deregulated in the disease have also been examined [27, 28]. Aberrant expression of both mRNA transcripts and microRNAs have been shown to be involved in glioma development and progression [29, 30]. One study performed miRNA expression profiling and found that differential expression of miRNAs in malignant astrocytomas may influence tumor proliferation and response to chemotherapy [31]. While a role for both mRNAs and miRNAs in glioma has been well studied, a role for lncRNAs in the disease has yet to be established. Identification of lncRNAs that are important for disease progression could result in new options for glioma prognosis and treatment [16, 32].

In this study, we performed lncRNA expression profiling using microarray technology. We identified thousands of lncRNAs that were differentially expressed in human glioma samples compared to peritumoral normal tissue. We selected 6 lncRNAs (ak125809, ak098473, uc002ehu.1, bc043564, NR_027322, and uc003qmb.2) with fold changes greater than 10 and with relatively high normalized intensities for qPCR validation in 40 pairs of




Figure 2. Cont. **B** – Pathway-associated down-regulated mRNAs are provided. **C** – Pathway-associated up-regulated mRNAs are provided

gliomas and their surrounding tissue. We hypothesize that these lncRNAs may be potential biomarkers in glioma. Additionally, the lncRNA HULC may be useful as a biomarker capable of being detected in plasma [12].

One of the 6 lncRNAs that was extensively examined in this study, ak125809, was upregulated

in tumor tissue. ak125809 is located on chr7 and has a length of 3854 nt. It is the natural anti-sense transcript of the gene encoding vasoactive intestinal peptide receptor 2 (VIPR2), also known as PACAP-R-3. VIPR2-mediated signal transduction results in glial cell proliferation, a process that is closely related to glioma formation and




Figure 3. LncRNA-mRNA co-expression network: nodes with red cycle represent lncRNAs, nodes without cycle represent mRNAs, straight lines represent interactions between genes, purple represents increased expression, and blue represents decreased expression. The size of the node represents the degree; the higher the degree, the more genes interact with the particular node in the network

repair following neurological damage. Ak125809 may have an indirect relation to glioma proliferation due to its regulation of VIPR2 expression [33]. The lncRNA BC043564, located on chr1 with a length of 2022 nt, is the exon sense-overlapping transcript of the *Kcn2a* gene, which encodes one subtype of proliferation-related voltage-gated potassium ion channel [34]. The lncRNA ak098473, located on chrX with a length of 1658 nt, is the natural antisense transcript of the gene encoding synaptotagmin-like protein 4 (SYTL4). The lncRNA uc002ehu.1 is located on chr17 and has a length of 643 nt. NR_027322 is located on chr10 with a length of 2427 nt. The biological

function of the latter three lncRNAs is currently unknown.

We identified several oncogene-related lncRNAs in our microarray analysis. BC007968 was found to be significantly upregulated; it is a natural antisense transcript of cyclin-dependent kinase 4 (CDK4), a Ser/Thr protein kinase. Expression of CDK4 and its associated proteins, including D-type cyclins, p16 (INK4a), and Rb, has been shown to be important for glioma tumorigenesis; moreover, disruption of the p16INK4a-Cdk4-Rb pathway has been shown to be useful for glioma therapy [35, 36]. BC007968 is very likely to interfere with the p16INK4a-Cdk4-Rb pathway as a CDK4 antisense

Table III. Degree was used to assess interactions in the lncRNA/mRNA network. This table is a collection of a series of key lncRNA/mRNAs

Gene symbol	Style	Regulation	Degree	Gene symbol	Style	Regulation	Degree
ENST00000425922	Up	LncRNA	13	LOC729991	Down	mRNA	15
ENST00000455568	Down	LncRNA	12	NUDCD1	Up	mRNA	15
ENST00000500949	Down	LncRNA	12	SNX32	Down	mRNA	14
uc001xvj.2	Up	LncRNA	12	TRAF3IP1	Down	mRNA	14
uc002ukq.2	Up	LncRNA	12	KCNK7	Down	mRNA	13
uc003qmb.2	Down	LncRNA	11	MORN1	Down	mRNA	13
ENST00000502715	Up	LncRNA	11	PALM2-AKAP2	Down	mRNA	13
ENST00000502999	Up	LncRNA	11	PMP2	Up	mRNA	13
NR_028344	Down	LncRNA	11	PXK	Down	mRNA	13
uc002pgz.1	Up	LncRNA	11	TXNRD3	Up	mRNA	13
ENST00000450820	Up	LncRNA	11	ZNF501	Up	mRNA	13




Figure 4. Comparison of microarray data and qPCR results. **A** – qPCR was used to verify expression of lncRNAs ak125809, ak098473, uc002ehu.1, bc043564, NR_027322, and uc003qmb.2. **B** – Distribution of lncRNA expression levels were provided. All six lncRNAs of ak125809, ak098473, uc002ehu.1, bc043564, NR_027322, and uc003qmb.2 were validated by qPCR analysis in the 40 paired glioma and peritumoral tissues. Each histogram represents the average fold change (T/N) with logarithmic conversion. Error bars are indicative of standard deviation. Distribution of lncRNA expression

transcript and would be predicted to regulate glioma formation. The lncRNA BC008580 was also upregulated in glioma. This is an exon sense-overlapping transcript of glial cell line-derived neurotrophic factor (GDNF). Recent work has shown that GDNF is overexpressed in glioma, where it likely contributes to increased proliferation and metastasis [37, 38]. It is possible that the lncRNA BC008580 contributes to glioma development via regulation of GDNF. Taken together, our data support a role for lncRNAs in glioma tumorigenesis.

Interaction or co-expression networks can be generated based on multiple assays, including co-immunoprecipitation experiments [39] and gene function analyses [40]. We generated an lncRNA-mRNA expression network based on data from our microarray expression profiling. We determined Pearson correlation coefficients and selected highly correlated lncRNA-mRNA pairs to establish the network. The network indicated which lncRNAs might be responsible for regulation of specific mRNAs. While mRNAs seemed to directly

influence tumorigenesis and disease malignancy, lncRNAs tended to have a more indirect effect, possibly via regulation of downstream mRNAs. Degree scores were assigned to molecules in the network, and those with the highest degree scores likely displayed the greatest extent of interaction. Through these analyses, we identified potential lncRNA-mRNA interactions. A better understanding of these interactions will likely help elucidate some key roles for lncRNAs in glioma.

We performed KEGG annotation of all mRNAs differentially expressed between glioma and normal tissue. Several pathways, including MAPK signaling, the Wnt pathway, p53 signaling, and the glioma pathway, were all found to be deregulated in the disease. Most of these pathways have been previously shown to influence glioma tumorigenesis and response to chemotherapy. For example, one study showed that down-regulation of Wnt2 and beta-catenin inhibited proliferation of malignant glioma [41]. Additionally, work by Johannesen *et al.* showed that activation of p53 by the MDM2 antagonist nutlin-3 improved sensitivity to chemotherapy treatment [42]. Of all pathways identified by KEGG analysis, the “Glioma” pathway (KEGG Pathway ID: hsa05214) particularly stood out [43–45]. Several genes belonging to this pathway, including PDGF, SHC, MDM2, and EGFR, have previously been shown to be important for glioma development [46, 47]. Many of these genes were also found to be differentially expressed in this study. Additionally, our lncRNA-mRNA co-expression network analysis showed that several of these genes correlate with particular lncRNAs. For example, *shc3* correlated with uc002ukz.1, enst00000473644, and uc010ayj.1 (correlation = 1, $p < 0.01$). *Pdgfra* correlated with enst00000502715 and NR_022011. *Mdm2* correlated with NR_027873 and uc.208- (correlation = 1, $p < 0.01$). We hypothesize that many of these lncRNAs regulate glioma formation by interfering with the expression of key factors in the “Glioma” pathway. We plan to test this hypothesis in future studies.

In this study, we identified differentially expressed lncRNAs in human glioma cases by both microarray profiling and qPCR validation. Importantly, data from both microarray and qPCR were consistent. We established an lncRNA-mRNA co-expression network to better understand the mechanisms of lncRNA action in the disease. We found lncRNA that may be associated with the pathogenesis of glioma, as a phenomenon, but the mechanism was not further explored, and whether it can be detected in blood and body fluids and become a biomarker is yet to be studied. The project team will further study this series of issues later. While this work provides a solid foundation, more research must be done to clearly

define the molecular roles of lncRNAs in glioma development.

Acknowledgments

This work was supported by the Starting Fund for Young Scientists of Nanjing Public Health Bureau (QYK11126).

Yi Ding, Xinfu Wang and Junchen Pan have equal contribute to this paper. Thanks to Mingjun Jis laboratory and their colleagues.

Conflict of interest

The authors declare no conflict of interest.

References

1. Patnaik A, Mishra SS, Senapati SB. Intraventricular glioblastoma multiforme mimicking meningioma and review of the literature. *Asian J Neurosurg* 2017; 12: 75-7.
2. Ohgaki H, Kleihues P. Epidemiology and etiology of gliomas. *Acta Neuropathol* 2005; 109: 93-108.
3. Zhong D, Ran J, Zhang X, et al. Syntenin is expressed in human gliomas and may correlate with tumor migration. *Arch Med Sci* 2015; 11: 1303-7.
4. Wilusz JE, Sunwoo H, Spector DL. Long noncoding RNAs: functional surprises from the RNA world. *Genes Dev* 2009; 23: 1494-504.
5. Zhao J, Sun BK, Erwin JA, Song JJ, Lee JT. Polycomb proteins targeted by a short repeat RNA to the mouse X chromosome. *Science* 2008; 322: 750-6.
6. Pandey RR, Mondal T, Mohammad F, et al. Kcnq1ot1 antisense noncoding RNA mediates lineage-specific transcriptional silencing through chromatin-level regulation. *Mol Cell* 2008; 32: 232-46.
7. Wang X, Arai S, Song X, et al. Induced ncRNAs allosterically modify RNA-binding proteins in cis to inhibit transcription. *Nature* 2008; 454: 126-30.
8. Beltran M, Puig I, Peña C, et al. A natural antisense transcript regulates Zeb2/Sip1 gene expression during Snail1-induced epithelial-mesenchymal transition. *Genes Dev* 2008; 22: 756-69.
9. Yu G, Yao W, Wang J, et al. lncRNAs expression signatures of renal clear cell carcinoma revealed by microarray. *PLoS One* 2012; 7: e42377.
10. Xu C, Yang M, Tian J, Wang X, Li Z. MALAT-1: a long non-coding RNA and its important 3' end functional motif in colorectal cancer metastasis. *Int J Oncol* 2011; 39: 169-75.
11. Birney E, Stamatoyannopoulos JA, Dutta A, et al. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 2007; 447: 799-816.
12. Panzitt K, Tschernatsch MM, Guelly C, et al. Characterization of HULC, a novel gene with striking up-regulation in hepatocellular carcinoma, as noncoding RNA. *Gastroenterology* 2007; 132: 330-42.
13. Yuan SX, Yang F, Yang Y, et al. Long noncoding RNA associated with microvascular invasion in hepatocellular carcinoma promotes angiogenesis and serves as a predictor for hepatocellular carcinoma patients' poor recurrence-free survival after hepatectomy. *Hepatology* 2012; 56: 2231-41.
14. Bracken AP, Helin K. Polycomb group proteins: navigators of lineage pathways led astray in cancer. *Nat Rev Cancer* 2009; 9: 773-84.

15. Calin GA, Liu CG, Ferracin M, et al. Ultraconserved regions encoding ncRNAs are altered in human leukemias and carcinomas. *Cancer Cell* 2007; 12: 215-29.
16. Wang P, Ren Z, Sun P. Overexpression of the long non-coding RNA MEG3 impairs in vitro glioma cell proliferation. *J Cell Biochem* 2012; 113: 1868-74.
17. Benetatos L, Vartholomatos G, Hatzimichael E. MEG3 imprinted gene contribution in tumorigenesis. *Int J Cancer* 2011; 129: 773-9.
18. Han L, Zhang K, Shi Z, et al. LncRNA profile of glioblastoma reveals the potential role of lncRNAs in contributing to glioblastoma pathogenesis. *Int J Oncol* 2012; 40: 2004-12.
19. Zhang X, Sun S, Pu JK, et al. Long non-coding RNA expression profiles predict clinical phenotypes in glioma. *Neurobiol Dis* 2012; 48: 1-8.
20. Guo Y, Vickers K, Xiong Y, et al. Comprehensive evaluation of extracellular small RNA isolation methods from serum in high throughput sequencing. *BMC Genomics* 2017; 18: 50.
21. Han B, Bu P, Meng X, Hou X. Microarray profiling of long non-coding RNAs associated with idiopathic pulmonary arterial hypertension. *Exp Ther Med* 2017; 13: 2657-66.
22. Shang D, Zheng T, Zhang J, Tian Y, Liu Y. Profiling of mRNA and long non-coding RNA of urothelial cancer in recipients after renal transplantation. *Tumour Biol* 2016; 37: 12673-84.
23. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 2001; 29: e45.
24. Draghici S, Khatri P, Tarca AL, et al. A systems biology approach for pathway level analysis. *Genome Res* 2007; 17: 1537-45.
25. Yi M, Horton JD, Cohen JC, Hobbs HH, Stephens RM. WholePathwayScope: a comprehensive pathway-based analysis tool for high-throughput data. *BMC Bioinformatics* 2006; 7: 30.
26. Kanehisa M, Goto S, Kawashima S, Okuno Y, Hattori M. The KEGG resource for deciphering the genome. *Nucleic Acids Res* 2004; 32: D277-80.
27. Yang H, Zhang QX, Pei DS, Xu F, Li Y, Yu RT. FK506-binding protein 5 inhibits proliferation and stimulates apoptosis of glioma cells. *Arch Med Sci* 2015; 11: 1074-80.
28. Hong Y, Shi Y, Shang C, Xue Y, Liu Y. Influence of far upstream element binding protein 1 gene on chemotherapy sensitivity in human U251 glioblastoma cells. *Arch Med Sci* 2016; 12: 156-62.
29. Vitucci M, Hayes DN, Miller CR. Gene expression profiling of gliomas: merging genomic and histopathological classification for personalised therapy. *Br J Cancer* 2011; 104: 545-53.
30. Shai R, Shi T, Kremen TJ, et al. Gene expression profiling identifies molecular subtypes of gliomas. *Oncogene* 2003; 22: 4918-23.
31. Rao SA, Santosh V, Somasundaram K. Genome-wide expression profiling identifies deregulated miRNAs in malignant astrocytoma. *Mod Pathol* 2010; 23: 1404-17.
32. Barsyte-Lovejoy D, Lau SK, Boutros PC, et al. The c-Myc oncogene directly induces the H19 noncoding RNA by allele-specific binding to potentiate tumorigenesis. *Cancer Res* 2006; 66: 5330-7.
33. Jaworski DM. Expression of pituitary adenylate cyclase-activating polypeptide (PACAP) and the PACAP-selective receptor in cultured rat astrocytes, human brain tumors, and in response to acute intracranial injury. *Cell Tissue Res* 2000; 300: 219-30.
34. Akhtar S, McIntosh P, Bryan-Sisneros A, Barratt L, Robertson B, Dolly JO. A functional spliced-variant of beta 2 subunit of Kv1 channels in C6 glioma cells and reactive astrocytes from rat lesioned cerebellum. *Biochemistry* 1999; 38: 16984-92.
35. Parsons DW, Jones S, Zhang X, et al. An integrated genomic analysis of human glioblastoma multiforme. *Science* 2008; 321: 1807-12.
36. Cancer Genome Atlas Research Network: Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature* 2008; 455: 1061-8.
37. Ng WH, Wan GQ, Peng ZN, Too HP. Glial cell-line derived neurotrophic factor (GDNF) family of ligands confer chemoresistance in a ligand-specific fashion in malignant gliomas. *J Clin Neurosci* 2009; 16: 427-36.
38. Golan M, Schreiber G, Avissar S. Antidepressants elevate GDNF expression and release from C(6) glioma cells in a beta-arrestin1-dependent, CREB interactive pathway. *Int J Neuropsychopharmacol* 2011; 14: 1289-300.
39. Smidtas S, Yartseva A, Schachter V, Kepes F. Model of interactions in biology and application to heterogeneous network in yeast. *C R Biol* 2006; 329: 945-52.
40. Nikiforova VJ, Willmitzer L. Network visualization and network analysis. *EXS* 2007; 97: 245-75.
41. Pu P, Zhang Z, Kang C, et al. Downregulation of Wnt2 and beta-catenin by siRNA suppresses malignant glioma cell growth. *Cancer Gene Ther* 2009; 16: 351-61.
42. Johannessen TC, Prestegarden L, Grudic A, Hegi ME, Tysnes BB, Bjerkvig R. The DNA repair protein ALKBH2 mediates temozolomide resistance in human glioblastoma cells. *Neuro Oncol* 2013; 15: 269-78.
43. Bansal K, Liang ML, Rutka JT. Molecular biology of human gliomas. *Technol Cancer Res Treat* 2006; 5: 185-94.
44. Kapoor GS, O'Rourke DM. Mitogenic signaling cascades in glial tumors. *Neurosurgery* 2003; 52: 1425-34; discussion 1434-25.
45. Hulleman E, Helin K. Molecular mechanisms in glioma genesis. *Adv Cancer Res* 2005; 94: 1-27.
46. Martinho O, Longatto-Filho A, Lambros MB, et al. Expression, mutation and copy number analysis of platelet-derived growth factor receptor A (PDGFRA) and its ligand PDGFA in gliomas. *Br J Cancer* 2009; 101: 973-82.
47. Ortensi B, Osti D, Pellegatta S, et al. Rai is a new regulator of neural progenitor migration and glioblastoma invasion. *Stem Cells* 2012; 30: 817-32.