

RESEARCH LETTER – Physiology & Biochemistry

Cysteine biosynthesis in *Lactobacillus casei*: identification and characterization of a serine acetyltransferase

Biljana Bogicevic¹, H el ene Berthoud¹, Reto Portmann¹, Tharmatha Bavan¹, Leo Meile² and Stefan Irmeler^{1,*}¹Agroscope, Institute for Food Sciences IFS, 3003 Berne, Switzerland and ²Laboratory of Food Biotechnology, Institute of Food Science and Nutrition, ETH Z urich, 8092 Z urich, Switzerland

*Corresponding author: Agroscope, Institute for Food Sciences IFS, Schwarzenburgstrasse 161, 3003 Berne, Switzerland. Tel: +41-58-463-81-55;

Fax: +41-58-463-82-27; E-mail: stefan.irmeler@agroscope.admin.ch**One sentence summary:** A gene that is annotated as homoserine succinyltransferase actually encodes a serine acetyltransferase in *Lactobacillus casei*.

Editor: Akihito Endo

ABSTRACT

In bacteria, cysteine can be synthesized from serine by two steps involving an L-serine O-acetyltransferase (SAT) and a cysteine synthase (CysK). While CysK is found in the publicly available annotated genome from *Lactobacillus casei* ATCC 334, a gene encoding SAT (*cysE*) is missing. In this study, we found that various strains of *L. casei* grew in a chemically defined medium containing sulfide as the sole sulfur source, indicating the presence of a serine O-acetyltransferase. The gene lying upstream of *cysK* is predicted to encode a homoserine trans-succinylase (*metA*). To study the function of this gene, it was cloned from *L. casei* FAM18110. The purified, recombinant protein did not acylate L-homoserine *in vitro*. Instead, it catalyzed the formation of O-acetyl serine from L-serine and acetyl-CoA. Furthermore, the plasmid expressing the *L. casei* gene complemented an *Escherichia coli cysE* mutant strain but not an *E. coli metA* mutant. This clearly demonstrated that the gene annotated as *metA* in fact encodes the SAT function and should be annotated as *cysE*.

Keywords: *Lactobacillus casei*; serine acetyltransferase; *cysE*; cysteine biosynthesis

INTRODUCTION

Lactobacillus casei is often present in the non-starter microbiota of cheese at the end of ripening (Beresford and Williams 2004). This implies that this species plays a significant role in proteolysis and flavor development during cheese ripening and that it could be used as adjunct culture with starter bacteria to improve ripening control and enhance flavor intensity. Knowledge of metabolic activities and pathways is advantageous in the selection of proper strains. The increasing number of newly sequenced genomes allows for the prediction of phylogenetic relatedness and metabolic pathways (Siezen *et al.* 2004; Liu *et al.*

2008). However, annotation of these genomes relies mostly on sequence and structural comparisons, which may lead to incorrect functional gene assignments and consequently to incorrect metabolic pathway predictions.

The biosynthesis of cysteine in cheese-inhabiting bacteria is of particular interest since cysteine is, unlike methionine, present in lower amounts in caseins (Farrell *et al.* 2004) and could therefore be a growth-limiting source. Furthermore, cysteine and methionine are precursors for the formation of odor-active volatile sulfur compounds during cheese ripening (Landaud, Helinck and Bonnarme 2008).

Received: 4 November 2015; Accepted: 21 December 2015

  FEMS 2016. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com




Figure 1. Biosynthesis of cysteine in *L. casei* based on genome sequence information. Genes encoding SAT and CBS have not yet been identified. Recently, we presented functional data for CysK, MetB, MalY and CGL (Irmeler et al. 2008, 2009; Bogicevic et al. 2012). A gene putatively encoding either SAT or MetA (shaded in gray) was studied in this report. SAT: serine acetyltransferase; CysK: cysteine synthase; MetA: homoserine O-succinyltransferase; MetB: cystathionine gamma-synthase; MalY: cystathionine beta-lyase; CGL: cystathionine gamma-lyase; CBS: cystathionine beta-synthase.

Cysteine can be synthesized from serine or homocysteine (Fig. 1). The biosynthesis from serine is well characterized in *Enterobacteriaceae* (Kredich 1996). First, serine is acetylated to O-acetyl-L-serine (OAS) by a serine acetyltransferase (SAT, EC 2.3.1.30), which is encoded by *cysE*. Second, cysteine synthase (EC 2.5.1.47) encoded by *cysK* replaces the acetyl group by sulfide to form cysteine. We have recently demonstrated that the orthologous *cysK* gene from *L. casei* FAM18110 actually synthesized cysteine from OAS and sulfide (Bogicevic et al. 2012). Surprisingly, the annotated genome of *L. casei* ATCC 334 apparently does not carry a gene encoding SAT, which is necessary for the formation of OAS (Makarova et al. 2006; Liu et al. 2008). In connection with the characterization of *cysK* from *L. casei*, we observed that the gene was co-transcribed with the coding sequence that lies upstream of *cysK* (Bogicevic et al. 2012). This gene is annotated as homoserine O-succinyltransferase (MetA, EC 2.3.1.46) in various *L. casei* genomes that are deposited in the GenBank database (Table S1, Supporting Information). MetA is an enzyme that catalyzes the acylation of homoserine (Fig. 1) and should therefore be involved in methionine biosynthesis (Born and Blanchard 1999).

In this paper, we provide functional data showing that *L. casei* synthesizes cysteine from serine. First, the growth of various *L. casei* strains with sulfide as the sulfur source was determined. Furthermore, the gene encoding the putative *metA* was cloned and the recombinant protein was used for enzymatic assays. Finally, complementation tests of *Escherichia coli* mutants were performed to confirm enzymatic results.

MATERIALS AND METHODS

Bacterial strains, plasmids, DNA manipulation and culture media

The strains and plasmids used in this study are listed in Table 1. Genomic and plasmid DNA were prepared as described previously (Bogicevic et al. 2012). *Escherichia coli* strains were grown at 37°C in LB broth (Sambrook, Fritsch and Maniatis 1989). When

necessary, ampicillin (amp) was added at a concentration of 100 µg mL⁻¹. *Lactobacillus casei* strains were grown at 30°C in MRS broth (de Man, Rogosa and Sharpe 1960).

The chemically defined medium (CDM) described by Christensen and Steele (2003) was used to study the dependency on various sulfur sources. CDM containing either sodium sulfide, L-cysteine or L-cysteine together with L-methionine at a final concentration of 0.7 mM was inoculated with *L. casei* (1% v/v). After incubation for 3 days at 30°C, the optical density at 600 nm (OD₆₀₀) of the cultures was determined using a spectrophotometer (LKB Biochrom 4050 Ultrospec II).

Cloning, expression and purification of His-tagged CysE

The orthologous *LSEI.0479* gene was amplified from genomic DNA of *L. casei* FAM18110 with the primer pair *LSEI.0479.F1/LSEI.0479.R* (Table 1) and cloned into the pEXP5-CT/TOPO plasmid vector (Life Technologies, Zug, Switzerland) according to the manufacturer's instructions. The plasmid containing the gene in the proper orientation was named pEXP5-CT/*cysE* and maintained in *E. coli* BL21(DE3). Expression of recombinant protein was performed as described previously (Bogicevic et al. 2012).

The recombinant protein was purified with Protino Ni-TED 1000 packed columns kit (Machery-Nagel, Düren, Germany) according to the manufacturer's instructions. Therefore, bacterial cells were suspended in 1 mL of LEW buffer provided in the kit and disrupted with glass beads (212–300 µm in diameter, 0.6 g) and an Omni Bead Ruptor 24 (Omni International, Kennesaw, US). After elution, the buffer of the purified protein was exchanged to 20 mM sodium phosphate using PD-10 columns (GE Healthcare, Little Chalfont, UK). Finally, the protein was concentrated with Ultracel 30K centrifugal filters (Millipore, Cork, Ireland). The concentration and purity of the purified protein was determined with the Qubit Protein Assay Kit (Life Technologies) and with sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), followed by colloidal Coomassie staining, respectively.

Table 1. Strains, plasmids and primers used in this study.

Strains	Genotype or relevant properties	Source, reference or purpose
<i>E. coli</i> BL21(DE3)	F ⁻ , <i>ompT hsdS_B(r_B⁻ m_B⁻) gal dcm</i> (DE3)	Invitrogen
JM39	F ⁺ , <i>cysE51 trf-8</i>	Jones-Mortimer (1968)
DL41	F ⁻ , λ ⁻ , <i>metA28</i>	Hendrickson, Horton and Lemaster (1990)
<i>L. casei</i> FAM8407, FAM 18101, FAM18108, FAM18110, FAM18121	Isolated from cheese	Agroscope Culture Collection
FAM18149, FAM18168, FAM18172 ATCC 334	Isolated from milk Genome sequenced strain, isolated from cheese	American Type Culture Collection
Plasmids		
pEXP5-CT/TOPO	<i>E. coli</i> cloning vector, Amp ^r , 2685 bp	Invitrogen
pEXP5-CT/ <i>cysE</i>	pEXP5-CT/TOPO containing <i>cysE</i> from <i>L. casei</i> FAM18110, 3498 bp	This study
pEXP5-NT/CALML3	Expression control plasmid, 3194 bp	Life Technologies
Primers		
	Sequence (5' to 3')	Used for
LSEL_0479_F1	ATG ^a GAAAAATCGCCACTCAAGAT	Cloning of <i>L. casei cysE</i> (this study)
LSEL_0479_R	ATGATTGTCATTGCGATGATCCG	Cloning of <i>L. casei cysE</i> (this study)

^aThe ORF of LSEL_0479 in *L. casei* ATCC 334 starts with TTG.

Enzyme assays

The acylation of L-serine, D-serine and L-homoserine was studied using a method based on mass spectrometry: for this, an enzymatic assay was performed in a syringe at RT containing 200 μL of 50 mM ammonium acetate (pH 7.4), 1 mM amino acid, 0.06 mM acyl-CoA (acetyl-CoA, propionyl-CoA, butyryl-CoA or succinyl-CoA) and recombinant enzyme. To test inhibition, 1 mM L-cysteine was added.

The reaction mixture was analyzed via continuous flow injection at 5 μL min⁻¹ into a high-resolution Bruker maXis 4G+ QTOF-mass spectrometer (Bruker Daltonics GmbH, Bremen, Germany) equipped with an electrospray ionization (ESI) interface. The mass spectrometer was operated in positive ion mode with a capillary voltage of 4 kV, an endplate offset of -500 V, a nebulizer pressure of 0.4 bar and a drying gas flow rate of 8 L min⁻¹ at 200°C.

Data acquisition, processing and reporting were achieved using Data Analysis 4.2 software (Bruker Daltonics GmbH, Bremen, Germany). The observed *m/z* window was in the range from 75 to 1250.

A photometric assay was used to determine the kinetic parameters. The release of coenzyme A from the acetylated form was measured with 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB). The reaction mixture contained 0.2 mL of 50 mM ammonium acetate buffer (pH 7.4), 1 mM DTNB, 1 mM EDTA, various concentrations of L-serine (0.1–10 mM) and various concentrations of acetyl-CoA (0.01–0.16 mM). The *K_M* for L-serine was determined with a fixed concentration of acetyl-CoA (0.06 mM), while the *K_M* for acetyl-CoA was determined with a fixed concentration of L-serine (1 mM). The reaction was initiated by the addition of purified recombinant protein and carried out at 25°C. The change of absorption at 412 nm was recorded at 5 min intervals for 60 min. *K_M* was determined by using the Hanes-Woolf

transformation (S versus S^{-1}), where v is the formation rate of CoA and S is the concentration of each substrate.

Complementation of *E. coli* auxotrophs

The plasmids pEXP5-CT/*cysE* and pEXP5-NT/CALML3 (Table 1) were used to transform the cysteine auxotroph *E. coli* JM39 and the methionine auxotroph *E. coli* DL41. Transformants were selected on LB/amp agar plates, checked for the presence of the plasmids and finally grown in LB/amp to yield stocks of the transformed bacteria. Transformed *E. coli* were grown in LB/amp medium for 6 h. After the determination of OD₆₀₀, the cells were harvested by centrifugation (3000 g, 10 min), washed once with 1X M9 salts and then resuspended in 1X M9 salts to a final OD₆₀₀ of 2. Transformed *E. coli* JM39 (50 μL) were plated on M9 minimal medium (Sambrook, Fritsch and Maniatis 1989) containing glucose (0.4%), IPTG (1 mM) and ampicillin (100 μg mL⁻¹) in the absence and presence of L-cysteine (1 mM). Transformed *E. coli* DL41 were plated as described above but in the absence and presence of L-methionine (1 mM). All plates were incubated at 37°C for 3 days.

RESULTS

Growth with various sulfur compounds

We studied the requirement for various sulfur sources by inoculating a CDM containing either sulfide, cysteine or methionine with various *L. casei* strains (Table 1). Medium that did not contain any of these sulfur compounds served as negative control. All strains tested grew in medium containing cysteine and methionine (Fig. 2), whereas no growth was observed in medium lacking the sulfur compounds. All strains also showed proper growth, when either sulfide or cysteine was




Figure 2. Bacterial growth of various *L. casei* strains in a CDM that contained neither sulfur containing amino acids nor sulfide (white bars). To study dependency on sulfur sources, sodium sulfide (striped bars), cysteine (gray bars) or cysteine and methionine (dark bars) was added. The data are represented as the mean \pm SD of three independently conducted experiments.

added as the sole sulfur source, except ATCC 334 and FAM18172 which failed to grow under both conditions.

Cloning of the putative *cysE* gene

The genome of six *L. casei* strains deposited in GenBank contains an 816-bp coding sequence, annotated as homoserine succinyltransferase, which lies upstream of *cysK* (Table S1, Supporting Information). A comparison of these genes showed a high nucleotide sequence identity (above 98%). Based on this information, we designed the primer pair LSEI_0479_F1/R (Table 1) to amplify the gene from *L. casei* FAM18110. A PCR product of approximately 800 bp was obtained and cloned into the expression vector pEXP5-CT/TOPO, thereby adding a 6 \times His tag to the C-terminus of the predicted protein.

The deduced amino acid sequence of the amplified gene shared a high sequence identity (above 98%) with the putative homoserine succinyltransferases from other *L. casei* strains (Fig. 3). Since we assumed that this gene in fact encodes a serine O-acetyltransferase, we call the gene *cysE* and the protein SAT in the remainder of this report. The nucleotide sequence of the *cysE* gene was deposited at GenBank under the accession number KU216159.

Heterologous expression of pEXP5-CT/*cysE* in *E. coli* BL21(DE3) yielded a soluble recombinant protein, which was purified by




Figure 3. Protein sequence alignment of the SAT protein from *L. casei* FAM18110 with the putative homoserine O-succinyltransferase from other *L. casei* strains. The sequences are labeled with the locus tag and the corresponding strains are given in brackets. Matching amino acids are represented as dots and amino acid changes are highlighted in gray.




Figure 4. ESI mass spectrum assay with recombinant SAT (3 μ g) from *L. casei*. Illustrated are the mass spectra of incubation of SAT with L-serine and acetyl-CoA (A), L-serine and succinyl-CoA (B), L-serine and propionyl-CoA (C) and L-serine and butyryl-CoA (D) after 20 min reaction time. The peak m/z 148.0, 162.0 and 206.0 correspond to O-acetylserine (OAS), O-propionylserine (OPS) and O-succinylserine (OSS), respectively. The formation of O-butyrylserine (m/z 176) was not observed. The two intense peaks at m/z 106.0 and 128.0 that are present in all spectra represent the protonated L-serine ($M + H$)⁺ and the serine sodium adduct ($M + Na$)⁺, respectively.

nickel affinity chromatography. The purified protein migrated as a single protein band with a molecular weight of 32 kDa in SDS-PAGE (expected 32.5 kDa; data not shown).

Substrate specificity and kinetic parameters of the putative CysE

To test whether the purified enzyme exhibited acylation activity, the enzyme was incubated together with various amino acids (L-serine, D-serine and L-homoserine) and acyl-CoAs (acetyl-CoA, propionyl-CoA, butyryl-CoA and succinyl-CoA). The enzymatic assay was continuously injected into a mass spectrometer. When the enzyme was incubated with L-serine and acetyl-CoA, a peak with an m/z of 148 appeared (Fig. 4A). This peak represents the protonated OAS ($M + H$)⁺. When succinyl-CoA or propionyl-CoA was used together with serine, the enzyme formed O-succinyl serine (m/z 206; Fig. 4B) and O-propionyl serine (m/z 162, Fig. 4C), respectively. The intensities of both peaks were lower than that for OAS, demonstrating that propionyl-CoA and succinyl-CoA were not the preferred acyl donor substrates. The formation of O-butyryl serine (m/z 176) was not observed when butyryl-CoA was used as the acyl donor (Fig. 4D). Further, no product formation was observed when D-serine or L-homoserine was used as acyl acceptors (data not shown).

When L-cysteine was added to the enzymatic reaction as a putative enzyme inhibitor, the peak corresponding to OAS (m/z 148.0) was still clearly detected by MS (Fig. S3, Supporting Information).

The K_M for the acetylation of serine was determined by a photometric assay that contained Ellman's reagent. This compound reacts with the sulfhydryl group of the released CoA, yielding a yellow color. Thus, we determined a K_M of 1.13 (± 0.13) mM and 0.021 (± 0.004) mM for serine and acetyl-CoA, respectively. Again, when L-serine was replaced by D-serine or L-homoserine, no color formation was observed (data not shown).

Functional analysis by genetic complementation

Each of the expression plasmids listed in Table 1 was introduced into the cysteine auxotroph *E. coli* JM39 (*cysE* mutant) and into the methionine auxotroph mutant *E. coli* DL41 (*metA* mutant). Transformation of *E. coli* JM39 with pEXP5-CT/*cysE* resulted in the growth of the strain in the absence of L-cysteine (Fig. 5). Transformed *E. coli* DL41 failed to grow in the absence of L-methionine in the culture medium (Fig. 5).

DISCUSSION

The genome of *L. casei* ATCC 334 was the first strain of this species to be sequenced (Makarova et al. 2006). With regard




Figure 5. Growth properties of transformed *E. coli* JM39 and *E. coli* DL41 bearing either plasmid pEXP5-CT/*cysE* (*cysE*) or pEXP5-NT/CALML3 (CALML3) on minimal medium (MM) agar plates. When indicated (right-hand side), the medium was supplemented with sulfur-containing amino acid.

to the two-step synthesis of cysteine from serine, the gene encoding cysteine synthase is found in the annotated genome, but a gene encoding the enzyme that provides the substrate OAS for the cysteine synthase is missing. This implies that *L. casei* cannot synthesize cysteine from serine. However, when we inoculated a CDM containing only sulfide as the sole sulfur source with various *L. casei* strains, we observed growth in seven out of nine strains. This indicated that *L. casei*, in general, possesses the genes necessary for the synthesis of cysteine from serine and sulfide. The two strains that did not grow with sulfide—ATCC 334 and FAM18172—also did not grow when we used cysteine as the sole sulfur source, but they did grow very well when cysteine and methionine were present (Fig. 2). This indicates that the pathway for the conversion of cysteine to methionine is inactive. Previously, we showed that two genes from *L. casei* FAM18168, named *metB* and *malY*, encode cystathionine synthase and cystathionine lyase, respectively (Irmler *et al.* 2008). We assume that both genes constitute the transsulfuration pathway which converts cysteine to homocysteine, the precursor of methionine. We sequenced both genes from *L. casei* FAM18172 and FAM18110 and compared the deduced amino acid sequences with the one of FAM18168 and ATCC 334 (Figs S1 and S2, Supporting Information). We found that ATCC 334 possesses a strain-specific glu-

tamine to proline amino acid substitution (Q169P) in *MalY*. A slight truncation of the *MetB* protein as a result from a nonsense mutation in the corresponding gene is specific for FAM18172. Whether these genetic changes render the strains auxotrophic for methionine needs further investigations.

With regard to the synthesis of cysteine from serine, we assumed that the *metA* gene that forms an operon with *cysK* in fact encodes a SAT protein. To study our hypothesis, we cloned the putative *metA* from *L. casei* FAM18110 and expressed it heterologously in *E. coli*. We found that the purified recombinant protein clearly acetylated L-serine. Slight acylation activity was also observed when propionyl-CoA and succinyl-CoA were used as substrates. The enzyme did not show acylation activity when D-serine or L-homoserine was used as acyl donors. Taken together with the complementation of an *E. coli* *cysE* mutant strain, this study clearly showed that the *L. casei* gene encodes SAT and therefore should be named *cysE* and not *metA*.

Liu *et al.* (2012) made a comprehensive *in silico* comparison of genes involved in sulfur metabolism. The authors describe that the genomes of other LAB species such as *L. acidophilus*, *L. delbrueckii* subsp. *bulgaricus*, *L. rhamnosus*, *L. helveticus*, *L. fermentum*, *L. kefirifaciens*, *L. reuteri* and *L. salivarius* possess a *cysK* gene but not a *cysE* gene. Moreover, *cysK* is co-localized with a

metA gene in the genomes of this species. Consequently, these species should not be able to synthesize cysteine from serine. The situation resembles the one of *L. casei* and considering the genetic organization we assume that *metA* in these LAB species could also encode a SAT. Our assumption could be clarified by studying growth of these bacteria in the presence and absence of various sulfur compounds.

It has been reported that SAT activity in various organisms is inhibited by cysteine (Johnson, Roderick and Cook 2005). We did not observe this kind of feedback inhibition with the *L. casei* SAT *in vitro*. The SATs from other bacteria, such as *E. coli*, *Salmonella enterica*, *Thermus thermophilus*, *Paracoccus denitrificans* and *Haemophilus influenzae*, have a K_M for serine of 0.11, 0.77, 0.013, 0.4 and 4.7 mM and for acetyl-CoA of 0.56, 0.1, 0.011, 0.1 and 0.7 mM, respectively (Kredich and Tomkins 1966; Burnell and Whatley 1977; Baecker and Wedding 1980; Kobayashi et al. 2004; Guan et al. 2008). The *L. casei* SAT showed affinity for both substrates within the same order of magnitude (K_M of 1.1 mM for serine and 0.02 mM for acetyl-CoA).

According to Johnson, Roderick and Cook (2005), acyltransferases can be divided into hexapeptide and non-hexapeptide proteins. This classification is based on the presence or absence of imperfect tandem repeats of a hexapeptide sequence. While MetA from *E. coli* and a homoserine acetyltransferase from *H. influenzae* belong to the non-hexapeptide acyltransferases, SATs generally possess this structural motif that can be found using InterProScan (IPR001451). We searched for these hexapeptide sequences in the *L. casei* SAT protein but did not detect this structural motif (data not shown). Therefore, the *L. casei* SAT is, to our knowledge, the first SAT to be described as a non-hexapeptide acyltransferase.

With the advent of next-generation sequencing, the amount of assembled and annotated genome data of microbes is rapidly accumulating. Currently, the annotation of genomes is mainly based on sequence and structural comparisons. This study shows that functional confirmations are very valuable, especially when metabolic pathways that are predicted from genome data seem to be incomplete.

SUPPLEMENTARY DATA

Supplementary data are available at FEMSLE online.

Conflict of interest. None declared.

REFERENCES

- Baecker PA, Wedding RT. Purification of serine acetyltransferase, a component of a multienzyme complex, by immunoabsorption and selective dissociation of the complex. *Anal Biochem* 1980;**102**:16–21.
- Beresford T, Williams A. The microbiology of cheese ripening. In: Fox PF, McSweeney PLH, Cogan TM et al. (eds). *Cheese: Chemistry, Physics and Microbiology*, Vol. 1. London: Elsevier Academic Press, 2004, 287–317.
- Bogicevic B, Berthoud H, Portmann R et al. CysK from *Lactobacillus casei* encodes a protein with O-acetylserine sulphydrylase and cysteine desulfurization activity. *Appl Microbiol Biot* 2012;**94**:1209–20.
- Born T, Blanchard JS. Enzyme-catalyzed acylation of homoserine: mechanistic characterization of the *Escherichia coli metA*-encoded homoserine transsuccinylase. *Biochemistry* 1999;**38**:14416–23.
- Burnell JN, Whatley FR. Sulphur metabolism in *Paracoccus denitrificans*. Purification, properties and regulation of serine transacetylase, O-acetylserine sulphydrylase and β -cystathionase. *Biochim Biophys Acta* 1977;**481**:246–65.
- Christensen JE, Steele JL. Impaired growth rates in milk of *Lactobacillus helveticus* peptidase mutants can be overcome by use of amino acid supplements. *J Bacteriol* 2003;**185**:3297–306.
- de Man JC, Rogosa M, Sharpe ME. A medium for the cultivation of lactobacilli. *J Appl Bacteriol* 1960;**23**:130–5.
- Farrell HM, Jimenez-Flores R, Bleck GT et al. Nomenclature of the proteins of cows' milk—sixth revision. *J Dairy Sci* 2004;**87**:1641–74.
- Guan R, Roderick SL, Huang B et al. Roles of histidines 154 and 189 and aspartate 139 in the active site of serine acetyltransferase from *Haemophilus influenzae*. *Biochemistry* 2008;**47**:6322–8.
- Hendrickson WA, Horton JR, Lemaster DM. Selenomethionyl proteins produced for analysis by multiwavelength anomalous diffraction (MAD): a vehicle for direct determination of three-dimensional structure. *EMBO J* 1990;**9**:1665–72.
- Irmeler S, Raboud S, Beisert B et al. Cloning and characterization of two *Lactobacillus casei* genes encoding a cystathionine lyase. *Appl Environ Microb* 2008;**74**:99–106.
- Irmeler S, Schaefer H, Beisert B et al. Identification and characterization of a strain-dependent cystathionine β/γ -lyase in *Lactobacillus casei* potentially involved in cysteine biosynthesis. *FEMS Microbiol Lett* 2009;**295**:67–76.
- Johnson CM, Roderick SL, Cook PF. The serine acetyltransferase reaction: acetyl transfer from an acylpantothienyl donor to an alcohol. *Arch Biochem Biophys* 2005;**433**:85–95.
- Jones-Mortimer MC. Positive control of sulphate reduction in *Escherichia coli*. Isolation, characterization and mapping of cysteineless mutants of *E. coli* K 12. *Biochem J* 1968;**110**:589–95.
- Kobayashi S, Masui R, Yokoyama S et al. A novel metal-activated L-serine O-acetyltransferase from *Thermus thermophilus* HB8. *J Biochem* 2004;**136**:629–34.
- Kredich NM. Biosynthesis of cysteine. In: Neidhardt FC, Curtiss IR, Ingraham JL et al. (eds). *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Washington, DC: ASM Press, 1996, 514–27.
- Kredich NM, Tomkins GM. The enzymic synthesis of L-cysteine in *Escherichia coli* and *Salmonella typhimurium*. *J Biol Chem* 1966;**241**:4955–65.
- Landaud S, Helinck S, Bonnarme P. Formation of volatile sulfur compounds and metabolism of methionine and other sulfur compounds in fermented food. *Appl Microbiol Biot* 2008;**77**:1191–205.
- Liu M, Nauta A, Francke C et al. Comparative genomics of enzymes in flavor-forming pathways from amino acids in lactic acid bacteria. *Appl Environ Microb* 2008;**74**:4590–600.
- Liu M, Prakash C, Nauta A et al. Computational analysis of cysteine and methionine metabolism and its regulation in dairy starter and related bacteria. *J Bacteriol* 2012;**194**:3522–33.
- Makarova K, Slesarev A, Wolf Y et al. Comparative genomics of the lactic acid bacteria. *P Natl Acad Sci USA* 2006;**103**:15611–6.
- Sambrook J, Fritsch EF, Maniatis T. *Molecular Cloning: A Laboratory Manual*. New York: Cold Spring Harbor Laboratory Press, 1989.
- Siezen RJ, van Enkevort FH, Kleerebezem M et al. Genome data mining of lactic acid bacteria: the impact of bioinformatics. *Curr Opin Biotech* 2004;**15**:105–15.