

COMMENTARY

Health information technology as a learning health system: Call for a national monitoring system

Tiago K. Colicchio¹  | Guilherme Del Fiol² | James J. Cimino¹¹Informatics Institute, University of Alabama at Birmingham, Birmingham, Alabama²Department of Biomedical Informatics, University of Utah, Salt Lake City, Utah**Correspondence**

Tiago K. Colicchio, Informatics Institute, University of Alabama at Birmingham, Birmingham, AL 35294.

Email: tcolicchio@uabmc.edu

Abstract

After over half a century of computer application development in medicine, the US health system has gone digital with an enthusiastic confidence for rapid improvements in care outcomes, especially those of quality of care, safety, and productivity. The bad news is that evidence for the justification of the hype around health information technology (HIT) is conflicting, and the expected benefits of a digital health system have not yet materialized. We propose a national system for monitoring HIT impact based on the paradigm of the learning health system (LHS): learning from practical experience through high-quality, ongoing monitoring of care outcomes. Our proposal aims at leveraging current de facto standard research data repositories used to support large-scale clinical studies by incorporating data needed for more robust HIT assessments and application of rigorous research designs that are now feasible on a large scale.

KEYWORDS

adoption, electronic health records, learning health systems, medical informatics applications, outcome assessment

1 | HEALTH INFORMATION TECHNOLOGY: EXPECTATIONS AND REALITY

After over half a century of computer application development in medicine, the US health system has gone digital. Most health care settings across the country have now adopted commercial electronic health record (EHR) systems with an enthusiastic confidence for rapid improvements in care outcomes, especially those of quality (eg, mortality and readmissions), safety (eg, medication errors and hospital-acquired infections), and productivity (eg, care cost and provider efficiency). By 2009, EHRs had been adopted by 12% of US hospitals; today, adoption is above 90%.¹ The bad news is that evidence for the justification of the hype around health information technology (HIT) is conflicting, and previous studies that have reported predominantly positive results are now being criticized for relying on weak research methods.^{2,3} Evidence of the safety risks of

HIT are starting to accumulate⁴ along with the need to finding definitive solutions, still resisted by current systems.⁵ Four years after going digital, the US health system is still the most expensive in the world and lags behind other developed countries in some important quality outcomes.⁶

HIT evaluations in which the EHR or its components were treated as interventions have served as the basis for the Meaningful Use (MU) program. However, most were single site studies, focused on home-grown systems that have been discontinued and, in most cases, the studies provided putative, non-generalizable outcomes.⁷ It is not surprising, then, that the expected benefits of a digitized health system have not yet materialized. However, as a result of nationwide adoption of a relatively small set of commercial EHRs, a large digital infrastructure is now available and can be leveraged to support more robust studies that were not feasible in the pre-MU era. In this paper, we discuss the potential benefits of a national monitoring system of HIT impact to

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Learning Health Systems* published by Wiley Periodicals, Inc. on behalf of the University of Michigan

facilitate comparison of consensus outcomes across large, diverse sets of health care organizations implementing similar HIT tools “in the wild.” Hopefully, such a monitoring system would increase our understanding of the full impact of HIT interventions on care outcomes.

2 | CURRENT STATE OF THE EVIDENCE PRODUCED BY PREVIOUS HIT EVALUATIONS

Care delivery in the United States is provided in fast-paced, distributed care settings under constant adaptation to ever increasing medical knowledge. In such environments, implementation of a newly adopted EHR or the new version of an installed EHR—a complex process that can last several months—will inevitably add to the complexity of several aspects of care. These implementations tend to be ongoing sociotechnical processes, in which improved or new functionality is continuously implemented and system errors are constantly addressed. Such a cycle has no end since people continuously adapt to the technology and vice versa in order to optimize their work,⁸ which results in users constantly facing a learning curve that can last from months to years.⁹ Despite the complex and ever changing environment in which HIT is implemented, most HIT evaluations assess their interventions with simple short-term pretest-posttest designs.¹⁰ Such methods are not capable of detecting time-sensitive effects¹¹ that are common to HIT interventions.¹² Due to the heterogeneity of systems used in the pre-MU era and the challenges associated with data collection across multiple organizations, most studies are site-specific and use a small set of nonconsensus measurements.^{3,10} The use of agreed-upon outcome measures that are shared among researchers is paramount to allow reproducibility of studies¹³ and the comparison of outcomes across studies of different sites by systematic reviews and meta-analyses.³ Furthermore, lessons learned from other service sectors demonstrate that IT adoption rarely produces positive results if not accompanied by complementary factors or investments.¹⁴ Despite these lessons, HIT evaluations rarely consider potential context-dependent factors that can affect their outcomes of interest.¹⁰

Finally, due to the high cost and complexity involved in EHR implementations and their ongoing maintenance, the definition of the implementation strategy, timeline, and settings to be implemented is naturally business-driven decisions; in such cases, opportunities for randomization of intervention and control settings are limited, confirmatory studies are also difficult to conduct, and so a more practical approach to assess HIT impact on care outcomes is needed.

As large-scale EHR implementations and their ongoing effects become ubiquitous, a robust national monitoring system can provide data needed to support future studies capable of comparing consensus outcomes across multiple settings using the same EHR system, generating a more agile, practice-based learning process.

3 | HIT IMPACT AND THE LEARNING HEALTH SYSTEM

The increasing evidence of medical errors and dependence on evidence-based guidelines generated by artificially controlled research have contributed to establishing a new vision for improving US health care, referred to as the “learning health system” (LHS).¹⁵ The LHS proposes a paradigm shift in which medical knowledge would be primarily acquired from practical experience in naturalistic settings. By accessing large data sets and applying novel evaluations, known as “practice-based evidence methods,” findings from traditional, expensive, and artificially controlled research methods such as randomized controlled trials can be augmented to develop more generalizable guidelines.

Understanding of the full impact of HIT interventions on care outcomes will require a paradigm analogous to the LHS: learning from practical experience through high-quality, ongoing monitoring and pragmatic studies. Comprehensive quasi-experimental methods for ongoing monitoring of HIT interventions¹⁶ and several pragmatic methods applicable to HIT research questions have been proposed recently.¹⁷ However, due to the constraints of large-scale studies, they have mostly been applied to single-site evaluations. By leveraging the digital infrastructure of the post-MU era, a comprehensive database of consensus outcomes could be created, which could then be studied on a scale larger than ever before using methods such as interrupted time-series designs with multiple control sites (for detection of ongoing effects) or pragmatic designs such as multisite clustered trials and stepped-wedge designs (in which the EHR or an EHR's component is the intervention).

As necessary improvements are added to current EHRs and users adapt either to the multifunctional system or to isolated components added over time, continuous, near real-time monitoring of HIT impact will be necessary to leverage future research. Care measures more likely to detect the impact of HIT interventions have been identified in the literature³ and combined with interviews and national surveys with subject-matter experts.¹⁸ These measures assess care outcomes of quality, productivity, and safety and have already been tested in a large-scale EHR implementation for identification of changes introduced immediately after the EHR go live or with long lasting effects.^{12,16} That evaluation detected a significant impact attributable to the commercial EHR implementation in 40 (98%) measures monitored.¹⁶ An example includes a significant increase in the emergency department length of stay of four hospitals; the effect was observed only in the intervention sites and lasted from 10 to 15 months. A retrospective assessment of the same implementation revealed several internal and external context-dependent factors that may have affected the longitudinally monitored outcomes, attesting to the fact that a full understanding of HIT impact will likely be reached only when such factors are considered.¹⁹ Several factors with the potential to affect care outcomes during HIT interventions can be quantitatively monitored with data readily available in electronic format such as provider-patient ratio, changes to health insurance coverage, intentional changes to the volume patients, or seasonal variations. If such

variables are collected from multiple organizations along with quality, safety and productivity outcomes, they can be analyzed as covariates or confounders to account for context-dependent factors recognized as paramount for HIT appraisals,¹⁷ but rarely reported in the literature.¹⁰

Challenges related to technical aspects of data access and concerns about privacy and publicly available sensitive data will have to be managed for monitoring to occur beyond individual institutions. However, several measures that will likely be monitored may already be publicly available through benchmarking databases such the Centers for Medicare and Medicaid Services (CMS) Hospital Compare or the Healthcare Effectiveness Data and Information Set (HEDIS).¹⁸ These reporting systems include multiple quality and safety outcomes capable of detecting HIT impact^{12,16} and could be combined with other measures from the National Quality Forum HIT safety measurements.²⁰ To minimize the safety risks of HIT on a national level, an EHR oversight program²¹ and patient safety goals²² will need to be included explicitly as part of a national monitoring agenda. To capture data, one possible approach is to use de facto standard research data repositories like the Observational Health Data Sciences and Informatics (OHDSI)²³ or the National Patient-Centered Clinical Research Network (PCORnet).²⁴ For example, PCORnet currently incorporates 13 clinical data research networks (CDRNs) that routinely collect care outcomes data (mostly quality and safety) from multiple organizations across the country to facilitate large-scale clinical studies. To form the platform for a national monitoring system of HIT impact, such infrastructure could be leveraged to incorporate data that are not currently captured by CDRNs, but would be needed for monitoring HIT interventions, such as productivity-related outcomes (eg, volume of visits, orders and hospitalizations), information about planned HIT interventions (eg, go live date of a new EHR component or a new EHR version), or data needed to monitor context-dependent variables (eg, local insurance/market changes, seasonal variations, and provider-patient ratio).¹⁹ The latter could be identified with guidance from implementation science frameworks such as the Consolidated Framework for Implementation Research (CFIR),²⁵ which covers context domains such as inner and outer setting and individual characteristics that may influence the effect of HIT interventions. Another challenge is the business nature of most HIT interventions, which may hamper the possibility of randomization in the case of pragmatic studies. Since most integrated care delivery systems now use similar commercial EHRs, one approach would be to form groups of clients and EHRs and organize a shared plan of implementation of new versions or system components, in order to orchestrate the randomization of these interventions.

Imagine a resource with data for several outcomes collected from multiple institutions on a regular basis or in real time depending on data availability. A dashboard built on top of such a resource would allow monitoring with statistical evaluation of longitudinal change patterns introduced by the implementation of a new EHR, a new EHR version, a new EHR component (eg, clinical decision support), outside applications, or system customization requests in a specific setting or group of settings spread across the nation. Each of these can be assessed as an individual intervention for identification of effects

observed immediately after its introduction or effects observed over longer periods. For example, if modifications to a function used by infection prevention specialists decreases the specialists' efficiency, their capacity to investigate suspected infections may be hampered, resulting in an unexpected apparent decrease in the rate of hospital-acquired infections. Studies using stepped-wedge designs²⁶ could be applied on subsequent implementations of the same product in different clients, confirming (or ruling out) a potential safety risk introduced by a new EHR version (infection rate could have decreased as a result of fewer cases investigated and not as a result of a real decrease in the number of infections). These effects could be shared with other institutions planning on implementing the same product or product component. Organizations not yet participating will pay their own way to join so that they can compare their systems with national experience and be informed about what to expect when they implement systems previously tested elsewhere, anticipating the need for organizational changes to prevent negative effects or to maximize positive ones. As mentioned above and demonstrated by previous HIT evaluations,^{12,16} several measures that will likely be relevant to enable the proposed monitoring system are already captured on a regular basis by most care delivery systems across the country, and so the cost to develop the data feed to populate repositories such as PCORnet and OHDSI is likely to be manageable. Such a cost will surely be paid back many times over potentially improving provider satisfaction and productivity, as well as preventing quality and safety hazards.

Our approach will further LHS goals by allowing accumulation of evidence of HIT impact from practical experience to reach the larger medical and informatics communities in near real time, as opposed to lengthy site-specific evaluations that, in addition to being less generalizable than what was desired, will be published long after their maximum usefulness, if they are published at all. The proposed monitoring system has the potential to lead us to a better understanding of the full impact of HIT on care outcomes and to avoid wasted time and money on another cycle of hundreds of studies that may not provide any valuable knowledge.

ACKNOWLEDGEMENT

This work was supported by research funds from the Informatics Institute of the University of Alabama at Birmingham.

CONFLICTS OF INTEREST

The authors have no competing interests to declare.

ORCID

Tiago K. Colicchio  <https://orcid.org/0000-0003-3405-1538>

REFERENCES

1. The Office of the National Coordinator for Health Information Technology [Internet]. Health IT Dashboard; [cited 2018 July 27]. Accessed from: <https://dashboard.healthit.gov/quickstats/quickstats.php>.
2. Rudin RS, Jones SS, Shekelle P, Hillestad RJ, Keeler EB. The value of health information technology: filling the knowledge gap. *Am J Manag Care*. 2020;4 [11 Spec No. 17]:eSP1-eSP8.

3. Colicchio TK, Facelli JC, Del Fiol G, Scammon DL, Bowes WA III, Narus SP. Health information technology adoption: understanding research protocols and outcome measurements for IT interventions in health care. *J Biomed Inform.* 2016;63:33-44. PMID: 27450990
4. Kim MO, Coiera E, Magrabi F. Problems with health information technology and their effects on care delivery and patient outcomes: a systematic review. *J Am Med Inform Assoc.* 2017;24(2):246-250.
5. Hingle S. Electronic health records: an unfulfilled promise and a call to action. *Ann Intern Med.* 2016;165(11):818-819.
6. Papanicolaos I, Woskie LR, Jha AK. Health care spending in the United States and other high-income countries. *JAMA.* 2018;319(10):1024-1039. PMID: 29536101
7. Colicchio TK, Cimino JJ, Del Fiol G. Unintended consequences of nationwide electronic health record adoption: challenges and opportunities in the post-meaningful use era. *J Med Internet Res.* 2019;21(6):e13313.
8. Harrison MI, Koppel R, Bar-Lev S. Unintended consequences of information technologies in health care—an interactive sociotechnical analysis. *J Am Med Inform Assoc.* 2007;14(5):542-549.
9. Abramson EL, Malhotra S, Osorio SN, et al. A long-term follow-up evaluation of electronic health record prescribing safety. *J Am Med Inform Assoc.* 2013;20(e1):e52-e58.
10. Jones SS, Rudin RS, Perry T, Shekelle PG. Health information technology: an updated systematic review with a focus on meaningful use. *Ann Intern Med.* 2014;160(1):48-54.
11. Lagarde M. How to do (or not to do) ... Assessing the impact of a policy change with routine longitudinal data. *Health Policy Plan.* 2012;27(1):76-83. PMID: 21278077
12. Colicchio TK, Fiol GD, Stoddard GJ, Narus SP. Evaluation of a systematic methodology to detect in near real-time performance changes during electronic health record system implementations: a longitudinal study. *AMIA Annu Symp Proc.* 2018;2017:595-604.
13. Ioannidis JPA. Why most published research findings are false. *PLoS Med.* 2005;2(8):e124. <https://doi.org/10.1371/journal.pmed.0020124>
14. Brynjolfsson E, Hitt LM. Beyond the productivity paradox. *Commun ACM.* 1998;41(8):49-55.
15. Olsen L, Aisner D, McGinnis JM. *Learning Healthcare System Workshop Summary, IOM Roundtable on Evidence-Based Medicine.* Washington: National Academies Press; 2007 [cited 2018 July 27]. Accessed from: <https://www.ncbi.nlm.nih.gov/books/NBK53494>.
16. Colicchio TK, Del Fiol G, Scammon DL, Facelli JC, Bowes WA, Narus SP. Comprehensive methodology to monitor longitudinal change patterns during EHR implementations: a case study at a large health care delivery network. *J Biomed Inform.* 2018;83:40-53.
17. Murray E, Hekler EB, Andersson G, et al. Evaluating digital health interventions: key questions and approaches. *Am J Prev Med.* 2016;51(5):843-851.
18. Colicchio TK, Fiol GD, Scammon DL, Bowes WA III, Facelli JC, Narus SP. Development and classification of a robust inventory of near real-time outcome measurements for assessing information technology interventions in health care. *J Biomed Inform.* 2017;73:62-75.
19. Colicchio TK, Borbolla D, Colicchio VD, et al. Looking behind the curtain: identifying factor affecting care outcomes during a large commercial EHR implementation. *eGEMs.* 2019;7(1):1-12.
20. National Quality Forum. Prioritization and Identification of Health IT Patient Safety Measures [Internet]. [Cited 2019 October 22]. Accessed from: <http://www.qualityforum.org/ProjectDescription.aspx?projectId=77689/>.
21. Singh H, Classen DC, Sittig DF. Creating an oversight infrastructure for electronic health record-related patient safety hazards. *J Patient Saf.* 2011;7(4):169-174.
22. Sittig DF, Classen DC, Singh H. Patient safety goals for the proposed Federal Health Information Technology Safety Center. *J Am Med Inform Assoc.* 2015;22(2):472-478.
23. Observational Health Data Sciences and Informatics - OHSDI [Internet]. [Cited 2018 August 20]. Accessed from: <https://www.ohdsi.org/>.
24. The National Patient-Centered Clinical Research Network - PCORnet [Internet]. [Cited 2018 August 20]. Accessed from: <https://pcor.net.org/>.
25. Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci.* 2009;4(1):50.
26. Westbrook JI, Li L, Raban MZ, et al. Stepped-wedge cluster randomised controlled trial to assess the effectiveness of an electronic medication management system to reduce medication errors, adverse drug events and average length of stay at two paediatric hospitals: a study protocol. *BMJ Open.* 2016;6(10):e011811. <https://doi.org/10.1136/bmjopen-2016-011811>