



Contents lists available at ScienceDirect

## Journal of Hand Surgery Global Online

journal homepage: [www.JHSGO.org](http://www.JHSGO.org)



### Original Research

## Evaluating the Impact of Social Deprivation on Mid-Term Outcomes Following Distal Radius Open Reduction Internal Fixation

Michelle Zeidan, MD, \* Andrew R. Stephens, BS, \* Chong Zhang, MS, † Angela P. Presson, PhD, †  
Nikolas H. Kazmers, MD, MSE \*

\* Department of Orthopaedics, University of Utah, Salt Lake City, UT

† Division of Public Health, University of Utah, Salt Lake City, UT



### ARTICLE INFO

#### Article history:

Received for publication March 28, 2021

Accepted in revised form June 1, 2021

Available online July 3, 2021

#### Key words:

Area deprivation index  
Biopsychosocial  
Distal radius fracture  
QuickDASH  
Social deprivation

**Purpose:** Although baseline biopsychosocial factors are associated with short-term patient-reported outcomes following distal radius fracture open reduction internal fixation (ORIF), their effect on mid-term outcomes is unclear. We aimed to evaluate the effect of social deprivation, previously established as a surrogate for depression, pain interference, and anxiety, on quick Disabilities of the Arm, Shoulder and Hand (QuickDASH) scores >1 year following distal radius ORIF.

**Methods:** Adult patients ( $\geq 18$  years) with an isolated distal radius fracture treated with ORIF by orthopedic trauma and hand surgeons at a single tertiary academic center over a 3-year period were included. Outcomes at  $\geq 1$  year postoperatively were evaluated using QuickDASH. Age, follow-up duration, area deprivation index (ADI; a measure of social deprivation), subspecialty (hand vs trauma), and postoperative alignment were assessed using linear regression with 95% confidence intervals after bootstrapping and a permutation test for P values to test for their association with the final QuickDASH score.

**Results:** Follow-up data were obtained for 98 of 220 (44.5%) patients at a mean of  $3.1 \pm 1.0$  years after surgery. Mean age and ADI were  $53.2 \pm 15.4$  years and  $26.8 \pm 18.7$ , respectively. Most fractures were intra-articular (67.3%), and 72.4% had acceptable postoperative alignment parameters, as defined by the American Academy of Orthopaedic Surgeons clinical practice guidelines. The mean QuickDASH score was  $13.0 \pm 16.5$ . There were no significant associations between the final QuickDASH score and any studied factor, including ADI, as determined using univariable analysis. Multivariable analysis showed no association between ADI and the final QuickDASH score, independent of age, sex, treating service, follow-up duration, and fracture alignment or pattern.

**Conclusions:** At mid-term follow up after distal radius ORIF, ADI did not correlate with QuickDASH scores, and the QuickDASH scores of the patients did not differ from those of the general population. However, our cohort mostly comprised patients with levels of deprivation below the national median. Although studies have shown that the short-term outcomes of distal radius ORIF are influenced by biopsychosocial factors, outcomes at the time of final recovery may not be associated with social deprivation.

**Type of study/level of evidence:** Prognostic IV.

Copyright © 2021, THE AUTHORS. Published by Elsevier Inc. on behalf of The American Society for Surgery of the Hand.  
This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Distal radius fractures represent 1 of the most commonly treated skeletal injuries.<sup>1</sup> Over time, the decision to treat these fractures with open reduction internal fixation (ORIF), rather than with casting, has been increasing.<sup>2,3</sup> Much of the research regarding the

outcomes of distal radius fracture ORIF has focused on postoperative complications and the restoration of anatomic alignment.<sup>1,4–6</sup>

Recent research, however, has focused on biopsychosocial factors that affect the outcomes of upper- and lower-extremity surgery.<sup>7–13</sup> More specifically, evidence has suggested that biopsychosocial factors such as depression, pain interference, catastrophizing, education level, and income level are associated with worse short-term ( $\leq 1$  year) outcomes following distal radius fractures.<sup>7,14–16</sup> Studies have shown that these factors and higher levels of social deprivation are also associated with lower levels of

**Declaration of interests:** No benefits in any form have been received or will be received related directly or indirectly to the subject of this article.

**Corresponding author:** Nikolas H. Kazmers, MD, MSE, Department of Orthopaedics, University of Utah, 590 Wakara Way, Salt Lake City, UT 84108.

**E-mail address:** [nkazmers@gmail.com](mailto:nkazmers@gmail.com) (N.H. Kazmers).

<https://doi.org/10.1016/j.jhsg.2021.06.002>

2589-5141/Copyright © 2021, THE AUTHORS. Published by Elsevier Inc. on behalf of The American Society for Surgery of the Hand. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

physical function.<sup>17–20</sup> However, the effect of biopsychosocial factors and social deprivation on mid- and long-term outcomes remains less clear. Because the management of distal radius fractures contributes a substantial amount of burden to the health care system, it is critical to understand which factors impact the outcomes.<sup>21,22</sup>

The purpose of this study was to evaluate the effect of social deprivation on patient-reported outcomes more than 1 year following distal radius ORIF. Specifically, we hypothesized that increased levels of social deprivation are associated with worse outcomes, determined using quick Disabilities of the Arm, Shoulder and Hand (QuickDASH), following distal radius ORIF.

## Materials and Methods

This study was approved by our institutional review board. Adult patients ( $\geq 18$  years of age) who sustained a distal radius fracture treated with ORIF by orthopedic trauma and hand surgeons between January 2014 and March 2019 were identified using current procedural terminology codes (25607, 25608, and 25609). The patients were excluded if they did not have radiographs at least 5 weeks after ORIF, sustained additional upper-extremity injuries at the same time as the distal radius fracture, were less than 18 years of age, had been treated nonsurgically, or had undergone the surgery less than 1 year prior.

QuickDASH was used as a measure of functional outcome in the patients. QuickDASH is an abbreviated version of the Disabilities of the Arm, Shoulder and Hand questionnaire, a valid measure of physical function in patients with upper-limb musculoskeletal disorders. QuickDASH is a shorter, 11-item questionnaire that has been shown to have the same validity and reliability as the Disabilities of the Arm, Shoulder and Hand questionnaire, with lesser burden on the patient.<sup>23,24</sup>

Age, sex, follow-up duration, subspecialty (hand vs trauma), intra-articular versus extra-articular fractures, and postoperative alignment parameter data were collected via a chart review. The postoperative alignment parameter data collected included radial height, ulnar variance, and volar tilt. Acceptable postoperative parameters were defined by the American Academy of Orthopaedic Surgeons (AAOS) criteria of a radial shortening of  $<3$  mm, a dorsal tilt of  $<10^\circ$ , or an intra-articular stepoff of  $<2$  mm.<sup>25</sup> Radial height outside of the AAOS criteria was defined as less than 8.6 mm, based on the average population's radial height.<sup>26</sup> The methods of measurement were based on those described by Mulders et al.<sup>27</sup> Patients who met the inclusion criteria were notified via email that they would be receiving a request to complete a QuickDASH questionnaire via email or phone call, and if they did not want to be contacted, they could notify the study investigators or decline the request. For responders, REDCap was used to collect QuickDASH scores at the time of final follow up.

The degree of social deprivation was measured using each patient's area deprivation index (ADI) (higher ADI represents more social deprivation; the possible range of scores span from 1 to 100). The ADI was developed by the Health Resources and Services Administration as a measure of social deprivation in a specific area based on 17 United States census-based factors, including education level, employment status, housing quality, the cost of living, and poverty level.<sup>28,29</sup> A patient's zip code is used to identify their ADI score. The ADI value used is reported as the national percentile. Based on data from a previous study, we expected the ADI to be associated with QuickDASH scores, where a 1-point increase in the ADI was associated with a 0.53-point (95% confidence interval: 0.26–0.80) increase in the QuickDASH score.<sup>30</sup> For the current study with 98 subjects, we had 80% power at a 0.05 significance

**Table 1**  
Patient Demographic and Baseline Characteristics

Variable	Value (n = 98)
Age (y)	53.2 (15.4)
Mean (SD)	
Sex	
Male	22 (22.4%)
Female	76 (77.6%)
Race	
Caucasian	94 (95.9%)
Hispanic	3 (3.1%)
Asian	1 (1%)
Area deprivation index	
Mean (SD)	26.8 (18.7)
Quartile 1 (range 1–25)	54 (55.1%)
Quartile 2 (range 25–50)	36 (36.7%)
Quartile 3 (range 51–75)	4 (4.1%)
Quartile 4 (range 76–100)	4 (4.1%)
Treating orthopedic subspecialty	
Hand	78 (79.6%)
Trauma	20 (20.4%)
Follow-up duration (y)	
Mean (SD)	3.1 (1.0)
Range	1–98
Alignment parameters outside of AAOS criteria	
Yes	27 (27.6%)
No	71 (72.4%)
Fracture type	
Extra-articular	32 (32.7%)
Intra-articular	66 (67.4%)

level to detect an effect size of 0.089 or greater in a multivariable regression model adjusting for 5 additional variables.

Associations between variables of interest and the final QuickDASH score were analyzed using univariable and multivariable linear regression analyses. Because the QuickDASH score distribution exhibited a notable right skew with a minimum value of 0, we used bias-corrected and accelerated 95% bootstrap confidence intervals and permutation test *P* values estimated from 3,000 samples. To obtain *P* values, on each iteration, we permuted QuickDASH values across the subjects, and the *P* values were estimated as the proportion of coefficients that were as extreme as or more extreme than the coefficients estimated for the original data. Regression coefficients were reported as mean changes in QuickDASH scores. The analyses were conducted using R, version 4.0 (R Core Team [2020]. R: A language and environment for statistical computing. R Foundation for Statistical Computing. <https://www.R-project.org/>). *P* values  $<.05$  were deemed significant. All the tests were 2-sided.

## Results

A total of 371 patients were identified using the current procedural terminology code query, and 220 met the inclusion criteria. Follow-up QuickDASH scores at least 1 year after surgery were obtained for 98 of 220 (44.5%) patients at a mean of  $3.1 \pm 1.0$  years after surgery. The mean age was  $53.2 \pm 15.4$  years, and 77.6% (76/98) of the patients were women (Table 1). Most fractures were intra-articular (67.3%), and 72.4% had acceptable postoperative AAOS alignment parameters. For patients with alignment parameters outside of the AAOS criteria, the majority had unacceptable radial height (23/27, 85.2%), whereas 2 (7.4%) patients had a dorsal angulation of  $>10^\circ$ , and 3 (11.1%) patients had an intra-articular stepoff of  $>2$  mm. The mean ADI percentile was  $26.8 \pm 18.7$  (interquartile range: 14.0–37.8), representing the bottom of the second quartile, and the mean QuickDASH score was  $13.0 \pm 16.5$ . The average QuickDASH score of patients with alignment parameters within the AAOS criteria was  $14.4 \pm 17.9$ , whereas the average QuickDASH score of patients with alignment parameters outside of

**Table 2**

Distal Radius Alignment Parameters for Patients With Parameters Within or Outside of AAOS Criteria

Value	All Patients	Alignment Parameters Within AAOS Criteria (n = 71)	Alignment Parameters Outside of AAOS Criteria (n = 27)
Radial height (mm)	10.1 ± 2.0	10.9 ± 1.7	8.1 ± 1.1
Volar angulation (°)	5.2 ± 6.5	5.9 ± 5.6	3.6 ± 8.5
Intra-articular stepoff (mm)	0.2 ± 0.6	0.1 ± 0.4	0.4 ± 0.9

the AAOS criteria was  $9.3 \pm 11.7$  ( $P > .05$ ). Most patients (95/98 or 96.9%) were treated with volar plate fixation, whereas the remaining patients were treated with a dorsal spanning plate. Table 2 presents data regarding the radial height, angulation, and stepoff for each group's AAOS alignment parameters.

In the univariable analysis (Table 3), no significant association was found between the final QuickDASH score and any studied factor, including ADI, final fracture alignment, type of fracture (intra- vs extra-articular), treating service, sex, or age. Similarly, in the multivariable analysis (Table 3), no association was found between ADI and the final QuickDASH score, independent of age, sex, treating service, fracture type, final fracture alignment, or follow-up duration.

## Discussion

The main finding of this study is that at an average of 3.1 years after distal radius ORIF in a cohort with an average ADI of  $26.8 \pm 18.7$  (interquartile range 14.0–37.8), the outcomes are not affected by social deprivation, as measured using the ADI. The ADI serves as a surrogate for a variety of biopsychosocial factors, with the prior demonstration of a significant correlation with depression, anxiety, and pain interference.<sup>20</sup> Furthermore, we found no association between the final QuickDASH score and any of the variables investigated, including age, fracture type, final alignment parameters, or sex.

There has been increasing focus on biopsychosocial factors that are associated with the patient-reported outcomes of upper- and lower-extremity surgeries.<sup>7–13</sup> In a recent study, Jayakumar et al suggested that factors such as depression, pain interference, and pain catastrophizing, which are associated with higher levels of social deprivation, are associated with worse early outcomes following distal radius fractures.<sup>7,17–20</sup> Notably, this study was specific to adult patients with distal radius fractures, although only 16% of the enrolled patients had undergone ORIF. With the caveat that the study population differed from that of the current study, the use of opioids and greater pain interference or pain catastrophizing within 1 week of injury were predictive of the results at 6–9 months after surgery. This must be interpreted in light of their evaluation of the early postoperative phase because other studies have suggested that continued recovery after distal radius fractures can be expected for 1 or more years after surgery.<sup>31–35</sup>

In addition, in a prospective study, Chung et al<sup>15</sup> found that in 66 patients, worse outcomes 1 year after distal radius ORIF, as measured using the Michigan Hand Outcome Questionnaire, were associated with increased age and lower income. In addition, there was no association with sex, fracture classification, radiographic parameters, or workers' compensation status. Additionally, in another prospective study, MacDermid et al<sup>36</sup> found that in 120 patients sustaining a distal radius fracture (with varying treatment options), injury compensation and lower levels of education were associated with worse outcomes at 6 months after fracture, as

measured using the patient-rated wrist evaluation. It should be noted that the ADI takes into account an area's average income and education level, and our results would argue that these factors might play a role in the short-term outcomes of distal radius fractures but may not actually have an effect on their mid- or long-term outcomes. In fact, in the current study, it was found that at a mean of 3.1 years following ORIF, the QuickDASH scores were actually similar to those of the general population.<sup>37</sup> This suggests that provided a long enough follow-up period, patients undergoing ORIF for a distal radius fracture function similar to the general population, regardless of social deprivation and other biopsychosocial factors that may be impactful in the early recovery phase.<sup>7</sup>

We also found that regardless of whether the patients' final radiographic parameters were within the AAOS criteria or not, there was no clinically relevant or statistically significant difference between each group's QuickDASH score.<sup>38</sup> There has been a large amount of research on the radiographic parameters that are the most important to restore after a distal radius fracture. Studies have suggested that malunions involving dorsal angulation, ulnar variance, and radial height tend to have worse outcomes.<sup>39–43</sup> However, multiple studies have also suggested that clinically important differences in the outcomes are not associated with anatomic radiographic reduction parameters and that other factors may play a more important role.<sup>27,44–52</sup> This study supports that near-anatomic reduction, especially at the time of long-term follow up, may provide no clinically important additional benefit. More studies are needed to evaluate whether a poor anatomic reduction leads to long-term issues due to potential changes in carpal stability and radiocarpal degenerative changes.

We also found that age had no impact on the QuickDASH scores at the time of final follow up. Therefore, despite how young or old a patient was, their outcome was essentially the same after distal radius ORIF. Older patients are thought to place less demand on their wrist and, therefore, usually treated with the traditional method of casting and closed reduction because it is thought to have the same final outcome as operative treatment.<sup>53–55</sup> However, there has been a growing trend toward fixing distal radius fractures in older adults rather than treating them conservatively because of the advent of volar locking plates.<sup>56</sup> Although studies have shown that ORIF may provide an improvement in the outcome scores and grip strength earlier, most patients have the same outcome at 1 year or more.<sup>56–59</sup>

There are several limitations to this study. The retrospective nature of this study introduces inherent biases based on the study's design. Furthermore, QuickDASH questionnaires were sent out to patients for response, and our response rate was 44.5%; thus, there was likely a large component of responder bias that could not be accounted for. Our results may have changed with a greater follow-up rate. Although some studies may call into question the validity of QuickDASH, multiple studies in the literature have supported the use of this patient-reported outcome measure for distal radius fractures, and it has been established as a primary outcome measure.<sup>23,60–65</sup> In addition, the generalizability of these results is somewhat limited. Our sample of patients included only patients treated with ORIF, and therefore, the result does not apply to patients treated with closed reduction and casting. In addition, the median patient age was 56 years, and the mean ADI was 26.8 (relatively low levels of social deprivation); so, these results may not apply to much older or younger patients as well as to more or less socially deprived areas. We acknowledge that although this ADI may have captured a group of less disadvantaged or deprived patients, the results of this study provide at least 1 data point indicating that perhaps biopsychosocial factors play less of a role in long-term outcomes. Further studies should continue to

**Table 3**

Univariable and Multivariable Regression Analyses With Bootstrapped Confidence Intervals for Correlation With QuickDASH Score\*

Variable†	Univariable Coefficient (95% CI)	P Value	Multivariable Coefficient (95% CI)	P Value
Age	0.09 (−0.11 to 0.25)	.45	0.20 (−0.05 to 0.55)	.10
Female versus male sex	−3.0 (−17.2 to 3.7)	.46	−3.5 (−17.2 to 6.7)	.44
ADI	0.11 (−0.05 to 0.43)	.23	0.14 (−0.04 to 0.42)	.14
Trauma versus hand service	3.2 (−3.5 to 12.6)	.46	4.2 (−2.7 to 12.6)	.33
Follow-up duration	2.4 (−0.45 to 6.2)	.14	2.5 (−0.32 to 6.1)	.13
Parameters outside AAOS criteria	−5.1 (−10.9 to 0.95)	.18	−5.1 (−12.3 to 0.73)	.20
Extra-articular versus intra-articular fractures	−3.6 (−10.0 to 1.7)	.30	−3.6 (−10.4 to 1.9)	.33

CI, confidence interval.

\* Reported as mean differences.

† Independent variables. The dependent variable is the QuickDASH score.

investigate this in more socially deprived patients. Last, we used the ADI as the biopsychosocial factor of interest in this study. Although studies have shown that factors such as depression and pain interference are associated with higher levels of social deprivation and lower levels of physical function, the true measures of each patient's level of depression, anxiety, or pain interference were not data points in this study.<sup>17–20</sup>

The results of this study in a relatively narrow cohort contribute to the growing literature that focuses on the psychological and socioeconomic factors that contribute to the outcomes of hand surgery and, more specifically, distal radius fracture ORIF. Prior studies have suggested that depression, anxiety, income, employment, and education levels impact outcomes up to 1 year after a distal radius fracture.<sup>66–68</sup> However, at mid-term follow up, we found that ADI deprivation, which has previously been established as a surrogate for depression, pain interference, and anxiety, was not associated with the functional outcomes.<sup>20</sup>

## Acknowledgments

This investigation was supported by the University of Utah Study Design and Biostatistics Center, with funding in part from the National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health, through Grant UL1TR002538 (formerly 5UL1TR001067-05, 8ULTR000105, and UL1RR025764).

## References

- Levin LS, Rozell JC, Pulos N. Distal radius fractures in the elderly. *J Am Acad Orthop Surg.* 2017;25(3):179–187.
- Mellstrand-Navarro C, Pettersson HJ, Tornqvist H, Ponzer S. The operative treatment of fractures of the distal radius is increasing: results from a nationwide Swedish study. *Bone Joint J.* 2014;96(7):963–969.
- Farner S, Malkani A, Lau E, Day J, Ochoa J, Ong K. Outcomes and cost of care for patients with distal radius fractures. *Orthopedics.* 2014;37(10):e866–e878.
- Chung KC, Cho HE, Kim Y, Kim HM, Shauer MJ. Assessment of anatomic restoration of distal radius fractures among older adults. *JAMA Network Open.* 2020;3(1):e1919433.
- Nelson GN, Stepan JG, Osei DA, Calfee RP. The impact of patient activity level on wrist disability after distal radius malunion in older adults. *J Orthop Trauma.* 2015;29(4):195–200.
- Clement ND, Duckworth AD, Court-Brown CM, McQueen MM. Distal radial fractures in the superelderly: does malunion affect functional outcome? *ISRN Orthop.* Published online March 4, 2014. doi:10.1155/2014/189803
- Jayakumar P, Teunis T, Vranceanu AM, Lamb S, Ring D, Gwilym S. Early psychological and social factors explain the recovery trajectory after distal radial fracture. *J Bone Joint Surg.* 2020;102(9):788–795.
- Wolfensberger A, Vuistiner P, Konzelmann M, Plomb-Holmes C, Léger B, Luthi F. Clinician and patient-reported outcomes are associated with psychological factors in patients with chronic shoulder pain. *Clin Orthop Relat Res.* 2016;474(9):2030–2039.
- Potter MQ, Wylie JD, Greis PE, Burks RT, Tashjian RZ. Psychological distress negatively affects self-assessment of shoulder function in patients with rotator cuff tears. *Clin Orthop Relat Res.* 2014;472(12):3926–3932.
- Ayers DC, Franklin PD, Ring DC. The role of emotional health in functional outcomes after orthopaedic surgery: extending the biopsychosocial model to orthopaedics. *J Bone Joint Surg.* 2013;95(21):e165.
- Kazmers NH, Hung M, Rane AA, Bounsanga J, Weng C, Tyser AR. Association of physical function, anxiety, and pain interference in nonshoulder upper extremity patients using the PROMIS platform. *J Hand Surg.* 2017;42(10):781–787.
- Rosenberger PH, Jokl P, Ickovics J. Psychosocial factors and surgical outcomes: an evidence-based literature review. *J Am Acad Orthop Surg.* 2006;14(7):397–405.
- Mavros MN, Athanasiou S, Gkegkes ID, Polyzos KA, Peppas G, Falagas ME. Do psychological variables affect early surgical recovery? *PLoS One.* 2011;6(5):e20306.
- Jayakumar P, Overbeek CL, Lamb S, et al. What factors are associated with disability after upper extremity injuries? A systematic review. *Clin Orthop Relat Res.* 2018;476(11):2190–2215.
- Chung KC, Kotsis SV, Kim HM. Predictors of functional outcomes after surgical treatment of distal radius fractures. *J Hand Surg.* 2007;32(1):76–83.
- Roh YH, Lee BK, Noh JH, Oh JH, Gong HS, Baek GH. Effect of anxiety and catastrophic pain ideation on early recovery after surgery for distal radius fractures. *J Hand Surg.* 2014;39(11):2258–2264.
- Okoroafor UC, Gerull W, Wright M, Guattary J, Sandvall B, Calfee RP. The impact of social deprivation on pediatric PROMIS health scores after upper extremity fracture. *J Hand Surg.* 2018;43(10):897–902.
- Court-Brown CM, Aitken SA, Duckworth AD, Clement ND, McQueen MM. The relationship between social deprivation and the incidence of adult fractures. *J Bone Joint Surg.* 2013;95(6):e32.
- Menon MR, Walker JL, Court-Brown CM. The epidemiology of fractures in adolescents with reference to social deprivation. *J Bone Joint Surg.* 2008;90(11):1482–1486.
- Wright MA, Adelani M, Dy C, O'Keefe R, Calfee RP. What is the impact of social deprivation on physical and mental health in orthopaedic patients? *Clin Orthop Relat Res.* 2019;477(8):1825–1835.
- Shauer MJ, Yin H, Banerjee M, Chung KC. Current and future national costs to medicare for the treatment of distal radius fracture in the elderly. *J Hand Surg.* 2011;36(8):1282–1287.
- Kazmers NH, Judson CH, Presson AP, Xu Y, Tyser AR. Evaluation of factors driving cost variation for distal radius fracture open reduction internal fixation. *J Hand Surg.* 2018;43(7):606–614.
- Beaton DE, Wright JG, Katz JN. Development of the QuickDASH: comparison of three item-reduction approaches. *J Bone Joint Surg.* 2005;87(5):1038–1046.
- Gummesson C, Ward MM, Atroshi I. The shortened Disabilities of the Arm, Shoulder and Hand questionnaire (QuickDASH): validity and reliability based on responses within the full-length DASH. *BMC Musculoskelet Disord.* 2006;7(1):1–7.
- American Academy of Orthopaedic Surgeons. The Treatment of Distal Radius Fractures - Guideline and Evidence Report. <https://www.aaos.org/globalassets/quality-and-practice-resources/distal-radius/distal-radius-fractures-clinical-practice-guideline.pdf>. Accessed February 28, 2020.
- Medoff RJ. Essential radiographic evaluation for distal radius fractures. *Hand Clin.* 2005;21(3):279–288.
- Mulders MA, Detering R, Rikli DA, Rosenwasser MP, Goslings JC, Schep NW. Association between radiological and patient-reported outcome in adults with a displaced distal radius fracture: a systematic review and meta-analysis. *J Hand Surg.* 2018;43(8):710–719.e715.
- Kind AJH, Buckingham WR. Making neighborhood-disadvantage metrics accessible—the neighborhood atlas. *N Engl J Med.* 2018;378(26):2456–2458.
- Kind AJ, Jencks S, Brock J, et al. Neighborhood socioeconomic disadvantage and 30-day rehospitalization: a retrospective cohort study. *Ann Intern Med.* 2014;161(11):765–774.
- van der Horst A, Potter J, Stephens A, et al. Prognostic factors affecting long-term outcomes after elbow dislocation: a longitudinal cohort study. Paper presented at: American Association for Hand Surgery Annual Meeting; February 7, 2021; Virtual.
- Brogren E, Hofer M, Petranek M, Dahlin LB, Atroshi I. Fractures of the distal radius in women aged 50 to 75 years: natural course of patient-reported outcome, wrist motion and grip strength between 1 year and 2–4 years after fracture. *J Hand Surg.* 2011;36(7):568–576.

32. Shimura H, Nimura A, Fujita K, Miyamoto T. Mid-term functional outcome after volar locking plate fixation of distal radius fractures in elderly patients. *J Hand Surg.* 2018;23(2):238–242.
33. Lalone E, MacDermid J, Grewal R, King G. Patient reported pain and disability following a distal radius fracture: a prospective study. *Open Orthop J.* 2017;11: 589–599.
34. Landgren M, Teurneau V, Abramo A, Geijer M, Tägil M. Intermediate-term outcome after distal radius fracture in patients with poor outcome at 1 year: a register study with a 2- to 12-year follow-up. *J Hand Surg.* 2019;44(1):39–45.
35. Van Leerdam RH, Huizing F, Termaat F, et al. Patient-reported outcomes after a distal radius fracture in adults: a 3–4 years follow-up. *Acta Orthopaedica.* 2019;90(2):129–134.
36. MacDermid JC, Donner A, Richards RS, Roth JH. Patient versus injury factors as predictors of pain and disability six months after a distal radius fracture. *J Clin Epidemiol.* 2002;55(9):849–854.
37. Aasheim T, Finsen V. The DASH and the QuickDASH instruments. Normative values in the general population in Norway. *J Hand Surg.* 2014;39(2):140–144.
38. Kazmers NH, Qiu Y, Yoo M, Stephens AR, Tyser AR, Zhang Y. The minimal clinically important difference of the PROMIS and QuickDASH instruments in a nonshoulder hand and upper extremity patient population. *J Hand Surg.* 2020;45(5):399–407.
39. Wilcke MK, Abbaszadegan H, Adolphson PY. Patient-perceived outcome after displaced distal radius fractures: a comparison between radiological parameters, objective physical variables, and the DASH score. *J Hand Ther.* 2007;20(4): 290–298.
40. Grewal R, MacDermid JC. The risk of adverse outcomes in extra-articular distal radius fractures is increased with malalignment in patients of all ages but mitigated in older patients. *J Hand Surg.* 2007;32(7):962–970.
41. Dario P, Matteo G, Carolina C, et al. Is it really necessary to restore radial anatomic parameters after distal radius fractures? *Injury.* 2014;45(suppl 6): S21–S26.
42. Batra S, Gupta A. The effect of fracture-related factors on the functional outcome at 1 year in distal radius fractures. *Injury.* 2002;33(6):499–502.
43. Fitoussi F, Ip WY, Chow SP. Treatment of displaced intra-articular fractures of the distal end of the radius with plates. *J Bone Joint Surg.* 1997;79(9): 1303–1312.
44. Anzarut A, Johnson JA, Rowe BH, Lambert RG, Blitz S, Majumdar SR. Radiologic and patient-reported functional outcomes in an elderly cohort with conservatively treated distal radius fractures. *J Hand Surg.* 2004;29(6): 1121–1127.
45. Plant CE, Parsons NR, Costa ML. Do radiological and functional outcomes correlate for fractures of the distal radius? *Bone Joint J.* 2017;99(3):376–382.
46. Ranjeet N, Estrella EP. Distal radius fractures: does a radiologically acceptable reduction really change the result? *J Clin Diagn Res.* 2012;6(8):1388–1392.
47. Finsen V, Rod O, Rod K, Rajabi B, Alm-Paulsen PS, Russwurm H. The relationship between displacement and clinical outcome after distal radius (Colles') fracture. *J Hand Surg.* 2013;38(2):116–126.
48. Martinez-Mendez D, Lizaur-Utrilla A, de Juan-Herrero J. Prospective study of comminuted articular distal radius fractures stabilized by volar plating in the elderly. *Int Orthop.* 2018;42(9):2243–2248.
49. Perugia D, Guzzini M, Civitenga C, et al. Is it really necessary to restore radial anatomic parameters after distal radius fractures? *Injury.* 2014;45(suppl 6): S21–S26.
50. Kumar S, Penematsa S, Sadri M, Deshmukh SC. Can radiological results be surrogate markers of functional outcome in distal radial extra-articular fractures? *Int Orthop.* 2008;32(4):505–509.
51. Wright TW, Horodyski M, Smith DW. Functional outcome of unstable distal radius fractures: ORIF with a volar fixed-angle tine plate versus external fixation. *J Hand Surg.* 2005;30(2):289–299.
52. Chung KC, Cho HE, Kim Y, Kim HM, Shauer MJ. Assessment of anatomic restoration of distal radius fractures among older adults: a secondary analysis of a randomized clinical trial. *JAMA Netw Open.* 2020;3(1):e1919433.
53. Chen Y, Chen X, Li Z, Yan H, Zhou F, Gao W. Safety and efficacy of operative versus nonsurgical management of distal radius fractures in elderly patients: a systematic review and meta-analysis. *J Hand Surg.* 2016;41(3):404–413.
54. Ju JH, Jin GZ, Li GX, Hu HY, Hou RX. Comparison of treatment outcomes between nonsurgical and surgical treatment of distal radius fracture in elderly: a systematic review and meta-analysis. *Langenbecks Arch Surg.* 2015;400(7): 767–779.
55. Song J, Yu AX, Li ZH. Comparison of conservative and operative treatment for distal radius fracture: a meta-analysis of randomized controlled trials. *Int J Clin Exp Med.* 2015;8(10):17023–17035.
56. Arora R, Lutz M, Deml C, Krappinger D, Haug L, Gabl M. A prospective randomized trial comparing nonoperative treatment with volar locking plate fixation for displaced and unstable distal radial fractures in patients sixty-five years of age and older. *J Bone Joint Surg.* 2011;93(23):2146–2153.
57. Bartl C, Stengel D, Bruckner T, Gebhard F. The treatment of displaced intra-articular distal radius fractures in elderly patients. *Dtsch Arztebl Int.* 2014;111(46):779–787.
58. Ochen Y, Peek J, van der Velde D, et al. Operative vs nonoperative treatment of distal radius fractures in adults: a systematic review and meta-analysis. *JAMA Netw Open.* 2020;3(4):e203497.
59. Landgren M, Abramo A, Geijer M, Kopylov P, Tägil M. Similar 1-year subjective outcome after a distal radius fracture during the 10-year-period 2003–2012. *Acta Orthop.* 2017;88(4):451–456.
60. Changulani M, Okonkwo U, Keswani T, Kalairajah Y. Outcome evaluation measures for wrist and hand—which one to choose? *Int Orthop.* 2008;32(1): 1–6.
61. Dowrick AS, Gabbe BJ, Williamson OD, Cameron PA. Does the Disabilities of the Arm, Shoulder and Hand (DASH) scoring system only measure disability due to injuries to the upper limb? *J Bone Joint Surg.* 2006;88(4):524–527.
62. Tsang P, Walton D, Grewal R, MacDermid J. Validation of the QuickDASH and DASH in patients with distal radius fractures through agreement analysis. *Arch Phys Med Rehabil.* 2017;98(6):1217–1222.
63. Stephens AR, Presson AP, McFarland MM, et al. Volar locked plating versus closed reduction and casting for acute, displaced distal radial fractures in the elderly: a systematic review and meta-analysis of randomized controlled trials. *J Bone Joint Surg.* 2020;102(14):1280–1288.
64. Schurko BM, Lechtig A, Chen NC, et al. Outcomes and complications following volar and dorsal osteotomy for symptomatic distal radius malunions: a comparative study. *J Hand Surg.* 2020;45(2):158.e1.
65. Fares AB, Childs BR, Polmear MM, Clark DM, Nesti LJ, Dunn JC. Dorsal bridge plate for distal radius fractures: a systematic review. *J Hand Surg.* Published online February 8, 2021. doi:10.1016/j.jhsa.2020.11.026
66. Truong JL, Doherty C, Suh N. The effect of socioeconomic factors on outcomes of distal radius fractures: a systematic review. *Hand.* 2018;13(5):509–515.
67. Yeoh JC, Pike JM, Slobogean GP, O'Brien PJ, Broekhuysen HM, Lefavre KA. Role of depression in outcomes of low-energy distal radius fractures in patients older than 55 years. *J Orthop Trauma.* 2016;30(5):228–233.
68. Ring D, Kadzielski J, Fabian L, Zurakowski D, Malhotra LR, Jupiter JB. Self-reported upper extremity health status correlates with depression. *J Bone Joint Surg.* 2006;88(9):1983–1988.