



Review Article

Stroke and diets – A review

Chin-Lon Lin^{a,b*}

^aDepartment of Cardiology, Dalin Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Chiayi, Taiwan, ^bDepartment of Internal Medicine, College of Medicine, Tzu Chi University, Hualien, Taiwan

Submission : 07-Jul-2020
Revision : 28-Jul-2020
Acceptance : 25-Aug-2020
Web Publication : 24-Feb-2021

INTRODUCTION

There was a significant increase in stroke burden in the world over the past 25 years, especially in developing countries [1]. Stroke was the second-most common cause of death and the third-most common cause of disability after ischemic heart disease worldwide [1]. Globally, the majority (approximately three fourth) of strokes are ischemic, followed by intracranial hemorrhage, subarachnoid hemorrhage, and cerebral venous thrombosis, etc., [2,3]. Stroke is also the most common cause of complex disability and independent risk for dementia [4]. The direct and indirect cost of stroke care is a considerable burden to both society and patients' families. Although some risk factors of stroke are nonmodifiable, such as older age and male sex, current known modifiable risk factors—hypertension, current smoking, high waist-to-hip ratio, diet, regular physical activity, diabetes mellitus, alcohol consumption, psychological factors, cardiac causes, ApoB/Apo1 ratio—together account for >90% of population attributable risk for stroke [2,5]. In addition, in the United States of America, out of the seven metrics of cardiovascular health, healthy diet score is the poorest (91.6% poor, National Health and Nutrition Examination Survey 2011–2012.) [6]; hence, diet is the worst health issues in the United States [7]. The objective of this review is to present an overview of the

ABSTRACT

Stroke was one of the most common causes of death and disability worldwide. In addition, there was a significant increase in the disease burden of stroke in the world over the past 25 years, especially in developing countries. We searched PubMed (National Library of Medicine, USA) and Embase (Elsevier) databases using “Stroke and diets” as strategy, and additional references were obtained from the selected articles. The risk factors of stroke include age, sex, and modifiable factors such as hypertension, smoking, diet, physical activity, diabetes mellitus, alcohol consumption, psychological factors, and cardiac causes—together account for >90% of the population attributable risk for stroke. They were discussed individually (Hypertension, Diabetes, Dyslipidemia, Gut microbiota, Nutrition), and the dietary modifications to reduce these risks were also presented. In conclusion, besides low salt intake, plant-based dietary patterns (especially vegetarian diets) that are rich in fruit, vegetables, whole grains, legumes, seeds, nuts, and dairy and low in meat, sweets, and alcohol significantly improved risk factors for stroke, and observational studies clearly demonstrated the stroke morbidity and mortality benefits. Thus, dietary intervention should be considered as an important strategy in the prevention and management of stroke.

KEYWORDS: Hypertension, Microbiota, Plant-based diet, Stroke, Vegetarian diet

risk factors of stroke and how various dietary practices impact them.

We searched PubMed (National Library of Medicine, USA) and Embase (Elsevier) databases, using “Stroke and diets” as a key for peer-reviewed observational studies published in the last 5 years until June 2020. An additional manual search was also performed on the references of the studies selected from these databases. We concentrated on systemic reviews or meta-analysis with statistically significant results. We summarized our findings as follows:

MODIFIABLE RISK FACTORS OF STROKE

Hypertension

The most important risk factor for stroke is hypertension, accounting for 47.9%, 45.7%, and 56.4% of the population attributable risk for overall, ischemic, and hemorrhagic strokes, respectively, in the INTERSTROKE study [2]. Diets rich in fruits, vegetables, whole grains, legumes, seeds, nuts, fish, and dairy and low in meat, sweets such as the Dietary Approaches to Stop Hypertension (DASH) diets [8], the


*Address for correspondence:

Dr. Chin-Lon Lin,
Department of Cardiology, Dalin Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, 2, Minsheng Road, Dalin, Chiayi, Taiwan.
E-mail: cllinmd@tzuchi.com.tw

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Lin CL. Stroke and diets – A review. Tzu Chi Med J 2021;33:238-42.

Access this article online	
Quick Response Code: 	Website: www.tcmjmed.com
	DOI: 10.4103/tcmj.tcmj_168_20

Mediterranean diets (MDs) [9], the Nordic diets [10], and Vegetarian diets [11] all have been demonstrated to significantly lower systolic and diastolic blood pressures [12] and hence confer protection against stroke.

The DASH diet (diet rich in fruits, vegetables, and low-fat dairy products and with reduced saturated and total fat) reduced systolic and diastolic blood pressures by 5.5 and 3.0 mm Hg, respectively, as compared with control and it is estimated that a population-wide reduction in systolic or diastolic blood pressures of the magnitude observed would reduce stroke by approximately 27% [8], which is a very large proportion in the prevention of stroke.

After 1 year on MD, with extra virgin olive oil and mixed nuts, participants had 4.0–4.3 mm Hg lower mean systolic, and 1.9 mm Hg lower mean diastolic blood pressures on 24-h monitor than the control diet group. This reduction translated into a 30% reduction (hazard ratio [HR] 0.69) of major cardiovascular events (myocardial infarction, stroke, or death from cardiovascular causes) and 42% reduction of stroke [9].

Adherence to a healthy Nordic diet (consists of mainly whole grain and rye bread, berries, apples and pears, fish, cabbage and cruciferous vegetables, root vegetables, low-fat dairy products, potatoes, and vegetable fats) was associated with a lower risk of stroke. The HR comparing high adherence with low adherence was 0.86 (95% confidence interval [CI] 0.76–0.98) for total stroke [13]. The 15% reduction in total stroke incidence is probably an underestimate because the control is low adherence rather than no adherence.

The vegetarian diet group was also demonstrated to have a lower prevalence of hypertension than the nonvegetarian group and risk reduction in hemorrhagic, ischemic and total stroke [14]. All these studies demonstrated the significant beneficial effects of plant-based diets on hypertension and hence the risks of stroke.

Diabetes

Type 2 diabetes is a global epidemic and a rapidly rising prevalence, especially in middle- and low-income countries [15]. Diabetic patients have high rates of macro- and micro-vascular complications, including stroke, myocardial infarction, major lower-extremity amputations, end-stage renal disease, and visual impairment [16]. Increases in consumption of calorie-dense foods, including fast foods, meats and other animal fats, highly refined grains, and sugar-sweetened beverages, are thought to play a critical role in the rising rates of type 2 diabetes worldwide [17]. On the other hand, healthy diet, regular physical activity, maintaining a normal body weight, and avoiding tobacco use are ways to prevent or delay the onset of type 2 diabetes [18]. Lifestyle intervention reduced the incidence of diabetes by >50% as compared with placebo; and was even significantly more effective than metformin [19]. Plant-based diets—i.e., eating patterns that emphasize legumes, whole grains, vegetables, fruits, nuts, and seeds and discourage most or all animal products, are especially potent in preventing type 2 diabetes and have been associated with much lower rates of obesity, hypertension, hyperlipidemia, cardiovascular mortality, and cancer [20]. There is a general

consensus that plant-based diets are cost-effective, low-risk interventions that are not only highly beneficial for preventing and treating type 2 diabetes and other risk factors such as obesity, hypertension, hyperlipidemia, and inflammation, and they may also reduce the number of medications needed to treat chronic diseases and lower major cardiovascular disease mortality rates [20,21]. Hence, plant-based diets are important in the prevention of stroke.

Dyslipidemia

Though the relationship between cholesterol levels and stroke has been much less clear than between cholesterol levels and coronary heart disease, recent studies provided proof that total (and low-density lipoprotein [LDL]) cholesterol-lowering, reduces stroke risk proportional to the extent of cholesterol reduction [22]. In addition, after an ischemic stroke or TIA with evidence of atherosclerosis, patients who had been treated to a LDL cholesterol level of <70 mg per deciliter had a lower risk of subsequent cardiovascular events than those who had been treated to LDL cholesterol of 90 mg to 110 mg per deciliter [23]. The Portfolio diet (a dietary portfolio of cholesterol-lowering foods such as soy protein, viscous fiber, plant sterols, and almonds), could significantly lower serum LDL-cholesterol concentrations (more than 20%) [24]. In addition, plant-based diets, such as The Mediterranean, DASH, Nordic, and the vegetarian diets, all have positive effects on various cardiometabolic risk factors (glycemic control, blood lipids, body weight and adiposity, and blood pressure) and cardiovascular disease outcomes such as myocardial infarction and stroke [25]. Thus, plant-based diets not only lower blood pressure, reduce weight, improve glycemic control, they also lower total and LDL-cholesterol, and confer protection against stroke.

Gut microbiota

The gut microbiota (GM) is the community of microorganisms (bacteria, archaea, fungi, viruses) colonizing the gastrointestinal tract. GM regulates various metabolic pathways in the host, including those involved in energy homeostasis, glucose, lipid, and bile acid metabolism. GM dysbiosis has been shown to be involved in the pathogenesis of diverse diseases, such as metabolic syndrome, cardiovascular diseases, celiac disease, inflammatory bowel disease, and neurological disorders [26]. Multiple factors modulate the composition of the microbiota and its activity. The individual microbiota pattern is influenced by antibiotic use, medication (nonsteroidal anti-inflammatory drugs, proton pump inhibitors, etc.), infections, and chronic stress. However, one of the major factors triggering GM establishment is diet. Plant-based diets, with adequate fruit and vegetable intake, rich in dietary fiber, with healthy fats (monounsaturated fatty acids [MUFAs] and polyunsaturated fatty acids [PUFAs]), and a predominance of plant-derived proteins seem to be the best way to promote GM diversity and activity [26]. Vegan or vegetarian diet results in a significantly lower stool pH and shift in the microbiota, while total cell numbers remain unaltered as compared with omnivore diet [27]. Intestinal microbiota metabolism of dietary L-carnitine, trimethylamine abundant in red meat, produces trimethylamine N-oxide (TMAO) and accelerates atherosclerosis. In addition, omnivorous subjects produce

significantly more TMAO than vegans/vegetarians following the ingestion of L-carnitine through a microbiota-dependent mechanism [28]. Carnitine from meat and choline from eggs not only provide substrates for but also shift gut microbial community to enhance the production of TMAO [29], an atherosclerotic factor that has been shown to induce platelet hyperreactivity and thrombosis [30] and increase stroke incidence [31]. Thus, plant-based diets can alter the composition of intestinal microbiota and confer the protection of cardiovascular diseases through interactions with GM.

Nutrition

Salt

The close relationship between dietary sodium intake and hypertension is widely recognized. The high sodium intake in the diet increases water retention and systemic peripheral resistance, alters the endothelial function, causes changes in the structure and function of large elastic arteries, and modification in autonomic neuronal modulation of the cardiovascular system. A reduction in dietary sodium not only decreases the blood pressure and the incidence of hypertension, also is associated with a reduction in morbidity and mortality from cardiovascular diseases [32,33]. Fresh vegetables and fruits are generally low in salt content, but salt is often added in the preparation and cooking of plant-based foods and should be avoided.

Fiber

Increased fiber intake in diets lowers blood pressure and serum cholesterol levels, improves glycemic control and insulin sensitivity, enhances weight loss, and benefits a number of gastrointestinal disorders, including gastroesophageal reflux disease, duodenal ulcer, diverticulitis, constipation, and hemorrhoids [34]. Individuals with high intakes of dietary fiber appear to be at significantly lower risk for developing coronary heart disease, stroke, hypertension, diabetes, obesity [31]. In a meta-analysis, greater dietary fiber intake is significantly associated with a lower risk of first stroke [35]. Typical plant-based foods have high contents of fiber and, therefore, beneficial for the prevention of stroke.

Vitamins

Plasma homocysteine is derived from the essential amino acid methionine and its metabolic pathways are catalyzed by enzymes requiring folic acid, Vitamin B6 and Vitamin B12 (cobalamin) as essential cofactors. Hence, the total plasma concentration of total homocysteine (tHcy) is influenced by blood concentrations of the B vitamins; folic acid (B9), cyanocobalamin (B12), and pyridoxine (B6) [36]. Systematic reviews showed a strong, positive and dose-related association between the plasma concentration of tHcy and the risk of stroke, independent of other vascular risk factors [37,38]. In fact, supplementation with B vitamins (folic acid, Vitamin B12, and Vitamin B6) lower blood concentrations of tHcy by 25% and reduces the relative risk of stroke overall by about 10% compared with placebo [36]. Vegetarians tend to have a low intake of Vitamin B12, which is found mainly in animal-based foods, and this raises plasma homocysteine, leading to an increase in the risk of stroke [36]. However, the combined effect of the supposed harm of high homocysteine and

the supposed benefits of low BP, glucose, and cholesterol (the unique aspect of vegetarian diets) on stroke incidence is unclear. In fact, cohort studies demonstrated that vegetarian diet is associated with a lower risk of ischemic and hemorrhagic strokes in a nonsmoking, non-drinking cohort despite higher plasma homocysteine levels [14]. Whether those on plant-based diets should receive vitamin B12 supplements remained controversial.

Nuts

Nuts have unique nutritional profiles. Nuts contain 43%–67% fat, 8%–22% protein, and only 1%–4% carbohydrate by weight and are abundant in unsaturated fatty acids (UFAs) (containing both monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA)) and only 4%–5% of saturated fatty acids. Nuts are also high in vitamins, minerals, fiber, and phytochemicals [39,40]. Recent meta-analyses found that nut consumption has beneficial effects on the cardiometabolic disease with reduced stroke incidence and mortality. It is mainly attributable to decreases in fasting glucose, total cholesterol, and LDL-C. Nuts, with its high contents of UFAs and protein are important ingredients in nutritionally balanced plant-based diets.

EPIDEMIOLOGICAL AND OBSERVATIONAL STUDIES

Various nutrients or individual foods have been studied in relation to stroke risk; whole-grain, fruit, and vegetable consumption has been found to be protective against cardiovascular mortality [41,42], and fish consumption is weakly inversely associated with the risk of stroke [43]. On the other hand, increased meat, especially processed red meat consumption, has been shown to correlate with increased stroke risk [44,45].

Dietary pattern analysis has emerged as an approach to examining the relationship between diet and the risk of chronic diseases. Instead of looking at individual nutrients or foods, pattern analysis examines the effects of overall diet and represents a broader picture of food and nutrient consumption, and maybe more predictive of disease risk than individual foods or nutrients [46]. It has also been demonstrated that the highest behavioral risk is incorrect nutrition for noncommunicable diseases, and that the choice of a balanced diet can reduce the risk of stroke [47].

After 1 year on MD supplemented with extra virgin olive oil and mixed nuts, the adjusted HRs for major cardiovascular events (myocardial infarction, stroke, or death from cardiovascular causes) were 0.70 (95% CI, 0.54–0.92, olive oil) and 0.72 (95% CI, 0.54–0.96, mixed nuts), versus the control group. Specifically, the diet also lowers the incidence of stroke by 42% and total cardiovascular mortality by 20% [9].

In the landmark “China study” [48], comparing the dietary patterns and mortality data from a large geographic area of rural China to that of the United States of America, demonstrated the link between the consumption of animal products (including dairy) and chronic illnesses such as coronary heart disease, diabetes, breast cancer, prostate cancer, and bowel cancer. The coronary artery disease mortality

rates in rural China were inversely associated with the frequency of intake of green vegetables and plasma erythrocyte MUFAs, but positively associated with salt, animal protein intake and negatively associated with plant protein, legume, and light-colored vegetable intake. The authors conclude that people who eat predominantly whole-food–avoiding animal products, including beef, pork, poultry, fish, eggs, cheese, and milk, and reducing their intake of processed foods and refined carbohydrates–will escape, reduce, or reverse the development of numerous diseases [49].

In a meta-analysis of the associations between different dietary patterns and the risk of stroke by pooling available data from 21 published studies, using a “healthy” diet scoring system on the food Frequency Questionnaire, the “healthy” dietary pattern was characterized by high intakes of vegetables, fruits, fish, low-fat milk, and whole grains. A decreased risk of stroke was demonstrated for the highest compared with the lowest categories of healthy dietary pattern (odds ratio = 0.77; 95% CI = 0.64–0.93; $P = 0.006$) [50] that translates into a 23% reduction of stroke risks.

The Adventist Health Study 2 [51] demonstrated that vegetarian groups (older, more educated, more likely to be married, drink less alcohol, smokeless, exercise more, and thinner) were associated with lower all-cause and cardiovascular mortality. The adjusted HR for all-cause mortality in all vegetarians combined versus nonvegetarians was 0.88 (95% CI, 0.80–0.97) a reduction of only 12% risk. However, they included vegan (no animal products, 7.6%), lacto-ovo–vegetarians (no animal meat, but consumes milk and eggs, 28.9%), pesco-vegetarians (consumes fish, 9.8%), semi-vegetarians (occasional meat consumption, 5.5%), in their vegetarian group. This will attenuate the true effects of vegetarian diets (meatless).

The DASH dietary pattern improves blood pressure and other cardiometabolic risk factors and is associated with decreased incidence of cardiovascular disease in people with and without diabetes [52]. Compared with lower adherence, higher adherence to the DASH diet was related to a reduced risk of developing stroke (RR 0.88, 95% CI 0.83–0.93), and there is a linear association of the DASH diet score with stroke (P for nonlinearity = 0.411), and each 4-points increment in the score conferred a risk reduction of 4% (RR 0.96, 95% CI 0.94–0.97) in total stroke events [53]. Similarly, the HR comparing high adherence to the Nordic diet with low adherence was 0.86 (95% CI 0.76–0.98) for total stroke [13]. The 12%–15% reduction in total stroke incidence is probably an underestimate because the control is low adherence rather than no adherence.

In a study comparing the stroke incidence of Buddhist volunteers, the vegetarian (meatless) diet groups (mostly ovo-lactovegetarians, few vegans) had a lower prevalence of hypertension than the nonvegetarian group and approximately, 60%, 60%, 50% risk reduction of hemorrhagic, ischemic and total stroke respectively. In addition, both the study and control groups were similar in their lifestyles (non-smoking, non-drinking, and active community volunteers). The only difference between the study and control group is the amount of meat (beef, pork, poultry, fish, etc.) they consumed; the

magnitude of stroke risk reduction underscores the importance of meat abstinence in the prevention of stroke [14].

All these observational studies demonstrated the significant benefits of plant-based diets, especially vegetarian diets (no meat), on the risk of stroke.

CONCLUSION

In summary, besides low salt intake, plant-based dietary patterns that are rich in fruit, vegetables, whole grains, legumes, seeds, nuts, and dairy and low in meat, sweets, and alcohol, such as the DASH, Nordic, Mediterranean, and vegetarian diets significantly lowered blood pressure, diabetes and other risk factors of stroke, especially ischemic stroke. Most epidemiological or observational studies confirm such effects. As discussed in this review, plant-based diets, especially vegetarian diets (meatless), have a significant impact on major risk factors of stroke, such as hypertension, diabetes, hyperlipidemia, and GM, and should be recognized as one of the most important strategies in the prevention and management of stroke.

Financial support and sponsorship

Nil.

Conflicts of interest

Dr. Chin-Lon Lin, an editorial board member at *Tzu Chi Medical Journal*, had no role in the peer review process or decision to publish this article. .

REFERENCES

1. Feigin VL, Norrving B, Mensah GA. Global burden of stroke. *Cir Res* 2017;120:439-48.
2. O'Donnell MJ, Chin SL, Rangarajan S, Xavier D, Liu L, Zhang H, et al. Global and regional effects of potentially modifiable risk factors associated with acute stroke in 32 countries (INTERSTROKE): A case-control study. *Lancet* 2016;388:761-75.
3. Hsieh FI, Chiou HY. Stroke: Morbidity, risk factors, and care in Taiwan. *J Stroke* 2014;16:59-64.
4. Pohjasvaara T, Erkinjuntti T, Ylikoski R, Hietanen M, Vataja R, Kaste M. Clinical determinants of poststroke dementia. *Stroke* 1998;29:75-81.
5. Ma YH, Leng XY, Dong Y, Xu W, Cao XP, Ji X, et al. Risk factors for intracranial atherosclerosis: A systematic review and meta-analysis. *Atherosclerosis* 2019;281:71-7.
6. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart disease and stroke statistics--2015 update: A report from the American Heart Association. *Circulation* 2015;131:e29-322.
7. Spence JD. Nutrition and Risk of Stroke. *Nutrients* 2019;11:647.
8. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med* 1997;336:1117-24.
9. Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, et al. Primary prevention of cardiovascular disease with a mediterranean diet supplemented with extra-virgin olive oil or nuts. *N Engl J Med* 2018;378:e34.
10. Brader L, Uusitupa M, Dragsted LO, Hermansen K. Effects of an isocaloric healthy Nordic diet on ambulatory blood pressure in metabolic syndrome: A randomized SYSDIET sub-study. *Eur J Clin Nutr* 2014;68:57-63.
11. Yokoyama Y, Nishimura K, Barnard ND, Takegami M, Watanabe M, Sekikawa A, et al. Vegetarian diets and blood pressure: A meta-analysis. *JAMA Intern Med* 2014;174:577-87.

12. Ndanuko RN, Tapsell LC, Charlton KE, Neale EP, Batterham MJ. Dietary patterns and blood pressure in adults: A systematic review and meta-analysis of randomized controlled trials. *Adv Nutr* 2016;7:76-89.
13. Hansen CP, Overvad K, Kyrø C, Olsen A, Tjønneland A, Johnsen SP, et al. Adherence to a healthy nordic diet and risk of stroke: A danish cohort study. *Stroke* 2017;48:259-64.
14. Chiu THT, Chang HR, Wang LY, Chang CC, Lin MN, Lin CL. Vegetarian diet and incidence of total, ischemic, and hemorrhagic stroke in 2 cohorts in Taiwan. *Neurology* 2020;94:e1112-21.
15. The World Health Organization Fact Sheets. Available from: <https://www.who.int/news-room/fact-sheets/detail/diabetes>. [Last accessed on 2020 Apr 25].
16. Caspersen CJ, Thomas GD, Boseman LA, Beckles GL, Albright AL. Aging, diabetes, and the public health system in the United States. *Am J Public Health* 2012;102:1482-97.
17. Ley SH, Hamdy O, Mohan V, Hu FB. Prevention and management of type 2 diabetes: Dietary components and nutritional strategies. *Lancet* 2014;383:1999-2007.
18. Yokoyama Y, Barnard ND, Levin SM, Watanabe M. Vegetarian diets and glycemic control in diabetes: A systematic review and meta-analysis. *Cardiovasc Diagn Ther* 2014;4:373-82.
19. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393-403.
20. McMacken M, Shah S. A plant-based diet for the prevention and treatment of type 2 diabetes. *J Geriatr Cardiol* 2017;14:342-54.
21. Tuso PJ, Ismail MH, Ha BP, Bartolotto C. Nutritional update for physicians: Plant-based diets. *Perm J* 2013;17:61-6.
22. Salvatore T, Morganti R, Marchioli R, De Caterina R. Cholesterol lowering and stroke: No longer room for pleiotropic effects of statins - Confirmation from PCSK9 inhibitor studies. *Am J Med* 2020;133:95-9.e6.
23. Amarenco P, Kim JS, Labreuche J, Charles H, Abtan J, Bèjot Y, et al. A comparison of two LDL cholesterol targets after ischemic stroke. *N Engl J Med* 2020;382:9.
24. Jenkins DJ, Kendall CW, Faulkner DA, Nguyen T, Kemp T, Marchie A, et al. Assessment of the longer-term effects of a dietary portfolio of cholesterol-lowering foods in hypercholesterolemia. *Am J Clin Nutr* 2006;83:582-91.
25. Kahleova H, Salas-Salvadó J, Rahelić D, Kendall CW, Rembert E, Sievenpiper JL. Dietary patterns and cardiometabolic outcomes in diabetes: A summary of systematic reviews and meta-analyses. *Nutrients* 2019;11:2209.
26. Moszak M, Szulińska M, Bogdański P. You are what you eat-the relationship between diet, microbiota, and metabolic disorders-A review. *Nutrients* 2020; 12:1096.
27. Zimmer J, Lange B, Frick JS, Sauer H, Zimmermann K, Schwiertz A, et al. A vegan or vegetarian diet substantially alters the human colonic faecal microbiota. *Eur J Clin Nutr* 2012;66:53-60.
28. Koeth RA, Wang Z, Levison BS, Buffa JA, Org E, Sheehy BT, et al. Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat Med* 2013;19:576-85.
29. Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, et al. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med* 2013;368:1575-84.
30. Zhu W, Gregory JC, Org E, Buffa JA, Gupta N, Wang Z, et al. Gut microbial metabolite TMAO enhances platelet hyperreactivity and thrombosis risk. *Cell* 2016;165:111-24.
31. Nie J, Xie L, Zhao BX, Li Y, Qiu B, Zhu F, et al. Serum trimethylamine N-Oxide concentration is positively associated with first stroke in hypertensive patients. *Stroke* 2018;49:2021-8.
32. Grillo A, Salvi L, Coruzzi P, Salvi P, Parati G. Sodium Intake and Hypertension. *Nutrients* 2019;11:1970.
33. He FJ, Tan M, Ma Y, MacGregor GA. Salt reduction to prevent hypertension and cardiovascular disease: JACC state-of-the-art review. *J Am Coll Cardiol* 2020;75:632-47.
34. Anderson JW, Baird P, Davis RH Jr, Ferreri S, Knudtson M, Koraym A, et al. Health benefits of dietary fiber. *Nutr Rev* 2009;67:188-205.
35. Threapleton DE, Greenwood DC, Evans CE, Cleghorn CL, Nykjaer C, Woodhead C, et al. Dietary fiber intake and risk of first stroke: A systematic review and meta-analysis. *Stroke* 2013;44:1360-8.
36. Hankey GJ. B vitamins for stroke prevention. *Stroke Vasc Neurol* 2018;3:51-8.
37. Homocysteine Studies Collaboration. Homocysteine and risk of ischemic heart disease and stroke: A meta-analysis. *JAMA* 2002;288:2015-22.
38. Wald DS, Law M, Morris JK. Homocysteine and cardiovascular disease: Evidence on causality from a meta-analysis. *BMJ* 2002;325:1202.
39. Alasalvar C, Bolling BW. Review of nut phytochemicals, fat-soluble bioactives, antioxidant components and health effects. *Br J Nutr* 2015;113 Suppl 2:S68-78.
40. Venkatchalam M, Sathe SK. Chemical composition of selected edible nut seeds. *J Agric Food Chem* 2006;54:4705-14.
41. Zong G, Gao A, Hu FB, Sun Q. Whole grain intake and mortality from all causes, cardiovascular disease, and cancer: A meta-analysis of prospective cohort studies. *Circulation* 2016;133:2370-80.
42. Hu D, Huang J, Wang Y, Zhang D, Qu Y. Fruits and vegetables consumption and risk of stroke: A meta-analysis of prospective cohort studies. *Stroke* 2014;45:1613-9.
43. Larsson SC, Orsini N. Fish consumption and the risk of stroke: A dose-response meta-analysis. *Stroke* 2011;42:3621-3.
44. Yang C, Pan L, Sun C, Xi Y, Wang L, Li D. Red meat consumption and the risk of stroke: A dose-response meta-analysis of prospective cohort studies. *J Stroke Cerebrovasc Dis* 2016;25:1177-86.
45. Kim K, Hyeon J, Lee SA, Kwon SO, Lee H, Keum N, et al. Role of total, red, processed, and white meat consumption in stroke incidence and mortality: A systematic review and meta-analysis of prospective cohort studies. *J Am Heart Assoc* 2017; 6:e005983.
46. Hu FB. Dietary pattern analysis: A new direction in nutritional epidemiology. *Curr Opin Lipidol* 2002;13:3-9.
47. Altobelli E, Angeletti PM, Rapacchietta L, Petrocelli R. Overview of meta-analyses: The impact of dietary lifestyle on stroke risk. *Int J Environ Res Public Health* 2019; 16:3582.
48. Campbell TC, Parpia B, Chen J. Diet, lifestyle, and the etiology of coronary artery disease: The Cornell China study. *Am J Cardiol* 1998;82:18T-21T.
49. Campbell TC, Junshi C. Diet and chronic degenerative diseases: Perspectives from China. *Am J Clin Nutr* 1994;59:1153S-61S.
50. Zhang X, Shu L, Si C, Yu X, Gao W, Liao D, et al. Dietary patterns and risk of stroke in adults: A systematic review and meta-analysis of prospective cohort studies. *J Stroke Cerebrovasc Dis* 2015;24:2173-82.
51. Orlich MJ, Singh PN, Sabaté J, Jaceldo-Siegl K, Fan J, Knutsen S, et al. Vegetarian dietary patterns and mortality in adventist health study 2. *JAMA Intern Med* 2013;173:1230-8.
52. Chiavaroli L, Vigiouliou E, Nishi SK, Blanco Mejia S, Rahelić D, Kahleová H, et al. DASH dietary pattern and cardiometabolic outcomes: An umbrella review of systematic reviews and meta-analyses. *Nutrients* 2019;11:338.
53. Feng Q, Fan S, Wu Y, Zhou D, Zhao R, Liu M, et al. Adherence to the dietary approaches to stop hypertension diet and risk of stroke: A meta-analysis of prospective studies. *Medicine (Baltimore)* 2018;97:e12450.