

RESEARCH ARTICLE

Nocturnal to Diurnal Switches with Spontaneous Suppression of Wheel-Running Behavior in a Subterranean Rodent

Patricia Tachinardi¹, Øivind Tøien², Veronica S. Valentinuzzi^{3*}, C. Loren Buck^{4*}, Gisele A. Oda^{1*}

1 Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil, **2** Institute of Arctic Biology, University of Alaska, Fairbanks, United States of America, **3** Centro Regional de Investigaciones Científicas y Transferencia Tecnológica (CRILAR), La Rioja, Argentina, **4** Department of Biological Sciences, University of Alaska, Anchorage, United States of America

* gaoda@ib.usp.br (GAO); Loren.Buck@nau.edu (CLB); vvalentinuzzi@crilar-conicet.gob.ar (VSV)



CrossMark
click for updates

OPEN ACCESS

Citation: Tachinardi P, Tøien Ø, Valentinuzzi VS, Buck CL, Oda GA (2015) Nocturnal to Diurnal Switches with Spontaneous Suppression of Wheel-Running Behavior in a Subterranean Rodent. PLoS ONE 10(10): e0140500. doi:10.1371/journal.pone.0140500

Editor: Eric M Mintz, Kent State University, UNITED STATES

Received: June 25, 2015

Accepted: September 25, 2015

Published: October 13, 2015

Copyright: © 2015 Tachinardi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (www.fapesp.br): grants 12/15767-2, 12/16362-6, 13/50482-1 to GAO; 12/23393-5 to PT, Conselho Nacional de Desenvolvimento Científico e Tecnológico (www.cnpq.br): grant 142215/2012-0 to PT, Consejo Nacional de Investigaciones Científicas y Técnicas (www.conicet.gov.ar): grant PIP-11420090100252 to VSV and the Agencia Nacional de Promoción Científica y Tecnológica (www.agencia).

Abstract

Several rodent species that are diurnal in the field become nocturnal in the lab. It has been suggested that the use of running-wheels in the lab might contribute to this timing switch. This proposition is based on studies that indicate feed-back of vigorous wheel-running on the period and phase of circadian clocks that time daily activity rhythms. Tuco-tucos (*Ctenomys aff. knighti*) are subterranean rodents that are diurnal in the field but are robustly nocturnal in laboratory, with or without access to running wheels. We assessed their energy metabolism by continuously and simultaneously monitoring rates of oxygen consumption, body temperature, general motor and wheel running activity for several days in the presence and absence of wheels. Surprisingly, some individuals spontaneously suppressed running-wheel activity and switched to diurnality in the respirometry chamber, whereas the remaining animals continued to be nocturnal even after wheel removal. This is the first report of timing switches that occur with spontaneous wheel-running suppression and which are not replicated by removal of the wheel.

Introduction

The tuco-tuco (*Ctenomys aff. knighti*), a South American subterranean rodent, is among the several mammals described as having discrepant activity timing between field and laboratory conditions [1–7]. Whereas they are active during the day in semi-natural field enclosures, all individuals are nocturnal under laboratory conditions, with or without access to running-wheels [8–10] suggesting that a fundamental feature of their natural environment is not reproduced in the laboratory. Both ecological and physiological studies indicate the critical role of daily energy balance in constraining the timing of activity, which is primarily determined by the circadian clock [7,11–17]. Thus, differences in energy demand between field and laboratory conditions could be the fundamental feature ultimately leading to inversion in the timing of daily activity [7,11,17].

mincyt.gob.ar: grant PICT 2011-1979 to VSV. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Subterranean rodents are excellent subjects to explore this proposition because their foraging activity in the field involves intense underground excavation, an intense energy demanding activity [18–23]. Moreover, the periodic day-time emergence of tuco-tucos to the surface [9,24] might indicate economy in thermoregulation costs, allowing allocation of the rest phase inside burrows during the coldest hours of the desert night [7,25].

Few studies have addressed continuous, 24h rhythms of metabolism in subterranean rodents [13, 26]. We set out to assess the daily patterns of metabolic rate and its interrelationships with body temperature (T_b) and activity in wild-caught tuco-tucos. We hypothesized that tuco-tucos would show higher metabolic rates, levels of activity and T_b during darkness under laboratory light/dark cycles. Furthermore, we predicted that availability of a running wheel in the respirometry chamber would modulate amplitude but not timing of metabolic rate, as shown before for T_b [10]. To accomplish our investigation, we monitored individuals continuously for oxygen consumption ($\dot{V}O_2$), T_b , general motor activity and wheel-running over several consecutive days. Measurement of $\dot{V}O_2$ occurred inside a respirometry chamber, which, to our surprise, revealed a novel association between wheel-running and switches in timing of activity.

Materials and Methods

Ethics statement

All procedures followed the guidelines of the American Society of Mammalogists for the use of wild mammals in research [27] and the U.S. National Institutes of Health Guide for the Care and Use of Laboratory Animals [28]. All experiments were performed in Anillaco and were authorized by the Environmental Department of La Rioja (permits 028–10 and 062–08) and approved by the Ethics Committees of the Biosciences Institute of the University of São Paulo, Brazil (permit 164/2012), the University of Alaska Anchorage’s Institutional Animal Care and Use Committee (405977–1) and of the Faculty of Veterinary Sciences of La Plata National University, Argentina (permit 29-2-12).

Animals

Tuco-tucos were trapped within a 3 km radius of the town of Anillaco (28° 48′ S; 66° 56′ W; 1350 m) in the ecoregion of the Monte Desert, Argentina. The animals were live-trapped within a 15km² area surrounding the laboratory, with buried traps constructed from a 25-cm long PVC plumbing pipe with a 7.5-cm outer diameter. The traps were set by opening a burrow beneath a fresh mound of soil and positioning the pipe horizontally along the floor of the tunnel. Because the animals sometimes plugged the traps with loose soil, the traps were checked every 1–2 h, cleaned and reset as needed. Nine adult individuals (140–220 g) were used, of which five were females and four were males. Because these animals are solitary, they were housed individually in plastic cages (53×29×27 cm) equipped with running wheels (23 cm diameter, 10 cm wide, 1 cm between bars). Food (grass, native plants, carrot, sweet potato, rabbit pellets, oat, sunflower seeds) was provided *ad libitum* and replaced daily at various times. Water was not offered because subterranean rodents do not drink free water [19].

Cages were placed inside light-tight boxes equipped with one incandescent red light bulb providing continuous dim red light (1–5 lux) to facilitate animal care, and one fluorescent bulb of 200–250 lux at cage lid level connected to a timing device. Unless specified otherwise, tuco-tucos were kept under an LD cycle with 12 hours of “darkness” (1–5 lux) followed by 12 hours of light (LD 12:12), with lights on at 07:00 AM (local time, GMT -3).

Relative humidity ranged from 30–60% and room temperature was maintained at $25\pm 2^\circ\text{C}$, which is within the thermoneutral zone of other *Ctenomys* species [29] (Tachinardi, unpublished). Data loggers (HOBO U10/003, Onset Computer Corporation, Bourne, MA) recorded room temperature and relative humidity every 15 minutes.

Monitoring of wheel-running, general activity and body temperature

Tuco-tucos were surgically implanted with temperature sensitive transponders (G2 E-Mitters, Mini-Mitter, Bend, OR) to allow for continuous monitoring of core T_b and gross motor activity. Animals were anaesthetized using either ketamine/acepromazine (200 and 20 mg/Kg, respectively) or isoflurane anaesthesia (3%–5% with oxygen). Transponders were inserted into the peritoneal cavity through 1.5–2 cm vertical midline incision (1 cm bellow the rib cage) and sutured with poliglicolic acid thread (for more surgical details, see [10]). All surgeries were completed at least eight weeks prior to initiation of experiments.

Each cage was placed above a receiver (ER 4000, Mini-Mitter, Bend, OR) and data were collected and processed using the software VitalView (Mini-Mitter, Bend, OR); averages of T_b and activity were recorded each five minutes. Wheel-running was recorded as total revolutions in each 5-min interval by the ArChron Data Acquisition System (Simonetta System, Universidad Nacional de Quilmes, Buenos Aires, Argentina).

Respirometry

Rates of O_2 consumption were measured by open-flow respirometry during February and March of 2013 and 2014. In 2013, we used a FoxBox (Sable Systems, Las Vegas, NV) and Molecular Sieve 3Å (8–12 mesh, Sigma-Aldrich, Saint Louis, MO) as a desiccant with O_2 measurement only. In 2014 we used the Field Metabolic System (Sable Systems, Las Vegas, NV) and a Nafion Dryer to remove moisture from the air [30]. Since $\dot{V}\text{O}_2$ data collected in the two years did not significantly differ (two-tailed t-test, $p>0.05$), we merged data from both years for further analysis.

During the experiments, animals were individually kept inside a respirometry chamber. This is the home cage (volume = 40L) with its wire lid replaced by a sealed clear acrylic lid, with fittings for in-flow and out-flow of air for the respirometry measures (S1 and S2 Figs). Outside air was pulled through the metabolic chamber at 450–650 mL/min, depending on the size of the animal. Before entering the chamber, outside air was passed through copper tubing (2m length) to facilitate equilibration of incurrent air temperature with air temperature of the animal room. Flow was generated by a vacuum pump and measured by a mass flow meter (part of the FoxBox System or the Mass Flow System-5, Sable Systems, Las Vegas, NV).

Excurrent air was drawn through Molecular Sieve 3Å or the Nafion dryer to remove moisture prior to measurements of gas concentrations. A subsample was passed through oxygen and carbon dioxide analyzers. The O_2 analyzer was calibrated with ambient air every hour. Averages of flow rate and $\text{O}_2\%$ were logged onto a computer each minute and corrected for baseline drift by linear interpolation using modified version of LabGraph [30].

Mass specific rate of oxygen consumption ($\text{mL g}^{-1} \text{h}^{-1}$) was calculated using the following equations [30,31]:

$$\dot{V}\text{O}_2 = [V_E * (F\text{IO}_2 - F\text{EO}_2) / (1 - F\text{IO}_2 * (1 - RQ))] / BM$$

\dot{V}_E = airflow exiting chamber (mL/min), $F\text{IO}_2$ = fraction of O_2 entering chamber, $F\text{EO}_2$ = fraction of O_2 exiting chamber, RQ = respiratory quotient (assumed to be 0.85, BM = body mass (Kg)).

Integrity of the respirometry system was tested before the 2014 trials using alcohol burns [30].

Sufficient food for at least three days was placed inside the chamber at the beginning of the experiment. For trials lasting more than three days, additional food was supplied during the experiment by quickly opening and re-sealing the chamber. Chamber temperature was $25 \pm 1^\circ\text{C}$, recorded every 15 minutes by a data logger (HOBO U10/003, Onset Computer Corporation, Bourne, MA). Animals were weighed before and after each trial.

Experiments

We performed continuous 5–9 day long respirometry trials for each animal, previously entrained by CE12:12, using two protocols. In the first ($N = 4$), respirometry trials were initiated without animal access to a running wheel and wheels were added on day three inside the chamber. In the second protocol ($N = 5$), trials started with a running wheel inside the chamber but removed on the third day. Activity and T_b were monitored continuously for at least 3 days before, during, and for 3 days after the respirometry trials.

Data analysis

Animal activity and T_b were firstly depicted in double-plotted actograms using the software El Temps (Díez-Noguera, Universitat de Barcelona, Spain, 1999). Actograms allowed visual estimation of phase and rhythmic pattern.

To quantify phase changes in different conditions, we used a modified version of the diurnality index (D) proposed by Hoogenboom et al. [32][6,17]:

$$D = \sum [(T_{L_i} - M) - (T_{D_i} - M)]_i / \sum [(T_{L_i} - M) + (T_{D_i} - M)]_i$$

where T_{L_i} and T_{D_i} correspond respectively to each T_b measure during the light and dark phase (only values above the mean were considered) and M corresponds to the mean T_b during light and dark. This index is symmetric around 0 and runs from -1 (no high T_b during the day) to +1 (high T_b only during the day). We used T_b to calculate the D-Index because it was a variable recorded throughout all conditions.

$\dot{V}O_2$ data are presented as means \pm SEM. We tested for the significance ($\alpha = 0.05$) of differences in variables under different conditions using one-way ANOVAs (for multiple group comparisons) or two-tailed Student's t-test (when only two conditions were compared). To test for significant associations among D-Indices and measured variables, we ran Pearson's product-moment correlation tests. All analyses were performed with R version 2.11.1 [33].

Results

Before the start of the respirometry trials, all animals displayed a nocturnal pattern with high T_b , general activity and wheel-running concentrated in the dark phase. When animals were placed into the respirometry chamber, some animals showed a radical and immediate change in their timing of peak $\dot{V}O_2$, T_b and general activity. While some ($N = 3$) remained clearly nocturnal (D-Indices < -0.4), the majority ($N = 6$) changed their rhythmic pattern and became either robustly diurnal (D-Indices > 0 ; $N = 3$) or did not show clear nocturnality/diurnality (D-Indices between -0.1 and 0.1; $N = 3$). Fig 1A and 1B are representative data of one animal that remained nocturnal and of one that switched to diurnality inside the respirometry chamber, respectively. The left sections display time series of the variables inside the respirometer (colored backgrounds) to highlight their amplitude changes. The right sections display the corresponding whole data set (including days outside the chamber) in actogram format to




Fig 1. Simultaneous measurements of daily rhythms in oxygen consumption ($\dot{V}O_2$), body temperature (T_b), gross motor and wheel-running activity of tuco-tucos. Left: temporal series collected when the animal was inside the respirometry chamber, with and without a running-wheel. Shaded areas indicate dark phases and white areas light phases. Right: actograms across experimental conditions. Pink and blue backgrounds indicate data from animals inside the respirometry chamber, with and without access to wheels, respectively. (A) Representative individual that did not switch phase inside the respirometry chamber. Pink line in the left figure indicates introduction of the wheel to the chamber. (B) Representative individual that switched from nocturnal to diurnal inside the respirometry chamber. There was a 7-day interval outside the respirometry chamber before the wheel introduction due to technical problems. Pink broken line in the right figure separates days with and without wheels. General conditions: LD12:12 (L = 200–250 lux), $25 \pm 2^\circ\text{C}$ and food *ad libitum*.

doi:10.1371/journal.pone.0140500.g001

highlight phase shifts. Because the respirometry measurement involved days with and without access to running wheels, blue and pink backgrounds indicate these two different conditions, respectively. As can be seen in Fig 1A, an animal that remained nocturnal showed highest values of all variables during the dark phase, including vigorous wheel-running episodes that appear clearly at the bottom of Fig 1A (left, blue background) and along the running-wheel actogram (right, blue background). On the other hand, Fig 1B shows data of an animal that switched to diurnality, as indicated by a shift of the highest values of variables to the light phase. This shift is more easily visualized in the right actograms, where timing of high amplitude T_b and motor activity switch to the light phase inside the colored sections, in dramatic contrast to the undisturbed patterns displayed in Fig 1A (right). Most notably, however, there was a total suppression of wheel-running activity in this animal, as indicated by no revolutions in the bottom of Fig 1B (left, blue background) and along the running-wheel actogram (right, blue background), again in contrast to the pattern exhibited in Fig 1A.

D-indices ranged from -0.98 to -0.39 when animals were outside the respirometry chamber. Inside the respirometry chamber, D-indices ranged from -0.23 to +0.29 in the absence of the running wheel and from -0.61 to +0.83 when the wheel was available (Fig 2). One individual




Fig 2. Variation of diurnality indices across the stages of the experiment. D-index for individual tuco-tucos (N = 9) along days outside and inside the respirometry chamber both with and without running-wheels. D-Index for each individual in the different conditions are connected by a line. Points in orange indicate the values for animal #143 which showed the highest discrepancy in D-Indices across conditions. Dashed line connects the values for animal #146, which was not submitted to the “chamber without wheel” condition.

doi:10.1371/journal.pone.0140500.g002

showed a particularly dramatic change in the D-Index, switching from -0.83 outside the chamber to +0.83 inside the chamber in the presence of the wheel (indicated by orange). Finally, data of one individual that was not measured without wheel was included, highlighted with a dashed line. Differences in D-Indices among the three conditions were statistically significant ($p < 0.001$).

$\dot{V}O_2$ followed the same rhythmic patterns as general activity and T_b (Fig 1). In addition to the daily variation, $\dot{V}O_2$ periodically peaked for episodes of more than one hour corresponding to bouts of high general activity and T_b . Mean $\dot{V}O_2$ of tuco-tucos was $1.305 \pm 0.073 \text{ mL g}^{-1} \text{ h}^{-1}$. Mean $\dot{V}O_2$ of females ($1.235 \pm 0.060 \text{ mL g}^{-1} \text{ h}^{-1}$, N = 5) and males ($1.384 \pm 0.151 \text{ mL g}^{-1} \text{ h}^{-1}$, N = 4) did not significantly differ ($p > 0.05$). In S1 Table, we present the mean values of $\dot{V}O_2$ and T_b for each individual, during days with and without access to running wheels.

Total amount of wheel-running revolutions per day was significantly reduced when animals were housed in the respirometry chamber ($p < 0.001$). Whereas all individuals completed > 5000 wheel revolutions/day outside the respirometry chamber, only one displayed such intense running while inside the chamber (Fig 3A). Daily amount of wheel-running correlated negatively with D-index ($r = -0.73$, $p < 0.001$) with the most strongly nocturnal animals displaying the greatest amount of wheel-running (Fig 3A). Lower wheel-running and associated phase inversion occurred both in the animals exposed to the wheel immediately upon being placed in the respirometry chamber and in those animals that were provided a wheel after three days in respirometry chamber. Despite the drastic decrease in wheel-running activity, mean daily general activity, T_b and $\dot{V}O_2$ did not differ significantly among conditions ($p > 0.05$) and neither correlated with D-Indexes ($p > 0.05$) (Fig 3B, 3C and 3D, and S1 Table). Finally, no effect was observed in association to the order of wheel/non-wheel stages, in our protocol.

Discussion

Despite showing day-time activity under field conditions, tuco-tucos consistently display nocturnal patterns when housed in the laboratory irrespective of access to running-wheels [8–10].




Fig 3. Wheel-running, mean T_b and mean Oxygen consumption of tuco-tucos in relation to diurnality indexes. Measurements for each individual ($N = 9$), across the stages of the experiment including days outside (white squares) and inside (black square) the respirometry chamber both with and without (grey squares) running-wheels. (A) Mean daily wheel-running levels are associated with nocturnality. (B) Mean body temperatures during each stage. There is no clear correlation with D-Indices. (C) Mean $\dot{V}O_2$ during each stage. There is no clear correlation with D-Indices. (D) Mean amount of general activity per day during each stage. There is no clear correlation with D-Indices.

doi:10.1371/journal.pone.0140500.g003

In the present study, we report the first displays of diurnality in the lab, which occurred exclusively during our respirometry experiment (Fig 1). Some individuals in the new environment of the sealed respirometry chamber completely suppressed running-wheel activity and switched to diurnality as revealed by T_b , $\dot{V}O_2$ and general activity rhythms; while others remained nocturnal as usual in the laboratory and continued to run on the wheel (Fig 3).

The search for the critical factors which trigger the nocturnality/diurnality switch observed in other species often converge upon the issue of the meaning of the running-wheel activity in the laboratory [34–37]. In some rodent species, all individuals are diurnal in the field whereas in the laboratory some become nocturnal while others remain diurnal. Interestingly, when offered unrestricted access to running wheels, some diurnal individuals become nocturnal (grass rats, *Arvicantha niloticus* [38]; degus, *Octodon degus* [39]; and mongolian gerbils, *Meriones unguiculatus* [4]). This phenomenon has been associated with a line of research devoted to investigating the “effect of” vigorous wheel-running on the period and phase of circadian activity rhythms [39–45]. Phase shifts in the free-running suprachiasmatic nuclei (SCN) due to “pulses” of intense running wheel activity are reported but are of very small magnitude [41,46]. Downstream from the SCN, however, wheel-running activity could act directly on the activity/inactivity signaling between the SCN and locomotor centers, as proposed by Kas and Edgar (1999). Their proposal was based on investigations of degus, a species that is known to switch phase from nocturnal to diurnal activity when provided access to a running wheel while in DD yet without any change to the basic free-running rhythm period.




Fig 4. Schematic view of different phase switch patterns associated with the presence of running wheels. Based on: [1,39,56,57] for *Octodon degus*; [2,38] for *Arvicanthis niloticus*; [3,58] for *Acomys russatus*; [4] for *Meriones unguiculatus*; [9,10] for *Ctenomys aff. knighti*. *For each species, field data were collected using different methods and do not necessarily reflect activity patterns of whole populations.

doi:10.1371/journal.pone.0140500.g004

The spontaneous suppression of wheel-running activity was displayed by all individuals that switched to diurnality (i.e., animals which showed D-Index>0.5) when exposed to the new environment of the respirometry chamber (Fig 3A). This phenomenon occurred in both of our trials in two consecutive years. It is noteworthy that general motor activity was maintained and switched to a diurnal pattern in all individuals that stopped running on the wheel (Fig 1). General activity levels did not change upon placement in the respirometry chamber, even in those animals that suppressed wheel-running. It is conceivable that this counterintuitive fact is due to the intensification of non-wheel as “digging-like” movements, intense grooming and other non-specific movements, which are generally observed when tuco-tucos are deprived of wheels.

Our finding of a phase inversion (nocturnal to diurnal) in tuco-tucos when housed within a respirometry chamber illustrates a novel association between running-wheels and timing of activity not observed in any of the previous work on degus, grass rats and Mongolian gerbils. In common with the above species, the greatest levels of activity are always associated with nocturnality (Fig 4). It has been proposed that a shift to nocturnality in response to elevated activity is associated to thermoregulation by consolidating the activity during a time of day when body temperature is naturally lower in diurnal species [39],[47–49]. However, this seems to be unlikely in tuco-tucos, based on our previous finding [10], that activity has a greater impact on body temperature during the dark phase, suggesting that the allocation of activity during the night would decrease, instead of enhance, heat loss.

Although phase inversion inside the respirometry chamber occurs concomitantly with suppression of wheel-running behavior, it is not “a response” to removal of the wheel (Fig 3) because when the wheel is removed from the respirometry chamber the nocturnal individuals do not switch to diurnality. Robust nocturnal patterns have been previously observed in 100% of 18 animals with wheels [8,9] and of 5 without wheels [9]. Additionally, wheel removal

without switch in activity phase in tuco-tucos has been reported before in 100% of 6 animals, during experiments performed in other contexts [10], done in the same laboratory, under the same standard conditions (housing, feeding schedules, temperature, humidity).

Interestingly, both the most extreme diurnal and the most extreme nocturnal D-indices (Fig 2) are associated with the animals having access to a wheel while in the respiratory chamber. Because most animals didn't actually use the wheel to run, this indicates that the wheel may also have a running-independent effect on the temporal pattern of activity. A small body of literature shows that the mere presence of the wheel in the environment can have behavioral and neurogenic effects [50]. Mice kept in an environment in which a locked wheel is present show less anxiety and enhanced fear memory than those kept in a cage without the wheel [51–52]. In our case, however, the suppression of running occurred spontaneously, in contrast to locking the wheel. This opens even more possibilities for future studies on running-independent effects of the wheel.

Diurnal/nocturnal switches in the laboratory occur so rapidly that it has been argued that this flexibility might represent an adaptive mechanism to sudden changes in the species' natural environment [38, 39]. The switches in activity timing in tuco-tucos were triggered when inserted into a respirometry chamber (Fig 4) where tuco-tucos face mild alteration of gas composition of the ambient air (< 1% decrease in O₂ and < 0.5% increase in CO₂). A survey of the literature of the effects of O₂ and CO₂ content of air on circadian patterns reveal mostly changes in amplitude, with rhythmic depression as a consequence of hypoxia or hypercapnia in rats [53, 54]. However, minute phase changes have been observed in free-running golden hamsters exposed to pulses of hypoxic air [55]. These studies involved more extreme hypoxia/hypercapnia than faced by our tuco-tucos in the respirometry chamber and they were conducted with non-subterranean animals. It is conceivable that tuco-tucos, which live in sealed underground tunnels, are able to detect even small changes in gas composition and/or humidity. Perceived changes in the gas composition of the environment could serve as a triggering mechanism to incite an alertness response needed for predator avoidance or tunnel maintenance and, possibly, lead to changes in the temporal pattern of activity, as suggested by our results in the sealed chamber.

Several interesting insights have emerged from our simultaneous measurements of the interconnected $\dot{V}O_2$, T_b, general motor and wheel-running rhythms. Our results clearly demonstrate that switches in timing of activity phase can occur concomitantly with spontaneous suppression of wheel-running. Apparently, in tuco-tucos activity timing and wheel-running associations appear in a reformulated perspective.

Supporting Information

S1 Fig. Respirometry chamber and schematic illustration of the experimental protocol. (A) Photography of the respirometry chamber without the running wheel. The chamber consists in a standard home cage with an acrylic lid with fittings to allow the airflow. The chamber was kept in a light-tight cabinet, which was the same used in the non-respirometry steps of the experiments. (B) Scheme of the experimental protocol. At first, the animal was kept in its home cage with access to a running-wheel. Then, it was placed in the respirometry chamber. One group was put in a chamber with running-wheel and the other in a chamber without a wheel. The group that started with the wheel would then have it removed, while the other would have the wheel added to the chamber. After the respirometry trials, measurements would continue in a standard home cage.

(DOCX)

S2 Fig. Scheme of the respirometry system.

(DOCX)

S1 Table. Summary of the variables measured under different conditions, for each individual. ¹In mL g⁻¹h⁻¹, represented as mean±SD. ²In °C. ³Mean total daily revolutions.

(DOCX)

Acknowledgments

The authors thank Danilo Flôres for valuable discussions and help during experimental measurements, Melanie Richter and Barbara Joos for help with the respirometry setup, Jose Paliza for technical assistance and Vinícius Dokkedal for assisting with data collection. All the reviewers for criticisms and suggestions that greatly improved the manuscript.

Author Contributions

Conceived and designed the experiments: PT VSV CLB GAO. Performed the experiments: PT VSV CLB. Analyzed the data: PT OT. Wrote the paper: PT CLB GAO. Designed the respirometry setup: OT.

References

1. Labyak SE, Lee TM, Goel N. Rhythm chronotypes in a diurnal rodent, *Octodon degus*. *Am J Physiol*. 1997; 273:R1058–R1066. PMID: [9321886](#)
2. Blanchong JA, Smale L. Temporal patterns of activity of the unstriped Nile rat, *Arvicanthis niloticus*. *J Mammal*. 2000; 81(2):595–599.
3. Levy O, Dayan T, Kronfeld-Schor N. The relationship between the golden spiny mouse circadian system and its diurnal activity: an experimental field enclosure and laboratory study. *Chron Int*. 2007; 24(4):599–613.
4. Weinert D, Weinandy R, Gatterman R. Photic and non-photic effects on the daily activity pattern of Mongolian gerbils. *Physiol and Behav*. 2007; 90:325–333.
5. Barak O, Kronfeld-Schor N. Activity rhythms and masking response in the diurnal fat sand rat under laboratory conditions. *Chron Int*. 2013; 30(9):1123–1134.
6. Daan S, Spoelstra K, Albrecht U, Schmutz I, Daan M, Daan B, et al. Lab mice in the field: unorthodox daily activity and effects of a dysfunctional circadian clock allele. *J of Biol Rhythms* 2011; 26(2):118–129.
7. Hut RA, Kronfeld-Schor N, van der Vinne V, De la Iglesia H. In search of a temporal niche: environmental factors. *Prog Brain Res*. 2012; 199:281–304. doi: [10.1016/B978-0-444-59427-3.00017-4](#) PMID: [22877672](#)
8. Valentinuzzi VS, Oda GA, Araújo JF, Ralph MR. Circadian pattern of wheel-running activity of a South American subterranean rodent (*Ctenomys cf knightii*). *Chronobiol Int*. 2009; 26(1):14–27. doi: [10.1080/07420520802686331](#) PMID: [19142755](#)
9. Tomotani BM, Flôres DEFL, Tachinardi P, Paliza JD, Oda GA, Valentinuzzi VS. Field and laboratory studies provide insights into the meaning of day-time activity in a subterranean rodent (*Ctenomys aff. knightii*), the tuco-tuco. *PLoS ONE*. 2012; 7, e37918. doi: [10.1371/journal.pone.0037918](#) PMID: [22649565](#)
10. Tachinardi P, Bicudo JEW, Oda GA, Valentinuzzi VS. Rhythmic 24 h variation of core body temperature and locomotor activity in a subterranean rodent (*Ctenomys aff. knightii*), the tuco-tuco. *PLoS ONE*. 2014; 9, e85674. doi: [10.1371/journal.pone.0085674](#) PMID: [24454916](#)
11. Hut RA, Pilon V, Boerema AS, Strijkstra AM, Daan S. Working for food shifts nocturnal mouse activity into the day. *PLoS ONE*. 2011; 6(3), e17527. doi: [10.1371/journal.pone.0017527](#) PMID: [21479166](#)
12. Daan S. Adaptive daily strategies in behavior. In: Aschoff J, editor. *Handbook of behavioral neurobiology vol 4—Biological rhythms*. New York: Plenum Press; 1981. pp. 275–298.
13. Kenagy GJ, Vleck D. Daily temporal organization of metabolism in small mammals: adaptation and diversity. In: Aschoff J, Daan S, Groos GA, editors. *Vertebrate circadian systems*. Berlin: Springer-Verlag; 1982. pp. 322–338.

14. Halle S. Ecological relevance of daily activity patterns. In: Halle S, Stenseth NC, editors. Activity patterns in small mammals. Berlin, Heidelberg: Springer-Verlag; 2000. pp. 67–90.
15. Ruf T, Heldmaier G. Djungarian hamsters—small graminivores with daily torpor. In: Halle S, Stenseth NC, editors. Activity patterns in small mammals. Berlin, Heidelberg: Springer-Verlag; 2000. Pp. 217–243.
16. Kronfeld-Schor N, Shargal E, Haim A, Zisapel N, Heldmaier G. Temporal partitioning among diurnally and nocturnally active desert mice: energy and water turnover costs. *J Thermal Biol.* 2001; 26:139–142.
17. Van der Vinne V, Sjaak JR, Gorter JA, Eijer WG, Sellix MT, Menaker M, et al. Cold and hunger induce diurnality in a nocturnal mammal. *PNAS.* 2014; 1413135111.
18. Lacey EA, Patton JL, Cameron GN. Life underground: the biology of subterranean rodents. Chicago: University of Chicago Press; 2000.
19. Buffenstein R. Ecophysiological responses of subterranean rodents to underground habitats. In: Lacey EA, Cameron G, Patton JL, editors. Life underground: the biology of subterranean rodents. Chicago: University of Chicago Press; 2000. pp. 183–226.
20. Luna F, Antinuchi CD, Busch C. Digging energetics in the South American rodent *Ctenomys talarum* (Rodentia, Ctenomyidae). *Can J Zool.* 2002; 80 (12):2144–2149.
21. Luna F, Antinuchi CD. Energetics and thermoregulation during digging in the rodent tuco-tuco (*Ctenomys talarum*). *Comp Biochem Physiol A* 2007; 146(4):559–564.
22. Luna F, Antenucci CD, Bozinovic F. Comparative energetics of the subterranean *Ctenomys* rodents: breaking patterns. *Physiol Biochem Zool.* 2009; 82(3):226–235. doi: [10.1086/597526](https://doi.org/10.1086/597526) PMID: [19327041](https://pubmed.ncbi.nlm.nih.gov/19327041/)
23. Perissinotti PP, Antenucci CD, Zenuto R, Luna F. Effect of diet quality and soil hardness on metabolic rate in the subterranean rodent *Ctenomys talarum*. *Comp Biochem Physiol A.* 2009; 154: 298–307.
24. Flôres DEFL, Tomotani BM, Tachinardi P, Oda GA, Valentinuzzi VS. Modeling natural photic entrainment in a subterranean rodent (*Ctenomys* aff. *knighti*), the tuco-tuco. *PLoS ONE* 2013; 8(7), e68243. doi: [10.1371/journal.pone.0068243](https://doi.org/10.1371/journal.pone.0068243) PMID: [23874562](https://pubmed.ncbi.nlm.nih.gov/23874562/)
25. Burda H, Sumner R, Begall S (2007) Microclimate in burrows of subterranean rodents—revisited. In: Begall S, Burda H, Schleich CE, editors. Subterranean rodents: News from underground. Heidelberg: Springer-Verlag; 2007. pp.21–23.
26. Riccio AP, Goldman BD. Circadian rhythms of body temperature and metabolic rate in naked mole-rats. *Physiol Behav.* 2000; 71:16–22.
27. Sikes RS, Gannon WL. The animal care and use committee of the American Society of Mammalogists. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *J Mamm.* 2011; 92(1):235–253.
28. National Research Council (US) Committee for the Update of the Guide for the Care and Use of Laboratory Animals. Guide for the Care and Use of Laboratory Animals. 8th edition. Washington (DC): National Academies Press (US). 2011; Available from: <http://www.ncbi.nlm.nih.gov/books/NBK54050/>. Accessed 2014 September 17.
29. Busch C. Metabolic rate and thermoregulation in two species of tuco-tuco, *Ctenomys talarum* and *Ctenomys australis* (Caviomorpha, Octodontidae). *Comp Biochem and Physiol Part A: Physiology* 1989; 93(2):345–347.
30. Tøien Ø. Automated open flow respirometry in continuous and long-term measurements: design and principles. *J Appl Physiol.* 2013; 114(8):1094–1107. doi: [10.1152/jappphysiol.01494.2012](https://doi.org/10.1152/jappphysiol.01494.2012) PMID: [23349455](https://pubmed.ncbi.nlm.nih.gov/23349455/)
31. Withers PC. Measurement of VO₂, VCO₂, and evaporative water loss with a flow-through mask. *J Appl Physiol.* 1977; 42:120–123. PMID: [833070](https://pubmed.ncbi.nlm.nih.gov/833070/)
32. Hoogenboom I, Daan S, Dallinga JH, Schoenmakers M. Seasonal change in the daily timing of behavior of the common vole, *Microtus arvalis*. *Oecologia* 1984; 61(1):18–31.
33. R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0. R-Project website. 2010; Available: <http://www.R-project.org>. Accessed 2014 September 17.
34. Mather JG. Wheel-running activity: a new interpretation. *Mamm Rev.* 1981; 11(1):41–51.
35. Sherwin CM. Voluntary wheel running: a review and novel interpretation. *Animal Behav.* 1998; 56(1):11–27.
36. Novak CM, Burghardt PR, Levine JA. The use of a running wheel to measure activity in rodents: relationship to energy balance, general activity, and reward. *Neurosci Biobehav Rev.* 2012; 36:1001–1014. doi: [10.1016/j.neubiorev.2011.12.012](https://doi.org/10.1016/j.neubiorev.2011.12.012) PMID: [22230703](https://pubmed.ncbi.nlm.nih.gov/22230703/)
37. Meijer JH, Robbers Y. Wheel running in the wild. *Proc Royal Soc B.* 2014; 281: 20140210.

38. Blanchong JA, McElhinny TL, Mahoney MM, Smale L. Nocturnal and diurnal rhythms in the unstriped Nile rat, *Arvicanthis niloticus*. *J Biol Rhythms* 1999; 14:364–377. PMID: [10511004](#)
39. Kas MJH, Edgar DM. A nonphotic stimulus inverts the diurnal–nocturnal phase preference in *Octodon degus*. *J of Neurosci*. 1999; 19 (1):328–333.
40. Yamada N, Shimoda K, Ohi K, Takahashi S, Takahashi K. Free-access to a running wheel shortens the period of free-running rhythm in blinded rats. *Physiol Behav*. 1988; 42:87–91. PMID: [3387483](#)
41. Reeb SG, Mrosovsky N. Effects of induced wheel running on the circadian activity rhythms of syrian hamsters: entrainment and phase response curve. *J Biol Rhythms* 1989; 4(1):39–48. PMID: [2519579](#)
42. Van Reeth O, Turek FW. Stimulated activity mediates phase shifts in the hamster circadian clock induced by dark pulses or benzodiazepines. *Nature* 1989; 339: 49–51. PMID: [2654641](#)
43. Edgar DM, Martin CE, Dement WC. Activity feedback to the mammalian circadian pacemaker: influence on observed measures of rhythm period length. *J Biol Rhythms* 1991; 6(3):185–199. PMID: [1773091](#)
44. Mrosovsky N. A non-photic gateway to the circadian clock of hamsters. In: Ciba Foundation Symposium 183. Circadian clocks and their adjustment. Chichester: Wiley. 1995; pp. 154–174.
45. Redlin U. Neural basis and biological function of masking by light in mammals: suppression of melatonin and locomotor activity. *Chronobiol Int*. 2001; 18(5):737–758. PMID: [11763983](#)
46. Reeb SG, Mrosovsky N. Large phase-shifts of circadian rhythms caused by induced running in a re-entrainment paradigm: the role of pulse duration and light. *J Comp Physiol A*. 1989; 165:819–825. PMID: [2810152](#)
47. Refinetti R. Rhythms of body temperature and temperature selection are out of phase in a diurnal rodents, *Octodon degus*. *Physiol and Behav*. 1996; 60(3):959–961.
48. Kenagy GJ, Nespolo RF, Vásquez RA, Bozinovic F. Daily and seasonal limits of time and temperature to activity of degus. *Rev Chilena Hist Nat*. 2002; 75:567–581.
49. Kenagy GJ, Vásquez RA, Barnes BM, Bozinovic F. Microstructure of activity bouts of degus in a thermally heterogeneous habitat. *J Mammal* 85(2): 260–267.
50. Bednarczyk MR, Hacker MC, Fortin-Nunez LC, Aumont A, Bergeron R, Fernandes KJ. Distinct stages of adult hippocampal neurogenesis are regulated by running and the running environment. *Hippocampus*. 2011; 21:1334–1347. doi: [10.1002/hipo.20831](#) PMID: [20623741](#)
51. Dubreucq S, Marsicano G, Chaouloff F. Emotional consequences of wheel running in mice: which is the appropriate control? *Hippocampus*. 2011; 21:239–242. doi: [10.1002/hipo.20778](#) PMID: [20232385](#)
52. Dubreucq S, Marsicano G, Chaouloff F. Duration and environment-dependent effects of repeated voluntary exercise on anxiety and cued fear in mice. *Behav Brain Res*. 2015; 282: 1–5. doi: [10.1016/j.bbr.2014.12.042](#) PMID: [25546723](#)
53. Mortola JP, Seifert EL. Hypoxic depression of circadian rhythms in adult rats. *J Appl Physiol*. 2000; 88:365–368. PMID: [10657999](#)
54. Bishop B, Silva G, Krasney J, Salloum A, Roberts A, Nakano H, et al. Circadian rhythms of body temperature and activity levels during 63h of hypoxia in the rat. *Am J Physiol*. 2000; 279:R1378–R1385.
55. Jarsky TM, Stephenson R. Effects of hypoxia and hypercapnia on circadian rhythms in the golden hamster (*Mesocricetus auratus*). *J Appl Physiol*. 2000; 89:2130–2138. PMID: [11090559](#)
56. Fulk GW. Notes on the activity, reproduction and social behavior of *Octodon degus*. *J Mammal*. 1976; 57:495–505.
57. Hagenauer MH, Lee TM. Circadian organization of the diurnal Caviomorph rodent, *Octodon degus*. *Biol Rhythm Res*. 2008; 39(3):269–289.
58. Cohen R, Smale L, Kronfeld-Schor N. Plasticity of circadian activity and body temperature rhythms in golden spiny mice. *Chronobiol Int*. 2009; 26:430–446. doi: [10.1080/07420520902820939](#) PMID: [19360488](#)