

Comprehensive analysis of risk factors associating with Hepatitis B virus (HBV) reactivation in cancer patients undergoing cytotoxic chemotherapy

W Yeo^{*,1}, B Zee¹, S Zhong¹, PKS Chan², W-L Wong¹, WM Ho¹, KC Lam¹ and PJ Johnson¹

¹Department of Clinical Oncology, Sir Y.K. Pao Centre for Cancer, Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, NT, Hong Kong;

²Department of Microbiology, Sir Y.K. Pao Centre for Cancer, Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, NT, Hong Kong

For cancer patients with chronic hepatitis B virus (HBV) infection, who receive cytotoxic chemotherapy, HBV reactivation is a well-described complication, which may result in varying degrees of liver damage. Several clinical features and the pre-chemotherapy HBV viral load have been suggested to be associated with an increased risk of developing the condition: (1) to assess the clinical and virological factors in a comprehensive manner and thereby identify those that are associated with the development of HBV reactivation; (2) to develop a predictive model to quantify the risk of HBV reactivation. In all, 138 consecutive cancer patients who were HBV carriers and undergoing chemotherapy were studied, of which 128 patients had sera available for real-time PCR HBV DNA measurement. They were followed up throughout their course of chemotherapy and the HBV reactivation rate was determined. The clinical and virological features between those who did and did not develop viral reactivation were compared. These included age, sex, baseline liver function tests, HBeAg status and viral load (HBV DNA) prior to the chemotherapy, and the use of specific cytotoxic agents. In all, 36 (26%) developed HBV reactivation. Multivariate analysis revealed pre-chemotherapy HBV DNA level, the use of steroids and a diagnosis of lymphoma or breast cancer to be significant factors. Based on real-time HBV DNA PCR assay, detectable baseline HBV DNA prior to the administration of cytotoxic chemotherapy, the use of steroids and a diagnosis of lymphoma or breast cancer are predictive factors for the development of HBV reactivation. A predictive model was developed from the current data, based on a logistic regression method.

British Journal of Cancer (2004) **90**, 1306–1311. doi:10.1038/sj.bjc.6601699 www.bjcancer.com

Published online 9 March 2004

© 2004 Cancer Research UK

Keywords: hepatitis B virus reactivation; chemotherapy; risk factors

For cancer patients who have chronic hepatitis B virus (HBV) infection, there is a high rate of hepatic complications during cytotoxic chemotherapy, and this has mainly been attributable to HBV reactivation. The condition is manifested with abnormal liver function tests that show a hepatic picture, and it is confirmed by raised levels of serum HBV DNA. The clinical spectrum ranges from asymptomatic hepatitis to fatal hepatic failure (Galbraith *et al*, 1975; Hoofnagle *et al*, 1982; Lok *et al*, 1991; Nokamura *et al*, 1996; Kumagai *et al*, 1997; Markovic *et al*, 1999; Yeo *et al*, 2000a). However, even in its mildest form with spontaneous recovery, a patient's prognosis from cancer may still be impaired from the disruption in chemotherapeutic administration with treatment delay, or premature termination of the anticancer therapy. The incidence of HBV reactivation in hepatitis B surface antigen (HBsAg) seropositive cancer patients undergoing cytotoxic chemotherapy has been reported to be 20% or higher (Lok *et al*, 1991; Nokamura *et al*, 1996; Kumagai *et al*, 1997; Markovic *et al*, 1999; Yeo *et al*, 2000a). No preventive measures have been proven to prevent or reduce the incidence of HBV reactivation, although

more recent reports have suggested that the prophylactic use of the antiviral agent lamivudine, prior to the start of chemotherapy, may reduce the occurrence of the condition (Rossi *et al*, 2001; Liao *et al*, 2002; Lim *et al*, 2002; Persico *et al*, 2002; Shibolet *et al*, 2002; Yeo *et al*, 2002). There has also been concern about the emergence of viral mutant as a result of lamivudine therapy, and, to date, limited data are available on the clinical impact of these mutants in immunosuppressed subjects. Limiting the use of the prophylactic antiviral to patients who are at the highest risk of developing viral reactivation may reduce the potential complication of mutant emergence, and at the same time be more cost-effective.

Despite the wide recognition of HBV reactivation, there is still no consensus as to the associated risk factors. Clinical features such as young age, male sex, the diagnosis of lymphoma and the use of anthracyclines and/or steroids as part of the anticancer therapy have been suggested (Lok *et al*, 1991; Yeo *et al*, 2000a), while virological factors such as HBeAg positivity (Yeo *et al*, 2000a, b) and, more recently, the pre-chemotherapy HBV viral load (Lau *et al*, 2002) have also been associated.

The objectives of this study were: (1) to assess the clinical and virological factors in a comprehensive manner in order to identify those that are associated with the development of HBV reactivation; and (2) to develop a predictive model to identify patients more likely to develop HBV reactivation.

*Correspondence: Dr W Yeo; E-mail: winnieyeo@cuhk.edu.hk

Received 8 August 2003; revised 11 November 2003; accepted 14 November 2003; published online 9 March 2004

PATIENTS AND METHODS

In all, 138 consecutive cancer patients who were chronic HBV carriers and planned for cytotoxic chemotherapy were consented and followed up throughout their course of treatment. The study was approved by the Clinical Research Ethics Committee of the Chinese University of Hong Kong.

Investigations

Prior to study entry, the following investigations were undertaken in all patients: hepatitis B s-antigen (HBsAg), hepatitis B e-antigen/antibody (HBeAg/anti-HBe) HBV DNA level (measured by commercially available Quantiplex HBV DNA assay (bDNA), Chiron, USA), complete blood picture (CBP, which included haemoglobin, red cell count, mean cell volume, white cell count and differential and platelet count), renal function tests (RFT, which included sodium, potassium, urea, creatinine and creatinine clearance), liver function test (LFT, which included total protein, albumin, total bilirubin, alanine transaminase, alkaline phosphatase) and clotting profile. In addition, one 3-ml serum sample was collected for later retrospective analysis of HBV DNA by a real-time PCR assay (Zhong *et al*, in press).

During the course of chemotherapy, on days 1 and 10 of each cycle, CBP, clotting profile, RFT and LFT were monitored together with clinical signs and symptoms. Monitoring of these parameters was continued for 8 weeks after completion of chemotherapy.

When a patient was found to have developed hepatitis (as defined below) during the course of chemotherapy, HBV DNA was performed (measured by Quantiplex bDNA assay), with immunoglobulin M (IgM) antibody to hepatitis A virus (IgM anti-HAV), HCV RNA, anti-HDV, ANA and other investigations as clinically indicated.

Hepatitis serology

Hepatitis A, B and D markers were detected by commercial enzyme immunoassays (Cobas Core Anti-HAV IgM EIA, Roche Diagnostics GmbH, Deutschland; Auszyme MC Dynamic (HBsAg), Abbott Laboratories, USA; Cobas Core Anti-HBc IgM EIA, Roche Diagnostics GmbH, Deutschland; MONOLISA HBe, Sanofi Diagnostics Pasteur, USA; Abbott Anti-Delta EIA, Abbott Laboratories, USA). HCV RNA was detected by reverse transcription polymerase chain reaction using primers 209, 211, 939 and 940 as previously described (Chan *et al*, 1992).

For routine use including establishment of the diagnosis of HBV reactivation, HBV DNA level was measured by using the branched DNA hybridisation assay (Quantiplex HBV DNA assay (bDNA), Chiron, USA), which has a lower detection limit of 0.7×10^6 genome equivalent ml⁻¹.

Real-time HBV DNA PCR assay

For the purpose of assessing the impact of viral load on HBV reactivation, we used the real-time PCR to quantify serum HBV DNA, which has a lower detection limit for HBV DNA of 2.9×10^3 g.e. ml⁻¹ (Zhong *et al*, in press). In total, 128 patients had sera available for real-time PCR analysis and these were tested in a single batch. Their serum samples were separated as soon as the blood was coagulated and stored at -80°C in RNase-free tubes. Isolation of nucleic acids was performed using the High Pure Viral Nucleic Acid Kit (Roche, Mannheim, Germany). For each isolation, 200 μl of serum was used resulting in 50 μl of nucleic acid extract. For HBV DNA quantification by real-time PCR, PCR amplification was performed with PCR primers located at regions that are universally conserved among the six HBV genotypes (A–F) corresponding to the HBV X gene. The oligonucleotide sequences of primers were: txs3 (1434–154), TCT CAT CTG CCG GAC CGT

GT; xas1 (1668–148), AAT TTA TGC CTA CAG CCT CC. A volume of 5 μl equivalent of serum was used for real-time amplification in a final 10 μl reaction volume, using $1 \times$ Fast Start SYBR Green I Master Mix (Roche), MgCl₂ (4 mM) and 0.3 μM concentration of each primer. The amplification procedure, utilising the LightCycler instrument (Roche), was as follows: 95°C for 10 min, followed by 45 cycles of 95°C for 15 s, 57°C for 10 s and 72°C for 12 s, followed by denaturation of amplification samples by slow increase of temperature (0.1°C s⁻¹) up to 95°C.

Definition of HBV reactivation

The following definitions, based on a definition of Lok *et al* (1991), and subsequently modified by us (Yeo *et al*, 2000a), were applied. 'Hepatitis' was defined as a threefold or greater increase in serum ALT level that exceeded the reference range (>58 iu l⁻¹) or an absolute increase of ALT to over 100 iu l⁻¹. 'HBV reactivation' was defined as either one of the following: using the Quantiplex b DNA assay, Chiron assay, an increase in HBV DNA levels of 10-fold or greater when compared with the baseline level, or an absolute increase of HBV DNA level that exceeded 1000×10^6 g.e. ml⁻¹ during chemotherapy, in the absence of clinical or laboratory features of acute infection with hepatitis A, C and delta virus or other systemic infections.

Comprehensive assessment of the clinical and virological factors in association with the development of HBV reactivation

Among the 128 patients who had sera available for the real-time PCR analysis for HBV DNA, the following factors were compared between those who did and those who did not develop viral reactivation: age, sex, tumour type, use of steroids, anthracyclines and 5-fluorouracil, pre-chemotherapy liver functions (albumin, total bilirubin and alanine transaminase), HBeAg status, HBV DNA using real-time PCR measurement.

Statistical methods

Univariate analysis for detecting significant prognostic factors for HBV reactivation was done using χ^2 test or Fisher's exact test. Prognostic factors for HBV reactivation were then determined by multivariate analysis using a stepwise logistic regression model based on a backward elimination procedure for model selection, with a significance level of <0.1 for a factor to enter into the logistic model; the same significance level was used to retain variables in the final model. A predictive model was obtained using the parameters of the final logistic model. The classification to the HBV reactivation group was based on the probability estimate from the logistic model, where an optimal cutoff point was determined using the ROC curve technique; the ROC curve refers to the receiver operating characteristic curve, which is a plot of the true-positive rate vs false-positive rate associated with the classification rules for all possible choices of critical values (Hanley, 1989).

RESULTS

In all, 36 patients were determined to have developed HBV reactivation (the 'reactivation' group) and 92 patients were determined not to have HBV reactivation (the 'non-reactivation' group) (Table 1).

There were 56 males and 72 females; 16 (28.6%) males and 20 (27.8%) females developed HBV reactivation ($P=1.000$). Between the 'reactivation' and the 'non-reactivation' groups, the median ages were 46.5 and 49.0 years, respectively ($P=0.265$). There were 39 patients with breast cancer, 29 with gastrointestinal

Table 1 Characteristics of the 128 HBsAg seropositive cancer patients undergoing cytotoxic chemotherapy

	Patients who had hepatitis B viral reactivation during chemotherapy	Patients who did not develop hepatitis B viral reactivation during chemotherapy	P-value
Total no. of patients	36	92	
Sex*			
Male	16 (28.6%)	40 (71.4%)	1.0000
Female	20 (27.8%)	52 (72.2%)	
Median age (years)	46.5	49	0.2654
Range	30–71	20–78	
Tumour type*			
Breast cancers	16 (41.0%)	23 (59.0%)	0.0041
Lymphomas	7 (58.3%)	5 (41.7%)	
Gastrointestinal cancers	2 (6.9%)	27 (93.1%)	
Head and neck cancers	5 (29.4%)	12 (70.6%)	
Lung cancers	3 (23.1%)	10 (76.9%)	
Other cancers	3 (16.6%)	15 (83.4%)	
Pretreatment (baseline) biochemistry			
Median ALT levels (normal <58 iu l ⁻¹)	35 (range: 12–77)	27 (range: 10–96)	0.1025
Median total bilirubin levels (normal <15 µmol l ⁻¹)	7 (range: 2–26)	7 (range: 1–108)	0.9639
Median albumin levels (normal >40 g l ⁻¹)	35 (range: 22–43)	36 (range: 20–47)	0.8421
Use of anthracyclines			
Yes	19 (46.3%)	22 (53.7%)	0.0029
No	17 (19.5%)	70 (80.5%)	
Use of steroids			
Yes	22 (39.3%)	34 (61.2%)	0.0174
No	14 (19.4%)	58 (80.6%)	
Use of fluorouracil			
Yes	14 (23.0%)	47 (77.0%)	0.2419
No	22 (32.8%)	45 (67.2%)	
HBeAg status*			
Positive	4 (26.7%)	11 (73.3%)	1.0000
Negative	32 (28.3%)	81 (71.7%)	
HBV DNA-real-time PCR assay*			
Detectable	31 (37.8%)	51 (62.2%)	0.0010
Undetectable	5 (10.9%)	41 (89.1%)	

*Fisher's exact two-sided test.

malignancies, 17 head and neck cancers, 13 lung cancers, 12 non-Hodgkin's lymphoma and 18 other malignancies. Among these patients, 16 who had breast cancers (41.0% of the patients with this tumour type, $P = 0.053$), seven non-Hodgkin's lymphoma (58.3%, $P = 0.037$), two gastrointestinal malignancies (6.9%, $P = 0.0041$), five head and neck (29.4%, $P = 1.000$), three lung cancers (23.1%, $P = 1.000$) and three (16.6%, $P = 0.396$) with other malignancies developed viral reactivation.

Of the 128 patients, 15 were HBeAg positive, of which four (26.7%) developed reactivation, while 32 of the 113 (28.3%) patients who were HBeAg negative/anti-HBe positive developed the condition ($P = 1.000$). Using the real-time PCR assay, 82 patients were detected to have HBV DNA, 31 (37.8%) of whom developed reactivation; in contrast, only five of the 41 (10.9%) patients who had undetectable HBV DNA developed the condition ($P = 0.001$).

There was no statistically significant difference in baseline liver function in terms of ALT, bilirubin and albumin between the two groups of patients. Out of 56 patients, 22 (39%) received steroids and developed reactivation; while 14 of 72 patients who did not

receive the agent developed the condition ($P = 0.017$). Out of 41 patients, 19 (46.34%) received anthracyclines and developed reactivation, while 17 of the 87 (19.54%) patients who had no anthracyclines developed the condition ($P = 0.0029$). With respect to 5-fluorouracil, 61 received the drug, among whom 14 (23%) developed reactivation, while 22 of 67 patients (33%) who did not receive the agent developed the condition ($P = 0.2419$).


On multivariate analysis, the factors that were significantly associated with a higher risk of developing HBV reactivation were detectable HBV DNA levels based on real-time PCR measurement ($P = 0.0003$, Odds ratio of 8.4 with 95% CI from 2.6 to 27.2), the use of steroid ($P = 0.0465$, OR of 2.7 with 95% CI from 1.0 to 7.2), a diagnosis of lymphoma ($P = 0.0419$, OR of 5.0 with 95% CI from 1.1 to 23.5) and breast cancer ($P = 0.0041$, OR of 4.2 with 95% CI from 1.6 to 11.0).

Predictive model development

A predictive model was determined using the final logistic regression model, as shown in Table 2. The following is the

Table 2 Logistic regression model

Parameter	P-value	Odds ratio	95% confidence interval	
			Lower	Upper
Intercept	-3.6589 (β_0)			
Lymphoma (x1)	1.6086 (β_1)	0.0419	1.06	23.53
Breast (x2)	1.4264 (β_2)	0.0041	1.57	11.02
Steroid (x3)	0.9939 (β_3)	0.0465	1.02	7.19
HBV PCR (x4)	2.1339 (β_4)	0.0003	2.63	27.15

**Figure 1** Logistic model with PCR_HBV, steroid, lymphoma and breast sites.

mathematical formula that can be programmed into a personal computer or programmable calculator:

$$\log\left(\frac{p_R}{1-p_R}\right) = \beta_0 + \beta_1x_1 + \beta_2x_2 + \beta_3x_3 + \beta_4x_4$$

where p_R is the probability of HBV reactivation, x_1 , x_2 , x_3 and x_4 are indicator variables for the presence of lymphoma (presence = 1, absence = 0), breast cancer (presence = 1, absence = 0), the use of steroids (presence = 1, absence = 0) and HBV virology load determined by PCR (detectable = 1; not detectable = 0). The parameters β_0 , β_1 , β_2 , β_3 , β_4 are parameters whose values are given in Table 2. We would classify a patient to the group with a higher likelihood of developing HBV reactivation if the estimated p_R from the above logistic model is higher than 0.3. A sensitivity of 75% and a specificity of 79.3% were obtained based on a cutoff point of 0.3 for the predictive probability of the logistic model. This optimal predictive probability cutoff point was determined using the ROC curve method (Figure 1).

DISCUSSION

Hepatitis B virus reactivation during cytotoxic chemotherapy is a particularly important clinical issue in areas of the world such as China, where chronic HBV infection is endemic. With the increasing incidence of neoplastic diseases (Lok *et al*, 1991), and the more widespread use of cytotoxic chemotherapy, the occurrence of HBV reactivation is likely to increase further.

In this study, the identified risk factors included detectable pre-chemotherapy HBV DNA load (using real-time PCR measurement), the use of steroids, a diagnosis of lymphoma or breast cancer, another suggested factor being the use of anthracyclines. Previous studies have reported factors including male sex (Liaw *et al*, 1987; Liang *et al*, 1990; Lok *et al*, 1991), younger age (Yeo

et al, 2000a), HBeAg positivity (Lok *et al*, 1991; Yeo *et al*, 2000a,b) and the presence of lymphoma (Yeo *et al*, 2000a). Although the reason remains unknown, male sex has consistently been reported to be a risk factor for the exacerbation of chronic HBV infection in cancer patients undergoing chemotherapy, as well as in non-cancer subjects (Liaw *et al*, 1987; Liang *et al*, 1990; Lok *et al*, 1991). Although HBeAg positivity in immunocompromised cancer patients appears to be a risk factor for HBV reactivation (Liang *et al*, 1990; Yeo *et al*, 2000b), this has not been found to be universally the case (Lok *et al*, 1991; Nokamura *et al*, 1996), and an increased risk may be partly attributed to the presence of the pre-core/core promoter HBV mutant (i.e. HBeAg negative/anti-HBe positive), which had been associated with severe fulminant hepatitis (Omata *et al*, 1991; Ehata *et al*, 1993; Liang *et al*, 1994; Nokamura *et al*, 1996; Steinberg *et al*, 2000; Yeo *et al*, 2000a,b). In addition, consistent with other reports, the baseline (pretreatment) liver function including ALT, total bilirubin and albumin levels did not appear to be associated with the development of HBV reactivation.

Certain limitations are noted in this study. Of the 82 patients found to have detectable HBV DNA on real-time PCR assay, 67 were HBeAg seronegative, and the existence of precore mutants was not analysed. In addition, the diagnosis of HBV reactivation based on HBV DNA measurement upon the development of hepatitis could be suboptimal in two aspects. First, the commercially available assay (Quantiplex bDNA assay) used was relatively less sensitive, with a lower detection limit of 0.7×10^6 g.e. ml⁻¹. Secondly, in the absence of serial monitoring HBV DNA, the true incidence of viral reactivation might have been underestimated, as the rise in HBV DNA might have preceded overt hepatitis in some cases (Yeo *et al*, 2001).

Several chemotherapeutic agents have been reported to be associated with the development of HBV reactivation in cancer patients. Apart from steroids and anthracyclines, other drugs that have been reported included vincristine, bleomycin, etoposide, methotrexate, actinomycin D, mercaptopurine, azauridine, chlorambucil, cytosine arabinoside, leucovorin, cisplatin and gemcitabine (Galbraith *et al*, 1975; Hoofnagle *et al*, 1982; Thung *et al*, 1985; Bird *et al*, 1989; Lau *et al*, 1989; Liang *et al*, 1990; Pinto *et al*, 1990; Lok *et al*, 1991; Soh *et al*, 1992; Nokamura *et al*, 1996; Wong *et al*, 1996; Yeo *et al*, 2000a). In the present study, it is interesting to find that the use of steroids was an associated risk factor on multivariate analysis. Patients with lymphoma have more frequently been reported to develop HBV reactivation (Lok *et al*, 1991; Omata *et al*, 1991; Nokamura *et al*, 1996; Yeo *et al*, 2000a), and a recent report on breast cancer patients receiving chemotherapy has shown the viral reactivation rate to be as high as 41% (Yeo *et al*, 2003). The type of treatment and the type of tumour may be inter-related factors. Since disease site confounded with the use of certain cytotoxic agents, a separate stepwise logistic regression model without disease site was used for further analysis. The logistic model without considering the disease site revealed that anthracyclines ($P = 0.0662$, OR 2.44, 95% CI from 0.94 to 6.32), as well as steroids ($P = 0.0557$, OR 2.56, 95% CI from 0.98 to 6.69) and detectable HBV DNA ($P = 0.0011$, OR of 6.25, 95% CI from 2.08 to 18.8) were significant factors. Steroids and anthracyclines are commonly used as part of the cyclophosphamide/adriamycin/vincristine/prednisolone (CHOP) regimen for patients with non-Hodgkin's lymphoma; they also constitute adriamycin/cyclophosphamide (AC) combination chemotherapy commonly used for breast cancer patients (where steroid is generally administered in the anti-emetic pre-medication). Indeed, anthracyclines have mainly been administered to patients with non-Hodgkin's lymphoma and breast cancer (Table 3). Although it is generally accepted that CHOP is an immunosuppressive regimen, only 4% of Western breast cancer patients receiving AC developed severe immunosuppression (Fisher *et al*, 1990). However, recent data from Chinese breast cancer patient population revealed the

corresponding figure to be as high as 77% (Ma *et al*, 2002). This implies that the degree of immunosuppression as a result of the treatment could have been a more significant factor than the type of malignancy *per se* in the development of HBV reactivation. This is further supported by one of the findings from the present study that revealed that patients with gastrointestinal malignancies, who undergo cytotoxic chemotherapy mainly consisting of less immunosuppressive agents (fluorouracil and folinic acid), have a lower risk of developing viral reactivation. Previous study has suggested that the type of chemotherapeutic regimens, rather than the individual cytotoxic agent, may be a more important factor (Yeo *et al*, 2000a). Thus, when the use of 5-fluorouracil was analysed in the present study, no association was found with HBV reactivation; while the agent was commonly coupled with folinic acid in gastrointestinal malignancies, it was also often used in combination with other cytotoxics that resulted in a variable degree of immunosuppression for non-gastrointestinal malignancies.

Using the real-time PCR assay, which has a lower cutoff value of 2.9×10^3 g.e. ml⁻¹, the present study has revealed that detectable HBV DNA load prior to chemotherapy was a significant predictive factor for viral reactivation. Although a recent study has demonstrated that a detectable HBV DNA may predict HBV reactivation in HBsAg-positive individuals (Lau *et al*, 2002), the finding, which was based on a small heterogeneous group of patients who underwent autologous hematopoietic cell transplantation, could not be generalised and applied to the majority of cancer patients who require conventional cytotoxic chemotherapy without transplantation.

Prophylactic treatment with antiviral nucleotide analogue lamivudine prior to the start of chemotherapy has been suggested to be effective in reducing the incidence of HBV reactivation (Rossi *et al*, 2001; Liao *et al*, 2002; Lim *et al*, 2002; Persico *et al*, 2002; Shibolet *et al*, 2002; Yeo *et al*, 2002). The use of the proposed predictive model may aid the identification of high-risk patients

REFERENCES

Bird GL, Smith H, Portmann B, Alexander GJ, Williams R (1989) Acute liver decompensation on withdrawal of cytotoxic and immunosuppressive therapy in hepatitis B carriers. *Q J Med* 73: 895–902

Chan SW, McOmish F, Holmes EC, Dow B, Peutherer JF, Follett E, Yap PL, Marsden H (1992) Analysis of a new hepatitis C virus type and its phylogenetic relationship to existing variants. *J Gen Virol* 73: 1131–1141

Ehata T, Omata M, Chuang WL, Yokosuka O, Ito Y, Hosoda K, Ohto M (1993) Mutations in the core nucleoside sequence of hepatitis B virus correlate with fulminant and severe hepatitis. *J Clin Invest* 91: 1206–1213

Fisher B, Brown AM, Dimitrov NV, Poisson R, Redmond C, Margolese RG, Bowman D, Wolmark M, Wickerham DL, Kardinal CG (1990) Two months of doxorubicin-cyclophosphamide with and without interval reinduction therapy compared with six months of cyclophosphamide, methotrexate and fluorouracil in positive-node breast cancer patients with tamoxifen-nonresponsive tumors: result from the National Surgical Adjuvant Breast and Bowel Project B-15. *J Clin Oncol* 8(9): 1483–1496

Galbraith RM, Eddleston AL, Williams R, Zuckerman AJ, Bagshawe KD (1975) Fulminating hepatic failure in leukemia and choriocarcinoma related to withdrawal of cytotoxic drug therapy. *Lancet* 2: 528–530

Hanley JA (1989) Receiver operating characteristic (ROC) methodology: the state of the art. *Crit Rev Diagn Imaging* 29: 307–335

Hoofnagle JH, Dusheiko GM, Schafer DF, Jones EA, Micetich KC, Young RC, Costa J (1982) Reactivation of chronic hepatitis B virus infection by cancer chemotherapy. *Ann Intern Med* 96: 447–449

Kumagai K, Takagi T, Nakamura S, Sawada U, Kura U, Kodama F, Shimano S, Kudoh I, Nakamura H, Sawada K, Ohnoshi T (1997) Hepatitis B virus carriers in the treatment of malignant lymphoma: an epidemiological study in Japan. *Ann Oncol* 8: 107–109

Lau GK, Leung YH, Fong DY, Au WY, Kwong YL, Lie A, Hou JL, Wen YM, Nanj A, Liang R (2002) High hepatitis B virus (HBV) DNA viral load as the most important risk factor for HBV reactivation in patients positive

Table 3 Tumour type treated with anthracyclines

	No. of patients with the tumour type	No. of patients treated with anthracyclines
Breast cancers	39	22 (56%)
Lymphomas	12	10 (83%)
Gastrointestinal cancers	29	0
Head and neck cancers	17	0
Lung cancers	13	0
Other cancers	18	9 (50%)

who stand to benefit most from the antiviral, which could in turn be administered in a most cost-effective manner.

We conclude that, in cancer patients who are positive for HBsAg and who undergo cytotoxic chemotherapy, high HBV viral load ($>2.9 \times 10^3$ g.e. ml⁻¹) prior to the administration of cytotoxic chemotherapy is a significant predictive factor for the development of HBV reactivation. Other factors that have been identified include the use of steroids and a diagnosis of lymphoma and breast cancer. The combination of baseline HBV viral load (using real-time PCR) and clinical features provide a more sensitive and specific means of identifying patients who are at risk of developing HBV reactivation, and who would benefit most from lamivudine prophylaxis. Although a predictive model has been formed to identify patients who are at risk of developing HBV reactivation in this study, there is a need to validate it in a separate cohort of patients before it can be safely applied in the clinical practice.

ACKNOWLEDGEMENTS

We thank research nurse Miss Jam Jun Lee and secretary Miss Candy Ho for their support in data collection.

for HBV surface antigen undergoing autologous hematopoietic cell transplantation. *Blood* 99(7): 2324–2330

Lau VYN, Lai CL, Lin HJ, Lok ASF, Liang RHS, Wu PC, Chan TK, Todd D (1989) Fatal reactivation of chronic hepatitis B virus infection following withdrawal of chemotherapy in lymphoma patients. *Q J Med* 73: 911–917

Liang RHS, Lok ASF, Lai CL, Chan TK, Todd O, Chiu EKW (1990) Hepatitis B infection in patients with lymphomas. *Hematol Oncol* 8: 261–270

Liang TL, Hasegawa K, Munoz SJ, Shapiro CN, Yoffe B, McMahon BJ, Feng C, Bei H, Alter MJ, Dienstag JL (1994) Hepatitis B virus pre-core mutation and fulminant hepatitis in the United States. *J Clin Invest* 93: 950–955

Liao CA, Lee CM, Wu HC, Wang MC, Lu SN, Eng HL (2002) Lamivudine for the treatment of hepatitis B virus reactivation following chemotherapy for non-Hodgkin's lymphoma. *Br J Haematol* 116(1): 166–169

Liaw YF, Tai DI, Chu CM, Pao CC, Chen TJ (1987) Acute exacerbation in chronic type of B hepatitis: comparison between HBeAg and antibody positive patients. *Hepatology* 7: 20–23

Lim LL, Wai CT, Lee YM, Kong HL, Lim R, Koay E, Lim SG (2002) Prophylactic lamivudine prevents hepatitis B reactivation in chemotherapy patients. *Aliment Pharmacol Ther* 16(11): 1939–1944

Lok ASF, Liang RHS, Chiu EKW, Wong KL, Chan TK, Todd D (1991) Reactivation of hepatitis B virus replication in patients receiving cytotoxic therapy. Report of a prospective study. *Gastroenterology* 100: 182–188

Ma B, Yeo W, Hui P, Ho WM, Johnson PJ (2002) Acute toxicity of adjuvant doxorubicin and cyclophosphamide for early breast cancer – a retrospective review of Chinese patients and comparison with a historic Western series. *Radiother Oncol* 62(2): 185–189

Markovic S, Drozina G, Vovk M, Fidler-Jenko M (1999) Reactivation of hepatitis B but not hepatitis C in patients with malignant lymphoma and

- immunosuppressive therapy. A prospective study in 305 patients. *Hepatology* 46(29): 2925–2930
- Nokamura Y, Motokura T, Fujita A, Yamashita T, Ogata E (1996) Severe hepatitis related to chemotherapy in hepatitis B virus carriers with hematologic malignancies. Survey in Japan, 1987–1991. *Cancer* 78: 2210–2215
- Omata M, Ehata T, Yokosuka O, Hosoda K, Ohto M (1991) Mutations in the pre-core region of hepatitis B virus DNA in patients with fulminant and severe hepatitis. *N Engl J Med* 324: 1699–1704
- Persico M, De Marino F, Russo GD, Severino A, Palmentieri B, Picardi M, Morante A, Rotloi B, Torella R, De Renzo A (2002) Efficacy of lamivudine to prevent hepatitis reactivation in hepatitis B virus-infected patients treated for non-Hodgkin lymphoma. *Blood* 99(2): 724–725
- Pinto PC, Hu E, Bernstein-Singer M, Pinter-Brown L, Govindarajan S (1990) Acute hepatic injury after withdrawal of immunosuppressive chemotherapy in patients with hepatitis B. *Cancer* 65: 878–884
- Rossi G, Pelizzari A, Motta M, Puoti M (2001) Primary prophylaxis with lamivudine of hepatitis B virus reactivation in chronic HBsAg carriers with lymphoid malignancies treated with chemotherapy. *Br J Haematol* 115(1): 58–62
- Shibolet O, Ilan Y, Gillis S, Hubert A, Shouval D, Safadi R (2002) Lamivudine therapy for prevention of immunosuppressive-induced hepatitis B virus reactivation in hepatitis B surface antigen carriers. *Blood* 100: 391–396
- Soh LT, Ang PT, Sng I, Chua EJ, Ong YW (1992) Fulminant hepatic failure in non-Hodgkin's lymphoma patients treated with chemotherapy. *Eur J Cancer* 28A: 1338–1339
- Steinberg JL, Yeo W, Zhong S, Chan JYH, Tam JS, Chan PKS, Leung NWY, Johnson PJ (2000) Hepatitis B virus reactivation in patients undergoing cytotoxic chemotherapy for solid tumours: precore/core mutations may play an important role. *J Med Virol* 60: 249–255
- Thung SN, Gerber MA, Klion F, Gilbert H (1985) Massive hepatic necrosis after chemotherapy withdrawal in a hepatitis B carrier. *Arch Intern Med* 145: 1313–1314
- Wong GC, Tan P, Goh YT, Ng HS, Lee LH (1996) Exacerbation of hepatitis in hepatitis B carriers following chemotherapy for haematological malignancies. *Annals of the Academy of Medicine, Singapore* 25: 500–503
- Yeo W, Chan PKS, Chan HLY, Mo FKF, Johnson PJ (2001) Hepatitis B virus reactivation during cytotoxic chemotherapy-enhanced viral replication precedes overt hepatitis. *J Med Virol* 65: 473–477
- Yeo W, Chan PKS, Hui P, Ho WM, Lam KC, Kwan WH, Zhong S, Johnson PJ (2003) Hepatitis B virus reactivation in breast cancer patients undergoing cytotoxic chemotherapy: a prospective study. *J Med Virol* 70: 553–561
- Yeo W, Chan PKS, Zee B, Ho WM, Lam KC, Hui P, Mok TSK, Chan ATC, Zhong S, Leung WT, Lei KIK, Johnson PJ (2002) Lamivudine prevents hepatitis B reactivation in HBV carriers undergoing cytotoxic chemotherapy – a prospective study. *Program/Proc Am Soc Clin Oncol* 21: 362a
- Yeo W, Chan PKS, Zhong S, Ho WM, Steinberg JL, Tam JS, Hui P, Leung NWY, Zee B, Johnson PJ (2000a) Frequency of hepatitis B virus reactivation in cancer patients undergoing cytotoxic chemotherapy: a prospective study of 626 patients with identification of risk factors. *J Med Virol* 62: 299–307
- Yeo W, Zhong S, Chan PKS, Ho WM, Wong HTM, Chan ASK, Johnson PJ (2000b) Sequence variations of precore/core and precore promoter regions of hepatitis B virus in patients with or without viral reactivation during cytotoxic chemotherapy. *J Viral Hepatitis* 7: 448–458
- Zhong S, Yeo W, Schroder C, Chan PKS, Wong WL, Ho WM, Mo F, Zee B, Johnson PJ (2004) High hepatitis B virus (HBV) DNA viral load is an important risk factor for HBV reactivation in breast cancer patients undergoing cytotoxic chemotherapy. *J Viral Hepatitis* 11: 55–59