



Living Well With Kidney Disease by Patient and Care-Partner Empowerment: Kidney Health for Everyone Everywhere

Canadian Journal of Kidney Health and Disease
Volume 8: 1–7
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2054358121995276
journals.sagepub.com/home/cjk



**Kamyar Kalantar-Zadeh¹, Philip Kam-Tao Li²,
Ekamol Tantisattamo³ , Latha Kumaraswami⁴,
Vassilios Liakopoulos⁵, Siu-Fai Lui⁶, Ifeoma Ulasi⁷,
Sharon Andreoli⁸, Alessandro Balducci⁹, Sophie Dupuis¹⁰,
Tess Harris¹¹ , Anne Hradsky¹⁰, Richard Knight¹²,
Sajay Kumar⁴, Maggie Ng¹³, Alice Poidevin¹⁰, Gamal Saadi¹⁴,
and Allison Tong¹⁵ for the World Kidney Day Steering Committee#**

Abstract

Living with chronic kidney disease (CKD) is associated with hardships for patients and their care-partners. Empowering patients and their care-partners, including family members or friends involved in their care, may help minimize burden and consequences of CKD-related symptoms to enable life participation. There is a need to broaden the focus on living well with kidney disease and re-engagement in life, including emphasis on patients being in control. The World Kidney Day (WKD) Joint Steering Committee has declared 2021 the year of “Living Well with Kidney Disease” in an effort to increase education and awareness on the important goal of patient empowerment and life participation. This calls for the development and implementation of validated patient-reported outcome measures to assess and address areas of life participation in routine care. It could be supported by regulatory agencies as a metric for quality care or to support labeling claims for medicines and devices. Funding agencies could establish targeted calls for research that address the priorities of patients. Patients with kidney disease and their care-partners should feel supported to live well through concerted efforts by kidney care communities including during pandemics. In the overall wellness program for kidney disease patients, the need for prevention should be reiterated. Early detection with prolonged course of wellness despite kidney disease, after effective secondary and tertiary prevention programs, should be promoted. WKD 2021 continues to call for increased awareness of the importance of preventive measures throughout populations, professionals, and policy makers, applicable to both developed and developing countries.

Abrégé

Vivre avec l'insuffisance rénale chronique (IRC) représente un fardeau pour les patients et leurs partenaires soignants, dont leurs familles et amis. Une façon d'alléger ce fardeau et de réduire les conséquences des symptômes de l'IRC sur la participation à la vie serait de favoriser une plus grande autonomie des patients et de leurs partenaires soignants. Pour ce faire, l'accent mis sur le bien vivre avec la maladie et sur la participation à la vie doit être élargi en privilégiant notamment de mettre les patients en contrôle. Pour accroître l'éducation et la sensibilisation sur la responsabilisation des patients et, ultimement, encourager leur participation à la vie, le Comité directeur bipartite de la Journée Mondiale du Rein (JMR) a déclaré 2021 l'année du « bien vivre avec une maladie rénale ». Une stratégie qui nécessite l'élaboration et la mise en œuvre de mesures validées des résultats signalés par le patient afin d'aborder la participation à la vie dans les soins courants. Cette stratégie pourrait être appuyée par les organismes de réglementation comme mesure de qualité des soins ou pour appuyer les demandes relatives à l'étiquetage des médicaments et des dispositifs. Les organismes subventionnaires pourraient quant à eux établir des appels d'offres à objectifs de recherche définis afin de répondre aux priorités des patients. Les patients et leurs partenaires soignants devraient se sentir soutenus, même en contexte de pandémie, par des efforts concertés des communautés de soins en santé rénale. Dans le cadre du program de bien-être global destiné aux patients atteints d'une néphropathie, il convient de réitérer la prévention. À la suite de programs efficaces de prévention secondaire et tertiaire, il convient également de promouvoir une détection précoce avec une évolution prolongée du bien-être malgré une maladie rénale. La JMR 2021 continue d'appeler à une plus grande sensibilisation des populations, des professionnels et des décideurs quant à l'importance de mesures préventives applicables tant dans les pays développés que dans les pays en développement.



Keywords

patient empowerment, care-partner, low-middle-income countries, health policy

Received November 9, 2020. Accepted for publication January 21, 2021.

Patient Priorities for Living Well: A Focus on Life Participation

Chronic kidney disease (CKD) its associated symptoms, and its treatment, including medications, dietary and fluid restrictions, and kidney replacement therapy can disrupt and constrain daily living, and impair the overall quality of life of patients and their family members. Consequently, this can also impact on treatment satisfaction and clinical outcomes.¹ Despite this, the past several decades have seen limited improvement in the quality of life of people with CKD.¹ To advance research, practice, and policy, there is an increasing recognition of the need to identify and address patient priorities, values, and goals.¹

Several regional and global kidney health projects have addressed on these important questions including the *Standardized Outcomes in Nephrology* (SONG) with more than 9000 patients, family members, and health professionals from over 70 countries.^{2,3} Across all treatment stages, including CKD, dialysis, and transplantation; SONG participating children; and adults with CKD consistently gave higher priority to symptoms and life impacts than health professionals.^{2,3} In comparison, health professionals gave higher priority to mortality and hospitalization than patients and

family members. The patient-prioritized outcomes are shown in Figure 1. Irrespective of the type of kidney disease or treatment stage, patients wanted to be able to live well, maintain their role and social functioning, protect some semblance of normality, and have a sense of control over their health and wellbeing.

Life participation, defined as the ability to do meaningful activities of life including, but not limited to, work, study, family responsibilities, travel, sport, social, and recreational activities, was established a critically important outcome across all treatment stages of CKD.^{1,2} The quotations from patients with kidney disease provided in Box 1 demonstrates how life participation reflects the ability to live well with CKD.⁴ According to the World Health Organization (WHO), participation refers to “involvement in a life situation.”⁵ This concept is more specific than the broader construct of quality of life. Life participation places the life priorities and values of those affected by CKD and their family at the center of decision making. The World Kidney Day Steering Committee calls for the inclusion of life participation, a key focus in the care of patients with CKD, to achieve the ultimate goal of living well with kidney disease. This calls for the development and implementation of validated patient-reported outcome measures, that could be used to assess and

¹The International Federation of Kidney Foundation–World Kidney Alliance, Division of Nephrology and Hypertension and Kidney Transplantation, University of California, Irvine, Orange, USA

²Department of Medicine and Therapeutics, Carol & Richard Yu PD Research Centre, Prince of Wales Hospital, The Chinese University of Hong Kong, Hong Kong

³Division of Nephrology, Hypertension and Kidney Transplantation, Department of Medicine, University of California Irvine School of Medicine, Orange, California, USA

⁴Tanker Foundation, Chennai, India

⁵Division of Nephrology and Hypertension, 1st Department of Internal Medicine, AHEPA Hospital, Aristotle University of Thessaloniki, Greece

⁶Hong Kong Kidney Foundation, The International Federation of Kidney Foundations–World Kidney Alliance, Hong Kong, China

⁷Renal Unit, Department of Medicine, College of Medicine, University of Nigeria, Ituku-Ozalla, Nigeria

⁸James Whitcomb Riley Hospital for Children, Indiana University School of Medicine, Indianapolis, USA

⁹Italian Kidney Foundation, Rome, Italy

¹⁰World Kidney Day Office, Brussels, Belgium

¹¹Polycystic Kidney Disease Charity and Kidney Patient, London, UK

¹²American Association of Kidney Patients, Tampa, FL, USA

¹³Hong Kong Kidney Foundation, China

¹⁴Nephrology Unit, Department of Internal Medicine, Faculty of Medicine, Cairo University, Giza, Egypt

¹⁵Sydney School of Public Health, The University of Sydney, New South Wales, Australia

Corresponding Authors:

Kamyar Kalantar-Zadeh, Division of Nephrology and Hypertension and Kidney Transplantation, University of California, Irvine, School of Medicine, Orange, USA.

Email: kkz@uci.edu

Philip Kam-Tao Li, Department of Medicine and Therapeutics, Prince of Wales Hospital, The Chinese University of Hong Kong, 30–32 Ngan Shing Street, Shatin, New Territories, Hong Kong, China.

Email: philipli@cuhk.edu.hk




Figure 1. Conceptual framework of “Living Well with Kidney Disease” based on patient centeredness and empowering patient with focus on effective symptom management and life participation.

Box 1. Quotations From Patients With CKD Related to Priorities for Living Well.

“I don’t want to think about dying from my disease. I want to be able to live well with my disease.”—Patient with CKD

“Life participation is most important because without it, you can’t do anything.”—Child with CKD

“Maybe it’s as simple as asking patients whether, how well they are able to participate in the life that they want to lead because it’s going to be different for different people”—Kidney transplant recipient

“Everyone has to face death, what I would like to have is a good quality of life rather than to face death.”—Kidney transplant recipient

“So, it doesn’t actually really matter what the numbers say, and some of my numbers should have suggested that I should be feeling a lot worse than what I actually was, it’s about how much I feel I can do and participate in my life and feel normal.”—Patient with CKD

“I’m still living. I get out of bed, and I’m still living and still breathing. As long as I can do that, I’m going to carry on and be positive because life is short.” Patient with CKD⁴

“I put life participation because I know that looking from the outside, I know [his kidney disease] stops [him] from thinking bigger. . . Although that’s really big, there’s this life that has to happen at the same time.”—Family member

“Amazed at comments from professional(sic) about travel, free time, etc they seem to think the mechanics of dialysis far more important. Dialysis is a treatment which keeps us alive to live a life, not just to wait for death.—Patient receiving dialysis

“I prefer to be above ground, then below ground. So why not enjoy life whilst being above ground.” Adam Martin^a

“Over the years, I have learned to worry less, control my emotions and not fear death. I keep my mind active. I follow the advice of the philosopher-emperor Marcus Aurelius to ‘love the hand that fate (has dealt me) and play it as (my) own’. Living well with CKD means to live the best life I can in the time I have available . . . Living well with CKD is the same as living well.”—Tess Harris^a

“While CKD brings me some limitations, I can maximize the possibility to live well. I kept working when I was doing hemodialysis. After transplant, I could live: study, work, travel, marry, have children, and service the community.”—Maggie Ng^a

Note. CKD = chronic kidney disease.

^aPersonal communication; quotations are identified by name with permission.

address areas of life participation in routine care. Monitoring of life participation could be supported by regulatory agencies as a metric for quality care or to support labeling claims

for medicines and devices. Funding agencies could establish targeted calls for research that address the priorities of patients, including life participation.

Table 1. Suggested Strategies for “Living Well With CKD” Using a Strengths-Based Approach.

Strengths-based approach	Suggested strategies
Build resilience	<ul style="list-style-type: none"> Identify or provide strategies and resources to manage stress and functioning when encountering challenges, adversity, and trauma (eg, commencement of dialysis)
Harness social connections	<ul style="list-style-type: none"> Facilitate connections with other patients to learn coping strategies and for support Support family members/caregivers
Build awareness and knowledge	<ul style="list-style-type: none"> Provide education (including practical advice) on diet and lifestyle modifications Understand, identify, and address the potential impacts of CKD (eg, cognitive function). Encourage patients to ask questions. Encourage the use of knowledge to empower and prepare for the future.
Facilitate access to support	<ul style="list-style-type: none"> Refer to allied health care professionals (eg, dietitian, social worker, mental health professionals, occupation therapists) Provide support that enables patient to participate in important life activities, for example, work.
Establish confidence and control in self-management	<ul style="list-style-type: none"> Support informed and shared decision-making (including dialysis, kidney transplantation, conservative or nondialytic care) Encourage patients to learn to “get in tune” with what works well for them and to voice any concerns, and work together to develop better management strategies to enable patients to feel better. Provide strategies to prevent or manage complications (eg, infection) Support open communication regarding goals, concerns, and priorities

Note. CKD = chronic kidney disease (not receiving kidney replacement therapy).

Patient Empowerment, Partnership and a Paradigm Shift Towards a Strengths-Based Approach to Care

Patients with CKD and their family members including care-partners should be empowered to achieve the health outcomes and life goals that are meaningful and important to them. The WHO defines patient empowerment as “a process through which people gain greater control over decisions or actions affecting their health,”⁶ which requires patients to understand their role, to have knowledge to be able to engage with clinicians in shared decision-making, skills, and support for self-management. For patients receiving dialysis, understanding the rationale for lifestyle change, having access to practical assistance and family support promoted patient empowerment, while feeling limited in life participation undermined their sense of empowerment.⁷

The World Kidney Day Steering Committee advocates for strengthened partnership with patients in the development, implementation, and evaluation of interventions for practice and policy settings, that enable patients to live well with kidney diseases. This needs to be supported by consistent, accessible, and meaningful communication. Meaningful involvement of patients and family members across the entire research process, from priority setting and planning the study through to dissemination and implementation, is now widely advocated.⁸ There have also been efforts, such as the *Kidney Health Initiative*, to involve patients in the development of drugs and devices to foster innovation.⁹

We urge for greater emphasis on a strengths-based approach as outlined in Table 1, which encompasses strategies to support patient resilience, harness social connections, build patient awareness and knowledge, facilitate access to

support, and establish confidence and control in self-management. The strengths-based approach is in contrast to the medical model where chronic disease is traditionally focussed on pathology, problems, and failures.¹⁰ Instead, the strengths-based approach acknowledges that each individual has strengths and abilities to overcome the problems and challenges faced, and requires collaboration and cultivation of the patient’s hopes, aspirations, interests and values. Efforts are needed to ensure that structural biases, discrimination, and disparities in the health care system also need to be identified, so all patients are given the opportunity to have a voice.

The Role of Care-Partner

A care-partner is often an informal caregiver who is also a family member of the patient with CKD.¹¹ They may take on a wide range of responsibilities including coordinating care (including transportation to appointments), administration of treatment including medications, home dialysis assistance, and supporting dietary management. Caregivers of patients with CKD have reported depression, fatigue, isolation, and also burden out. The role of the care-partner has increasingly become more important in CKD care given the heightened complexity in communicative and therapeutic options including expansion of telemedicine under COVID-19 pandemic and given the goal to achieve higher life expectancy with CKD.¹² The experience of caring for a partially incapacitated family member with progressive CKD can represent a substantial burden on the care-partner and may impact family dynamics. Not infrequently, the career goals and other occupational and leisure aspects of the life of the care-partner are affected because of CKD care partnership, leading to

care-partner overload and burnout. Hence, the above-mentioned principles of life participation need to equally apply to care-partners as well as all family members and friends involved in the CKD care.

Living With Kidney Disease in Low Income Regions

In low and lower-middle income countries (LICs and LMICs) including in sub-Saharan Africa, South East Asia, and Latin America, patient's ability to self-manage or cope with chronic disease vary but may often be influenced by internal factors including spirituality, belief system, and religiosity, and external factors including appropriate knowledge of the disease, poverty, family support system, and one's grit and social relations network. The support system comprising healthcare providers and caregivers plays a crucial role as most patients rely on them in making decisions, and for the necessary adjustments in their health behavior.¹³ In LIC regions, where there are often a relatively low number physicians and even lower number of kidney care providers per population especially in rural areas, a stepwise approach can involve local and national stakeholders including both non-governmental organizations and government agencies by: (1) extending kidney patient education in rural areas, (2) adapting telehealth technologies if feasible to educate patients and train local community kidney care providers, and (3) implementing effective retention strategies for rural kidney health providers including adapting career plans and competitive incentives.

Many patients in low resource settings present in very late stage needing to commence emergency dialysis.¹⁴ The very few fortunate ones to receive kidney transplantation may acquire an indescribable chance to normal life again, notwithstanding the high costs of immunosuppressive medications in some countries. For some patients and care-partners in low income regions, spirituality and religiosity may engender hope, when ill they are energized by the anticipation of restored health and spiritual wellbeing. For many patients, informing them of a diagnosis of kidney disease is a harrowing experience both for the patient (and caregivers) and the healthcare professional. Most patients present to kidney physicians (usually known as "renal physicians" in many of these countries) with trepidations and apprehension. It is rewarding therefore to see the patient's anxiety dissipate after reassuring him or her of a diagnosis of simple kidney cysts, urinary tract infection, simple kidney stones, solitary kidneys, and so on, that would not require extreme measures like kidney replacement therapy. Patients diagnosed with glomerulonephritis who have appropriate characterization of their disease from kidney biopsies and histology; who receive appropriate therapies and achieve remission are relieved and are very grateful. Patients are glad to discontinue dialysis following resolution of AKI or acute on CKD.

Many CKD patients who have residual kidney function appreciate being maintained in relatively healthy state with conservative measures, without dialysis. They experience renewed energy when their anemia is promptly corrected using erythropoiesis stimulating agents. They are happy when their peripheral oedema resolves with treatment. For those on maintenance hemodialysis who had woeful stories from emergency femoral cannulations, they appreciate construction of good temporary or permanent vascular accesses. Many patients in low resource settings present in very late stage needing to commence emergency dialysis. Patients remain grateful on waking from uremic coma or recovering from recurrent seizures when they commence dialysis.

World Kidney Day 2021 Advocacy

World Kidney Day 2021 theme on "Living Well with Kidney Disease" is deliberately chosen to have the goals to redirect more focus on plans and actions toward achieving patient-centered wellness. "Kidney Health for Everyone, Everywhere" with emphasis on patient-centered wellness should be a policy imperative that can be successfully achieved if policy makers, nephrologists, health care professionals, patients, and care partners place this within the context of comprehensive care. The requirement of patient engagement is needed. WHO in 2016 put out an important document on patient empowerment (WHO 2016):

Patient engagement is increasingly recognized as an integral part of health care and a critical component of safe people-centred services. Engaged patients are better able to make informed decisions about their care options. In addition, resources may be better used if they are aligned with patients.

priorities and this is critical for the sustainability of health systems worldwide. Patient engagement may also promote mutual accountability and understanding between the patients and health care providers. Informed patients are more likely to feel confident to report both positive and negative experiences and have increased concordance with mutually agreed care management plans. This not only improves health outcomes, but also advances learning and improvement, while reducing adverse events. In the ISN Community Film Event at World Congress of Nephrology (WCN) 20 (ISN Community Film Event 2020), it is good to see a quote in the film from patients: "Tell me. I will forget; Show me. I will remember; Involve me. I will understand." ISN Global Kidney Policy Forum 2019 included a patient speaker Nicki Scholes-Robertson from New Zealand: "Culturally appropriate and sensitive patient information and care are being undertaken in New Zealand to fight inequities in kidney health, especially in Maori and other disadvantaged communities."

World Kidney Day 2021 would like to promote to the policy makers on increasing focus and resources on both drug and

nondrug programs in improving patient wellness. Examples include funding for erythropoiesis stimulating agents and anti-pruritic agents for managing anemia and itchiness respectively, just name but a few.^{15,16} Home dialysis therapies have been consistently found to improve patient autonomy and flexibility, quality of life in a cost-effective manner, enhancing life participation. Promoting home dialysis therapies should tie in with appropriate “assisted dialysis” programs to reduce patient and care partner fatigue and burnout. Also examples like self-management programs, cognitive behavioral therapy and group therapies for managing depression, anxiety, and insomnia should be promoted before resorting to medications.¹⁷ The principle of equity recognizes that different people with different levels of disadvantage require different approaches and resources to achieve equitable health outcomes. The kidney community should push for adapted care guidelines for vulnerable and disadvantaged populations. Involvement of primary care and general physicians especially in LICs and LMICs would be useful in improving the affordability and access to services through the public sector in helping the symptom management of CKD patients and improve their wellness. In the overall wellness program for kidney disease patients, the need for prevention should be reiterated. Early detection with prolonged course of wellness despite kidney disease, after effective secondary prevention program, should be promoted.¹⁸ Prevention of CKD progression can be attempted by lifestyle and diet modifications such as a plant-dominant low protein diet and by means of effective pharmacotherapy including administration of sodium-glucose transport protein 2 (SGLT2) inhibitors.¹⁹ WKD 2021 continues to call for increased awareness of the importance of preventive measures throughout populations, professionals, and policy makers, applicable to both developed and developing countries.¹⁸

Conclusion

Effective strategies to empower patients and their care-partners strive to pursue the overarching goal of minimizing burden of CKD-related symptoms in order to enhance patient satisfaction, health-related quality of life and life participation. World Kidney Day 2021 theme on “Living Well with Kidney Disease” is deliberately chosen to have the goals to redirect more focus on plans and actions toward achieving patient-centered wellness. Notwithstanding the COVID-19 pandemic that had overshadowed many activities in 2020 and beyond, the World Kidney Day Steering Committee has declared 2021 the year of “Living well with Kidney Disease” in an effort to increase education and awareness on the important goal of effective symptom management and patient empowerment. Whereas the World Kidney Day continues to emphasize the importance of effective measures to prevent kidney disease and its progression,¹⁸ patients with preexisting kidney disease and their care-partners should feel supported to live well through concerted efforts by kidney

care communities and other stakeholders throughout the world even during a world shattering pandemic as COVID-19 that may drain many resources.²⁰ Living well with kidney disease is an uncompromisable goal of all kidney foundations, patient groups, and professional societies alike, to which the International Society of Nephrology and the International Federation of Kidney Foundation World Kidney Alliance are committed at all times.

Ethics Approval and Consent to Participate

Permission to quote personal communication has been received from participants.

Consent for Publication

Not applicable.

Availability of Data and Materials

Not applicable.

Authors' Note

Members of the World Kidney Day Steering Committee are: Philip Kam Tao Li, Kamyar Kalantar-Zadeh, Sharon Andreoli, Alessandro Balducci, Sophie Dupuis, Latha Kumaraswami, Vassilios Liakopoulos, Siu-Fai Lui, Gamal Saadi, and Ifeoma Ulasi.

Declaration of Conflicting Interests

The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: K.K.-Z. reports honoraria from Abbott, Abbvie, ACI Clinical, Akebia, Alexion, Amgen, Ardelyx, Astra-Zeneca, Aveo, BBraun, Cara Therapeutics, Chugai, Cytokinetics, Daiichi, DaVita, Fresenius, Genentech, Haymarket Media, Hospira, Kabi, Keryx, Kissei, Novartis, Pfizer, Regulus, Relypsa, Resverlogix, Dr Schaefer, Sandoz, Sanofi, Shire, Vifor, UpToDate, and ZS-Pharma. P.K.T.L. reports personal fees from Fibrogen and Astra-Zeneca. G.S. reports personal fees from Multicare, Novartis, Sandoz, and Astra-Zeneca. V.L. reports nonfinancial support from Genesis Pharma. ET, LK, SL, IU, SA, AB, SD, TH, AH, RK, SK, MN, AP, and AT have declared that no conflict of interest exists.

This article was originally published in *Kidney International*, volume 99, pages 278-284, Copyright World Kidney Day Steering Committee (2021), and reprinted concurrently in several journals. The articles cover identical concepts and wording but vary in minor stylistic and spelling changes, detail, and length of manuscript in keeping with each journal's style. Any of these versions may be used in citing this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Ekamol Tantisattamo  <https://orcid.org/0000-0003-0883-6892>

Tess Harris  <https://orcid.org/0000-0003-3659-7438>

References

1. Tong A, Manns B, Wang AYM, et al. Implementing core outcomes in kidney disease: report of the Standardized Outcomes in Nephrology (SONG) implementation workshop. *Kidney Int.* 2018;94(6):1053-1068. doi:10.1016/j.kint.2018.08.018.
2. Carter SA, Gutman T, Logeman C, et al. Identifying outcomes important to patients with glomerular disease and their caregivers. *Clin J Am Soc Nephrol.* 2020;15(5):673-684. <https://www.ncbi.nlm.nih.gov/pubmed/32354728>.
3. Hanson CS, Craig JC, Logeman C, et al. Establishing core outcome domains in pediatric kidney disease: report of the Standardized Outcomes in Nephrology-Children and Adolescents (SONG-KIDS) consensus workshops. *Kidney Int.* 2020;98:553-565. <https://www.ncbi.nlm.nih.gov/pubmed/32628942>.
4. Gonzalez AM, Gutman T, Lopez-Vargas P, et al. Patient and caregiver priorities for outcomes in CKD: a multinational nominal group technique study. *Am J Kid Dis.* 2020;76:679-689. doi:10.1053/j.ajkd.2020.03.022.
5. World Health Organization. Towards a common language for functioning, disability and health. Geneva, Switzerland: World Health Organization; 2002. <http://www.who.int/classifications/icf/training/icfbeginnersguide.pdf>.
6. World Health Organization. Health promotion glossary. WHO/HPR/HEP/981; 1998. <https://www.who.int/healthpromotion/about/HPR%20Glossary%201998.pdf>.
7. Baumgart A, Manera KE, Johnson DW, et al. Meaning of empowerment in peritoneal dialysis: focus groups with patients and caregivers. *Nephrol Dial Transplant.* 2020;35:1949-1958. <https://www.ncbi.nlm.nih.gov/pubmed/32712672>.
8. Patient-Centered Outcomes Research Institute. The value of engagement. <https://www.pcori.org/about-us/our-programs/engagement/public-and-patient-engagement/value-engagement>. Published 2018. Accessed September 1, 2020.
9. Bonventre JV, Hurst FP, West M, Wu I, Roy-Chaudhury P, Sheldon M. A technology roadmap for innovative approaches to kidney replacement therapies: a catalyst for change. *Clin J Am Soc Nephrol.* 2019;14(10):1539-1547. <https://www.ncbi.nlm.nih.gov/pubmed/31562182>.
10. Ibrahim N, Michail M, Callaghan P. The strengths based approach as a service delivery model for severe mental illness: a meta-analysis of clinical trials. *BMC Psychiatry.* 2014;14:243. <https://www.ncbi.nlm.nih.gov/pubmed/25189400>.
11. Parham R, Jacyna N, Hothi D, Marks SD, Holttum S, Camic P. Development of a measure of caregiver burden in paediatric chronic kidney disease: the Paediatric Renal Caregiver Burden Scale. *Journal of Health Psychology.* 2016;21:193-205. <http://www.ncbi.nlm.nih.gov/pubmed/24648321>.
12. Subramanian L, Kirk R, Cuttitta T, et al. Remote management for peritoneal dialysis: a qualitative study of patient, care partner, and clinician perceptions and priorities in the United States and the United Kingdom. *Kidney Med.* 2019;1(6):354-365. <https://www.ncbi.nlm.nih.gov/pubmed/32734216>.
13. Angwenyi V, Aantjes C, Kajumi M, De Man J, Criel B, Bunders-Aelen J. Patients experiences of self-management and strategies for dealing with chronic conditions in rural Malawi. *PLoS ONE.* 2018;13(7):e0199977. <https://www.ncbi.nlm.nih.gov/pubmed/29965990>.
14. Ulasi II, Ijoma CK. The enormity of chronic kidney disease in Nigeria: the situation in a teaching hospital in South-East Nigeria. *J Trop Med.* 2010;2010:501957. <https://www.ncbi.nlm.nih.gov/pubmed/20613945>.
15. Spinowitz B, Pecoits-Filho R, Winkelmayr WC, et al. Economic and quality of life burden of anemia in patients with CKD on dialysis: a systematic review. *J Med Econ.* 2019;22(6):593-604. <https://www.ncbi.nlm.nih.gov/pubmed/30813807>.
16. Sukul N, Speyer E, Tu C, et al. Pruritus and patient reported outcomes in non-dialysis CKD. *Clin J Am Soc Nephrol.* 2019;14(5):673-681. <https://www.ncbi.nlm.nih.gov/pubmed/30975656>.
17. Gregg LP, Hedayati SS. Pharmacologic and psychological interventions for depression treatment in patients with kidney disease. *Curr Opin Nephrol Hypertens.* 2020;29(5):457-464. <https://www.ncbi.nlm.nih.gov/pubmed/32701597>.
18. Li PK, Garcia-Garcia G, Lui SF, et al. Kidney health for everyone everywhere—from prevention to detection and equitable access to care. *Kidney Int.* 2020;97(2):226-232. <https://www.ncbi.nlm.nih.gov/pubmed/31980067>.
19. Kalantar-Zadeh K, Li PK. Strategies to prevent kidney disease and its progression. *Nat Rev Nephrol.* 2020;16(3):129-130. <https://www.ncbi.nlm.nih.gov/pubmed/32005966>.
20. Kalantar-Zadeh K, Wightman A, Liao S. Ensuring choice for people with kidney failure—dialysis, supportive care, and hope. *N Engl J Med.* 2020;383(2):99-101. <https://www.ncbi.nlm.nih.gov/pubmed/32640129>.