

## Research Article

# Diverse Effect of Vitamin C and N-Acetylcysteine on Aluminum-Induced Eryptosis

Ali Reza Zangeneh <sup>1</sup>, Mohammad Ali Takhshid <sup>2</sup>, Reza Ranjbaran <sup>2</sup>,  
Mahsa Maleknia <sup>1</sup> and Mohammad Hassan Meshkibaf <sup>1</sup>

<sup>1</sup>Department of Clinical Biochemistry, Fasa University of Medical Sciences, Fasa, Iran

<sup>2</sup>Diagnostic Laboratory Sciences and Technology Research Center, Faculty of Paramedical Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

Correspondence should be addressed to Mohammad Hassan Meshkibaf; meshkibaf.m@fums.ac.ir

Received 21 October 2020; Revised 16 December 2020; Accepted 31 December 2020; Published 12 January 2021

Academic Editor: Saad Tayyab

Copyright © 2021 Ali Reza Zangeneh et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Purpose.** The role of oxidative stress in Aluminum (Al)-induced apoptotic effects has been investigated and suicidal death of erythrocytes, eryptosis, is characterized by cell shrinkage and phosphatidylserine externalization (PSE) at the surface of the erythrocyte cell membrane. Eryptosis is stimulated by an increase in cytosolic  $Ca^{2+}$  concentration and reactive oxygen species (ROS). This *ex vivo* study was conducted to evaluate the effect of well-known antioxidants including vitamin C (vit C) and N-acetylcysteine (NAC), against Al-induced hemolysis and eryptosis. **Methods.** Isolated erythrocytes from the healthy volunteers were partitioned into various groups (6 replicates/group) and treated by various concentrations of Al (3–100  $\mu$ M) in the presence and absence of vit C (0.6 mM) and NAC (1 mM). After 24 hours of treatment, hemolysis was determined from hemoglobin levels in the supernatant. Flowcytometric methods were applied to measure PSE, cell shrinkage,  $Ca^{2+}$  content, and ROS abundance using annexin V-binding, forward scatter, Fluo<sub>3</sub>-fluorescence, and DCFDA dependent fluorescence, respectively. Reduced glutathione (GSH) was measured by the ELISA method. **Results.** The results showed that a 24 hours' exposure of the erythrocytes to Al (10–100  $\mu$ M) significantly increased hemolysis in a dose and  $Ca^{2+}$ -dependent manner. Al also dramatically decreased forward scatter. The percentage of PSE cells, Fluo<sub>3</sub>-fluorescence, and DCFDA fluorescence were increased by Al. Furthermore, cotreatment with NAC inhibited the effect of Al on hemolysis, eryptosis, and ROS production. Vit C decreased Al-induced ROS production. However, increased Al-induced eryptosis. There were no significant changes in glutathione after the ALCL<sub>3</sub> treatment. **Conclusions.** Al-induced eryptosis and hemolysis through triggering oxidative stress, while NAC could diverse this effect. In contrast, vit C might intensify Al-induced eryptosis at particular doses through a less known mechanism.

## 1. Introduction

Due to high abundance in the Earth's crust and widespread use in daily life, the exposure of humans with Aluminum (Al) through occupational exposure, foods, drinking water, drugs, and cosmetic products has been increased [1]. Human and animal studies have depicted that Al led to pathological alteration in the structure and function of the various tissues. In the brain, the adverse influences of Al on the synthesis of neurotransmitters and synaptic transmission, posttranslational modification, degradation of proteins, and expression of genes have been demonstrated [2, 3]. Al could also inhibit

the mitochondrial electron transport chain and energy production in the liver. Furthermore, Al administration caused a significant decrease in the activity of antioxidant enzymes and the induction of oxidative stress in the hepatocytes [4]. Disturbances in the metabolism of phosphate and calcium in the bone and induction of osteomalacia [5], degeneration of the renal tubular cells [6, 7], and adverse effects on reproductive tissues were other reported toxic effects of Al [8]. Al has also revealed toxicity for the hematological system [9]. Inhibition of hemoglobin biosynthesis [10], alteration in membrane integrity, and stimulation of eryptosis [11, 12] were among possible

mechanisms that may have a role in the Al-induced anemia. Eryptosis is defined as suicidal cell death of the erythrocytes and has similar hallmarks to apoptosis in nucleated cells. In this phenomenon, activation of membrane  $\text{Ca}^{2+}$  channels and increase in cytoplasmic  $\text{Ca}^{2+}$  led to several modifications including cell shrinkage and translocation of phosphatidylserine (PS) from intracellular to extracellular leaflet of the cell membrane. Despite the cytoplasmic  $\text{Ca}^{2+}$  activity, eryptosis could be stimulated as a result of increase in ceramide, oxidative stress, energy depletion, and activated caspases as well. Externalization of PS mediated phagocytosis of erythrocyte by the macrophages and their clearance from the blood circulation can lead to anemia [13, 14]. The role of enhanced eryptosis in the pathogenesis of anemia associated disorders including sickle cell disease, thalassemia, and glucose-phosphate dehydrogenase deficiency have been demonstrated as well [15]. Lead [16], mercury [17], and gold [18] are among metal ions that can induce eryptosis. Although several mechanisms may underlie the initiation of eryptosis, an increase in oxidative stress is among the most cited mechanisms. One of the pieces of evidence in this issue is the protective impacts of several synthetic and natural antioxidants on this phenomenon [19]. Vitamin C (vit C) is an antioxidant vitamin that is generally used for ameliorating the adverse effects of toxins [20], cancer [21], and other conditions associated with enhanced-oxidative stress. It has been revealed that vit C can ameliorate the toxic effects of Al [22–24]. To the best of our knowledge, the effect of vit C supplementation against Al-induced hemolysis and eryptosis has not been studied. Therefore, the present study was designed to explore the possibility of Al-induced eryptosis, which might be ameliorated by vit C or N-acetylcysteine (NAC). To this end, the probable protective effect of vit C and NAC on  $\text{AlCl}_3$  induced hemolysis and eryptosis-related characteristics including PS exposure and oxidative stress was determined on the erythrocytes.

## 2. Materials and Methods

**2.1. Chemicals.** Ringer solution (pH 7.4) was prepared from NaCl (125 mM), KCl (5 mM),  $\text{MgSO}_4$  (1 mM),  $\text{CaCl}_2$  (1 mM), glucose (5 mM). In  $\text{Ca}^{2+}$ -free Ringer,  $\text{CaCl}_2$  (1 mM) was substituted by a similar amount of EGTA (1 mM).  $\text{AlCl}_3$  and other chemical reagents were purchased from Sigma, USA.

**2.2. Assay of Hemolysis.** The ex vivo hemolysis assay was conducted on the erythrocytes obtained from healthy volunteers (6 replicates/group) whom informed consent. The study has been approved by the ethics committee of Fasa University of Medical Sciences. In brief, erythrocytes (0.4% hematocrit) were incubated at  $37^\circ\text{C}$  for 24 hours in the absence (control group) or presence of  $\text{AlCl}_3$  (3–100  $\mu\text{M}$ ). The samples were centrifuged (3 min at  $400 \times g$ , RT) and the supernatants were harvested. Thereafter, the absorbance of hemoglobin (Hb) in the supernatants was determined at 405 nm. 100% hemolysate in hemolysis tests was prepared by mixing the erythrocyte solution with distilled water. As a

measure of hemolysis, the percent of hemolysis in each tube is calculated by dividing the absorbance of the test tube by the absorbance of 100% hemolysis [25].

**2.3. Flowcytometric Assay for Annexin V-PE Binding and Forward Scatter.** Flowcytometry was used to quantify the percent of Annexin V-PE positive erythrocyte and also to measure forward scatter [25]. In brief, after 24 hours of incubation, cells were washed in Ringer solution. Erythrocytes were then stained with Annexin V-PE (EXBIO Praha, Czech Republic) at 1:500 dilutions. After 15 min of incubation in the dark, samples were quantified using flowcytometric analysis (FACS Calibur from Becton Dickinson; Heidelberg, Germany). Cells were analyzed by forward scatter and annexin-V-fluorescence intensity was measured in fluorescence channel FL-2 with an excitation wavelength of 488 nm and an emission wavelength of 580 nm.

**2.4. Measurement of Intracellular  $\text{Ca}^{2+}$ .** Intracellular  $\text{Ca}^{2+}$  was measured using Fluo<sub>3</sub>/AM flowcytometry [25]. In brief, erythrocytes were washed in Ringer solution and then loaded with Fluo<sub>3</sub>/AM (Calbiochem; Bad Soden, Germany) in Ringer solution containing 5 mM  $\text{CaCl}_2$  and 2  $\mu\text{M}$  Fluo<sub>3</sub>/AM. The cells were incubated at  $37^\circ\text{C}$  for 20 min and washed twice in Ringer solution containing 5 mM  $\text{CaCl}_2$ . After that, the Fluo<sub>3</sub>/AM loaded erythrocytes, which were resuspended in 200  $\mu\text{l}$ , Ringer. Ionomycin (1.0  $\mu\text{M}$  for 30 min) was used as a positive control, though. Moreover,  $\text{Ca}^{2+}$ -dependent fluorescence intensity was determined in fluorescence channel FL-1.

**2.5. Reactive Oxygen Species (ROS) Assay.** 2', 7'-dichlorodihydrofluorescein diacetate (DCFDA, Sigma) was used in order to determine the role of oxidative stress in Al-induced eryptosis fluorescent ROS assay [25]. In this assay, DCFDA was oxidized in the presence of ROS and becomes green fluorescent. In brief, erythrocytes were incubated with  $\text{AlCl}_3$  (100  $\mu\text{M}$ ) in the presence and absence of vit C (0.6 mM) or NAC (1 mM). After the treatment, 150  $\mu\text{l}$  suspension of erythrocytes was washed in Ringer solution and stained with DCFDA (10  $\mu\text{M}$  in Ringer solution) at  $37^\circ\text{C}$  for 30 min in the dark. After washing with the Ringer solution, the DCFDA-stained erythrocytes were resuspended in 200  $\mu\text{l}$  Ringer solution and DCFDA fluorescence intensity was measured in the FL-1 channel at an excitation wavelength of 488 nm and an emission wavelength of 530 nm on a FACS Calibur (BD). The geomean of the DCFDA dependent fluorescence was then determined as well.

**2.6. Determination of Reduced Glutathione (GSH).** The level of erythrocytes glutathione (GSSG and GSH) was evaluated using Glutathione Assay Kit (Cayman Chemicals, IBL Hamburg, Hamburg, Germany) according to the manufacturer's protocol. In brief, erythrocytes were washed twice in PBS, incubating for 24 h at  $37^\circ\text{C}$  in Ringer solution in the absence or presence of different concentrations of  $\text{Al}^{3+}$  (3–100  $\mu\text{M}$ ). Following that, the erythrocytes were then

washed in PBS. Then, 50  $\mu\text{l}$  of the erythrocyte pellet was lysed in 200  $\mu\text{l}$  distilled water, centrifuged at  $14,000 \times g$ , and 150  $\mu\text{l}$  of the supernatant was deproteinated by adding 150  $\mu\text{l}$  metaphosphoric acid (10%). GSSG and GSH were then measured with the Glutathione Assay Kit (measured via ELISA reader within the wavelength of 405–414 nm).


**2.7. Statistical Analyses.** SPSS statistical software (SPSS, Chicago, IL, USA, version 21) was applied to analyze the data. Normal distribution of data was checked using the Shapiro–Wilk test ( $P < 0.05$ ). Nonparametric Kruskal–Wallis test was used for the comparison of data between control and experimental groups. The data are represented as mean  $\pm$  standard deviation (SD) of at least three independent experiments.  $P < 0.05$  was considered to be statistically significant.

### 3. Results

**3.1. The Effect of Al on Erythrocytes Hemolysis.** To explore the effect of Al on hemolysis, erythrocytes were treated with  $\text{AlCl}_3$  (3–100  $\mu\text{M}$ ) for 24 hours and the percentage of hemolysis was determined from the absorbance of hemoglobin in the supernatant. As illustrated in Figure 1, the treatment of erythrocytes with various concentrations, from 10 to 100  $\mu\text{M}$ , of  $\text{AlCl}_3$  increased the level of hemolysis in a dose-dependent manner compared to that of the control group, while 3  $\mu\text{M}$  concentration of  $\text{AlCl}_3$  had no significant effects on hemolysis compared to control group.

To explore whether the entry of extracellular  $\text{Ca}^{2+}$  to the erythrocytes was a prerequisite for Al-induced hemolysis or not, erythrocytes were incubated for 24 hours with various concentrations of  $\text{AlCl}_3$  in the presence of 1 mM of EGTA, as a calcium chelating compound. As can be seen in Figure 1, EGTA significantly decreased the effect of  $\text{AlCl}_3$  on hemolysis to that of control cells. Thus, Al may induce hemolysis by the stimulation of extracellular  $\text{Ca}^{2+}$  entry. Further experiments were performed to examine the possible impacts of antioxidants including vit C and NAC, on Al-induced hemolysis. To this end, erythrocytes were treated with  $\text{AlCl}_3$  in the absence and presence of vit C (0.6 mM) or NAC (1 mM). As demonstrated in Figure 2, NAC treatment decreased the level of Al-induced hemolysis significantly (Figure 2), suggesting the possible role of oxidative stress in Al-induced hemolysis. Although, cotreatment with vit C partially decreased the percent of hemolysis in Al-treated cells; no significant effects were observed compared to that of the cells treated with  $\text{AlCl}_3$  alone. vit C at the higher concentrations (>1 mM) significantly increased the level of hemolysis compared to the control group.

**3.2. The Effect of Vit C and NAC on Al-Induced Eryptosis.** Eryptosis is characterized by three main hallmarks including membrane blebbing, cell shrinkage, and PS externalization. As can be seen in Figure 3(b), a 24 hours' exposure to  $\text{AlCl}_3$  induced membrane blebbing in the erythrocytes. Furthermore,  $\text{AlCl}_3$  (30 and 100  $\mu\text{M}$ )



**FIGURE 1:** Effect of  $\text{AlCl}_3$  on hemolysis in the absence and presence of EGTA.  $\text{AlCl}_3$  increased hemolysis in a dose-dependent manner. EGTA (1 mM) decreased the effect of Al on erythrocyte hemolysis. Data are mean  $\pm$  SD of at least five independent experiments. Nonparametric Kruskal–Wallis test was used for the statistical analyses.  $P < 0.05$  was considered as significant statistical differences between the group.  $P$  values compared to the control untreated control group: \* $P = 0.028$ , \*\* $P = 0.002$ , \*\*\* $P = 0.0001$ , and  $P$  values compared to corresponding  $\text{AlCl}_3$ -treated groups: ## $P = 0.027$ , ### $P = 0.045$ .

increased the percentage of annexin V positive cells compared to the control group (Figures 3(c) and 3(d)), suggesting PS externalization by  $\text{AlCl}_3$ .  $\text{AlCl}_3$  also significantly decreased the forward scatter compared to that of the control group ( $P < 0.001$ ). All these pieces of evidence strongly suggest that  $\text{AlCl}_3$  induced eryptosis in the erythrocytes. NAC cotreatment (1 mM) blunted the effect of Al (100  $\mu\text{M}$ ) on PS externalization (Figure 4(b)). Unexpectedly, treatment with vit C alone significantly increased PSE compared to the control group. Furthermore, cotreatment of vit C with Al significantly augmented the effects of Al on the induction of PS externalization.

**3.3. The Effect of Al on Intracellular Co-Ca<sup>2+</sup> Concentration Using the Fluo<sub>3</sub>-Fluorescence Method.** An increase in intracellular  $\text{Ca}^{2+}$  concentration could trigger eryptosis. In this study, the Fluo<sub>3</sub> flow cytometry method was used in order to test whether Al could alter the intracellular concentration of  $\text{Ca}^{2+}$ . As can be seen in Figure 5, ionomycin (1  $\mu\text{M}$ ), as a positive control, increased the percent of Fluo<sub>3</sub>-fluorescence cells. Furthermore, incubation of erythrocytes for 24 hours with  $\text{AlCl}_3$  at 30 and 100  $\mu\text{M}$  concentrations increased the percent of Fluo<sub>3</sub> fluorescence cells significantly compared to the control group. However,  $\text{AlCl}_3$  at the concentration of 3 and 10  $\mu\text{M}$  had no significant effects on intracellular  $\text{Ca}^{2+}$ .




FIGURE 2: Effect of vitamin C (+vit C) and n-acetyl cysteine (+NAC) on Al-induced hemolysis. AlCl<sub>3</sub> increase hemolysis in a dose-dependent manner. NAC (1 mM) decreased Al-induced hemolysis while vit C (0.6 mM) had no significant effects on Al-induced hemolysis. Data are mean  $\pm$  SD of at least five independent experiments. Nonparametric Kruskal–Wallis was used for the statistical analyses.  $P < 0.05$  was considered as significant statistical differences between the group.  $P$  values compared to the untreated control group (\* $P = 0.028$ , \*\* $P = 0.002$ , \*\*\* $P = 0.0001$ ), significant differences compared to corresponding AlCl<sub>3</sub>-treated groups (# $P = 0.031$ , ## $P = 0.035$ , ### $P = 0.038$ ).

**3.4. The Effect of Al on Reactive Oxygen Species Production Using DCFDA Fluorescence Method.** Reactive oxygen species (ROS) can cause eryptosis. Therefore, the effects of Al on ROS production and possible ameliorating effects of vit C and NAC on this phenomenon were estimated by DCFDA flowcytometry. As illustrated in Figure 6, a 24 hours' exposure to 100  $\mu$ M concentration of AlCl<sub>3</sub> significantly increased the DCFDA fluorescence of erythrocytes. Moreover, treatment of erythrocytes with AlCl<sub>3</sub> (100  $\mu$ M) in the presence of vit C (0.6 mM) and NAC (1 mM) significantly decreased DCFDA flowcytometry, suggesting a possible role of oxidative stress in the Al-induced eryptosis.

**3.5. The Effect of Al on Glutathione Level in the Erythrocytes.** As depicted in Figure 7, a 24 hours' exposure to 100  $\mu$ M concentration of AlCl<sub>3</sub> had no significant effects on total glutathione, GSH, GSSG, and GSSG/GSH ratio compared to the control group.

## 4. Discussion

The morphological and biochemical alteration occurred following the 24 h exposure of erythrocytes with Al including membrane blebbing, cell shrinkage, increase in the PS externalization, increase in intracellular Ca<sup>2+</sup> level, and elevation of ROS levels, in order to provide clear evidence that Al can trigger eryptosis of erythrocytes in a dose-dependent manner. In addition, we have found that NAC attenuated Al-induced accumulation of ROS and inhibited the Al-induced eryptosis and hemolysis. Although vit C has decreased Al-induced accumulation of ROS, it would surprisingly have failed to reduce Al-induced hemolysis and also increased Al-induced eryptosis of the erythrocytes.

The critical role of oxidative stress in the induction of eryptosis has been demonstrated in numerous studies [15]. However, contradicting results have been reported about the effects of Al on the oxidative stress status of erythrocytes. While Niemoeller et al. [11] has demonstrated that oxidative stress did not involve Al-induced eryptosis, Vota et al. [12]




FIGURE 3: AlCl<sub>3</sub> induced membrane blebbing, PS externalization, and cell shrinkage in the erythrocytes. Micrographs showed normal shape erythrocytes in the control group (a) and membrane blebbing following treatment of erythrocytes with Ringer solution containing AlCl<sub>3</sub> (30 μM) (b). Incubation with AlCl<sub>3</sub> (30 and 100 μM) increased annexin V positive cells (c) and (d). Forward scatter in the erythrocytes ((e) and (f)) following exposure for 24 hours to Ringer solution (control; grey area), AlCl<sub>3</sub> (100 μM; black line).

have represented the chronic exposure of erythrocytes with AlCl<sub>3</sub> which increased the ROS production and decreased the GSH content of the erythrocytes, suggesting the role of oxidative stress in Al-induced effects. In the present investigation, we observed no changes in GSH level and GSH/GSSG following exposure of erythrocyte for 24 hours with Al, while flowcytometric analyses, using DCFDA fluorescence, showed that the ROS content could increase the following Al exposure. Meanwhile, the findings showed that NAC could inhibit the effects of Al on ROS accumulation, eryptosis, and hemolysis, suggesting that oxidative stress

may be a downstream mechanism in the Al-induced hemolysis and eryptosis. Plasma membranes of erythrocytes have a high content of polyunsaturated fatty acid [26]; hence, they are highly sensitive to oxidative stress and lipid peroxidation which disrupts the membrane integrity of erythrocytes and results in hemolysis [27]. The inhibition of Al-induced hemolysis by NAC suggested that observed Al-induced hemolysis may be attributed to excessive ROS production and oxidative stress.

It is well-known that vit C keeps cell components including cell membranes against ROS either directly or




FIGURE 4: The effect of vit C and NAC on Al-induced PS externalization using Annexin V-PE flowcytometric methods. (a) Histogram of PS externalization of erythrocytes following exposure for 24 hours to Ringer solution (control; grey area), AlCl<sub>3</sub> (100 μM; black line), AlCl<sub>3</sub> (100 μM) and vit C (0.6 mM; dashed line), AlCl<sub>3</sub> (100 μM) and NAC (1.0 mM; dotted line). (b) Arithmetic means ± SD of the percentage of erythrocyte with enhanced PS externalization following treatment with AlCl<sub>3</sub> in the presence and absence of vit C (0.6 mM) and NAC (1 mM) in Ringer solution for 24 hours. \* and \*\*  $P < 0.05$  significant difference compared to the AlCl<sub>3</sub>-treated group.




FIGURE 5: The effects of various concentrations of AlCl<sub>3</sub> on intracellular Ca<sup>2+</sup> levels in the erythrocytes. (a) Histogram of Fluo<sub>3</sub>-fluorescence of the erythrocytes following exposure for 24 hours to Ringer solution without (control; grey area) and with the presence of 30 μM (black line) and 100 μM of AlCl<sub>3</sub> μM (dotted line) and 1 μM of ionomycin (positive control; dashed line). (b) Arithmetic means ± SD of the percentage of erythrocyte with enhanced Fluo<sub>3</sub>-fluorescence following treatment with various concentrations of AlCl<sub>3</sub> and ionomycin (1 μM), in Ringer solution for 24 hours. \*, \*\*, and \*\*\*  $P < 0.001$  significant difference compared to control groups.

indirectly via the regeneration of oxidized vitamin E. However, contradicting results have been reported for the effects of vit C on erythrocytes eryptosis and hemolysis. In a recent study, Shan et al. showed that vit C attenuated H<sub>2</sub>O<sub>2</sub>-induced eryptosis in the erythrocytes of human glucose 6-phosphate deficient patients through inhibiting

ROS accumulation [28], whereas other studies demonstrated the hemolytic effects of vit C [29]. In agreement with the well-known antioxidant effects of vit C, the data from the current study revealed that vit C decreased ROS production induced by Al. However, vit C induced PS externalization by itself and caused a drastic increment in




FIGURE 6: The effects of AlCl<sub>3</sub> on the formation of reactive oxygen species. (a) Original histogram of DCFDA fluorescence in erythrocytes following exposure for 24 hours to Ringer solution (control, grey areas), AlCl<sub>3</sub> (100 μM, black lines) AlCl<sub>3</sub> (100 μM) in the presence of vit C (0.6 mM, dotted lines) and NAC (1 mM, dashed lines). (b) Geo means ± SD of DCFDA fluorescence in erythrocytes after a 24 hours' treatment with AlCl<sub>3</sub> (100 μM) in the presence of vit C (AlCl<sub>3</sub> + vit C) and NAC (AlCl<sub>3</sub> + NAC). \*\*\*P < 0.001 compared to the control group, #P < 0.01 compared to the AlCl<sub>3</sub> group, and \*\*P < 0.05 compared to the AlCl<sub>3</sub> group.




FIGURE 7: The effects of AlCl<sub>3</sub> on erythrocyte glutathione level. Total glutathione, reduced glutathione (GSH), and oxidized glutathione (GSSG) were determined in erythrocytes following 24 hours' exposure to AlCl<sub>3</sub> (100 μM). No significant differences were found between the control and AlCl<sub>3</sub>-treated group.

Al-induced eryptosis. The exact mechanism by which vit C induced PS externalization has not been clear yet. One possible explanation may be autooxidation of vit C in the presence of Al<sup>3+</sup> which was associated with the production of cytotoxic substances [21]. Moreover, vit C can readily be oxidized to dehydroascorbate (DHA), which is readily transported into erythrocytes by glucose transporter1 (Glut1), abundant in the membrane of erythrocytes. Intracellular DHA is rapidly converted to vit C by GSH dependent reaction. The depletion of GSH eventually

induces oxidative stress in the erythrocytes [21]. Further investigations will be needed to understand the interaction between Al<sup>3+</sup> and vit C.

In addition to ROS production and induction of oxidative stress, it has been demonstrated that disturbance in Ca<sup>2+</sup> hemostasis (involved in Al-mediated toxicity) and calcium channel blocker was useful for the amelioration of Al's toxic effects [30]. The findings of the current studies clearly demonstrated that Al increased the accumulation of Ca<sup>2+</sup> in the erythrocytes while calcium ion chelating agent and EGTA (1 mM) have attenuated the Al-induced hemolysis. Thus the observed effects of Al may be secondary to Ca<sup>2+</sup> entry from the extracellular space to erythrocytes, which is known to trigger eryptosis.

### 5. Conclusion

In conclusion, the investigation revealed that Al-induced eryptosis and hemolysis of erythrocytes, increased in ROS production, and Ca<sup>2+</sup> entry to erythrocytes which were underlying mechanisms in Al-induced effects. NAC ameliorated Al-induced hemolysis and eryptosis through its antioxidant effects. On the other hand, vit C did not have similar protective effects on eryptosis. At the same time, it has increased eryptosis at particular doses and suggested that ex vivo vit C was not a suitable compound against Al-induced hemolytic and eryptotic effect. On the contrary, NAC would be a better choice to exert a protective effect on cellular eryptosis.

### Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

## Conflicts of Interest

The authors declare no conflicts of interest.

## Acknowledgments

The authors would like to acknowledge the Fasa University of Medical Sciences and the Shiraz University of Medical Sciences for providing facilities and laboratory equipment for this study.

## References

- [1] K. Klotz, W. Weistenhofer, F. Neff, A. Hartwig, C. van Thriel, and H. Drexler, "The health effects of aluminum exposure," *Neurobiology of Aging*, vol. 114, pp. 653–659, 2017.
- [2] Z. Wu, Y. Du, H. Xue, Y. Wu, and B. Zhou, "Aluminum induces neurodegeneration and its toxicity arises from increased iron accumulation and reactive oxygen species (ROS) production," *Neurobiology of Aging*, vol. 33, p. 199, 2012.
- [3] J. Saberzadeh, R. Arabsolghar, and M. A. Takhshid, "Alpha synuclein protein is involved in Aluminum-induced cell death and oxidative stress in PC12 cells," *Brain Research*, vol. 1635, pp. 153–160, 2016.
- [4] S. Han, J. Lemire, V. P. Appanna, C. Auger, Z. Castonguay, and V. D. Appanna, "How aluminum, an intracellular ROS generator promotes hepatic and neurological diseases: the metabolic tale," *Cell Biology and Toxicology*, vol. 29, no. 2, pp. 75–84, 2013.
- [5] D. Chappard, P. Bizot, G. Mabileau, and L. Hubert, "Aluminum and bone: review of new clinical circumstances associated with Al<sup>3+</sup> deposition in the calcified matrix of bone," *Morphologie*, vol. 100, no. 329, pp. 95–105, 2016.
- [6] F. Geyikoglu, H. Türkez, T. O. Bakir, and M. Cicek, "The genotoxic, hepatotoxic, nephrotoxic, haematotoxic and histopathological effects in rats after aluminium chronic intoxication," *Toxicology and Industrial Health*, vol. 29, no. 9, pp. 780–791, 2013.
- [7] Y. Belaïd-Nouira, H. Bakhta, Z. Haouas, I. Flehi-Slim, and H. Ben Cheikh, "Fenugreek seeds reduce aluminum toxicity associated with renal failure in rats," *Nutrition Research and Practice*, vol. 7, no. 6, pp. 466–474, 2013.
- [8] E. Cheraghi, A. Golkar, K. Roshanaei, and B. Alani, "Aluminium-induced oxidative stress, apoptosis and alterations in testicular tissue and sperm quality in wistar rats: ameliorative effects of curcumin," *International Journal of Fertility & Sterility*, vol. 11, no. 3, pp. 166–175, 2017.
- [9] C. C. Willhite, N. A. Karyakina, R. A. Yokel et al., "Systematic review of potential health risks posed by pharmaceutical, occupational and consumer exposures to metallic and nanoscale aluminum, aluminum oxides, aluminum hydroxide and its soluble salts," *Critical Reviews in Toxicology*, vol. 44, pp. 1–80, 2014.
- [10] C.-Y. Lin, W.-C. Hsiao, C.-J. Huang, C.-F. Kao, and G.-S. W. Hsu, "Heme oxygenase-1 induction by the ROS-JNK pathway plays a role in aluminum-induced anemia," *Journal of Inorganic Biochemistry*, vol. 128, pp. 221–228, 2013.
- [11] N. V. Kiedaisch, P. Dreischer, T. Wieder, and F. Lang, "Stimulation of eryptosis by aluminium ions," *Journal of Cellular Biochemistry*, vol. 217, pp. 168–175, 2006.
- [12] D. M. Vota, R. L. Crisp, A. B. Nesse, and D. C. Vittori, "Oxidative stress due to aluminum exposure induces eryptosis which is prevented by erythropoietin," *Journal of Cellular Biochemistry*, vol. 113, no. 5, pp. 1581–1589, 2012.
- [13] L. Repsold and A. M. J. Eryptosis, "An erythrocyte's suicidal type of cell death," *Journal of Cellular Biochemistry*, vol. 2018, 2018.
- [14] M. MischitellJemaà, M. Jemaà, F. Lang, C. Faggio, and F. Lang, "Ca<sup>2+</sup> entry, oxidative stress, ceramide and suicidal erythrocyte death following diosgenin treatment," *Cellular Physiology and Biochemistry*, vol. 39, no. 4, pp. 1626–1637, 2016.
- [15] E. Lang and F. L. Triggers, "Inhibitors, mechanisms, and significance of eryptosis: the suicidal erythrocyte death," *BioMedical Research Institute*, vol. 2015, 2015.
- [16] I. C. Aguilar-Dorado, G. Hernandez, M. A. Quintanar-Escorza, M. Maldonado-Vega, M. Rosas-Flores, and J. V. Calderon-Salinas, "Eryptosis in lead-exposed workers," *Cellular Physiology and Biochemistry*, vol. 281, pp. 195–202, 2014.
- [17] A. Officioso, K. Alzoubi, F. Lang, and C. Manna, "Hydroxytyrosol inhibits phosphatidylserine exposure and suicidal death induced by mercury in human erythrocytes: possible involvement of the glutathione pathway," *Food Chemical Toxicology*, vol. 89, pp. 47–53, 2016.
- [18] M. Sopjani, M. Foller, and F. Lang, "Gold stimulates Ca<sup>2+</sup> entry into and subsequent suicidal death of erythrocytes," *Toxicology*, vol. 244, pp. 271–279, 2008.
- [19] A. Fazio, M. Briglia, C. Faggio, K. Alzoubi, and F. Lang, "Oxaliplatin induced suicidal death of human erythrocytes," *Cellular Physiology and Biochemistry*, vol. 37, no. 6, pp. 2393–2404, 2015.
- [20] M. A. Takhshid, A. R. Tavasuli, Y. Heidary, M. Keshavarz, and H. Kargar, "Protective effect of vitamins e and C on endosulfan-induced reproductive toxicity in male rats," *Iran Journal of Medical Science*, vol. 37, pp. 173–180, 2012.
- [21] J. Du, J. Cullen, and G. R. Buettner, "Ascorbic acid: chemistry, biology and the treatment of cancer," *Cellular Physiology and Biochemistry*, vol. 1826, pp. 443–457, 2012.
- [22] O. J. Olajide, E. O. Awson, I. T. Gbadamosi et al., "Ascorbic acid ameliorates behavioural deficits and neuropathological alterations in rat model of Alzheimer's disease," *Environmental Toxic Pharmacology*, vol. 50, pp. 200–211, 2017.
- [23] M. I. Yousef, "Aluminium-induced changes in hemato-biochemical parameters, lipid peroxidation and enzyme activities of male rabbits: protective role of ascorbic acid," *Toxicology*, vol. 199, pp. 47–57, 2004.
- [24] F. Gultekin, N. Delibas, S. Yasar, and I. Kilinc, "In Vivo changes in antioxidant systems and protective role of melatonin and a combination of vitamin C and vitamin E on oxidative damage in erythrocytes induced by chlorpyrifos-ethyl in rats," *Toxicology*, vol. 75, pp. 88–96, 2001.
- [25] A. Al Mamun Bhuyan, S. Nussle, H. Cao, S. Zhang, and F. Lang, "Simvastatin, a novel stimulator of eryptosis, the suicidal erythrocyte death," *Cell Physiology Biochemistry*, vol. 43, pp. 492–506, 2017.
- [26] P. Koehrer, S. Saab, O. Berdeaux et al., "Erythrocyte phospholipid and polyunsaturated fatty acid composition in diabetic retinopathy," *PLoS ONE*, vol. 9, 2014.
- [27] K. B. Pandey and S. I. Rizvi, "Markers of oxidative stress in erythrocytes and plasma during aging in humans," *PLoS ONE*, vol. 3, pp. 2–12, 2010.
- [28] F. Shan, R. Yang, T. Ji, and F. Jiao, "Vitamin C inhibits aggravated eryptosis by hydrogen peroxide in glucose-6-phosphated dehydrogenase deficiency," *Cell Physiology Biochemistry*, vol. 39, pp. 1453–1462, 2016.
- [29] Z. Zhang, E. Lee, J. Sudderth et al., "Glutathione depletion, pentose phosphate pathway activation, and hemolysis in



erythrocytes protecting cancer cells from vitamin C-induced oxidative stress,” *Journal of Biology Chemistry*, vol. 28, pp. 22861–22867, 2016.

- [30] J. Saberzadeh, M. Omrani, and M. A. Takhshid, “Protective effects of nimodipine and lithium against aluminum-induced cell death and oxidative stress in PC12 cells,” *Iran Journal of Basic Medical Science*, vol. 19, pp. 1251–1257, 2016.