

# Ammonium [(1*S*)-(endo,*anti*)]-(−)-3-bromocamphor-8-sulfonate

Muhammad Athar Abbasi,<sup>a,‡</sup> Aziz-ur-Rehman,<sup>a</sup> Mehmet Akkurt,<sup>b,\*</sup> Muhammad Jahangir,<sup>a</sup> Seik Weng Ng<sup>c</sup> and Islam Ullah Khan<sup>a</sup>

<sup>a</sup>Department of Chemistry, Government College University, Lahore 54000, Pakistan,

<sup>b</sup>Department of Physics, Faculty of Arts and Sciences, Erciyes University, 38039

Kayseri, Turkey, and <sup>c</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: akkurt@erciyes.edu.tr


Received 4 June 2010; accepted 14 June 2010

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.027;  $wR$  factor = 0.065; data-to-parameter ratio = 16.5.

In the title molecular salt,  $\text{NH}_4^+\cdot\text{C}_{10}\text{H}_{14}\text{BrO}_4\text{S}^-$ , the norbornane skeleton of the anion is composed of two five-membered rings in envelope conformations and a six-membered ring with one Br atom, one carbonyl O atom and a methyl group held in a boat conformation by a bridging methylene group. Short intramolecular  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{Br}$  interactions occur. In the crystal, the component ions are linked by intermolecular  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds.

## Related literature

For further synthetic details, see: Smith *et al.* (2008). For other structures with the norbornane skeleton, see: Jauch *et al.* (1992); Ustabaş *et al.* (2006); Ersanlı *et al.* (2005). For the use of 3-bromocamphor-8-sulfonic acid and its ammonium salts as chiral auxiliaries for the optical resolution of enantiomeric amines through diastereomeric salt formation, see: Bálint *et al.* (1999); Pellati *et al.* (2010); Roy *et al.* (2009); Zhao *et al.* (2002). For puckering parameters, see: Cremer & Pople (1975).



<sup>‡</sup> Additional corresponding author, e-mail: atrabbasi@yahoo.com.

## Experimental

### Crystal data

$\text{NH}_4^+\cdot\text{C}_{10}\text{H}_{14}\text{BrO}_4\text{S}^-$

$M_r = 328.22$

Monoclinic,  $P2_1$

$a = 7.2449$  (2) Å

$b = 7.0049$  (1) Å

$c = 13.2428$  (3) Å

$\beta = 104.704$  (1)°

$V = 650.06$  (3) Å<sup>3</sup>

$Z = 2$

Mo  $K\alpha$  radiation

$\mu = 3.33$  mm<sup>-1</sup>

$T = 296$  K

$0.42 \times 0.14 \times 0.11$  mm

### Data collection

Bruker Kappa APEXII CCD diffractometer

Absorption correction: refined from  $\Delta F$  (XABS2; Parkin *et al.*, 1995)

$T_{\min} = 0.336$ ,  $T_{\max} = 0.711$

2775 measured reflections

2775 independent reflections

2586 reflections with  $I > 2\sigma(I)$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.065$

$S = 1.03$

2775 reflections

168 parameters

5 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.35$  e Å<sup>-3</sup>

$\Delta\rho_{\min} = -0.47$  e Å<sup>-3</sup>

Absolute structure: Flack (1983), 1155 Freidel pairs

Flack parameter:  $-0.021$  (7)

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O1}^{\text{i}}$	0.92 (3)	1.92 (3)	2.835 (4)	173 (3)
$\text{N1}-\text{H2N}\cdots\text{O2}^{\text{ii}}$	0.90 (3)	2.05 (3)	2.899 (3)	157 (3)
$\text{N1}-\text{H3N}\cdots\text{O2}$	0.92 (3)	1.97 (3)	2.887 (3)	176 (3)
$\text{N1}-\text{H4N}\cdots\text{O3}^{\text{iii}}$	0.92 (3)	1.93 (3)	2.827 (3)	167 (4)
$\text{C4}-\text{H4B}\cdots\text{Br1}$	0.97	2.71	3.221 (3)	113
$\text{C8}-\text{H8A}\cdots\text{O2}$	0.96	2.44	3.104 (3)	126
$\text{C10}-\text{H10}\cdots\text{O1}^{\text{i}}$	0.98	2.49	3.451 (4)	167

Symmetry codes: (i)  $x, y - 1, z$ ; (ii)  $-x + 2, y - \frac{1}{2}, -z + 1$ ; (iii)  $-x + 1, y - \frac{1}{2}, -z + 1$ .

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999), PARST (Nardelli, 1983) and PLATON (Spek, 2009).

The authors are grateful to the Higher Education Commission for financial support to purchase the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5484).

## References

- Bálint, J., Egri, G., Fogassy, E., Böcskei, Z., Simon, K., Gajáry, A. & Friesz, A. (1999). *Tetrahedron Asymmetry*, **10**, 1079–1087.
- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Ersanlı, C. C., Çoruh, U., Hökelek, T., Vázquez-López, E. M. & Daştan, A. (2005). *Acta Cryst.* **E61**, o263–o265.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.

- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Jauch, J., Laderer, H. & Walz, L. (1992). *Acta Cryst.* **C48**, 1246–1248.
- Nardelli, M. (1983). *Comput. Chem.* **7**, 95–98.
- Parkin, S., Moezzi, B. & Hope, H. (1995). *J. Appl. Cryst.* **28**, 53–56.
- Pellati, F., Cannazza, G. & Benvenuti, S. (2010). *J. Chromatogr. A*, **1217**, 3503–3510.
- Roy, B. N., Singh, G. P., Srivastava, D., Jadhav, H. S., Saini, M. B. & Aher, P. (2009). *Org. Process Res. Dev.* **13**, 450–455.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Smith, E. D., Vinson, N. A., Zhong, D., Berrang, B. D., Catanzaro, J. L., Thomas, J. B., Navarro, H. A., Gilmour, B. P., Deschamps, J. & Carroll, F. I. (2008). *Bioorg. Med. Chem.* **16**, 822–829.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Ustabaş, R., Çoruh, U., Yavuz, M., Salamci, E. & Vázquez-López, E. M. (2006). *Acta Cryst.* **E62**, o1149–o1150.
- Zhao, M. M., McNamara, J. M., Ho, G.-J., Emerson, K. M., Song, Z. J., Tschaen, D. M., Brands, K. M. J., Dolling, U.-H., Grabowski, E. J. J. & Reider, P. J. (2002). *J. Org. Chem.* **67**, 6743–6747.

**supplementary materials**

*Acta Cryst.* (2010). E66, o1707-o1708 [ doi:10.1107/S1600536810022804 ]

## Ammonium [(1*S*)-(endo,anti)]-(-)-3-bromocamphor-8-sulfonate

M. A. Abbasi, Aziz-ur-Rehman, M. Akkurt, M. Jahangir, S. W. Ng and I. U. Khan

### Comment

3-Bromocamphor-8-sulfonic acid and its ammonium salts have extensively been used as chiral auxiliaries for the optical resolution of a number of enantiomeric amines through diastereomeric salt formation (Bálint *et al.*, 1999; Zhao *et al.*, 2002; Roy *et al.*, 2009; Pellati *et al.*, 2010).

In the bicyclo[2.2.1]heptane (norbornane) skeleton of the title compound, (I), (Fig. 1), the two five-membered rings have envelope conformations, with atom C2 displaced by 0.365 (3) Å from the C2–C6 plane [the puckering parameters (Cremer & Pople, 1975) are  $Q_2 = 0.573$  (3) Å and  $\varphi_2 = 5.3$  (3)°] and by 0.397 (3) Å from the C2/C3/C6/C9/C10 plane [the puckering parameters:  $Q_2 = 0.615$  (3) Å and  $\varphi_2 = 181.6$  (3)°] and the six-membered ring (C3–C6/C9/C10) adopts a boat conformation by the puckering parameters  $Q_T = 0.970$  (3) Å,  $\theta = 92.03$  (18)° and  $\varphi = 357.34$  (19)°.

In (I), the C—C single-bond lengths range from 1.491 (5) to 1.575 (4) Å, with a mean value of 1.535 (4) Å. In the bicyclo[2.2.1]heptane fragment, the angles between planes A (C3/C2/C6), B (C3–C6) and C (C3/C6/C9/C10) are as follows: A/B= 53.65 (19)°, A/C= 58.14 (18)° and B/C= 68.22 (13)°.

In the crystal, adjacent molecules of (I) are linked by intermolecular N—H···O and C—H···O hydrogen bonds (Table 1, Fig. 2).

### Experimental

3-Bromocamphor-8-sulfonic acid ammonium salt was prepared by modification in the reported method (Smith *et al.*, 2008). 3-Bromocamphor-8-sulfonic acid (1 g) was dissolved in 15 ml of ethanol and then 6 ml of NH<sub>3</sub> solution were added. The mixture was stirred until a clear solution was observed (about 20 min). The solution was slowly concentrated on water bath to half the volume over a 2 h period. The concentrate was allowed to crystallize undisturbed for 48 h. The resulting colourless prisms of (I) were carefully separated by filtration and washed with three 0.5-ml portions of petroleum ether.

### Refinement

In the ammonium ion, H atoms bound to N atoms were located in difference Fourier maps and their positional parameters were refined freely using a *DFIX* instruction [N—H = 0.93 (3) Å] in *SHELXL97*, with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{N})$ . H atoms bound to C atoms were placed in idealized positions and refined using a riding model with C—H = 0.96, 0.97 and 0.98 Å for CH<sub>3</sub>, CH<sub>2</sub> and CH, respectively.  $U_{\text{iso}}(\text{H})$  values were set at  $1.5U_{\text{eq}}(\text{C})$  for the methyl groups, and  $1.2U_{\text{eq}}(\text{C})$  for other H atoms.

## Figures




Fig. 1. View of (I) with displacement ellipsoids drawn at the 30% probability level.





Fig. 2. The crystal packing of (I) viewed down the *b*-axis. The hydrogen-bonds are drawn as a dashed lines. H-atoms not involved in hydrogen bonds have been omitted for clarity.

## Ammonium [(1*S*)-(endo,*anti*)]-(*-*)-(3-bromo-1,7- dimethyl-2-oxobicyclo[2.2.1]heptan-7-yl)methanesulfonate

### Crystal data



$M_r = 328.22$

Monoclinic,  $P2_1$

Hall symbol: P 2yb

$a = 7.2449 (2) \text{ \AA}$

$b = 7.0049 (1) \text{ \AA}$

$c = 13.2428 (3) \text{ \AA}$

$\beta = 104.704 (1)^\circ$

$V = 650.06 (3) \text{ \AA}^3$

$Z = 2$

$F(000) = 336$

$D_x = 1.677 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3356 reflections

$\theta = 2.9\text{--}28.3^\circ$

$\mu = 3.33 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Prism, colourless

$0.42 \times 0.14 \times 0.11 \text{ mm}$

### Data collection

Bruker Kappa APEXII CCD  
diffractometer

Radiation source: sealed tube  
graphite

$\varphi$  and  $\omega$  scans

Absorption correction: part of the refinement model

( $\Delta F$ )

(*XABS2*; Parkin *et al.*, 1995; quadratic fit to  $\sin(\theta)/\lambda$  -  
18 parameters)

$T_{\min} = 0.336$ ,  $T_{\max} = 0.711$

2775 measured reflections

2775 independent reflections

2586 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.000$

$\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 3.3^\circ$

$h = -8 \rightarrow 9$

$l = 0 \rightarrow 17$

Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.027$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.065$	$w = 1/[\sigma^2(F_o^2) + (0.033P)^2 + 0.1814P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
2775 reflections	$(\Delta/\sigma)_{\max} < 0.001$
168 parameters	$\Delta\rho_{\max} = 0.35 \text{ e } \text{\AA}^{-3}$
5 restraints	$\Delta\rho_{\min} = -0.47 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1155 Freidel pairs Flack parameter: $-0.021 (7)$

Special details

**Geometry.** Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement on  $F^2$  for ALL reflections except those flagged by the user for potential systematic errors. Weighted  $R$ -factors  $wR$  and all goodnesses of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The observed criterion of  $F^2 > \sigma(F^2)$  is used only for calculating  $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.09450 (4)	0.00017 (4)	0.27520 (3)	0.0440 (1)
S1	0.65799 (9)	0.68509 (9)	0.36429 (5)	0.0242 (2)
O1	0.7216 (3)	0.8793 (3)	0.3541 (2)	0.0434 (8)
O2	0.8199 (3)	0.5560 (3)	0.39714 (16)	0.0332 (6)
O3	0.5283 (3)	0.6697 (4)	0.42972 (17)	0.0448 (8)
O4	0.2268 (4)	0.0517 (4)	0.0568 (2)	0.0596 (10)
C1	0.5311 (4)	0.6201 (4)	0.2339 (2)	0.0277 (8)
C2	0.4754 (4)	0.4086 (4)	0.2154 (2)	0.0213 (7)
C3	0.3597 (3)	0.3263 (4)	0.2899 (2)	0.0223 (7)
C4	0.1802 (4)	0.4500 (4)	0.2647 (2)	0.0272 (8)
C5	0.1351 (4)	0.4758 (5)	0.1462 (2)	0.0352 (9)
C6	0.3067 (4)	0.3804 (4)	0.1148 (2)	0.0310 (9)
C7	0.3318 (6)	0.4411 (6)	0.0112 (2)	0.0519 (13)
C8	0.6514 (4)	0.2949 (4)	0.2067 (2)	0.0316 (9)
C9	0.2719 (4)	0.1690 (5)	0.1248 (2)	0.0342 (9)
C10	0.3097 (4)	0.1283 (4)	0.2418 (2)	0.0298 (8)

## supplementary materials

---

N1	0.8032 (3)	0.1906 (4)	0.4952 (2)	0.0311 (7)
H1A	0.41550	0.69600	0.21450	0.0330*
H1B	0.60920	0.65480	0.18710	0.0330*
H3	0.42880	0.32500	0.36380	0.0270*
H4A	0.20390	0.57210	0.30010	0.0330*
H4B	0.07600	0.38650	0.28480	0.0330*
H5A	0.12590	0.61000	0.12770	0.0420*
H5B	0.01620	0.41320	0.11210	0.0420*
H7A	0.22080	0.40620	-0.04250	0.0780*
H7B	0.44200	0.37920	-0.00150	0.0780*
H7C	0.34880	0.57700	0.01080	0.0780*
H8A	0.74270	0.29220	0.27340	0.0470*
H8B	0.70750	0.35440	0.15630	0.0470*
H8C	0.61420	0.16680	0.18500	0.0470*
H10	0.42240	0.04610	0.26290	0.0360*
H1N	0.775 (5)	0.096 (4)	0.445 (2)	0.0470*
H2N	0.914 (4)	0.170 (6)	0.543 (2)	0.0470*
H3N	0.806 (5)	0.305 (4)	0.461 (3)	0.0470*
H4N	0.706 (4)	0.195 (6)	0.528 (3)	0.0470*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0329 (2)	0.0300 (2)	0.0673 (2)	-0.0085 (1)	0.0093 (1)	0.0087 (2)
S1	0.0199 (3)	0.0209 (3)	0.0303 (3)	-0.0028 (2)	0.0036 (2)	-0.0024 (3)
O1	0.0492 (13)	0.0217 (11)	0.0526 (15)	-0.0096 (9)	0.0007 (11)	-0.0030 (9)
O2	0.0246 (9)	0.0298 (11)	0.0393 (12)	0.0024 (7)	-0.0029 (8)	-0.0036 (8)
O3	0.0304 (10)	0.0674 (17)	0.0391 (13)	-0.0101 (11)	0.0133 (9)	-0.0175 (12)
O4	0.0552 (15)	0.060 (2)	0.0580 (16)	-0.0138 (12)	0.0042 (12)	-0.0336 (13)
C1	0.0241 (13)	0.0248 (14)	0.0298 (15)	-0.0018 (11)	-0.0010 (11)	0.0010 (11)
C2	0.0168 (12)	0.0239 (13)	0.0226 (13)	-0.0016 (10)	0.0041 (10)	-0.0026 (10)
C3	0.0170 (11)	0.0207 (12)	0.0274 (14)	-0.0020 (9)	0.0022 (10)	-0.0004 (10)
C4	0.0184 (11)	0.0235 (15)	0.0403 (16)	0.0017 (9)	0.0084 (10)	-0.0020 (11)
C5	0.0222 (12)	0.0368 (19)	0.0406 (16)	0.0024 (13)	-0.0030 (11)	0.0017 (14)
C6	0.0239 (13)	0.0409 (17)	0.0248 (15)	-0.0040 (12)	-0.0002 (11)	-0.0024 (12)
C7	0.056 (2)	0.070 (3)	0.0247 (17)	-0.0134 (18)	0.0011 (15)	0.0033 (15)
C8	0.0206 (13)	0.0359 (16)	0.0382 (17)	-0.0007 (12)	0.0074 (11)	-0.0104 (13)
C9	0.0198 (12)	0.0392 (17)	0.0405 (16)	-0.0051 (12)	0.0018 (11)	-0.0130 (14)
C10	0.0213 (12)	0.0212 (13)	0.0444 (17)	-0.0006 (10)	0.0039 (11)	-0.0016 (11)
N1	0.0266 (12)	0.0334 (13)	0.0329 (13)	0.0019 (11)	0.0066 (10)	0.0030 (11)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Br1—C10	1.945 (3)	C6—C7	1.491 (4)
S1—O1	1.454 (2)	C6—C9	1.514 (4)
S1—O2	1.457 (2)	C9—C10	1.530 (4)
S1—O3	1.435 (2)	C1—H1B	0.9700
S1—C1	1.797 (3)	C1—H1A	0.9700
O4—C9	1.201 (4)	C3—H3	0.9800

N1—H1N	0.92 (3)	C4—H4B	0.9700
N1—H2N	0.90 (3)	C4—H4A	0.9700
N1—H3N	0.92 (3)	C5—H5A	0.9700
N1—H4N	0.92 (3)	C5—H5B	0.9700
C1—C2	1.539 (4)	C7—H7A	0.9600
C2—C3	1.558 (4)	C7—H7B	0.9600
C2—C6	1.575 (4)	C7—H7C	0.9600
C2—C8	1.532 (4)	C8—H8B	0.9600
C3—C4	1.527 (4)	C8—H8C	0.9600
C3—C10	1.531 (4)	C8—H8A	0.9600
C4—C5	1.530 (4)	C10—H10	0.9800
C5—C6	1.558 (4)		
O1—S1—O2	111.00 (13)	C3—C10—C9	102.4 (2)
O1—S1—O3	113.46 (15)	S1—C1—H1B	108.00
O1—S1—C1	104.17 (14)	S1—C1—H1A	108.00
O2—S1—O3	111.96 (14)	C2—C1—H1A	108.00
O2—S1—C1	107.84 (13)	C2—C1—H1B	108.00
O3—S1—C1	107.92 (14)	H1A—C1—H1B	107.00
H3N—N1—H4N	109 (3)	C10—C3—H3	114.00
H2N—N1—H4N	109 (3)	C4—C3—H3	114.00
H1N—N1—H3N	107 (3)	C2—C3—H3	115.00
H1N—N1—H4N	108 (3)	C5—C4—H4A	111.00
H1N—N1—H2N	113 (3)	C3—C4—H4B	111.00
H2N—N1—H3N	111 (3)	C3—C4—H4A	111.00
S1—C1—C2	116.52 (19)	C5—C4—H4B	111.00
C1—C2—C8	108.8 (2)	H4A—C4—H4B	109.00
C1—C2—C6	111.8 (2)	H5A—C5—H5B	109.00
C1—C2—C3	114.6 (2)	C4—C5—H5A	111.00
C6—C2—C8	110.7 (2)	C4—C5—H5B	111.00
C3—C2—C6	93.6 (2)	C6—C5—H5A	111.00
C3—C2—C8	116.6 (2)	C6—C5—H5B	111.00
C4—C3—C10	108.9 (2)	H7B—C7—H7C	109.00
C2—C3—C4	102.5 (2)	C6—C7—H7C	109.00
C2—C3—C10	100.4 (2)	C6—C7—H7A	110.00
C3—C4—C5	103.9 (2)	C6—C7—H7B	109.00
C4—C5—C6	104.2 (2)	H7A—C7—H7B	109.00
C7—C6—C9	114.9 (3)	H7A—C7—H7C	109.00
C2—C6—C7	119.5 (3)	C2—C8—H8B	109.00
C2—C6—C5	102.8 (2)	C2—C8—H8A	109.00
C5—C6—C9	103.6 (2)	H8A—C8—H8C	109.00
C2—C6—C9	99.2 (2)	C2—C8—H8C	109.00
C5—C6—C7	114.5 (3)	H8A—C8—H8B	109.00
O4—C9—C6	128.6 (3)	H8B—C8—H8C	110.00
O4—C9—C10	125.2 (3)	C9—C10—H10	109.00
C6—C9—C10	106.3 (2)	Br1—C10—H10	109.00
Br1—C10—C3	116.24 (18)	C3—C10—H10	109.00
Br1—C10—C9	111.62 (19)		
O1—S1—C1—C2	169.7 (2)	C2—C3—C4—C5	39.1 (3)



## supplementary materials

O2—S1—C1—C2	51.7 (2)	C10—C3—C4—C5	-66.7 (3)
O3—S1—C1—C2	-69.4 (3)	C2—C3—C10—Br1	-159.47 (17)
S1—C1—C2—C3	54.3 (3)	C2—C3—C10—C9	-37.5 (3)
S1—C1—C2—C6	159.2 (2)	C4—C3—C10—Br1	-52.3 (3)
S1—C1—C2—C8	-78.2 (3)	C4—C3—C10—C9	69.7 (3)
C1—C2—C3—C4	61.2 (3)	C3—C4—C5—C6	-5.6 (3)
C1—C2—C3—C10	173.5 (2)	C4—C5—C6—C2	-29.3 (3)
C6—C2—C3—C4	-54.8 (2)	C4—C5—C6—C7	-160.5 (3)
C6—C2—C3—C10	57.5 (2)	C4—C5—C6—C9	73.6 (3)
C8—C2—C3—C4	-170.0 (2)	C2—C6—C9—O4	-144.1 (3)
C8—C2—C3—C10	-57.8 (3)	C2—C6—C9—C10	34.8 (3)
C1—C2—C6—C5	-67.8 (3)	C5—C6—C9—O4	110.3 (4)
C1—C2—C6—C7	60.4 (4)	C5—C6—C9—C10	-70.9 (3)
C1—C2—C6—C9	-174.0 (2)	C7—C6—C9—O4	-15.4 (5)
C3—C2—C6—C5	50.6 (2)	C7—C6—C9—C10	163.5 (3)
C3—C2—C6—C7	178.7 (3)	O4—C9—C10—Br1	-54.7 (4)
C3—C2—C6—C9	-55.7 (2)	O4—C9—C10—C3	-179.8 (3)
C8—C2—C6—C5	170.7 (2)	C6—C9—C10—Br1	126.4 (2)
C8—C2—C6—C7	-61.1 (4)	C6—C9—C10—C3	1.3 (3)
C8—C2—C6—C9	64.5 (3)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N1—H1N $\cdots$ O1 <sup>i</sup>	0.92 (3)	1.92 (3)	2.835 (4)	173 (3)
N1—H2N $\cdots$ O2 <sup>ii</sup>	0.90 (3)	2.05 (3)	2.899 (3)	157 (3)
N1—H3N $\cdots$ O2	0.92 (3)	1.97 (3)	2.887 (3)	176 (3)
N1—H4N $\cdots$ O3 <sup>iii</sup>	0.92 (3)	1.93 (3)	2.827 (3)	167 (4)
C4—H4B $\cdots$ Br1	0.97	2.71	3.221 (3)	113
C8—H8A $\cdots$ O2	0.96	2.44	3.104 (3)	126
C10—H10 $\cdots$ O1 <sup>i</sup>	0.98	2.49	3.451 (4)	167

Symmetry codes: (i)  $x, y-1, z$ ; (ii)  $-x+2, y-1/2, -z+1$ ; (iii)  $-x+1, y-1/2, -z+1$ .

Fig. 1




Fig. 2

