



ELSEVIER

Contents lists available at ScienceDirect

SSM -Population Health

journal homepage: www.elsevier.com/locate/ssmph

Review Article

Spirituality, religiosity, aging and health in global perspective: A review

Zachary Zimmer^{a,b,*}, Carol Jagger^c, Chi-Tsun Chiu^d, Mary Beth Ofstedal^e, Florencia Rojo^a, Yasuhiko Saito^f^a University of California, San Francisco, USA^b Mount Saint Vincent University, Canada^c Newcastle University, UK^d Academia Sinica, Taiwan^e University of Michigan, USA^f Nihon University, Japan

ARTICLE INFO

Article history:

Received 15 January 2016

Received in revised form

16 April 2016

Accepted 18 April 2016

Keywords:

Aging

Global aging

Health expectancy

Older adults

Mindfulness

Mortality

Religion

Spirituality

ABSTRACT

Persistent population aging worldwide is focusing attention on modifiable factors that can improve later life health. There is evidence that religiosity and spirituality are among such factors. Older people tend to have high rates of involvement in religious and/or spiritual endeavors and it is possible that population aging will be associated with increasing prevalence of religious and spiritual activity worldwide. Despite increasing research on religiosity, spirituality and health among older persons, population aging worldwide suggests the need for a globally integrated approach. As a step toward this, we review a subset of the literature on the impact of religiosity and spirituality on health in later life. We find that much of this has looked at the relationship between religiosity/spirituality and longevity as well as physical and mental health. Mechanisms include social support, health behaviors, stress and psychosocial factors. We identify a number of gaps in current knowledge. Many previous studies have taken place in the U.S. and Europe. Much data is cross-sectional, limiting ability to make causal inference. Religiosity and spirituality can be difficult to define and distinguish and the two concepts are often considered together, though on balance religiosity has received more attention than spirituality. The latter may however be equally important. Although there is evidence that religiosity is associated with longer life and better physical and mental health, these outcomes have been investigated separately rather than together such as in measures of health expectancy. In conclusion, there is a need for a unified and nuanced approach to understanding how religiosity and spirituality impact on health and longevity within a context of global aging, in particular whether they result in longer healthy life rather than just longer life.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

Introduction	374
Religiosity and spirituality in the context of global aging and increasing longevity	374
Defining religiosity and spirituality for the purpose of global research	375
Connecting religiosity and spirituality to health of older persons	375
Potential mechanisms	376
Social support	377
Behaviors	377
Stress	377
Psychosocial responses	377
Future directions	377
Longitudinal data	377
Cross-national studies	378

* Correspondence to: Department of Social and Behavioral Sciences, University of California, 3333 California Street, Suite 455, San Francisco, CA 94118, USA.
E-mail address: zacharyzimmer@ucsf.edu (Z. Zimmer).

Health expectancy as opposed to life expectancy	378
Conclusion	378
Acknowledgments	378
References	379

Introduction

There has been growth in the volume of scientific studies that explore the connection between religion, spirituality and health. Earlier reviews have established that while associations are not universal, on balance, studies indicate salutary benefits arising from religious and spiritual involvement across a number of health outcomes (Hummer, Ellison, Rogers, Moulton, & Romero, 2004; Koenig, 2009; Koenig, King, & Carson, 2012; Marks, 2005; Seeman, Dubin, & Seeman, 2003). The current article is an essay that examines a selection of literature that derives from several notions: religiosity and spirituality is pervasive globally; health benefits that may stem from religious and spiritual involvement could be important for the future of global population health, and; this is particularly the case given realities of global population aging. The dialog to follow therefore identifies religiosity and spirituality as key components of health within the context of global aging and expanding life expectancy. The essay covers several topics. It begins by characterizing population aging and increasing longevity. It then defines religiosity and spirituality within a global context. Evidence is provided that links religiosity and spirituality to health of older persons. Potential mechanisms of these links are then discussed. This leads into a dialog on some potential future directions, followed by a concluding statement.

Religiosity and spirituality in the context of global aging and increasing longevity

Population aging is a term that refers to increasing numbers and proportion of older persons within populations. Population

Table 1
Percent responding in the affirmative to specific questions on the World Values Survey, 2010–2014, in the thirteen most highly populated countries covered by the 2015 release wave, by age.

	All ages		Under 60		60 and older	
	Religious ^a	Think ^b	Religious ^a	Think ^b	Religious ^a	Think ^b
China	12.9	50.2	12.3	52.5	16.6 [*]	37.5 [*]
India	78.3	66.4	78.6	66.8	75.8	61.7
USA	67.9	77.8	64.7	78.7	77.6 [*]	75.1
Brazil	81.2	79.9	79.7	81.1	87.8 [*]	74.3 [*]
Nigeria	95.8	92.4	96.0	92.4	94.3	94.3
Pakistan	99.8	82.7	99.7	82.7	100	82.7
Russia	61.2	71.1	60.8	70.6	62.5	73.1
Japan	25.4	79.2	22.5	80.1	30.1 [*]	77.8
Mexico	74.7	80.8	73.6	81.8	84.1 [*]	72.0 [*]
Philippines	80.8	92.0	80.1	92.8	83.9	87.9 [*]
Germany	50.9	69.9	44.9	66.7	64.1 [*]	76.9 [*]
Turkey	85.0	88.5	83.9	88.8	93.3 [*]	86.9
Thailand	33.0	81.4	30.9	81.2	48.6 [*]	76.6
Mean across countries	65.1	76.8	63.7	78.2	70.7 [*]	75.1
Standard deviation	27.2	11.5	27.9	11.4	25.5	14.0

^{*} Difference between under 60 and 60+ significant at $p < 0.05$.

^a The full question is: 'Do you consider yourself to be a religious person?'

^b The full question is: 'Do you think about/contemplate upon (sometimes or often) the meaning or purpose of life?'

aging is being experienced throughout the world (UNFPA & HelpAge International, 2012). According to United Nations estimates, the population aged 60 and older in 2015 was about 900 million, representing about 12% of the global population (United Nations, 2015). Given medium level growth projections, this number is expected to pass 2 billion by 2050, which at that time will represent close to 22% of the global population. Related to this is increasing longevity with people in many parts of the world living to much older ages than has ever been the case (Vaupel & Kistowski, 2005). With few exceptions, these changes are happening everywhere, in every global region (Zimmer & McDaniel, 2013). The impacts of population aging, including rising health care costs and formal and informal health care needs, are being shared across societies thousands of miles apart, rich and poor, with different cultures, languages, structures of government, family values, and economic systems. Many of the consequences of population aging are common across human societies in all corners of the world.

Given the ubiquity of population aging and increasing longevity, common ways in which good health in old age can be promoted is now of paramount concern to health professionals, researchers and policy makers worldwide (World Health Organization, 2015). While population aging is a recent phenomenon, religiosity, which evidence suggests may be a common factor associated with health, is a long-standing one that similarly extends to all corners of the world (George, Larson, Koenig, & McCullough, 2000; O'Brien, Palmer, & Barrett, 2007). When expanded to include meditative and contemplative activity, the desire to seek a meaning to life or the desire for a transcendent connection, it is clear that expressions of religiosity and spirituality is widespread across individuals living in human societies, across regions with differing ideologies, orientations and practices.

To illustrate we provide in Table 1 results from the 2015 release of the World Values Survey (WVS) (data collected between 2010 and 2014) for major countries with different religious traditions and degrees of secularization (World Values Survey Association, 2014). The countries we show are chosen because they are the most populated ones covered by the WVS. Together they represent 60% of the global population. Shown is the percent within each country that respond to questionnaire items by saying they consider themselves to be a religious person (labeled 'Religious') and the percent saying that they sometimes or often think about or contemplate upon the meaning of life (labeled 'Think'). The latter question is not a direct measure of spirituality, but, given common definitions of spirituality (Koenig, 2012) it does suggest reflection upon abstract and intangible things that relate to a purpose of life and therefore reflects a degree of spiritual-like thinking. Moreover, earlier versions of the WVS included additional questions about meditation and prayer, and the item on thinking and contemplating meaning of life correlates highly with these other measures that can also be linked to spiritualness. The first two columns compare these items for the total sample 18 and older. The next four columns show the percentages for those under age 60 and for those age 60 and older. Statistically significant differences across age groups are indicated by an asterisk.

The percent that say they consider themselves to be religious varies widely with lows of 12.9% in China, 25.4% in Japan and 33.0% in Thailand, and highs of 99.8% in Pakistan, 95.8% in Nigeria and

85.0% in Turkey. But, when it comes to the percent that frequently think about the meaning of life, there is less variation and the percentages in the more secular countries are quite high. In China, Japan, and Thailand, where a clear minority of persons report themselves as being religious, 50.2%, 79.2% and 81.4% respectively report thinking about the meaning or purpose of life. The average across these thirteen countries is higher and the standard deviation is lower for the item about meaning of life than the item about religiosity. While the different ways in which people respond to these two questions across countries is clearly a function of normative beliefs and practices, the point we make is that some form of spiritual-like thinking as it is ordinarily defined is quite common among people in countries around the world characterized by different dominant religious traditions and different degrees of secularism.

There are age differences in Table 1. Except for two countries, older persons are more likely to report they are religious, and in most cases, the difference is statistically significant. Differences tend to be non-significant within countries that report very high rates of being religious, such as in Nigeria and Pakistan. When it comes to the percent reporting that they think about the meaning of life, differences across age groups are less noticeable. China is an exception. In the USA, the percent that say they think about the meaning of life sometimes or often is 78.7% for those under 60 and 75.1% for those 60 and older, a difference that is not statistically significant. Differences in thinking about meaning of life also run in both directions. In Mexico younger persons are significantly more likely to think about the meaning of life, whereas in Germany it is older persons that are more likely.

It is because population aging and religiosity are both pervasive global phenomenon that the potential salutary effect of religiosity and spirituality is so important for global health. Evidence suggests that older persons tend to be more religious than younger ones; an association that has proven robust in both cross-sectional and longitudinal data (Argue, Johnson, & White, 1999; Levin, Taylor, & Chatters, 1994). Some of this could be a function of cohort differences, with the current older generation coming from a background and a time where religion was valued to a greater degree and thus they carry those values into old age (Wilhelm, Rooney, & Tempel, 2007). However, there is other evidence to suggest that people become more involved with religion and their sense of spirituality magnifies with age (Moberg, 2005; Wink & Dillon, 2001). Therefore, as the world grows older, it is quite possible that it will grow more religious and spiritual. It is not hyperbole to say that population aging coupled with increasing longevity will be among the most important trends driving population health and health care costs around the world over the next several decades (Mayhew, 1999; Zimmer & McDaniel, 2013). Therefore, how religiosity and spirituality associate to health among older persons will be critical for determining the health of these aging older populations and ultimately for health policies adopted by the societies in which older persons live.

Defining religiosity and spirituality for the purpose of global research

Distinguishing the difference between religiosity and spirituality has not been easy for any single culture let alone cross-nationally, and centuries of thinking have yet to provide standard delineations of the two concepts (Fetzer Institute, 2003; Hill et al., 2000; Idler et al., 2003; Koenig et al., 2012; MacKinlay, 2002; Ortiz & Langer, 2002). One difficulty is that the two concepts are related and have common characteristics (Seybold & Hill, 2001); both can involve personal transformation and the search for an ultimate truth. A religious person often defines themselves as being

spiritual, so that spirituality often encompasses religiosity, but is often defined in broader terms (Holmes, 2002). Religion around the world generally associates with specific foundational principles that are organized around distinct systems of beliefs, practices and rituals that take place within communities of participants. Spirituality is more difficult to define, particularly across cultures, as its characteristics are not easily agreed upon, and it can mean different things for different individuals in different places (Koenig et al., 2012). More so, spirituality conveys the notion of a personal search relating to things sacred and transcendent (Crowther, Parker, Achenbaum, Larimore, & Koenig, 2002). While levels of religiosity around the world are often measured in direct ways, for instance with respect to frequency of activity, such as ritualistic practice and attendance at places of worship, spirituality must be measured in more indirect personal experiential terms, such as the search for meaning, peace and personal fulfillment, contemplation about meaning of life, and the feeling of a personal relationship to a higher power (Zinnbauer et al., 1997). Religious activity is normally thought to take place within or in connection to formal institutions while spiritual activity, taking on a more personal tonality, can take place within and equally outside of formal institutions. This does not mean to say that religion is necessarily a public endeavor. Eastern religions, such as Shintoism as practiced in Japan and Buddhism as practiced across Southeast and South Asia, involve a greater degree of private worship than do the religions more commonly practiced in the United States and Europe (Hidajat, Zimmer, Saito, & Lin, 2013).

In many ways, religiosity is easier to quantify than spirituality. Because of the abstraction involved with the concept, survey research often includes indirect indicators of spirituality, such as the item referred to in Table 1. This type of measure demonstrates the broadness of spirituality. In some settings, many more people say that they reflect upon the meaning of life, meditate or participate in ritualistic activities rather than say they are religious. Most research around the world to date has focused on the link between religion and health as opposed to spirituality and health (Lucchetti, Lucchetti, & Koenig, 2011). Yet, spirituality and religiosity are often considered together, particularly when incorporating measures relating to meaning of life (Krause, 2003).

Another ambiguity in distinguishing religiosity from spirituality is that both can involve contemplative activity. These will differ across contexts around the world. Religious activities can involve prayer, chanting, singing and listening to chants, songs and sermons. More recently, research on spirituality and health has been extended to consider the impacts of mindfulness and non-denominational meditation (Davidson et al., 2003). While prayer may be considered a conversation, often internal, with a divine power or a higher self, meditation and mindfulness may be thought of as an attempt at deep concentration with a particular focus, such as on the breath, or an attempt at being acutely aware of one's feelings, thoughts and experiences (Shapiro & Carlson, 2009). Again, like religion and spirituality, there are close connections between prayer and mindfulness meditation and both are thought to elicit physical and mental relaxation (Ivanovski & Malhi, 2007).

Connecting religiosity and spirituality to health of older persons

A connections between religiosity or spirituality and health has been recognized for decades if not centuries (Dearmer, 1909; Hiltner, 1943; Koenig, 2012; Koenig et al., 2012; Lavretsky, 2010). Scientific dialog on the topic accelerated in the 1980s and 1990s with influential studies that focused on mental health (Bergin, 1983; Koenig, 1989, 1998). Religion has specifically been found to

exert positive impacts on disability and depressive outcomes of older persons (Idler & Kasl, 1997a; Lavretsky, 2010). Recent years have witnessed proliferation of research on mechanisms that link religiosity to health outcomes (Ellison & Levin, 1998; Hummer et al., 2004; Krause, 2008, 2009, 2011; Krause & Hayward, 2014; Larson, Swyers, & McCullough, 1998).

The impact of religion and spirituality on the health of older persons specifically has been a growing concern and there are a number of empirical and theoretical reviews on the topic (Krause, 1997, 2004; Levin & Chatters, 2008). For older populations, an increasing focus on how the association is manifest has been driven by not only concern for elderly health and related policy implications but also, as referenced above, increasing human longevity, the growth in the size of the older population globally, and the relationship between aging and spirituality (Krause, 2006). Demographic research has for one been critically involved in linking religiosity and longer life (Clark, Friedman, & Martin, 1999; Dupre, Franzese, & Parrado, 2006; Hummer, Benjamins, Ellison, & Rogers, 2010; Hummer, Rogers, Nam, & Ellison, 1999; Koenig et al., 1999; McCullough, Hoyt, Larson, Koenig, & Thoresen, 2000; Musick, House, & Williams, 2004; Strawbridge, Cohen, Shema, & Kaplan, 1997; Sullivan, 2010; Zhang, 2008). Hummer et al. (1999) provided evidence, using the U.S. National Health Interview Survey, that those that do not attend religious services have close to twice the risk of dying as opposed to those that do. Unfortunately, demographic research has almost exclusively focused on frequency of service attendance as an indicator of religiosity and, with few exceptions (noted later in this review), considered only U.S. data (Hummer et al., 2004).

Statistical correlation between religiosity, spirituality and health has been found with other health outcomes (Larson et al., 1998; Levin & Vanderpool, 1991). These include cardiovascular conditions common among older persons such as heart disease, blood pressure, cholesterol, myocardial infarction and stroke (Armstrong, Van Merwyk, & Coates, 1977; Brown, 2000; Sørensen, Danbolt, Lien, Koenig, & Holmen, 2011; Walsh, 1998). Benefits of religiosity and spirituality pre and post cardiac event have been observed (Berntson, Norman, Hawkey, & Cacioppo, 2008; Oxman, Freeman, & Manheimer, 1995). Salutary effects have been seen with respect to disability and functional limitation, kidney function, cirrhosis, emphysema, chronic pain, cancer and self-rated overall health (Gillum, 2006; Hank & Schaan, 2008; Hidajat et al., 2013; Idler & Kasl, 1997a, 1997b; Koenig et al., 2012; Kune, Kune, & Watson, 1993; Larson et al., 1998; Roff et al., 2006; Thege, Pilling, Székely, & Kopp, 2013). Religious attendance has been shown to buffer the need for and length of hospitalization (Koenig et al., 1992, 1998).

There is also strong evidence of a connection between religiosity and mental health (Musick, Traphagan, Koenig, & Larson, 2000). The mental health side of the literature has been more likely to recognize broader spirituality measures as predictors (Richards & Bergin, 2005). Religiosity and spirituality has been shown to relate to depressive and anxiety outcomes, particularly among older persons (Bergin, 1983; Braam, Beekman, & van Tilburg, 1999; Koenig, 1989, 2013, 1998; Lucchetti et al., 2011; Mueller, Plevak, & Rummans, 2001; Seybold & Hill, 2001). Some of the literature has indicated that religiosity can moderate the association between physical health conditions and depression through coping mechanisms, and can increase recovery time from a number of health disorders (Koenig et al., 1992). Cognition has been shown to benefit from religiosity (Corsentino, Collins, Sachs-Ericsson, & Blazer, 2009).

While many studies combine concepts of spirituality and religiosity, spirituality can also represent a separate component of the literature. As noted, spiritual attitudes are assessed using different types of indicators and may be more challenging to measure

(George et al., 2000; Miller & Thoresen, 2003; Schlehofer, Omoto, & Adelman, 2008). Atchley (2009) has devoted a book to the topic of spirituality and aging separate from religiosity, much of which illustrates connections between spirituality and well-being. Like religiosity, those that have focused on spirituality have generally found strong associations with both mental and physical health (Ballew, Hannum, Gaines, Marx, & Parrish, 2012; Crowther et al., 2002; Liu, Schieman, & Jang, 2011; Rippentrop, Altmaier, Chen, Found, & Keffala, 2005; Saad, Hatta, & Mohamad, 2010; Tan, Wutthilert, & O'Connor, 2011). Much of this literature concerns older persons (Kirby, Coleman, & Daley, 2004; Lavretsky, 2010; Reeder, 1991).

Expanded to include meditative and contemplative activities, there is the possibility of additional associations that have been underexplored (Jantos & Kiat, 2007). Prayer can impact stress and coping, and it has been shown to be related to happiness and feelings of general well-being (Boehnlein, 2008; Poloma & Pendleton, 1989, 1991). A mindfulness practice is increasingly being seen as effective in controlling stress and anxiety (Koszycki, Bengler, Shlik, & Bradwejn, 2007; Ledesma & Kumano, 2009; Roth & Creaser, 1997). Older persons benefit from such activity (Ernst et al., 2008). Studies have indicated mindfulness meditation positively impacts a host of physical health outcomes (Cahn & Polich, 2006; Davidson et al., 2003; Fang et al., 2010; Jevning, Wallace, & Beidebach, 1992; Solberg et al., 2000; Young & Taylor, 1998). However, very little research has examined the impact of prayer and meditation on demographic outcomes like life or healthy life expectancy.

While the preponderance of evidence thus far has linked religiosity and health in a positive way, effects are not universally beneficial. Certain aspects and measures of religiosity have been shown to be disadvantages to health. For instance, religion has been found to induce anxiety when it fosters psychologically harmful feelings such as guilt and shame, when it promotes adverse attitudes towards outsiders or when it encourages unquestioning devotion and obedience or beliefs that step toward 'fundamentalism' (Chatters, 2000; Nooney & Woodrum, 2002). It is harmful to rely on faith healing as a substitute for medical care (Pargament et al., 2001). Unfavorable interactions with fellow church members were identified in one study as leading to depressive outcomes among clergy and older church-goers (Krause, Ellison, & Wulff, 1998). Religious doubt, or a feeling of uncertainty towards religious beliefs has been associated with negative mental health outcomes (Krause & Wulff, 2004). There may also be causation and sampling issues in studies of religiosity and health that warrant attention. Depressed individuals may drop out of religious activities making samples biased (Maselko, Hayward, Hanlon, Buka, & Meador, 2012). In this case, mental health affects religious activity rather than the other way around. For older populations with high prevalence of disability, attending religious services requires a certain level of physical ability. This might suggest that attendance, which is the single most frequently used measure of religiosity and the one most often linked to health outcomes, could bias associations more so than other measures, like private practice, which may be the activity of choice for chronically ill and those near death (Yeager et al., 2006).

Potential mechanisms

Much of the literature indicating a favorable connection between religiosity, spirituality and health is descriptive in nature, meaning that the research empirically demonstrates a relationship, but the mechanisms driving the relationship are unexplained. This is particularly the case for research that connects religious attendance and mortality (Hummer et al., 2004).

Research that does examine mechanisms suggest they are complex and multi-faceted. Yet, recent research is uncovering several that appear to be consistent across cultures and countries: social support; behavior; stress reduction; and psychosocial indicators.

Social support

Religious institutions worldwide, such as churches, temples and mosques are often a focal point for interaction, exchange and support. They can play a role in integrating families and providing encouragement for an active family life. A place of worship is often a location where social interaction happens, friends meet, families gather, and supportive activities take place. A place of worship is often a focal point for volunteer and community building work (Wilson & Musick, 1997). Neal Krause has explored the concept of spiritual support, by which he refers to social support that is geared towards enhancing the spiritual and religious experience (Krause, 2002). Spiritual support can involve tangible exchanges such as help and advice in applying spiritual principles and emotional assistance in spiritual pursuits. In addition, in many societies around the world and across religious denominations, places of worship play a central role in organizing the community, providing a place to gather to celebrate events or mourn losses. In the latter case, the place of worship can play a supportive role particularly important for older persons who, in some countries, will be frequently living alone, could be lonely and may be experiencing frequent losses.

Behaviors

Religious institutions have long been viewed as organizations that promote healthy behaviors (Hill, Ellison, Burdette, & Musick, 2007; Powell, Shahabi, & Thoresen, 2003; Strawbridge, Shema, Cohen, & Kaplan, 2001; Takyi, 2003; Thege et al., 2013). Individuals who are religious are often committed to healthy lifestyles, and many religious orientations around the world possess negative views about tobacco, alcohol consumption, drugs and non-marital risky sex. Recent evidence suggests spirituality is associated with the popularity of activities such as meditation and yoga for the maintenance of health (Ross & Thomas, 2010). Related, some have referenced the ability of religion to imbue a sense of self-regulation and self-control (McCullough & Willoughby, 2009).

Stress

Perhaps the largest body of literature on mechanisms relates to the impact of religiosity and spirituality on stress and coping with stressful life events. To some, a chief function of religion is to assist in the face of adversity (Pargament, 1997; Pargament & Raiya, 2007). This is important for older persons who often cope with losses of loved ones, loss of mental and physical function and the propinquity of their own mortality (McFadden, 1995; Rogers, 1976). By prompting a meaning to life, religion has been shown to bolster life satisfaction, optimism and self-esteem, all of which act to reduce the impact of stressful life events (Krause, 2003; Krause, Ellison, Shaw, Marcum, & Boardman, 2001; Pargament, Koenig, & Perez, 2000; Ryff & Singer, 1998). There is also a link between the social support that can be provided through religious activity and the reduction of stress (Nooney & Woodrum, 2002).

Psychosocial responses

Some of the most profound advances in mechanisms have come by way of the testing of psychosocial mediators. Forgiveness for instance has been examined as a health determinant both within and external to religion (Enright & North, 1998; Thoresen,

Harris, & Luskin, 2000). By promoting forgiveness religion has been shown to reduce feelings of hostility and anger, which is beneficial for health (Bono & McCullough, 2004). According to some, one of the primary aims of faith is to promote virtues such as humility, compassion, gratitude, wisdom and altruism (Lundberg, 2010). Across a series of studies, Krause has attempted to establish structural links between religiosity and health through these types of psychosocial mechanisms. In a recent publication, Krause and Hayward (2014) show that people who go to church more frequently are more likely to receive a type of support that relates to a sense of belonging that promotes self-ratings of health. In an earlier study (Krause, 2011), church attendance is linked to what is called human needs, such as self-transcendence, control and coping, social participation and development of intimate relationships.

These psychosocial concepts may be difficult to define and measure and the links between them are intricate and at times difficult to observe empirically. Each potential mechanism may be considered as providing a piece of the theoretical puzzle. Krause (2011) asserts that past research represents a disheveled or fractured literature, in need of grander perspective to connect the multidimensional parts. The literature is replete with potential mediators and their connection is often difficult to pinpoint. It is also important to note that across the many studies we reviewed and that adjust and control for multiple intervening mechanisms, there is still a residual impact of religiosity on health. In other words, the impact of religiosity and spirituality on health has yet to be fully explained, especially across global contexts and traditions.

Future directions

Population aging has considerable consequences for global health. Given the potential for religiosity and spirituality to impact on the health of older persons, there is clearly an advantage to public health that can be gained by better understanding the connection between spirituality, religiosity and health in old age across denominational and ritualistic backgrounds and ways of thinking about religious practice. As Hummer et al. (2010) note, future work is needed to determine the strength of the association, the mechanisms through which religiosity and spirituality impact on health across different societies, and the subpopulations for whom religious involvement may or may not be influential. While there has been a burgeoning of scholarly research on religion, spirituality and health, there remain challenges to address, including some that are substantive and some that are methodological (Moberg, 2005; Sloan, Bagiella, & Powell, 1999). Below we identify three contemporary challenges facing the scientific community interested in this topic: longitudinal data; cross-national studies; using health expectancy as an outcome.

Longitudinal data

Studies that link religiosity and mortality provide some of the most robust evidence of positive influences. By definition, such research incorporates longitudinal data. Outside of this, the literature is still weighted toward cross-sectional data, and many of the mortality studies involve a single observation of religiosity or spirituality followed by monitoring of mortality without interim observations. Moreover, connections between religion or spirituality and health transitions, meaning changes in health states while alive, have been mostly absent. This limits ability to make causal connections. Further, while religiosity and spirituality can impact upon health, health, especially among older persons, can make religious and spiritual involvement more or less accessible

or desired. As noted above, in a practical sense, disability typical in older ages can limit one's ability to transport from home to a place of worship. In contrast, life threatening disease may intensify the desire to find a meaning to life and increase the appeal of religious or spiritual involvement. Further, while research has connected many dimensions of early life exposure to later life health and mortality (Elo & Preston, 1992; Galobardes, Smith, & Lynch, 2006; Smith, Mineau, Garibotti, & Kerber, 2009; Temby & Smith, 2013), early life exposure to religion and spiritual thinking on late life health is missing from this dialog.

Cross-national studies

The field of religiosity, spirituality, global aging and health at present relies very much on data from the United States and other developed countries (Hank & Schaan, 2008; Krause & Booth, 2004; Lalive d'Épinay & Spini, 2004). A consideration of the developing world, where most of global population aging is taking place, where most of the world's old people live, and where health care costs are likely to increase most dramatically as a result, exist but has been more limited (e.g., Benjamins, 2007; Chokkanathan, 2013; Hidajat et al., 2013; Kodzi, Gyimah, Emina, & Ezeh, 2011; Nelson, 2000; Reyes-Ortiz, Pelaez, Koenig, & Mulligan, 2007; Zeng, Gu, & George, 2011; Zhang, 2008). There are many benefits of examining associations in non-western settings. Religious orientations differ around the world and, as such, not much is known about whether and to what extent the association persists across boundaries characterized by different ways of expressing religious conviction. Moreover, how different individuals practice religiosity varies widely across countries (Koenig, 2001). When contemplative and meditative pursuits are included, the notion of practice can become murkier. It can be difficult to tease out the impacts of religion versus spirituality versus meditation, and effects of public versus private expressions (Helm, Hays, Flint, Koenig, & Blazer, 2000). As mentioned earlier, Buddhism is organized around individualistic reflective principles much more so than western orientations (Batchelor, 1987; Lutz, Slagter, Dunne, & Davidson, 2008). Buddhists do engage in collective rituals, such as chanting, and visiting temples with family members during specific festivals. But, much more so than in western traditions, visitors to the temple engage in activities that promote meditative absorption, which in Buddhism is viewed as an attempt to cultivate a state of mindfulness. Referring back to Table 1, this could explain why such a minority of Thais, who tend to be practicing Buddhists, report that they are religious but a majority report regularly thinking about the meaning or purpose of life.

Health expectancy as opposed to life expectancy

An underlying goal within medical and other health sciences is to help people live long lives and the increasing longevity being experienced globally is a key indicator of improvement in the human condition. Thus, longevity itself can be a proxy for health. As such, as the global population increasingly lives to older ages it is often assumed that individuals are living both longer and healthier lives. However, the notion that longer life necessarily translates into a healthier old-age is often untested (Crimmins, Hayward, & Saito, 1994; Jagger, 2006). The issue has indeed led to some spirited debate amongst scholars (Fries, 2003; Jagger, 2000; Nusselder, 2003). Recent evidence now indicates that there are at least parts of the world where longer life has not brought about a healthier old-age (World Health Organization, 2015).

In this respect, an important concept is that of health expectancy, or amount of time that individuals within a population can expect to live in relative health. Health expectancy has become a widely used tool for comparing health across populations (Chiu,

2013; Jagger et al., 2009; Robine, Jagger, & Romieu, 2001; Saito, Qiao, & Jitapunkul, 2003). It allows for the quantification of differences in health in terms that are not only easily understood, with reference to number of years of life and percent of life to be lived in healthy states, but also can be translated into future health care needs. Although researchers have recognized the importance of religiosity and spirituality as a determinant of longevity, there is now a need to determine whether religiosity and spirituality associate with extra years of life being lived in good or poor health. That is, whether religiosity and spirituality contribute globally to healthy life expectancy as well as total life expectancy.

Conclusion

Several authors have noted that research on religiosity or spirituality and health can be controversial and critics of religiosity as a determinant of health can be outspoken (Hummer, 2005; Lavretsky, 2010). It can be considered by some to be 'unscientific' to consider religiosity among other health determinants, despite evidence cited above. Yet, there does appear to be an intensification of spiritual affectations with aging. At the same time, the global population is getting older and living to increasingly advanced ages. Now for the first time in human history, most people in the world can expect to live into their 60s (World Health Organization, 2015). These facts strongly point toward a requirement and even an obligation on the part of the scientific community to explore the connection between religiosity, spirituality and health in order to more fully understand the determinants of quality of life in old age and in so doing suggest ways for improving human health and the human condition.

The current essay has its limitations. More systematic reviews on religion and health exist elsewhere (Hummer et al., 2004; Koenig, 2009; Marks, 2005; McCullough et al., 2000; Seeman et al., 2003). Rather than doing a review that would involve empirical assessments of a set of literature, we have sought to focus on a particular subset that facilitates a discussion on religiosity and spirituality within the context of global aging. Therefore, we have chosen to examine literature that not only speaks to issues of religiosity and spirituality among older persons in different regions of the world, but also touches on the global aging taking place and the implications of this for overall health. While the lack of a systematic strategy herein results in selection bias, we believe that we have provided an honest and balanced look at the literature that links religion, spirituality and health among older persons.

The field of religiosity, spirituality and health is still growing. In the current essay we have suggested that advancements are needed in order to fully appreciate the impact of religiosity and spirituality on health within the context of global aging. These include examining related concepts of meditation and mindfulness, using longitudinal data more frequently to strengthen causal connections, conducting cross-national studies since religious and spiritual orientations, values, beliefs and practices differ worldwide, and examining a broader slate of health outcomes that are relevant for aging populations, which include contrasting changes in total life years versus years in states of healthfulness.

Acknowledgments

This work is funded by the John Templeton Foundation Grant number 57521 "Linking spirituality and religiosity to life and health expectancy: A global comparative study." The authors wish to thank four anonymous reviewers for their helpful comments on an earlier version.

References

- Argue, A., Johnson, D. R., & White, L. K. (1999). Age and religiosity: Evidence from a three-wave panel analysis. *Journal for the Scientific Study of Religion*, 38, 423–435.
- Armstrong, B., Van Merwyk, A. J., & Coates, H. (1977). Blood pressure in Seventh-day adventist vegetarians. *American Journal of Epidemiology*, 105, 444–449.
- Atchley, R. C. (2009). *Spirituality and aging*. Baltimore: Johns Hopkins University Press.
- Ballew, S. H., Hannum, S. M., Gaines, J. M., Marx, K. A., & Parrish, J. M. (2012). The role of spiritual experiences and activities in the relationship between chronic illness and psychological well-being. *Journal of Religion and Health*, 51, 1386–1396.
- Batchelor, S. (1987). *The jewel in the lotus: A guide to the buddhist traditions of Tibet*. London: Wisdom Publications.
- Benjamins, M. R. (2007). Predictors of preventive health care use among middle-aged and older adults in Mexico: The role of religion. *Journal of Cross-Cultural Gerontology*, 22, 221–234.
- Bergin, A. E. (1983). Religiosity and mental health: A critical reevaluation and meta-analysis. *Professional Psychology: Research and Practice*, 14, 170–184.
- Berntson, G. G., Norman, G. J., Hawley, L. C., & Cacioppo, J. T. (2008). Spirituality and autonomic cardiac control. *Annals of Behavioral Medicine*, 35, 198–208.
- Psychiatry and religion: The convergence of mind and spirit In: J. K. Boehlein (Ed.), Washington, DC: American Psychiatric Pub.
- Bono, G., & McCullough, M. E. (2004). Religion, forgiveness, and adjustment in older adulthood In: K. W. Schaie, N. Krause, & A. Booth (Eds.), *Religious influences on health and well-being in the elderly* (pp. 163–186). New York: Springer.
- Braam, A. W., Beekman, A. T., & van Tilburg, W. (1999). Religion and depression in later life. *Current Opinion in Psychiatry*, 12, 471–475.
- Brown, C. M. (2000). Exploring the role of religiosity in hypertension management among African Americans. *Journal of Health Care for the Poor and Underserved*, 11, 19–32.
- Cahn, B. R., & Polich, J. (2006). Meditation states and traits: EEG, ERP, and neuroimaging studies. *Psychological Bulletin*, 132, 180–211.
- Chatters, L. M. (2000). Religion and health: Public health research and practice. *Annual Review of Public Health*, 21, 335–367.
- Chiu, C. T. (2013). *A cross-national comparison of health expectancy: Japan, the United States and Taiwan* (Ph.D. dissertation). Austin TX: University of Texas.
- Chokkanathan, S. (2013). Religiosity and well-being of older adults in Chennai, India. *Aging Mental Health*, 17, 880–887.
- Clark, K. M., Friedman, H. S., & Martin, L. R. (1999). A longitudinal study of religiosity and mortality risk. *Journal of Health Psychology*, 4, 381–391.
- Corsentino, E. A., Collins, N., Sachs-Ericsson, N., & Blazer, D. G. (2009). Religious attendance reduces cognitive decline among older women with high levels of depressive symptoms. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 64, 1283–1289.
- Crimmins, E. M., Hayward, M. D., & Saito, Y. (1994). Changing mortality and morbidity rates and health status and life expectancy of the older population. *Demography*, 31, 159–175.
- Crowther, M. R., Parker, M. W., Achenbaum, W. A., Larimore, W. L., & Koenig, H. G. (2002). Rowe and Kahn's model of successful aging revisited positive spirituality – The forgotten factor. *The Gerontologist*, 42, 613–620.
- Davidson, R. J., Kabat-Zinn, J., Schumacher, J., Rosenkranz, M., Muller, D., Santorelli, S. F., et al. (2003). Alterations in brain and immune function produced by mindfulness meditation. *Psychosomatic Medicine*, 65, 564–570.
- Dearmer, P. (1909). *Body and soul: An enquiry into the effect of religion upon health, with a description of Christian works of healing from the New Testament to the present day*. New York: EP Dutton & Company.
- Dupre, M. E., Franzese, A. T., & Parrado, E. A. (2006). Religious attendance and mortality: Implications for the black–white mortality crossover. *Demography*, 43, 141–164.
- Ellison, C. G., & Levin, J. S. (1998). The religion–health connection: Evidence, theory, and future directions. *Health Education and Behaviors*, 25, 700–720.
- Elo, I. T., & Preston, S. H. (1992). Effects of early life conditions on adult mortality: A review. *Population Index*, 58, 186–212.
- Enright, R. D., & North, J. E. (1998). *Exploring forgiveness*. Madison, WI: University of Wisconsin Press.
- Ernst, S., Welke, J., Heintze, C., Zöllner, A., Kiehne, S., Schwantes, U., et al. (2008). Effects of mindfulness-based stress reduction on quality of life in nursing home residents: a feasibility study. *Forschende Komplementärmedizin/Research in Complementary Medicine*, 15, 74–81.
- Fang, C. Y., Reibel, D. K., Longacre, M. L., Rosenzweig, S., Campbell, D. E., & Douglas, S. D. (2010). Enhanced psychosocial well-being following participation in a mindfulness-based stress reduction program is associated with increased natural killer cell activity. *The Journal of Alternative and Complementary Medicine*, 16, 531–538.
- Fetzer Institute (2003). *Multidimensional measurement of religiousness/spirituality for use in health research: A report of a national working group*. Kalamazoo, MI: Fetzer Institute.
- Fries, J. F. (2003). Measuring and monitoring success in the compression of morbidity. *Annals of Internal Medicine*, 139, 455–459.
- Galobardes, B., Smith, G. D., & Lynch, J. W. (2006). Systematic review of the influence of childhood socioeconomic circumstances on risk for cardiovascular disease in adulthood. *Annals of Epidemiology*, 16, 91–104.
- George, L. K., Larson, D. B., Koenig, H. G., & McCullough, M. E. (2000). Spirituality and health: What we know, what we need to know. *Journal of Social and Clinical Psychology*, 19, 102–116.
- Gillum, R. F. (2006). Frequency of attendance at religious services, overweight, and obesity in American women and men: The Third National Health and Nutrition Examination Survey. *Annals of Epidemiology*, 16, 655–660.
- Hank, K., & Schaan, B. (2008). Cross-national variations in the correlation between frequency of prayer and health among older Europeans. *Research on Aging*, 30, 36–54.
- Helm, H. M., Hays, J. C., Flint, E. P., Koenig, H. G., & Blazer, D. G. (2000). Does private religious activity prolong survival? A six-year follow-up study of 3851 older adults. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 55, M400–M405.
- Hidajat, M., Zimmer, Z., Saito, Y., & Lin, H.-S. (2013). Religious activity, life expectancy, and disability-free life expectancy in Taiwan. *European Journal of Ageing*, 10, 1–8.
- Hill, P. C., Pargament, K. I., Hood, R. W., McCullough, M. E., Swyers, J. P., Larson, D. B., et al. (2000). Conceptualizing religion and spirituality: Points of commonality, points of departure. *Journal for the Theory of Social Behaviour*, 30, 51–77.
- Hill, T. D., Ellison, C. G., Burdette, A. M., & Musick, M. A. (2007). Religious involvement and healthy lifestyles: Evidence from the survey of Texas adults. *Annals of Behavioral Medicine*, 34, 217–222.
- Hiltner, S. (1943). *Religion and health*. New York: Macmillan.
- Holmes III, U. T. (2002). *Spirituality for ministry*. Harrisburg, PA: Morehouse Publishing.
- Hummer, R. A. (2005). Commentary: Understanding religious involvement and mortality risk in the United States: Comment on Bagiella, Hong, and Sloan. *International Journal of Epidemiology*, 34, 452–453.
- Hummer, R. A., Benjamins, M. R., Ellison, C. G., & Rogers, R. G. (2010). Religious involvement and mortality risk among pre-retirement aged US adults In: C. G. Ellison, & R. A. Hummer (Eds.), *Religion, families, and health: Population-based research in the United States* (pp. 273–291). New Brunswick, NJ: Rutgers University Press.
- Hummer, R. A., Ellison, C. G., Rogers, R. G., Moulton, B. E., & Romero, R. R. (2004). Religious involvement and adult mortality in the United States: Review and perspective. *Southern Medical Journal*, 97, 1223–1230.
- Hummer, R. A., Rogers, R. G., Nam, C. B., & Ellison, C. G. (1999). Religious involvement and US adult mortality. *Demography*, 36, 273–285.
- Idler, E. L., & Kasl, S. V. (1997a). Religion among disabled and nondisabled persons I: Cross-sectional patterns in health practices, social activities, and well-being. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 52, S294–S305.
- Idler, E. L., & Kasl, S. V. (1997b). Religion among disabled and nondisabled persons II: Attendance at religious services as a predictor of the course of disability. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 52, S306–S316.
- Idler, E. L., Musick, M. A., Ellison, C. G., George, L. K., Krause, N., Ory, M. G., et al. (2003). Measuring multiple dimensions of religion and spirituality for health research conceptual background and findings from the 1998 general social survey. *Research on Aging*, 25, 327–365.
- Ivanovski, B., & Malhi, G. S. (2007). The psychological and neurophysiological concomitants of mindfulness forms of meditation. *Acta Neuropsychiatrica*, 19, 76–91.
- Jagger, C. (2000). Compression or expansion of morbidity – What does the future hold. *Age and Ageing*, 29, 93–94.
- Jagger, C. (2006). Can we live longer, healthier lives? In: Y. Zeng (Ed.), *Longer life and healthy aging* (pp. 7–22). New York: Springer.
- Jagger, C., Gillies, C., Moscone, F., Cambois, E., Van Oyen, H., Nusselder, W., et al. (2009). Inequalities in healthy life years in the 25 countries of the European Union in 2005: A cross-national meta-regression analysis. *The Lancet*, 372, 2124–2131.
- Jantos, M., & Kiat, H. (2007). Prayer as medicine: How much have we learned? *Medical Journal of Australia*, 186, 51–53.
- Jevning, R., Wallace, R., & Beidebach, M. (1992). The physiology of meditation: A review. A wakeful hypometabolic integrated response. *Neuroscience Biobehavioral Reviews*, 16, 415–424.
- Kirby, S. E., Coleman, P. G., & Daley, D. (2004). Spirituality and well-being in frail and nonfrail older adults. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 59, P123–P129.
- Kodzi, I. A., Gyimah, S. O., Emina, J., & Ezech, A. C. (2011). Religious involvement, social engagement, and subjective health status of older residents of informal neighborhoods of Nairobi. *Journal of Urban Health*, 88, 370–380.
- Koenig, H. G. (1989). Research on religion and mental health in later life: A review and commentary. *Journal of Geriatric Psychiatry*, 23, 23–53.
- Koenig, H. G. (2001). Religion, spirituality, and medicine: How are they related and what does it mean? *Mayo Clinic Proceedings*, 76, 1189–1191.
- Koenig, H. G. (2009). Research on religion, spirituality, and mental health: A review. *Canadian Journal of Psychiatry*, 54, 283–291.
- Koenig, H. G. (2012). Religion, spirituality, and health: The research and clinical implications. *International Scholarly Research Network*, 2012, 1–33.
- Koenig, H. G. (2013). *Is religion good for your health?: The effects of religion on physical and mental health*. New York: Routledge.
- Handbook of religion and mental health In: H. G. Koenig (Ed.), San Diego, CA: Elsevier.
- Koenig, H. G., Cohen, H. J., Blazer, D. G., Pieper, C., Meador, K. G., Shelp, F., et al. (1992). Religious coping and depression among elderly, hospitalized medically ill men. *The American journal of psychiatry*, 149, 1693–1700.

- Koenig, H. G., Hays, J. C., Larson, D. B., George, L. K., Cohen, H. J., McCullough, M. E., et al. (1999). Does religious attendance prolong survival? A six-year follow-up study of 3968 older adults. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 54, 370–376.
- Koenig, H. G., King, D., & Carson, V. B. (2012). *Handbook of religion and health*. New York: Oxford University Press.
- Koenig, H. G., Larson, D. B., Hays, J. C., McCullough, M. E., George, L. K., Branch, P. S., et al. (1998). Religion and the survival of 1010 hospitalized veterans. *Journal of Religion and Health*, 37, 15–30.
- Koszycski, D., Benger, M., Shlik, J., & Bradwejn, J. (2007). Randomized trial of a meditation-based stress reduction program and cognitive behavior therapy in generalized social anxiety disorder. *Behaviour Research and Therapy*, 45, 2518–2526.
- Krause, N. (1997). Religion, aging, and health: Current status and future prospects. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 52, S291–S293.
- Krause, N. (2002). Church-based social support and health in old age exploring variations by race. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 57, S332–S347.
- Krause, N. (2003). Religious meaning and subjective well-being in late life. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 58, S160–S170.
- Krause, N. (2004). Religion, aging, and health: Exploring new frontiers in medical care. *Southern Medical Journal*, 97, 1215–1222.
- Krause, N. (2006). Exploring the stress-buffering effects of church-based and secular social support on self-rated health in late life. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 61, 332–347.
- Krause, N. (2008). *Aging in the church: How social relationships affect health*. West Conshohocken, PA: Templeton Foundation Press.
- Krause, N. (2009). Meaning in life and mortality. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 64, 517–527.
- Krause, N. (2011). Religion and health: Making sense of a disheveled literature. *Journal of Religion and Health*, 50, 20–35.
- Krause, N., & Booth, A. (2004). *Religious influences on health and well-being in the elderly*. New York: Springer Publishing Company.
- Krause, N., Ellison, C. G., Shaw, B. A., Marcum, J. P., & Boardman, J. D. (2001). Church-based social support and religious coping. *Journal for the Scientific Study of Religion*, 40, 637–656.
- Krause, N., Ellison, C. G., & Wulff, K. M. (1998). Church-based emotional support, negative interaction, and psychological well-being: Findings from a national sample of Presbyterians. *Journal for the Scientific Study of Religion*, 37, 725–741.
- Krause, N., & Hayward, D. R. (2014). Religious involvement, practical wisdom, and self-rated health. *Journal of Aging and Health*, 26, 540–558.
- Krause, N., & Wulff, K. M. (2004). Religious doubt and health: Exploring the potential dark side of religion. *Sociology of Religion*, 65, 35–56.
- Kune, G. A., Kune, S., & Watson, L. F. (1993). Perceived religiousness is protective for colorectal cancer: Data from the Melbourne Colorectal Cancer Study. *Journal of the Royal Society of Medicine*, 86, 645–647.
- Lalivé d'Épinay, C., & Spini, D. (2004). Religion and health: a European perspective. In: K. W. Schaie, N. Krause, & A. Booth (Eds.), *Religious influence on health and well-being in the elderly* (pp. 44–58). New York: Springer.
- Larson, D. B., Swyers, J. P., & McCullough, M. E. (1998). *Scientific research on spirituality and health: A report based on the scientific progress in spirituality conferences*. Washington, DC: National Institute for Healthcare Research.
- Lavretsky, H. (2010). Spirituality and aging. *Aging Health*, 6, 749–769.
- Ledesma, D., & Kumano, H. (2009). Mindfulness-based stress reduction and cancer: A meta-analysis. *Psycho-Oncology*, 18, 571–579.
- Levin, J. S., & Chatters, L. M. (2008). Religion, aging, and health: Historical perspectives, current trends, and future directions: Public health. *Journal of Religion, Spirituality Aging*, 20, 153–172.
- Levin, J. S., Taylor, R. J., & Chatters, L. M. (1994). Race and gender differences in religiosity among older adults: Findings from four national surveys. *Journal of Gerontology*, 49, S137–S145.
- Levin, J. S., & Vanderpool, H. Y. (1991). Religious factors in physical health and the prevention of illness. *Prevention in Human Services*, 9, 41–64.
- Liu, E. Y., Schieman, S., & Jang, S. J. (2011). Religiousness, spirituality, and psychological distress in Taiwan. *Review of Religious Research*, 53, 137–159.
- Lucchetti, G., Lucchetti, A. G. L., Badan-Neto, A. M., Peres, P. T., Peres, M. F., Moreira-Almeida, A., et al. (2011a). Religiousness affects mental health, pain and quality of life in older people in an outpatient rehabilitation setting. *Journal of Rehabilitation Medicine*, 43, 316–322.
- Lucchetti, G., Lucchetti, A. L., & Koenig, H. G. (2011b). Impact of spirituality/religiosity on mortality: Comparison with other health interventions. *EXPLORE: The Journal of Science and Healing*, 7, 234–238.
- Lundberg, C. D. (2010). *Unifying truths of the world's religions: Practical principles for living and loving in peace*. New Fairfield, CT: Heavenlight Press.
- Lutz, A., Slagter, H. A., Dunne, J. D., & Davidson, R. J. (2008). Attention regulation and monitoring in meditation. *Trends in Cognitive Sciences*, 12, 163–169.
- MacKinlay, E. (2002). The spiritual dimension of caring: Applying a model for spiritual tasks of ageing. *Journal of Religious Gerontology*, 12, 151–166.
- Marks, L. (2005). Religion and bio-psycho-social health: A review and conceptual model. *Journal of Religion and Health*, 44, 173–186.
- Maselko, J., Hayward, R. D., Hanlon, A., Buka, S., & Meador, K. (2012). Religious service attendance and major depression: A case of reverse causality? *American Journal of Epidemiology*, 175, 576–583.
- Mayhew, L. (1999). *Health and welfare services expenditure in an aging world*. Laxenburg, Austria: International Institute for Applied Systems Analysis.
- McCullough, M. E., Hoyt, W. T., Larson, D. B., Koenig, H. G., & Thoresen, C. (2000). Religious involvement and mortality: A meta-analytic review. *Health Psychology*, 19, 211–222.
- McCullough, M. E., & Willoughby, B. L. B. (2009). Religion, self-regulation and self-control: Associations, explanations and implications. *Psychological Bulletin*, 135, 69–93.
- McFadden, S. H. (1995). Religion and well-being in aging persons in an aging society. *Journal of Social Issues*, 51, 161–175.
- Miller, W. R., & Thoresen, C. E. (2003). Spirituality, religion, and health: An emerging research field. *American Psychologist*, 58, 24–35.
- Moberg, D. O. (2005). Research in spirituality, religion, and aging. *Journal of Gerontological Social Work*, 45, 11–40.
- Mueller, P. S., Plevak, D. J., & Rummans, T. A. (2001). Religious involvement, spirituality, and medicine: Implications for clinical practice. *Mayo Clinic Proceedings*, 76, 1225–1235.
- Musick, M. A., House, J. S., & Williams, D. R. (2004). Attendance at religious services and mortality in a national sample. *Journal of Health and Social Behavior*, 45, 198–213.
- Musick, M. A., Traphagan, J. W., Koenig, H. G., & Larson, D. B. (2000). Spirituality in physical health and aging. *Journal of Adult Development*, 7, 73–86.
- Nelson, J. K. (2000). *Enduring identities: The guise of Shinto in contemporary Japan*. Honolulu, HI: University of Hawaii Press.
- Nooney, J., & Woodrum, E. (2002). Religious coping and church-based social support as predictors of mental health outcomes: Testing a conceptual model. *Journal for the Scientific Study of Religion*, 41, 359–368.
- Nusselder, W. J. (2003). Compression of morbidity. In: J.-M. Robine, C. Jagger, C. D. Mathers, E. M. Crimmins, & R. M. Suzman (Eds.), *Determining Health expectancies* (pp. 35–57). Hoboken, NY: John Wiley & Sons.
- O'Brien, J., Palmer, M., & Barrett, D. B. (2007). *The atlas of religion*. Berkeley: University of California Press.
- Ortiz, L. P., & Langer, N. (2002). Assessment of spirituality and religion in later life: Acknowledging clients' needs and personal resources. *Journal of Gerontological Social Work*, 37, 5–21.
- Oxman, T. E., Freeman, D. H., & Manheimer, E. D. (1995). Lack of social participation or religious strength and comfort as risk factors for death after cardiac surgery in the elderly. *Psychosomatic Medicine*, 57, 5–15.
- Pargament, K. I. (1997). *The psychology of religion and coping: Theory, practice and research*. New York: Guilford.
- Pargament, K. I., Koenig, H. G., & Perez, L. M. (2000). The many methods of religious coping: Development and initial validation of the RCOPE. *Journal of Clinical Psychology*, 56, 519–543.
- Pargament, K. I., Koenig, H. G., Tarakeshwar, N., & Hahn, J. (2001). Religious struggle as a predictor of mortality among medically ill elderly patients: A 2-year longitudinal study. *Archives of Internal Medicine*, 161, 1881–1885.
- Pargament, K. I., & Raiya, H. A. (2007). A decade of research on the psychology of religion and coping: Things we assumed and lessons we learned. *Psyke & Logos*, 28, 742–766.
- Poloma, M. M., & Pendleton, B. F. (1989). Exploring types of prayer and quality of life: A research note. *Review of Religious Research*, 31, 46–53.
- Poloma, M. M., & Pendleton, B. F. (1991). The effects of prayer and prayer experiences on measures of general well-being. *Journal of Psychology and Theology*, 19, 71–83.
- Powell, L. H., Shahabi, L., & Thoresen, C. E. (2003). Religion and spirituality: Linkages to physical health. *American Psychologist*, 58, 36–52.
- Reed, P. G. (1991). Spirituality and mental health in older adults: Extant knowledge for nursing. *Family & Community Health*, 14, 14–25.
- Reyes-Ortiz, C. A., Pelaez, M., Koenig, H. G., & Mulligan, T. (2007). Religiosity and self-rated health among Latin American and Caribbean elders. *The International Journal of Psychiatry in Medicine*, 37, 425–443.
- Richards, P. S., & Bergin, A. E. (2005). *A spiritual strategy for counseling and psychotherapy*. Washington, DC: American Psychological Association.
- Rippentrop, A. E., Altmaier, E. M., Chen, J. J., Found, E. M., & Keffala, V. J. (2005). The relationship between religion/spirituality and physical health, mental health, and pain in a chronic pain population. *Pain*, 116, 311–321.
- Robine, J.-M., Jagger, C., & Romieu, I. (2001). Disability-free life expectancies in the European Union countries: Calculation and comparisons. *Genus*, 42, 89–101.
- Roff, L. L., Klemmack, D. L., Simon, C., Cho, G. W., Parker, M. W., Koenig, H. G., et al. (2006). Functional limitations and religious service attendance among African American and white older adults. *Health and Social Work*, 31, 246–255.
- Rogers, T. (1976). Manifestations of religiosity and the aging process. *Religious Education*, 71, 405–415.
- Ross, A., & Thomas, S. (2010). The health benefits of yoga and exercise: A review of comparison studies. *The Journal of Alternative and Complementary Medicine*, 16, 3–12.
- Roth, B., & Creaser, T. (1997). Mindfulness meditation-based stress reduction: Experience with a bilingual inner-city program. *The Nurse Practitioner*, 22, 150–178.
- Ryff, C. D., & Singer, B. (1998). The contours of positive human health. *Psychological Inquiry*, 9, 1–28.
- Saad, Z. M., Hatta, Z. A., & Mohamad, N. (2010). The impact of spiritual intelligence on the health of the elderly in Malaysia. *Asian Social Work and Policy Review*, 4, 84–97.
- Saito, Y., Qiao, X., & Jitapunkul, S. (2003). Health expectancy in Asian countries. In: J.-M. Robine, C. Jagger, C. D. Mathers, E. M. Crimmins, & R. M. Suzman (Eds.), *Determining health expectancies* (pp. 289–318). Hoboken, NJ: John Wiley & Sons.

- Schlehofer, M. M., Omoto, A. M., & Adelman, J. R. (2008). How do "religion" and "spirituality" differ? Lay definitions among older adults. *Journal for the Scientific Study of Religion*, 47, 411–425.
- Seeman, T. E., Dubin, L. F., & Seeman, M. (2003). Religiosity/spirituality and health: A critical review of the evidence for biological pathways. *American Psychologist*, 58, 53–63.
- Seybold, K. S., & Hill, P. C. (2001). The role of religion and spirituality in mental and physical health. *Current Directions in Psychological Science*, 10, 21–24.
- Shapiro, S. L., & Carlson, L. E. (2009). *The art and science of mindfulness: Integrating mindfulness into psychology and the helping professions*. Washington, DC: American Psychological Association.
- Sloan, R. P., Bagiella, E., & Powell, T. (1999). *Reliness: Integrating mindfulness into psychology and the helping professions*. *The Lancet*, 353. Washington, DC: American Psychological Association, spirituality, and medicine.
- Smith, K. R., Mineau, G. P., Garibotti, G., & Kerber, R. (2009). Effects of childhood and middle-adulthood family conditions on later-life mortality: Evidence from the Utah Population Database, 1850–2002. *Social Science and Medicine*, 68, 1649–1658.
- Solberg, E., Ingjier, F., Ekberg, O., Holen, A., Standal, P., & Vikman, A. (2000). Blood pressure and heart rate during meditation. *Journal of Psychosomatic Research*, 48, 283–287.
- Sørensen, T., Danbolt, L. J., Lien, L., Koenig, H. G., & Holmen, J. (2011). The relationship between religious attendance and blood pressure: The HUNT study, Norway. *The International Journal of Psychiatry in Medicine*, 42, 13–28.
- Strawbridge, W. J., Cohen, R. D., Shema, S. J., & Kaplan, G. A. (1997). Frequent attendance at religious services and mortality over 28 years. *American Journal of Public Health*, 87, 957–961.
- Strawbridge, W. J., Shema, S. J., Cohen, R. D., & Kaplan, G. A. (2001). Religious attendance increases survival by improving and maintaining good health behaviors, mental health, and social relationships. *Annals of Behavioral Medicine*, 23, 68–74.
- Sullivan, A. R. (2010). Mortality differentials and religion in the United States: Religious affiliation and attendance. *Journal for the Scientific Study of Religion*, 49, 740–753.
- Takyi, B. K. (2003). Religion and women's health in Ghana: Insights into HIV/AIDS preventive and protective behavior. *Social Science Medicine*, 56, 1221–1234.
- Tan, H., Wutthilert, C., & O'Connor, M. (2011). Spirituality and quality of life in older people with chronic illness in Thailand. *Progress in Palliative Care*, 19, 177–184.
- Temby, O. F., & Smith, K. R. (2013). The association between adult mortality risk and family history of longevity: The moderating effects of socioeconomic status. *Journal of Biosocial Science*, 45, 1–14.
- Thege, B. K., Pilling, J., Székely, A., & Kopp, M. S. (2013). Relationship between religiosity and health: Evidence from a post-communist country. *International Journal of Behavioral Medicine*, 20, 477–486.
- Thoresen, C. E., Harris, A. H., & Luskin, F. (2000). *Forgiveness and health: An unanswered question*. New York: Guilford Press.
- UNFPA, & HelpAge International (2012). *Ageing in the twenty-first century: A celebration and a challenge*. New York: United Nations Population Fund and Help Age International.
- United Nations (2015). *World population prospects: The 2015 revision*. New York: United Nations.
- Vaupel, J. W., & Kistowski, K.Gv (2005). Broken limits to life expectancy. *Ageing Horizons*, 3, 6–13.
- Walsh, A. (1998). Religion and hypertension: Testing alternative explanations among immigrants. *Behavioral Medicine*, 24, 122–130.
- Wilhelm, M. O., Rooney, P. M., & Tempel, E. R. (2007). Changes in religious giving reflect changes in involvement: Age and cohort effects in religious giving, secular giving, and attendance. *Journal for the Scientific Study of Religion*, 46, 217–232.
- Wilson, J., & Musick, M. (1997). Who cares? Toward an integrated theory of volunteer work. *American Sociological Review*, 62, 694–713.
- Wink, P., & Dillon, M. (2001). Religious involvement and health outcomes in late adulthood In: T. G. Plante, & A. C. Sherman (Eds.), *Faith and health: Psychological perspectives* (pp. 75–106). New York: Guilford Press.
- World Health Organization (2015). *World report on ageing and health*. Geneva: World Health Organization.
- World Values Survey Association (2014). *Wave 6 2010–2014 OFFICIAL AGGREGATE v.20150418. Aggregate File Producer: Asep/JDS*. Madrid, Spain: World Values Survey Association.
- Yeager, D. M., Gleib, D. A., Au, M., Lin, H.-S., Sloan, R. P., & Weinstein, M. (2006). Religious involvement and health outcomes among older persons in Taiwan. *Social Science Medicine*, 63, 2228–2241.
- Young, J. D.-E., & Taylor, E. (1998). Meditation as a voluntary hypometabolic state of biological estivation. *Physiology*, 13, 149–153.
- Zeng, Y., Gu, D., & George, L. K. (2011). Association of religious participation with mortality among Chinese old adults. *Research on Aging*, 33, 51–83.
- Zhang, W. (2008). Religious participation and mortality risk among the oldest old in China. *Journal of Gerontology: Social Sciences*, 63, 293–297.
- Zimmer, Z., & McDaniel, S. (2013). *Global ageing in the twenty-first century: An introduction. Global ageing in the twenty-first century: Challenges, opportunities and implications* (pp. 1–12). Surrey: Ashgate.
- Zinnbauer, B. J., Pargament, K. I., Cole, B., Rye, M. S., Butter, E. M., Belavich, T. G., et al. (1997). Religion and spirituality: Unfuzzifying the fuzzy. *Journal for the Scientific Study of Religion*, 36, 549–564.