

Outcomes in patients with glenoid labral lesions: a cohort study

Marc Zughaib, Christopher B Robbins, Bruce S Miller, Joel J Gagnier

To cite: Zughaib M, Robbins CB, Miller BS, *et al*. Outcomes in patients with glenoid labral lesions: a cohort study. *BMJ Open Sport Exerc Med* 2017;**2**:e000209. doi:10.1136/bmjsem-2016-000209

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjsem-2016-000209>).

Accepted 31 January 2017

ABSTRACT

Background/aim: For patients presenting with glenoid labral pathologies, there is little information on how operative interventions affect long-term outcomes and health-related quality of life (HRQoL). This study evaluated outcomes in operative and non-operative patients presenting with labral tears versus labral degenerations.

Methods: Participants completed a pain Visual Analogue Scale (VAS), the Veterans RAND 12-item Health Survey, Western Ontario Rotator Cuff Index, American Shoulder and Elbow Surgeons instrument, Scapular Assistance Test (SAT), Shoulder Activity Level, and Single Assessment Numeric Evaluation at baseline and at 6-month, 12-month and 2-year follow-ups. χ^2 and Student's t-test were used to test the differences between categorical and continuous variables. Analysis of variance investigated the differences between groups, and linear regression analyses explored the relationship of baseline characteristics with outcome scores.

Results: After 2 years, the operative cohort (n=68) significantly improved in all measures. The non-operative cohort (n=55) showed significant improvements in all scores except the mental component summary (MCS) and pain VAS. Labral tear patients (n=52) within the operative group (n=28) significantly improved in all measures except MCS. Non-operative labral tear patients (n=24) indicated significant improvements in all measures except MCS, VAS and SAT. Labral degeneration patients (n=71) within the operative group (n=27) significantly improved in all measures except MCS and SAT. Non-operative labral degeneration patients (n=44) indicated significant improvements in all measures except the physical component summary, MCS, VAS and SAT.

Conclusion: Patients who were surgically treated for labral tears or degenerations had significantly improved outcomes and HRQoL scores after 2 years compared with the non-operative cohort.

INTRODUCTION

The anatomic configuration of the shoulder joint (glenohumeral joint) is often compared with a golf ball on a tee. The labrum is a fibrocartilaginous ring that attaches to the bony rim of the glenoid fossa,¹ doubling the depth of the glenoid fossa to help provide stability.² The superior labrum attaches to the superior glenoid

Key messages

What are the new findings?

- This study highlights the impact of glenoid labral lesions in patients with rotator cuff tears on health-related quality-of-life outcomes.
- Patients with rotator cuff tear with labral pathologies tend to do worse on outcomes than those without labral pathologies.

How might it impact clinical practice in the near future?

- This study could help guide clinicians on the importance of labral pathologies in predicting outcomes in patients with rotator cuff tear.

by loose connective fibres and provides attachment for the long head of the biceps tendon, forming the biceps anchor. The anterior superior labrum is attached to the middle and inferior glenohumeral ligaments, and the inferior labrum is attached firmly to the glenoid by inelastic fibrous tissue.³

Glenoid labral lesions can often lead to significant discomfort and restriction during daily living activities, as well as various sporting activities.⁴ The prevalence of these tears can vary between 6% in the general population and 35% in the sporting population.^{5 6} The high prevalence of shoulder dysfunction is associated with high societal cost and patient burden. In 2013, a reported 86 690 work-related shoulder injuries and illnesses involving days away from work occurred in the United States.⁷ In 2005, the treatment measures of shoulder pain accumulated to an estimated annual cost of \$39 billion.⁸ Zhang *et al*⁹ reported the highest incidence of repair is in those aged 20–29 years and 40–49 years. In addition, a significant gender difference exists, with men having three times higher incidence of repair.⁹

Glenoid labral pathology can be classified based on location and morphology. With respect to location, the labrum can be



CrossMark

Department of Orthopaedic Surgery, University of Michigan, Ann Arbor, Michigan, USA

Correspondence to

Dr Joel J Gagnier; jgagnier@med.umich.edu

divided into six areas (superior, posterosuperior, posteroinferior, inferior, anteroinferior and anterosuperior) or alternatively into clock-face 'time zones' (1–12 o'clock) (figures 1 and 2).¹⁰ Labral pathology can be classified morphologically as torn, degenerated or blunted. Diagnosis of these pathologies can be

optimally performed using magnetic resonance arthrography (MRA). Studies have shown the sensitivity of MRA to be between 82% and 100%, with a specificity of between 71% and 98%.^{11–14}

Tears of the superior labrum account for 80%–90% of labral pathology in the stable shoulder.¹⁵ These lesions were first described in 1990 by Snyder *et al*¹⁶ as superior labrum anterior posterior (SLAP) tears. Originally SLAP tears were classified into four types (I–IV). Subsequently, lesion types V–IX were included in the classification as well.^{17 18} The most common mechanisms of injury include chronic, repetitive microtrauma secondary to overhead throwing-type movement or an acute fall onto the outstretched hand.¹⁹

Treatment options for patients presenting with glenoid labral pathology include both surgical and non-surgical interventions. Previous work determined that the population incidence of SLAP repairs in years 2002 and 2009 increased from 3.54 to 10.89 per 100 000.²⁰ Multiple reports document successful outcomes with operative repair of SLAP tears.^{5 16 21–28} Non-operative repair of SLAP lesions has also been associated with successful outcomes.²⁹ Edwards *et al* reported significant improvements in pain, function and health-related quality of life (HRQoL). The HRQoL measure is increasingly used within orthopaedic research, as clinicians increasingly look beyond strictly functional outcomes to physical, psychological and social factors.³⁰




Figure 1 Labral division: six areas. Figure reproduced from Mohana-Borges *et al*¹⁰ with tentative permission of the American Roentgen Ray Society.




Figure 2 Labral division: time zones. For both shoulders, 12- to 6-o'clock position faces anteriorly, and 6- to 12-o'clock position faces posteriorly. Figure reproduced from Mohana-Borges *et al*¹⁰ with tentative permission of the American Roentgen Ray Society.

Unfortunately, there is not much information on the HRQoL associated with operative interventions of labral pathology. The incentive for this shift from a biomedical model to a psychosocial model has been influenced by research addressing the relationship to HRQoL and successful outcomes within a variety of orthopaedic disciplines, including labral and rotator cuff disease.^{31–33} It is the intent of this research to examine the outcomes and HRQoL between operative and non-operative groups in a cohort of patients with labral pathology. The research will also compare patients presenting with labral tears with those presenting with labral degenerations.

METHODS

Study design

A cohort sample of 123 patients presenting with glenoid labral pathologies (tears or degenerations) was identified retrospectively by sampling consecutive patients from July 2012 to January 2015 from a pragmatic prospective Shoulder Registry cohort.

Inclusion/exclusion criteria

The following were the inclusion criteria: (1) greater than 18 years of age, (2) labral tear of any size and (3) the first tear of the affected shoulder. The following were the exclusion criteria: (1) a labral tear in which complete footprint coverage is not possible (for the surgical group); and (2) history of prior surgery, fracture, dislocation or infection of the affected shoulder.

Data collection

Study approval was obtained from the University of Michigan Institutional Review Board (no HUM00056320). Participants provided baseline

demographic information and completed several previously validated paper-based outcome measures. Data collected included the pain Visual Analogue Scale (VAS), the Veterans RAND 12-Item Health Survey (VR-12), Western Ontario Rotator Cuff Index (WORC), American Shoulder and Elbow Surgeons (ASES) instrument, Scapular Assistance Test (SAT), Shoulder Activity Level (SAL), and Single Assessment Numeric Evaluation (SANE). Participants were tracked for operative versus non-operative intervention and completed the same measures at baseline and at 6-month, 12-month and 2-year follow-ups (excluding SAL, which was collected at baseline only).

Measures

The primary outcome measure used was the pain VAS, which is self-completed by the respondent. The respondents are asked to place a line perpendicular to the VAS line at the point that represents their pain intensity. Using a ruler, the score is determined by measuring the distance (millimetre) on the 10 cm line between the 'no pain' anchor and the patient's mark, providing a range of scores from 0 to 100.³⁴ A higher score indicates greater pain intensity.

The VR-12 Health Survey was derived from the RAND 36-item Short-Form Health Survey (SF-36).^{35–38}

The VR-12 is a self-administered health survey used to measure HRQoL and estimate disease burden. This instrument comprises 12 items corresponding to eight principal physical and mental health domains. These domains include general health perceptions, physical functioning, role limitations due to physical and emotional problems, bodily pain, energy-fatigue, social functioning, and mental health. The 12 items are then summarised into two scores, a 'physical component summary' (PCS) score and a 'mental component summary' (MCS) score.

The WORC index is a valid, reliable and responsive HRQoL self-assessment measurement tool for shoulder disease.^{39–40} A recently published study determined the WORC to have the highest ratings among all shoulder instruments.⁴¹ The items included in the instrument were determined based on a literature review, along with discussions among patients, orthopaedic surgeons, sports and family physicians, and physiotherapists. The WORC includes 21 questions encompassing five domains. The domains include pain and physical symptoms (six questions), sports and recreation (four questions), work function (four questions), social function (four questions) and emotional function (three questions). In addition to these questions, a 100 mm VAS is included. Scores range from 0 to 100, with a higher score indicating worse shoulder function.

The ASES shoulder outcome score was developed in 1994 in the United States by the research committee of the ASES.⁴² The purpose was to measure shoulder function through a physician-determined and self-assessment questionnaire. The instrument is divided

into three domains: pain, instability and activities of daily living. These domains includes several yes/no questions and a VAS. A 4-point ordinal scale rates 10 questions, with a maximum score for the self-assessment portion of 30. This score is converted to a 100-point scale (100 is the best score). The construct validity, internal consistency and reliability have been shown to be good.^{43–48}

The SAT instrument is presumed to indirectly measure the function of the scapular rotators.⁴⁹ In this test, the examiner pushes laterally and upward on the inferior medial border of the scapula to simulate the serratus anterior/lower trapezius force couple as the patient elevates the arm. According to Kibler,⁵⁰ the SAT is considered positive if the manual assistance diminished or abolished the symptoms of impingement. The subject rates the pain felt while performing the movement with the assistance of the examiner on an 11-point scale. A reduction of two or more points on the 11-point numeric pain rating scale during assisted elevation, compared with elevation without assistance, is considered a positive test. Previous work has shown that a reduction of 2 points on an 11-point numeric pain rating scale represents a clinically important difference.⁵¹ This instrument has been shown to provide moderate inter-rater reliability.⁵²

The SAL instrument generates a score based on the patient's greatest level of activity during the previous year. The most recent level of activity may be limited by other factors such as weather, conflicting commitments, or complicating illness or injury.⁵³ The patient's greatest activity level during a 1-year period is likely to be the most accurate.⁵⁴ A reasonable interpretation of the numeric activity score is high, ≥ 16 (mean+SD); average, 7–15; and low, ≤ 6 (mean–SD).

The SANE rating is determined by the patient's written response to the following question: 'How would you rate your shoulder today as a percentage of normal (0% to 100% scale, with 100% being normal)?' SANE correlates well with similar outcome scores; correlation coefficients of 0.46–0.69 have been found for SANE and ASES scores.⁵⁵

Statistical analysis

A sample-size calculation was performed ($n=54$, $\alpha=0.05$, $\beta=0.10$, effect size=0.913) using a hypothesised difference in change in pain scores in operative versus non-operative interventions (G*Power v3.1.9.2). Descriptive and inferential analyses were conducted using SAS v. 9.4. χ^2 analysis and Student's t-test were used to examine the differences between categorical and continuous variables at baseline and at 2-year follow-up. Analysis of variance was used to test the differences between groups. Data were then stratified based on labral pathology. Methods were repeated for each level of stratification.

Simple linear regression analyses were performed on statistically significant baseline characteristics within

each strata of labral pathology. These outcome measures for patients with labral tear included pain VAS and ASES scores at baseline. For labral degeneration patients, models were created using age and ASES scores at baseline. Change in pain VAS from baseline to 2-year follow-up was used as the outcome variable for the models.

RESULTS

Participant demographics consisted of 123 patients. Of these, 82 were male and 41 were female (table 1). There were 68 participants in the non-operative group and 55 in the operative group. Comparing baseline demographics and clinical characteristics indicated a significant difference in age ($p=0.01$), with non-operative patients being older. Among baseline scores, there were significant differences in pain VAS ($p=0.02$) and baseline ASES score ($p=0.01$), with both baseline scores indicating 'worse' scores for the operative patients. No significant differences were noted between groups for baseline PCS, MCS, WORC, SAT, SANE and SAL scores.

Paired sample t-tests within the operative group indicate significant improvements between baseline and 2-year scores for pain VAS (mean difference (MD) = -38.51 , $p<0.01$), PCS (MD = 8.75 , $p<0.01$), MCS (MD = 3.49 , $p=0.05$), ASES (MD = 35.53 , $p<0.01$), WORC (MD = -826.2 , $p<0.01$), SAT (MD = 9.82 , $p<0.01$) and SANE (MD = 53.08 , $p<0.01$) (table 2). Paired sample t-tests within the non-operative group indicate significant improvements between baseline and 2-year scores for PCS (MD = 3.53 , $p=0.02$), ASES (MD = 10.63 , $p<0.01$), WORC (MD = -295.3 , $p<0.01$), SAT (MD = 7.95 , $p=0.02$) and SANE (MD = 21.89 , $p<0.01$). When comparing the difference of the 2-year means of outcome measures between groups, only MCS ($p=0.81$) and SAT ($p=0.1$) did not display a significant improvement.

Labral tear patient demographics consisted of 52 patients. Of these, 36 were male and 16 were female (table 3). There were 28 patients in the operative group and 24 in the non-

operative group. Comparing baseline demographics and clinical characteristics revealed no significant differences. Among baseline outcome scores, there were differences in pain VAS ($p=0.04$) and ASES scores ($p=0.05$). Both baseline outcome scores indicated 'worse' for the operative patients. No significant differences were noted between groups for baseline PCS, MCS, WORC, SAT, SANE and SAL scores.

Paired sample t-tests for patients with labral tear within the operative group indicate significant improvements between baseline and 2-year follow-up scores for pain VAS (MD = -43.08 , $p<0.01$), PCS (MD = 10.81 , $p<0.01$), ASES (MD = 35.39 , $p<0.01$), WORC (MD = -891.74 , $p<0.01$), SAT (MD = 11.41 , $p=0.03$) and SANE (MD = 55.08 , $p<0.01$) (table 4). Paired sample t-tests within the non-operative group

for patients with labral tear indicate significant improvements between baseline and 2-year follow-up scores for the PCS (MD = 6.89 , $p=0.02$), ASES (MD = 7.38 , $p<0.01$), WORC (MD = -277.76 , $p=0.04$) and SANE (MD = 15.55 , $p=0.03$). When comparing the difference of the 2-year means of outcome measures between groups, only PCS ($p=0.07$), MCS ($p=0.77$) and SAT ($p=0.07$) did not display a significant improvement.

Labral degeneration patient demographics consisted of 71 patients. Of these, 46 were male and 25 were female (table 5). There were 27 patients in the operative group and 44 in the non-operative group. Comparing

baseline demographics and clinical characteristics indicated a significant difference in age ($p=0.01$), with non-operative patients being older. No significant differences in outcome measures were noted between groups.

Paired sample t-tests for patients with labral degeneration within the operative group ($n=27$) indicated significant improvements between baseline and 2-year follow-up scores for pain VAS (MD = -33.94 , $p<0.01$), PCS (MD = 6.55 , $p=0.01$), ASES (MD = 35.71 , $p<0.01$), WORC (MD = -757.06 , $p<0.01$) and SANE (MD = 51.08 , $p<0.01$) (table 6). Paired sample t-tests within the non-operative group ($n=44$) for patients with labral degeneration indicate significant improvements between baseline and 2-year follow-up scores for ASES (MD = 12.72 , $p=0.01$), WORC (MD = -305.59 , $p=0.02$) and SANE (MD = 26.41 , $p<0.01$). When comparing the difference of the 2-year means of outcome measures between the two groups, only MCS ($p=0.51$) and SAT ($p=0.44$) did not display a significant improvement.

In our linear regression analyses for patients with labral tear using change in pain VAS score from baseline to 2-year follow-up as the response variable, we found baseline pain VAS ($\beta=-0.718$, $p<0.01$) and ASES scores ($\beta=0.597$, $p=0.03$) were significant predictor variables. Also, in the linear regression analyses for labral degeneration, we found baseline ASES scores ($\beta=0.663$, $p=0.02$) to be significant predictors of change in pain.

DISCUSSION

Many published studies have demonstrated good to excellent outcomes of repairs of labral pathology. However, orthopaedic research is increasingly assessing results using HRQoL measures. In this particular patient population, the two groups were fairly homogenous regarding variables related to the injury, such as type of labral pathology, cause of injury, side of injury and symptom duration. However, the operative group was younger and had significantly worse baseline scores for pain VAS and ASES. These differences could be attributed to the guidelines and recommendations provided to

Table 1 Demographics, clinical features and baseline outcome scores

	Operative (n=55)			Non-operative (n=68)			p Value
	Mean/n	SD/%	95% CI	Mean/n	SD/%	95% CI	
Age*	58.51	9.11	56.05 to 60.97	63.09	10.06	60.65 to 65.52	0.0100
Gender†							0.3694
Male	39	70.91		43	63.24		
Female	16	29.09		25	36.76		
Body mass index*	28.63	5.31	27.19 to 30.07	28.54	5.78	27.14 to 29.94	0.9272
Cause of injury†							0.3694
Known	39	70.91		43	63.24		
Unknown	16	29.09		25	36.76		
Labral pathology†							0.0813
Tear	28	50.91		24	35.29		
Degeneration	27	49.09		44	64.71		
Side of injury†							0.2655
Right	31	56.36		45	66.18		
Left	24	43.64		43	33.82		
Symptom duration†							0.4084
<1 year	21	38.18		31	45.59		
>1 year	34	61.82		27	54.41		
Diabetes†							0.4529
Yes	7	12.73		12	17.65		
No	48	87.27		56	82.35		
Smoking†							0.7559
Yes	4	7.27		4	5.88		
No	51	92.73		64	94.12		
Pain VAS*	54.31	26.26	46.99 to 61.62	42.57	26.93	35.95 to 49.19	0.0191
Baseline PCS*	39.11	9.02	36.14 to 42.07	40.03	8.66	37.49 to 42.58	0.6308
Baseline MCS‡	48.50	11.24	44.81 to 52.19	51.47	10.22	48.47 to 54.47	0.2909
Baseline ASES*	50.63	21.30	44.64 to 56.62	60.73	20.67	55.34 to 66.12	0.0132
Baseline WORC*	1184.30	455.20	1058.80 to 1309.80	1061.90	453.90	947.60 to 1176.20	0.1512
SAT‡	79.20	26.08	71.55 to 86.86	70.23	33.10	61.28 to 79.18	0.1726
SANE‡	30.94	24.04	24.32 to 37.57	37.34	22.88	31.67 to 43.01	0.0819

*p values calculated using t-tests.

†p values calculated using Wilcoxon signed-rank tests.

‡p values calculated using chi-square tests.

Continuous data presented as mean, SD. Categorical data presented as n (%).

ASES, American Shoulder and Elbow Surgeons; MCS, mental component summary; PCS, physical component summary; SAL, Shoulder Activity Level; SANE, Single Assessment Numeric Evaluation; SAT, Scapular Assistance Test; VAS, Visual Analogue Scale; WORC, Western Ontario Rotator Cuff Index.

physicians for determining whether a patient should elect operative or non-operative treatment. The patient who elected operative intervention experienced significantly improved long-term HRQoL outcomes and function scores. This trend held true

after stratification by labral pathology, as operative patients consistently reported more favourable outcome scores after the 2-year follow-up period.

In this patient population, pain VAS was significantly higher in the operative group at baseline. However,

Table 2 Mean differences of outcome measures from baseline to 2-year follow-up within and between operative and non-operative groups

	Operative (n=55)						Non-operative (n=68)						
	n	Baseline mean	2-year mean	Mean difference	95% CI	Within group p value*	n	Baseline mean	2-year mean	Mean difference	95% CIs	Within group p value*	Between group p value†
Pain VAS‡	36	47.43	8.92	-38.51	-29.89 to -47.14	<0.0010	48	42.74	33.48	-9.26	0.61 to -19.13	0.0652	<0.0001
PCS‡	29	41.31	50.06	8.75	11.56 to 5.94	<0.0001	41	40.00	43.52	3.53	6.59 to 0.47	0.0248	0.0002
MCS§	29	51.02	54.51	3.49	7.77 to -0.79	0.0511	41	51.06	53.24	2.18	4.74 to -0.39	0.0879	0.8108
ASES‡	32	54.72	90.25	35.53	43.43 to 27.63	<0.0001	41	61.7317	72.37	10.63	16.86 to 4.41	0.0013	<0.0001
WORC‡	37	1112.16	285.95	-826.20	-678.00 to -974.40	<0.0001	46	1006.52	711.22	-295.30	-121.80 to -468.80	0.0013	<0.0001
SAT§	34	79.68	89.50	9.82	19.35 to 0.29	0.0034	38	72.61	80.55	7.95	16.32 to -0.43	0.0205	0.0981
SANE§	36	29.81	82.89	53.08	63.05 to 43.11	<0.0001	48	36.03	57.92	21.89	30.89 to 12.88	<0.0001	<0.0001

*p values for mean difference from baseline to 2-year mean.

†p values for difference between 2-year operative mean and 2-year non-operative mean.

‡p values calculated using t-tests.

§p values calculated using Wilcoxon signed-rank tests.

ASES, American Shoulder and Elbow Surgeons; MCS, mental component summary; PCS, physical component summary; SANE, Single Assessment Numeric Evaluation; SAT, Scapular Assistance Test; VAS, Visual Analogue Scale; WORC, Western Ontario Rotator Cuff Index.

Table 3 Labral tear demographics, clinical features and baseline outcome scores

	Operative (n=28)			Non-operative (n=24)			p Value
	Mean (n)	SD (%)	95% CI	Mean (n)	SD (%)	95% CI	
Age*	57.89	9.27	54.30 to 61.49	59.33	11.74	54.38 to 64.29	0.6232
Gender†							0.3303
Male	21	75.00		15	53.57		
Female	7	25.00		9	46.43		
Body mass index*	27.24	5.46	25.12 to 29.36	27.80	6.32	25.13 to 30.47	0.7310
Cause of injury†							0.6250
Known	21	75.00		18	64.29		
Unknown	7	25.00		6	35.71		
Side of injury†							0.5479
Right	14	50.00		14	58.33		
Left	14	50.00		10	41.67		
Symptom duration†							0.6856
<1 year	9	32.14		9	37.50		
>1 year	19	67.86		15	62.50		
Diabetes‡							0.1103
Yes	1	3.57		4	14.29		
No	27	96.43		20	85.71		
Smoking‡							0.8408
Yes	3	10.71		3	10.71		
No	25	89.29		25	89.29		
Pain VAS*	57.36	27.71	45.92 to 68.80	40.58	27.29	29.06 to 52.11	0.0381
Baseline PCS*	36.83	7.89	33.03 to 40.64	37.90	10.45	32.70 to 43.10	0.7271
Baseline MCS‡	48.57	11.85	42.86 to 54.29	51.38	10.01	46.41 to 56.36	0.5136
Baseline ASES*	51.27	22.33	42.25 to 60.29	63.90	20.37	54.63 to 73.18	0.0510
Baseline WORC*	1209.60	444.80	1033.70 to 1385.60	1042.90	476.80	831.50 to 1254.30	0.2126
SAT‡	79.45	26.12	67.87 to 91.04	62.81	38.03	45.50 to 80.12	0.2431
SANE‡	27.94	22.10	19.01 to 36.87	39.50	24.71	29.07 to 49.93	0.0754
SAL*	11.21	8.60	8.61 to 13.81	9.76	4.51	7.45 to 12.08	0.3925

*p values calculated using t-tests.

†p values calculated using χ^2 tests.

‡p values calculated using Wilcoxon signed-rank tests.

Continuous data presented as mean, SD. Categorical data presented as n (%).

ASES, American Shoulder and Elbow Surgeons; MCS, mental component summary; PCS, physical component summary; SAL, Shoulder Activity Level; SANE, Single Assessment Numeric Evaluation; SAT, Scapular Assistance Test; VAS, Visual Analogue Scale; WORC, Western Ontario Rotator Cuff Index.

there was evidence to support the efficacy of surgical intervention as these patients electing operative treatment displayed significant improvements from baseline to 2-year follow-up across all outcome measures. While the non-operative group also reported significant improvements in outcome measures, the operative group exhibited larger MDs from baseline to

follow-up, as well as higher scores associated with better outcomes. Additionally, the reported raw 2-year follow-up scores were significantly higher/improved when compared with the non-operative group except for MCS and SAT.

Although numerous studies explicitly examine the functional outcomes following labrum repair,^{5 16 22-24}

Table 4 Labral tear mean differences of outcome measures from baseline to 2-year follow-up within and between operative and non-operative groups

	Operative (n=28)					Non-operative (n=24)							
	n	Baseline mean	2-year mean	Mean difference	95% CI	Within group p value*	n	Baseline mean	2-year mean	Mean difference	95% CIs	Within group p value*	Between group p value†
Pain VAS‡	18	50.81	7.72	-43.08	-31.71 to -54.45	<0.0001	20	41.10	33.10	-8.00	4.89 to -20.89	0.2094	0.0014
PCS‡	15	38.30	49.10	10.81	14.16 to 7.45	<0.0001	17	37.31	44.20	6.89	12.37 to 1.42	0.0167	0.0726
MCS§	15	50.82	54.01	3.19	10.02 to -3.65	0.1876	17	50.69	52.80	2.11	5.97 to -1.75	0.4586	0.7659
ASES‡	18	57.78	93.17	35.39	45.24 to 25.54	<0.0001	16	61.56	68.94	7.38	14.21 to 0.54	0.0363	0.0002
WORC‡	19	1139.74	248.00	-891.74	-701.00 to -1082.50	<0.0001	17	1003.76	726.00	-277.76	-21.81 to -533.70	0.0352	0.0027
SAT§	16	80.03	91.44	11.41	23.47 to -0.66	0.0305	17	66.68	73.94	7.27	20.70 to -6.17	0.2490	0.0670
SANE§§	18	28.58	83.67	55.08	64.43 to 45.73	<0.0001	20	39.75	55.30	15.55	27.69 to 3.41	0.0319	0.0012

*p values for mean difference from baseline to 2-year mean.

†p values for difference between 2-year operative mean and 2-year non-operative mean.

‡p values calculated using t-tests.

§p values calculated using Wilcoxon signed-rank tests.

§§p values calculated using Wilcoxon signed-rank tests. ASES, American Shoulder and Elbow Surgeons; MCS, mental component summary; PCS, physical component summary; SANE, Single Assessment Numeric Evaluation; SAT, Scapular Assistance Test; VAS, Visual Analogue Scale; WORC, Western Ontario Rotator Cuff Index.

Table 5 Labral degeneration demographics, clinical features and baseline outcome scores

	Operative (n=27)			Non-operative (n=44)			p Value
	Mean/n	SD/%	95% CI	Mean/n	SD/%	95% CI	
Age*	59.15	9.07	55.56 to 62.73	65.14	8.48	62.56 to 67.72	0.0064
Gender†							0.7952
Male	18	66.67		28	63.64		
Female	9	33.33		16	36.36		
Body mass index*	30.07	4.84	28.15 to 31.99	28.94	5.50	27.26 to 30.61	0.3815
Cause of injury†							0.4098
Known	18	66.67		25	56.82		
Unknown	9	33.33		19	43.18		
Side of injury†							0.5126
Right	17	62.96		31	70.45		
Left	10	37.04		13	29.55		
Symptom duration†							0.6492
<1 year	15	55.56		22	50		
>1 year	12	44.44		22	50		
Diabetes†							0.6779
Yes	6	22.22		8	18.18		
No	21	77.78		36	81.82		
Smoking†							0.7235
Yes	1	3.70		1	2.27		
No	26	96.30		43	97.73		
Pain VAS*	51.48	25.02	41.58 to 61.38	43.70	27.00	35.29 to 52.11	0.2338
Baseline PCS*	41.38	9.70	36.70 to 46.05	41.36	7.23	38.61 to 44.11	0.9937
Baseline MCS‡	48.43	10.91	43.17 to 53.69	51.52	10.53	47.51 to 55.53	0.4231
Baseline ASES*	49.96	20.61	41.45 to 58.47	58.97	20.89	52.11 to 65.84	0.0973
Baseline WORC*	1158.00	473.10	966.90 to 1349.10	1072.10	446.80	931.10 to 1213.10	0.4562
SAT‡	78.98	26.58	68.01 to 89.95	74.81	29.32	64.58 to 85.04	0.4518
SANE‡	33.83	25.86	23.60 to 44.06	36.07	21.96	29.14 to 43.01	0.5185
SAL*	10.69	3.86	8.63 to 12.74	11.76	5.01	9.48 to 14.04	0.4816

*p values calculated using t-tests.

†p values calculated using χ^2 tests.

‡p values calculated using Wilcoxon signed-rank tests.

Continuous data presented as mean, SD. Categorical data presented as n (%).

ASES, American Shoulder and Elbow Surgeons; MCS, mental component summary; PCS, physical component summary; SAL, Shoulder Activity Level; SANE, Single Assessment Numeric Evaluation; SAT, Scapular Assistance Test; VAS, Visual Analogue Scale; WORC, Western Ontario Rotator Cuff Index.

²⁷ few studies look at the relationship between labral pathology and quality of life. The HRQoL outcomes for participants in the present study support previous research that has been conducted with non-operative patients. While the present study used the VR-12 as the HRQoL measure, studies that have used different

HRQoL measures have reported similar results. Edwards *et al*²⁹ found in non-operative patients that the SF-36, ASES and VAS scores all improved at final follow-up.

VR-12 (MCS and PCS) scores were not significantly different at baseline between operative and non-

Table 6 Labral degeneration mean differences of outcome measures from baseline to 2-year follow-up within and between operative and non-operative groups

	Labral degeneration (n=71)												
	Operative (n=27)					Non-operative (n=44)							
	n	Baseline mean	2-year mean	Mean difference	95% CI	Within group p value*	n	Baseline mean	2-year mean	Mean difference	95% CIs	Within group p value*	Between group p value†
Pain VAS‡	18	44.06	10.11	-33.94	-20.13 to -47.76	<0.0001	28	43.91	33.75	-10.16	4.73 to -25.05	0.1729	0.0027
PCS‡	14	44.54	51.09	6.55	11.30 to 1.79	0.0108	24	41.90	43.04	1.14	4.68 to -2.39	0.5096	0.0012
MCS§	14	51.24	55.06	3.82	9.81 to -2.18	0.2349	24	51.33	53.55	2.22	5.90 to -1.46	0.1394	0.5065
ASES‡	14	50.79	86.50	35.71	50.22 to 21.21	0.0001	25	61.84	74.56	12.72	22.25 to 3.19	0.0110	0.0523
WORC‡	18	1083.06	326.00	-757.06	-512.10 to -1002.10	<0.0001	29	1008.14	702.55	-305.59	-63.00 to -548.20	0.0154	0.0102
SAT§	18	79.36	87.78	8.42	24.08 to -7.24	0.0798	21	77.40	85.90	8.50	20.14 to -3.14	0.0650	0.4425
SANE§	18	31.03	82.11	51.08	69.86 to 32.31	0.0002	28	33.38	59.79	26.41	39.56 to 13.26	0.0001	0.0186

*p values for mean difference from baseline to 2-year mean.

†p values for difference between 2-year operative mean and 2-year non-operative mean.

‡p values calculated using Wilcoxon signed-rank tests.

§p values calculated using Wilcoxon signed-rank tests.

ASES, American Shoulder and Elbow Surgeons; MCS, mental component summary; PCS, physical component summary; SANE, Single Assessment Numeric Evaluation; SAT, Scapular Assistance Test; VAS, Visual Analogue Scale; WORC, Western Ontario Rotator Cuff Index.

operative groups. Despite observing an increased MCS score in both operative and non-operative groups, there was no statistical difference between 2-year MCS scores of either treatment group. Throughout this study, MCS consistently did not display significant improvements over time. This phenomenon is not fully understood, but can possibly be attributed to the nature of a repaired labrum. Previous research involving overhead athletes receiving non-operative treatments of SLAP tears found return to sport difficult to achieve, despite improvement in pain scores²⁹: 'Although 71% of patients returned to the same or better levels of participation, only 10 of the 15 overhead athletes (67%) returned to their overhead sport at the same or higher level'.²⁹ Additionally, Cohen *et al*²¹ found that only 14 of 29 patients (48%) in their series could return to pre-participation levels after SLAP repair, and only 71% of patients were satisfied. This may suggest that patients in our population experienced less pain and higher PCS scores, but did not report increased MCS scores due to disappointing results when returning to sport or work.

Stratifying by labral pathology allowed a deeper look into the relationship between treatment and HRQoL. Among patients with labral tears, the two groups were fairly homogenous except for the operative group presenting significantly worse baseline scores for pain VAS and ASES. As previously mentioned, this difference is likely the result of physician guidelines and recommendations for patients presenting with higher levels of pain. In this population, only the operative group reported a significant improvement in pain VAS scores. Both groups displayed significant improvement in PCS scores from baseline to 2-year follow-up, with operative patients reporting a nearly statistically greater improvement in PCS scores. The efficacy of operative treatment of labral tears has been shown again, as operative patients displayed larger improvement in pain VAS, ASES, WORC and SANE scores when compared with non-operative patients.

Patients presenting with labral degenerations reported improvements in outcome scores. However, it is important to note non-significant improvements in pain VAS and PCS scores in patients receiving non-operative treatment. Operative patients reported significant improvements in these scores, further supporting surgical treatment of labral pathologies. While it may seem intuitive for a physician to treat a less severe lesion with a conservative approach, this stratified non-operative group displayed the smallest MD in scores from baseline to 2-year follow-up.

Study limitations include single-centre patient information, as well as smaller sample sizes after stratification. Additionally, the patient population was derived from patients presenting with rotator cuff pathology. Despite controlling for this variable, it may prove useful to investigate the effects of treatment on patients presenting specifically with labral pathology.

Future research may include additional long-term trials evaluating HRQoL in patients choosing operative and non-operative treatment modalities.

The present study provides important information on expected outcomes in patients who receive operative or non-operative treatment for labral pathologies, and may help guide decisions on when to consider intervention types based on these criteria. Operative groups in this cohort consistently exhibited larger MDs for all outcome measures, and higher scores associated with better outcomes. This study demonstrates that patients who received operative treatment of labral pathology (either a tear or degeneration) significantly improved HRQoL and outcome scores after 2 years when compared with non-operative treatment.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Wilk KE, Reinold MM, Dugas JR, *et al*. Current concepts in the recognition and treatment of superior labral (SLAP) lesions. *J Orthop Sports Phys Ther* 2005;35:273–91.
2. Perry J. Anatomy and biomechanics of the shoulder in throwing, swimming, gymnastics, and tennis. *Clin Sports Med* 1983;2:247–70.
3. Robinson G, Ho Y, Finlay K, *et al*. Normal anatomy and common labral lesions at MR arthrography of the shoulder. *Clin Radiol* 2006;61:805–21.
4. Vander Kraats R, Doss A. Glenoid labral tear follow up case series on ultrasound guided autologous platelet rich plasma in conjunction with a progressive rehabilitation program. *F1000Res* 2012;1:68.
5. Funk L, Snow M. SLAP tears of the glenoid labrum in contact athletes. *Clin J Sport Med* 2007;17:1–4.
6. Handelberg F, Willems S, Shahabpour M, *et al*. SLAP lesions: a retrospective multicenter study. *Arthroscopy* 1998;14:856–62.
7. U.S Department of Labor. Bureau of Labor Statistics. Nonfatal occupational injuries and illnesses requiring days away from work. 2014. <http://www.bls.gov/news.release/pdf/osh2.pdf> (accessed 16 Dec 2015).
8. Johnson MP, Crossley KL, O'Neil ME, *et al*. Estimates of direct health care expenditures among individuals with shoulder dysfunction in the United States. APTA combined sections meeting abstract no. PL8. *J Orthop Sports Phys Ther* 2005;35:A4.
9. Zhang AL, Kreulen C, Ngo SS, *et al*. Demographic trends in arthroscopic SLAP repair in the United States. *Am J Sports Med* 2012;40:1144–7.
10. Mohana-Borges AV, Chung CB, Resnick D. Superior labral anteroposterior tear: classification and diagnosis on MRI and MR arthrography. *AJR Am J Roentgenol* 2003;181:1449–62.
11. Bencardino JT, Beltran J, Rosenberg ZS, *et al*. Superior labrum anterior-posterior lesions: diagnosis with MR arthrography of the shoulder. *Radiology* 2000;214:267–71.
12. Dinauer PA, Flemming DJ, Murphy KP, *et al*. Diagnosis of superior labral lesions: comparison of noncontrast MRI with indirect MR arthrography in unexercised shoulders. *Skeletal Radiol* 2007;36:195–202.
13. Herold T, Hente R, Zorger N, *et al*. Indirect MR-arthrography of the shoulder-value in the detection of SLAP-lesions. *Rofo* 2003;175:1508–14.
14. Waldt S, Burkart A, Lange P, *et al*. Diagnostic performance of MR arthrography in the assessment of superior labral anteroposterior lesions of the shoulder. *AJR Am J Roentgenol* 2004;182:1271–8.
15. Clavert P. Glenoid labrum pathology. *Orthop Traumatol Surg Res* 2015;101(1 Suppl):S19–24.

16. Snyder SJ, Banas MP, Karzel RP. An analysis of 140 injuries to the superior glenoid labrum. *J Shoulder Elbow Surg* 1995;4:243–8.
17. Maffet MW, Gartsman GM, Moseley B. Superior labrum-biceps tendon complex lesions of the shoulder. *Am J Sports Med* 1995;23:93–8.
18. Solomon DJ, Levine WN. SLAP tears: pearls and pitfalls in diagnosis and management. *Sports Med Update* 2011;2–6.
19. Rowbotham EL, Grainger AJ. Superior labrum anterior to posterior lesions and the superior labrum. *Semin Musculoskelet Radiol* 2015;19:269–76.
20. Vogel LA, Moen TC, Macaulay AA, et al. Superior labrum anterior-to-posterior repair incidence: a longitudinal investigation of community and academic databases. *J Shoulder Elbow Surg* 2014;23:e119–26.
21. Cohen DB, Coleman S, Drakos MC, et al. Outcomes of isolated type II SLAP lesions treated with arthroscopic fixation using a bioabsorbable tack. *Arthroscopy* 2006;22:136–42.
22. Field LD, Savoie FH. Arthroscopic suture repair of superior labral detachment lesions of the shoulder. *Am J Sports Med* 1993;21:783–90.
23. Nam EK, Snyder SJ. The diagnosis and treatment of superior labrum, anterior and posterior (SLAP) lesions. *Am J Sports Med* 2003;31:798–810.
24. O'Brien SJ, Allen AA, Coleman SH, et al. The trans-rotator cuff approach to SLAP lesions: technical aspects for repair and a clinical follow-up of 31 patients at a minimum of 2 years. *Arthroscopy* 2002;18:372–7.
25. Pagnani MJ, Speer KP, Altchek DW, et al. Arthroscopic fixation of superior labral lesions using a biodegradable implant: a preliminary report. *Arthroscopy* 1995;11:194–8.
26. Resch H, Golser K, Thoeni H, et al. Arthroscopic repair of superior glenoid labral detachment (the SLAP lesion). *J Shoulder Elbow Surg* 1993;2:147–55.
27. Rhee YG, Lee DH, Lim CT. Unstable isolated SLAP lesion: clinical presentation and outcome of arthroscopic fixation. *Arthroscopy* 2005;21:1099.e1–1099.e7.
28. Yoneda M, Hirooka A, Saito S, et al. Arthroscopic stapling for detached superior glenoid labrum. *J Bone Joint Surg Br* 1991;73:746–50.
29. Edwards SL, Lee JA, Bell JE, et al. Nonoperative treatment of superior labrum anterior posterior tears: improvements in pain, function, and quality of life. *Am J Sports Med* 2010;38:1456–61.
30. Cho CH, Song KS, Hwang I, et al. Does rotator cuff repair improve psychologic status and quality of life in patients with rotator cuff tear? *Clin Orthop Relat Res* 2015;473:3494–500.
31. Ayers DC, Franklin PD, Ring DC. The role of emotional health in functional outcomes after orthopaedic surgery: extending the biopsychosocial model to orthopaedics: AOA critical issues. *J Bone Joint Surg Am* 2013;95:e165–1–7.
32. Chung SW, Park JS, Kim SH, et al. Quality of life after arthroscopic rotator cuff repair: evaluation using SF-36 and an analysis of affecting clinical factors. *Am J Sports Med* 2012;40:631–9.
33. Mohtadi NG, Hollinshead RM, Sasyniuk TM, et al. A randomized clinical trial comparing open to arthroscopic acromioplasty with mini-open rotator cuff repair for full-thickness rotator cuff tears: disease-specific quality of life outcome at an average 2-year follow-up. *Am J Sports Med* 2008;36:1043–51.
34. Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison of six methods. *Pain* 1986;27:117–26.
35. Hays RD, Sherbourne CD, Mazel RM. The RAND 36-Item Health Survey 1.0. *Health Econ* 1993;2:217–27.
36. Kazis LE, Miller DR, Skinner KM, et al. Applications of methodologies of the Veterans Health Study in the VA Healthcare system: conclusions and summary. *J Ambul Care Manage* 2006;29:182–8.
37. Kazis LE, Miller DR, Skinner KM, et al. Patient-reported measures of health: The Veterans Health Study. *J Ambul Care Manage* 2004;27:70–83.
38. Kazis LE, Selim A, Rogers W, et al. Dissemination of methods and results from the veterans health study: final comments and implications for future monitoring strategies within and outside the veterans healthcare system. *J Ambul Care Manage* 2006;29:310–9.
39. Kirkley A, Griffin S, Dainty K. Scoring systems for the functional assessment of the shoulder. *Arthroscopy* 2003;19:1109–20.
40. Lopes AD, Ciconelli RM, Carrera EF, et al. Validity and reliability of the Western Ontario Rotator Cuff Index (WORC) for use in Brazil. *Clin J Sport Med* 2008;18:266–72.
41. Huang H, Grant JA, Miller BS, et al. A systematic review of the psychometric properties of patient-reported outcome instruments for use in patients with rotator cuff disease. *Am J Sports Med* 2015;43:2572–82.
42. Richards RR, An KN, Bigliani LU, et al. A standardized method for the assessment of shoulder function. *J Shoulder Elbow Surg* 1994;3:347–52.
43. Angst F, Goldhahn J, Drerup S, et al. Responsiveness of six outcome assessment instruments in total shoulder arthroplasty. *Arthritis Rheum* 2008;59:391–8.
44. Beaton D, Richards RR. Assessing the reliability and responsiveness of 5 shoulder questionnaires. *J Shoulder Elbow Surg* 1998;7:565–72.
45. Beaton DE, Richards RR. Measuring function of the shoulder. A cross-sectional comparison of five questionnaires. *J Bone Joint Surg Am* 1996;78:882–90.
46. Bot SD, Terwee CB, van der Windt DA, et al. Clinimetric evaluation of shoulder disability questionnaires: a systematic review of the literature. *Ann Rheum Dis* 2004;63:335–41.
47. Kocher MS, Horan MP, Briggs KK, et al. Reliability, validity, and responsiveness of the American Shoulder and Elbow Surgeons subjective shoulder scale in patients with shoulder instability, rotator cuff disease, and glenohumeral arthritis. *J Bone Joint Surg Am* 2005;87:2006–11.
48. Michener LA, McClure PW, Sennett BJ. American Shoulder and Elbow Surgeons standardized shoulder assessment form, patient self-report section: reliability, validity, and responsiveness. *J Shoulder Elbow Surg* 2002;11:587–94.
49. Borstad JD, Ludewig PM. The effect of long versus short pectoralis minor resting length on scapular kinematics in healthy individuals. *J Orthop Sports Phys Ther* 2005;35:227–38.
50. Kibler WB. The role of the scapula in athletic shoulder function. *Am J Sports Med* 1998;26:325–37.
51. Farrar JT, Young JP, LaMoreaux L, et al. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain* 2001;94:149–58.
52. Rabin A, Irrgang JJ, Fitzgerald GK, et al. The intertester reliability of the Scapular Assistance Test. *J Orthop Sports Phys Ther* 2006;36:653–60.
53. Brophy RH, Beauvais RL, Jones EC, et al. Measurement of shoulder activity level. *Clin Orthop Relat Res* 2005;439:101–8.
54. Marx RG, Stump TJ, Jones EC, et al. Development and evaluation of an activity rating scale for disorders of the knee. *Am J Sports Med* 2001;29:213–8.
55. Williams GN, Gangel TJ, Arciero RA, et al. Comparison of the single assessment numeric evaluation method and two shoulder rating scales. outcomes measures after shoulder surgery. *Am J Sports Med* 1999;27:214–21.