

Natural Selection Mediated Association of the Duffy (*FY*) Gene Polymorphisms with *Plasmodium vivax* Malaria in India

Anita Chittoria¹, Sujata Mohanty², Yogesh Kumar Jaiswal³, Aparup Das^{1*}

1 Division of Genomics and Bioinformatics, National Institute of Malaria Research, New Delhi, India, **2** Department of Biotechnology, Jaypee Institute of Information Technology, Noida, Uttar Pradesh, India, **3** School of Studies in Biotechnology, Jiwaji University, Gwalior, Madhya Pradesh, India

Abstract

The Duffy (Fy) antigens act as receptors for chemokines as well as for *Plasmodium vivax* to invade human RBCs. A recent study has correlated the occurrence of the *FY*A* allele of Duffy gene with decreased susceptibility to vivax malaria, but no epidemiological correlation between the distribution of *FY*A* allele and incidences of vivax malaria has been established so far. Furthermore, if such correlations exist, whether natural selection has mediated the association, is an important question. Since India is highly endemic to *P. vivax* malaria with variable eco-climatic and varying vivax malaria epidemiology across different regions, such a question could well be answered in Indians. For this, we have genotyped the *FY* gene at the –33rd and the 125th nucleotide positions in 250 Indians sampled from six different zonal plus one tribal population covering the whole of India and studied possible correlations with eco-climatic and vivax malaria incidences. No *FY*O* allele was found, however, both the *FY*A* and *FY*B* alleles forming *FY*A/FY*A*, *FY*A/FY*B* and *FY*B/FY*B* genotypes were widely distributed among Indians. Five out of seven population samples significantly deviated from the Hardy-Weinberg equilibrium expectation, and two alleles (*FY*A* and *FY*B*) and the homozygote genotype, *FY*B/FY*B* were clinally distributed over the population coordinates. Furthermore, vivax malaria incidences over the past five years were significantly negatively and positively associated with the frequencies of the *FY*A* and *FY*B* alleles, respectively. The Northern Indians were highly differentiated from the other zonal population samples at the *FY* gene, as evidenced from the reconstructed Neighbor-Joining phylogenetic tree. The results specify the role of natural selection in the distribution of *FY* gene polymorphism in India. Furthermore, the hypotheses on the part of the *FY*A* allele in conferring protection to vivax malaria could be validated following population genetic studies in a vivax malaria epidemiological setting, such as India.

Citation: Chittoria A, Mohanty S, Jaiswal YK, Das A (2012) Natural Selection Mediated Association of the Duffy (*FY*) Gene Polymorphisms with *Plasmodium vivax* Malaria in India. PLoS ONE 7(9): e45219. doi:10.1371/journal.pone.0045219

Editor: Ana Paula Arez, Instituto de Higiene e Medicina Tropical, Portugal

Received: June 1, 2012; **Accepted:** August 14, 2012; **Published:** September 21, 2012

Copyright: © 2012 Chittoria et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: AC is recipient of Junior and Senior Research Fellowships from the Council of Scientific and Industrial Research (CSIR), New Delhi, India. AD thanks the Indian Council of Medical Research (ICMR) for intramural funding support. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: aparup@mrindia.org

Introduction

The human Duffy blood-group (Fy) antigens are major determinants for the invasion of malaria parasite *Plasmodium vivax* in RBCs and are encoded by Duffy (*FY*) gene which is characterized by three different alleles at the genetic level; *FY*A*, *FY*B* and *FY*O* [1–3]. While the *FY*A* and *FY*B* alleles could be identified by a missense nucleotide mutation (G125A) and present the common Fy(a+b–), Fy(a–b+) and Fy(a+b+) serological phenotypes, the *FY*O* allele (absence of the Fy antigen) gives rise to Fy(a–b–) phenotype due to T-33C mutation at the promoter region of the *FY* gene [4]. Since the T-33C mutation stops the expression of the *FY* gene and thus blocks the red blood cell invasion of the human malaria parasite *Plasmodium vivax*; the *FY*O* allele is at or near to fixation in most of the sub-Saharan African populations, with very rare occurrence in populations outside Africa - indicating the role of positive natural selection for the *FY*O* allele in humans [5,6]. This argument is in agreement with the observation of resistance to *P. vivax* infection in humans homozygous to the *FY*O* allele [7] with some exceptions [8]. Very

recently, the *FY*A* allele has been shown to significantly diminish binding of *P. vivax* Duffy binding protein in comparison to *FY*B*, hence it is argued that individuals possessing the *FY*A* allele are in reduced risk to *P. vivax* malaria [9]. However, no correlation between the allele frequency and vivax malaria epidemiology has yet been established in any endemic country setting. Now-a-days, Duffy blood group is majorly determined by DNA sequencing of the *FY* gene [10], which enables in identification of any of the three different genotypes (*FY*A/FY*A*, *FY*A/FY*B* and *FY*B/FY*B*) in a single individual. Population data from multiple individuals further enables in determination of the frequency of three different alleles (*FY*A*, *FY*B* and *FY*O*) which could be utilized in evolutionary inference of the *FY* gene [5,6]. Since eco-climatic factors are highly variable in India (large longitudinal and latitudinal transects) with complex malaria epidemiology due to both *P. vivax* and *P. falciparum* infections [11] and Indians are genetically heterogeneous with complex population history [12,13,14]; we were interested to determine (i) the variable distribution of *FY* gene polymorphisms and genetic differentiations among populations living in different eco-climatic conditions, and

(ii) possible correlation between *FY* gene polymorphisms and vivax malaria epidemiology by population genetic analyses in India. Although previous studies have indicated the distribution of different serological phenotypes of the Fy antigen in different local populations in India [15,16,17,18], neither any survey on the distribution of different *FY* alleles representing the whole of India has been conducted, nor any correlation of *FY* polymorphisms to eco-climatic and/or vivax malaria epidemiology has been established. In addition, the results on the role of the *FY*A* allele to reduced risk of malaria in comparison to the *FY*B* allele [9], has not been tested/reconfirmed in any field vivax malaria epidemiological settings so far. We propose that no better place other than India could serve as the model field to understand evolution of such host-parasite interaction [19], with vivax malaria as an example.

Materials and Methods

In order to catch most of the polymorphisms of the *FY* gene and to meaningfully conduct population genetic survey, we have divided the whole of India into six distinct zones; *viz.*, North India (NI), Central India (CI), West India (WI), South India (SI), East India (EI) and North-East India (NEI). Further, since many tribes live in different Indian states, we have included samples from three Indian tribes; Juango, Bonda and Kutia Kandha inhabiting in Odisha state. The details of the Indian states included under each zone and the number of individuals sampled from each state are provided in the Table S1. In total, 250 healthy and unrelated Indians (including 28 tribal individuals) representing almost all the political states of India were sampled. For positive control, blood samples from two Africans (Cameroonians colleagues of the authors working in India) were also analyzed. From each individual, two milliliter of intravenous blood was collected in heparinized vacutainers and stored at -20°C until DNA extraction was performed.

We have selected the 1096 bp region of the *FY* gene located in the human chromosome 1 (Figure S1), comprising the two main polymorphic sites which govern the *FY*O* allele (T-33C) in the promoter region and the *FY*A/FY*B* alleles (G125A) in the exon-2. For ease in PCR amplification and automated DNA sequencing, we have divided the 1096 bp region into two overlapping sequences (Figure S1) employing two primer pairs (Table S2). Genomic DNA from blood sample of each individual was extracted and PCR amplification reactions were carried out following standard protocols in a final volume of 20 μl . Five microliters of amplified PCR product of each fragment were run on 2% agarose gel to check the quality of amplification. If a single band without any primer-dimer was present, the amplicons were considered for DNA sequencing. For this, the PCR amplicons were purified with Exonuclease-I and Shrimp Alkaline Phosphatase (Exo-Sap, Fermentas, Life Sciences) following standard protocol. Sequencing reactions were followed with Big Dye Terminator (BDT) ready reaction mix as per the Applied Biosciences (ABI) protocol and DNA sequencing was performed in an ABI 3730XL DNA Analyzer (in-house facility of NIMR). Each DNA fragment was sequenced in both the forward and reverse directions ($2\times$ coverage) and the resulted DNA sequences were assembled and edited using the SeqMan and EditSeq computer programs (DNASTAR, Madison, WI, USA) for each individual. The DNA sequence chromatograms were carefully visually inspected for the occurrence of single/double peaks in the sequence chromatogram at both the T-33C and G125A nucleotide positions. Altogether, 252 DNA sequences (250 Indian, two African) were aligned using the MegAlign computer program

(DNASTAR) following Clustal-W algorithm to identify Single Nucleotide Polymorphisms (SNPs) at these two nucleotide positions (T-33C and G125A). For instance, detection of the C nucleotide in a single peak at the -33^{rd} position indicates the presence of *FY*O* homozygote, both C and T in double peak as heterozygote genotype and the T nucleotide in a single peak indicates the absence of *FY*O* genotype. Similarly, detection of the G and A nucleotides (in single peaks) at the 125^{th} position indicate homozygosity for *FY*A* and *FY*B*, respectively, whereas presence of double peaks of both G and A nucleotides at this position categorize the individual as a heterozygote (*FY*A/FY*B*). Following these approaches, each of the 250 Indians and the two Africans were independently genotyped and frequencies of each allele for the seven Indian population samples were calculated based on the genotype data for population genetic analyses (see below).

The genotype data (*FY*A/FY*A*, *FY*A/FY*B* and *FY*B/FY*B*) from each of the seven zonal population samples were analyzed by Chi-square Test to determine if the population samples deviate from the expectation under the Hardy-Weinberg (H-W) equilibrium. Genetic differentiation between pairs of population samples were determined by two test statistics; (i) Fisher's Exact Test of differentiation using the Arelquine computer program (<http://cmpg.unibe.ch/software/arlquin3>), and (ii) Nei's genetic distance between a pair of population sample (as measured by *D*) [20] using the POPGENE computer program (<http://www.ualberta.ca/~fyeh/>). In order to visualize genetic interrelationships among different Indian population samples, the *D* matrix was used to obtain a Neighbour-Joining (NJ) population phylogenetic tree using the Phylip computer program (<http://evolution.gs.washington.edu/phylip>) and visualized with the online computer program Drawgram (http://www.phylogeny.fr/version2_cgi/one_task.cgi?task_type=drawgram) following a similar approach as described by Das and co-workers [21] and Gupta and co-workers [22]. In order to discern if ecological parameters (that change with population coordinates) influence the distribution patterns of the two *FY* alleles, frequencies of the two different *FY* alleles were correlated with both latitude and longitude of the population samples (determined from the centrally placed city of each zone, see Supplementary Table S1) independently by calculating the Pearson's correlation coefficient (*r*). Furthermore, since the *FY*A* allele has been shown to provide protection against vivax malaria in comparison to the *FY*B* allele [9], and epidemiology of *P. vivax* malaria varies in India, we have correlated the extent of vivax malaria for each Indian state (<http://www.nvbdc.gov.in/>) with the frequencies of both the *FY*A* and *FY*B* alleles and with three *FY* genotypes by calculating the *r* values independently. Data in percent frequencies were converted through Arcsine Transformation before employing in any of the above statistical analyses.

Ethics Statement

The Institutional ethics committee of the National Institute of Malaria Research (NIMR), New Delhi, India has approved the study and written informed consent from each blood donor has been obtained.

Results and Discussion

In none of the 250 Indians, the C nucleotide at the -33^{rd} promoter region of the *FY* gene could be found (no *FY*O* allele), indicating no Duffy-negative individual in the presently analyzed samples from India. The results are not quite surprising, as some earlier studies in India employing serological techniques had identified only the tribal Indians dwelling in Andaman and

Nicobar Islands and Madhya Pradesh as Duffy-negative [16,17]. On the other hand, studies following serological techniques, no Duffy-negative individual could be found in tribes inhabiting in Rajasthan and Nagaland states [15,18]. Non-occurrence of Duffy negative Indians, especially the three tribes presently analyzed (Juango, Bonda, Kutia Kandha inhabiting in Odisha state) indicates that the Duffy-negative Indians might be restricted to particular tribes and locations in India. More number of tribal individuals populating in different Indian states would be needed to establish the prevalence of the *FY*O* allele in Indians in general, and in indigenous Indians in particular. However, both the African samples employed as positive control in the present study were found to be Duffy-negative [as determined by the presence of C nucleotide instead of a T nucleotides at the -33^{rd} promoter region of the *FY* gene]; one individual with single peak of the C nucleotide (homozygous) and the other with two peaks of both T and C nucleotides (heterozygous), indicating the flawlessness of DNA sequencing in unraveling the Duffy genotypes in the present study.

For the 125th nucleotide position, both the A (*FY*B*) and the G (*FY*A*) nucleotides were found to be segregated in all the six Indian zonal population samples and also in the tribes. Thus, each Indian could be characterized into one of the three genotypes at this nucleotide position; either as one of the two homozygotes (*FY*A/FY*A* and *FY*B/FY*B*) or as a heterozygote (*FY*A/FY*B*). In total, out of 250 Indians, 53 were homozygous for *FY*B* genotype; 130 were homozygous for *FY*A* and the rest 67 were heterozygotes (*FY*A/FY*B*). Similarly, both the African individuals were found to be *FY*B* homozygous. Distribution of the three different genotypes (*FY*A/FY*A*, *FY*B/FY*B* and *FY*A/FY*B*) and allele (*FY*A* and *FY*B*) frequencies in each population sample are detailed in Table 1. In general, majority of Indians (52% of the total samples) were found to be homozygous to *FY*A*, while Indians possessing either the *FY*B* homozygote or the *FY*A/FY*B* heterozygote were in 21.2% and 26.8%, respectively. Correspondingly, in each population sample, the *FY*A* homozygote was found to be highest in number than the corresponding *FY*B* homozygote and the heterozygote. Moreover, the observation of a comparatively lesser number of heterozygotes (*FY*A/FY*B*) indicates no evidence of heterozygous advantage (heterosis), except for the SI and OT population samples, where fair number of heterozygous individuals were found to be present. It is well known that heterosis is maintained by balancing selection in many natural populations for functionally relevant genes in humans [23,24]. Absence of an adequate number of heterozygous individuals thus specifies that the *FY* gene does not evolve following the balancing selection model of natural selection in India. This observation further implicates the role of some other form of natural selection (e.g. positive natural selection) or population demography on the observed pattern of distribution of different alleles (and genotypes) of the *FY* gene. To this respect, observations of statistically significant deviation from the H-W equilibrium expectation in five (NI, NEI, CI, WI and EI) of the seven samples (with less number of heterozygotes, Table 1) indicate demographic disequilibrium or positive natural selection (also see below) in majority of the Indian samples. However, the observed demographic disequilibrium (excess of homozygotes) might not have been caused due to inbreeding, as each zonal population sample in the present study is composed of unrelated individuals originating from different Indian states (Supplementary Table S1). Thus, the excess of *FY*A* homozygote that is essentially responsible for deviation from H-W expectation in the majority of zonal population samples points towards the role of positive natural selection in maintaining the *FY*A* alleles in high frequency in India (also see below). Such

an observation seems to be unique to India thus far, as in Brazilians (another vivax malaria endemic country), no significant departure from the H-W expectation at the *FY* gene could be observed [25]. Thus, the differences in the pattern of evolution of the *FY* gene between India and Brazil might be due to differential evolutionary pressures at the *FY* gene and variable host-parasite interaction mechanisms prevailing in the two vivax malaria eco-epidemiological settings.

In order to estimate genetic differentiations between different zonal populations and to perform other population genetic studies involving the *FY* gene polymorphism in India, we first calculated the frequencies of the two alleles (*FY*A* and *FY*B*) from the genotype data for each population. As expected, in all the seven samples, the frequency of the *FY*A* allele was higher than the *FY*B*, indicating high prevalence of *FY*A* allele in Indians (Table 1). This is interesting, as the *FY*A* allele (derived, conferring protection to *P. vivax* infection) [9] was found to dominate over the *FY*B* (ancestral, providing vivax malaria susceptibility) [26] allele in India. In order to measure genetic differentiation between population pairs at the *FY* gene, we utilized the allele frequency data and calculated the Nei's *D* [19] (Table S3) and Fisher's Exact Test. While the *D* values were utilized to reconstruct a population phylogenetic tree [21,22] (see below); the Fisher's Exact Test statistic yielded a total of only three statistically significant cases of pair-wise genetic differentiation, always involving the NI population sample (NI vs. CI = 0.014; NI vs. EI = 0.009 and NI vs. NEI = 0.005), indicating that the northern Indians are well differentiated from individuals inhabiting in the central, eastern and northeastern India.

Since the genotype frequency distribution data indicated towards the role of positive natural selection in evolution of the *FY* gene (see above), to strengthen our hypothesis we were interested to discern whether eco-climatic conditions of the population sample have influenced the frequency distribution of the two different alleles (*FY*A* and *FY*B*) of the *FY* gene in India. Although no statistically-significant correlation (with respect to the *r* values) could be obtained between the frequencies of the two alleles (*FY*A* and *FY*B*) and the latitude, statistically significant *r* values (in opposite direction for the two alleles) could be obtained with longitude (Table 2). In particular, the ancestral *FY*B* allele was found to be negatively correlated, and the *FY*A* was positively associated with the longitude in India (Table 2). Since eco-climatic conditions of different locations change with the latitude and longitude [12], the distribution patterns of the two *FY* alleles seem to follow strict eco-climatic trend in India as a result of local adaptation by natural selection. It is known that the *FY*A* allele have almost reached to fixation in Southeast Asia [9], and the present results indicate that its frequency decreases as we move in westward directions in India. Considering India as a part of the ancestral range of *P. vivax* [22], and the incidences of vivax malaria decreases as we move in eastward direction (Figure 1), it can be argued that the resistance to *P. vivax* infection has been well-evolved in the eastern, central and northeastern Indians as reflected by the high prevalence of the *FY*A* allele in those regions (Figure 1). This hypothesis is further substantiated by the fact that *P. vivax* populations from these areas in India have undergone significant population size reduction in the past [22]. However, all these regions are highly endemic to *P. falciparum* malaria [11] with high rates of mixed parasite infections [27]. Such complex malaria epidemiology in India [11], especially the decreased incidences of *P. vivax* infection over *P. falciparum* in the past years [19], presents two principal hypotheses to be tested; (i) outperformance of *P. falciparum* over *P. vivax* and (ii) evolving host resistance to vivax malaria. While testing of the first hypotheses requires separate

Table 1. Details of locations and numbers of samples, average vivax malaria incidences over five years period (2007–2011), Hardy Weinberg Expectations and allele and genotype frequencies of the FY gene in Indians.

| Locations and no. of samples | Central city | Latitude/Longitude | # <i>P. vivax</i> malaria (%) | Observed no. of genotypes | | | HWE (χ^2) | Allele frequency (%) | |
|------------------------------|--------------|--------------------|-------------------------------|---------------------------|-----------|-----------|------------------|----------------------|------|
| | | | | FY*B/FY*B | FY*A/FY*A | FY*A/FY*B | | FY*A | FY*B |
| North India (50) | Haridwar | 29°57'N/78°10'E | 97 | 18 | 22 | 10 | 17.85*** | 54 | 46 |
| North-East India (32) | Guwahati | 26°11'N/91°44'E | 41 | 7 | 24 | 1 | 26.12*** | 77 | 23 |
| East India (36) | Ranchi | 23°21'N/85°20'E | 49 | 5 | 22 | 9 | 4.23* | 74 | 26 |
| Odisha Tribes (28) | Bhubaneswar | 20°16'N/85°50'E | 49 | 5 | 12 | 11 | 0.73 | 63 | 37 |
| Central India (40) | Bhopal | 23°15'N/77°25'E | 46 | 6 | 24 | 10 | 5.56** | 72 | 28 |
| West India (32) | Ahmedabad | 23°02'N/77°35'E | 77 | 9 | 16 | 7 | 9.35** | 59 | 41 |
| South India (32) | Chennai | 13°05'N/80°16'E | 73 | 3 | 14 | 15 | 0.01 | 67 | 33 |

*P<0.05,
**P<0.01,
***P<0.001.

HWE = Hardy Weinberg Expectations.

Average yearly proportions of *P. vivax* malaria in India. Source: www.nvbdc.gov.in; The total number of *P. falciparum* malaria cases (as provided in the website) for each state under different zones were deducted from the total malaria cases for each year (2007–2011) to obtain the total number of *P. vivax* malaria cases. The data on *P. vivax* might include a few number of *P. malariae* infections, in the EI samples.
doi:10.1371/journal.pone.0045219.t001

experimental set ups, the results from the present study could be able to demonstrate evolving acquired immunity in Indians against *P. vivax* infection in general, and in the central, eastern and northeastern Indians, in particular. This proposition could possibly explain the observed declining trend of vivax malaria in these three zones in India [19], although other unseen factors might also be responsible. Furthermore, dominance of the FY*A homozygotes (75% in NEI, 60% in CI and 56% in EI) in these three zones (Central, Eastern and North Eastern) suggests that individuals homozygous to FY*A might be more capable of resisting to *P. vivax* malaria infection than the heterozygotes. Moreover, tight positive latitudinal clines for both the FY*A and FY*B homozygotes (Table 2) indicate the role of natural selection through local adaptation of these two homozygotes [28,29]. Whether such clines are solely adaptive for both the genotypes could not be ascertained in the present study, but it might be predicted that the cline for the FY*A/FY*A genotype over latitude might be a reflection of its increasing trends towards the northern Indian populations due to local adaptation by natural selection and the cline observed for the FY*B homozygote may be historically maintained due to its ancestral background [26].

The results of the present study provide us with opportunities to test if the high prevalence of FY*A allele that is maintained by the influence of positive natural selection (see above) has, by any means, impacted the vivax malaria epidemiology in India. For this purpose, we obtained the number of vivax malaria incidences for each year separately for the last five years (2007–2011) and also five years' combined data (2007–2011) for each Indian state [courtesy; National Vector Borne Disease Control Program, (www.nvbdc.gov.in), New Delhi, India] and collated for six zonal population samples (separately for each year and the combined data for all the five years). The five-year combined state-wise data on vivax malaria majorly fall in three different categories; high (80–100% of malaria cases due to *P. vivax* infection), medium (60–79%) and low (0–59%) (Figure 1). Accordingly, the whole of India could be divided into three distinct zones based on vivax malaria endemicity (high, medium and low) (Figure 1). Further, data on the frequency of the two alleles (FY*A and FY*B) and number of the three genotypes (FY*A/FY*A, FY*A/FY*B and FY*B/FY*B) were also assembled separately for these three zones and plotted as two different types of pie charts (for allele frequencies and for genotypes) in the respective zones (Figure 1). In addition, Pearson's

Table 2. Pearson's correlation coefficient (*r*) between the alleles and genotypes frequencies of the FY gene with vivax malaria incidences over five years period (2007–2011) and population coordinates (latitude and longitude) in India.

| Alleles and genotypes of the FY gene | | Incidences of <i>Plasmodium vivax</i> malaria in India | | | | | | Population coordinates | |
|--------------------------------------|-----------|--|--------|--------|--------|--------|------------|------------------------|-----------|
| | | 2007 | 2008 | 2009 | 2010 | 2011 | (2007-11)# | Latitude | Longitude |
| Alleles | FY*A | -0.81* | -0.83* | -0.91* | -0.95* | -0.95* | -0.92* | 0.01 | 0.67* |
| | FY*B | 0.83* | 0.86* | 0.92 | 0.96* | 0.96* | 0.94* | 0.04 | -0.65* |
| Genotypes | FY*A/FY*A | -0.09 | 0.04 | -0.02 | -0.22 | -0.2 | -0.2 | 0.68 | 0.21 |
| | FY*B/FY*B | 0.72* | 0.89** | 0.85* | 0.7* | 0.71* | 0.79* | 0.78* | -0.32 |
| | FY*A/FY*B | 0.38 | 0.2 | 0.21 | 0.35 | 0.34 | 0.31 | -0.59 | -0.41 |

*P<0.05,
**P<0.01.

Five years average incidences of *P. vivax* malaria in India.

doi:10.1371/journal.pone.0045219.t002




Figure 1. Map of India showing partitions of different zones (NI, NEI, EI, CI, WI and SI) considered for sampling of population in the present study. Three different blue colour shades represent three different vivax malaria transmission areas (high, medium and low). The frequencies of the alleles (FY^*A and FY^*B) and genotypes (FY^*A/FY^*A , FY^*B/FY^*B and FY^*A/FY^*B) were indicated as pi-charts in the three respective regions in each of these three areas.

doi:10.1371/journal.pone.0045219.g001

correlation coefficient (r) values were calculated between the exact incidences of vivax malaria and the frequencies of the two FY alleles (FY^*A and FY^*B) as well as the number of three different genotypes (Table 2). As indicated in Table 2, several cases of statistically significant r values were obtained with such genetic and epidemiological correlation analyses, involving both the yearly as well as the five-year combined vivax malaria data in India. Interestingly, statistically significant correlations for both the FY^*A and FY^*B alleles were observed over all the five years (2007–2011), and also for the combined five year data. However, the trends were in opposite directions (negative correlation for FY^*A and positive for FY^*B) (Table 2). For the genotypes, the FY^*B homozygote presented statistically significant positive correlations with vivax malaria cases in all the five years independently (and also with the five-year combined data), corroborating the fact that individuals of this particular genotype (FY^*B/FY^*B) may be highly susceptible to vivax malaria. However, no such association over the years could be observed for either the FY^*A/FY^*A homozygote or the FY^*A/FY^*B heterozygote (Table 2). However, considering the FY^*A genotype lowers the risk to vivax malaria than the FY^*B allele [9], and field epidemiological data on vivax malaria in India reveal negative association with the frequency distribution of the FY^*A alleles and positive association with the FY^*B alleles (see

above), it can be ascertained that the FY^*A allele might be playing some role in conferring protection to vivax malaria, and is maintained under positive natural selection in India. Whether such an association exists in other vivax malaria endemic countries remains to be seen, as this type of study has never been conducted in any other malaria epidemiological setting yet. On the contrary, the observed statistically significant positive association between vivax malaria and FY^*B allele frequency was mainly due to high prevalence of the FY^*B homozygote in places where vivax malaria incidences are high (e.g. NI population sample) (Table 1). To this respect, it was interesting to know if genetic relatedness among different Indian zonal samples fits well with the observed differential frequency distribution of the FY^*A and FY^*B alleles by reconstruction of a NJ population phylogenetic tree (Figure 2) from genetic distance matrix [21,22]. Interestingly, the NJ tree produced three different clades (Figure 2), exactly representing the three distinct zones based on differential incidences of vivax malaria (high, medium and low) (Figure 1). The northern Indian (NI) sample was clearly separated from the rest six population samples in the NJ tree (Figure 2), suggesting the fact that this population sample (with the highest incidences of both vivax malaria and FY^*B alleles) is in fact genetically distinct from the rest of six Indian population samples. This observation further

corroborated with (i) the observations on the statistically significant Fisher's Exact Test between the NI and three other population samples (see above) and (ii) the partitioning of whole of India based of differential incidences of vivax malaria (Figure 1). The results from a diverse array of population genetic analyses thus point towards correspondence between the genetic (*FY* gene) and epidemiological (incidences of *P. vivax* infection) aspects in Indians.

In conclusion, the results of the present population genetic study indicate the role of natural selection in the variable distribution of two different alleles of the *FY* gene (*FY*A* and *FY*B*) in Indians. The significant departure from H-W expectation in five out of seven samples justifies this contention. In addition, a clear-cut negative correlation with vivax malaria epidemiology in Indians with the *FY*A* allele frequency distribution points towards the *FY*A* homozygote genotype being under strong influence of positive natural selection. In view of the fact that the derived *FY*A* allele confers resistance to vivax malaria [9], its tight negative correlation with incidences of vivax malaria indicates evolving resistance to vivax malaria in Indians. However, the prevalence of *FY*B* allele in all zones (although in a lower frequency than the *FY*A* allele) suggests that Indians are still susceptible to *P. vivax* infection. Since it is argued that the ideal populations to test PVDBPII-based vaccine are with genetic background where combinations of *FY*A* and *FY*B* alleles exist [9], India can be the model field for trial of such a vaccine against *P. vivax* infection. Considering both the falciparum and vivax malaria are prevalent in almost all over India [11,19] and the *FY* polymorphisms have

been subjected to malarial selection shortly before known *P. falciparum* protective mutation [30,31], comparative genetic studies of genes responsible for susceptibility to both falciparum and vivax malaria in common populations could not only inform evolution of host-parasite interaction mechanisms in malaria but also help in devising novel control measures for both forms of malaria in India.

Supporting Information

Figure S1 Schematic diagram of the of 1096 bp DNA fragment of the Duffy gene.

(TIF)

Table S1 Details of the sample location, size and vivax malaria proportions for each Indian state that are used to form six different zones for the present study.

(RTF)

Table S2 Description of primer sequences and annealing temperatures for PCR amplification and sequencing of the 1096 bp DNA fragment.

(RTF)

Table S3 Pairwise-population genetic distance as calculated using Nei's genetic distance (Nei's *D*, 1972).

(RTF)

Acknowledgments

The authors are highly thankful to all blood donors and to Prof. A. P. Dash, former Director of the National Institute of Malaria Research (NIMR) for facilities and encouragements. We thank Dr. M. R. Ranjit of the Regional Medical Research Centre (RMRC), Bhubaneswar for his kind support in sampling the tribal populations of Odisha, to the Director, National Vector Borne Disease Control Programme (NVBDCP), New Delhi, for the annual malaria incidence data in the present analyses and to Dr. Ana Paula Arez and three anonymous reviewers for constructive comments on an earlier version of the manuscript. SM thanks the Vice Chancellor, Jaypee Institute of Information Technology, Noida for facilities and encouragements. This paper bears the NIMR publication screening committee approval number 011/2012.

Author Contributions

Conceived and designed the experiments: AD AC. Performed the experiments: AC. Analyzed the data: AC SM YKJ AD. Contributed reagents/materials/analysis tools: AD AC. Wrote the paper: AC SM YKJ AD.

References

1. Chaudhuri A, Polyakova J, Zbrzezna V, Williams K, Gulati S, et al. (1993) Cloning of glycoprotein D cDNA, which encodes the major subunit of the Duffy blood group system and the receptor for the Duffy blood group system and the receptor for the Plasmodium vivax malaria parasite. Proc Natl Acad Sci USA 90: 10793–10797.
2. Iwamoto S, Li J, Ormi T, Ikemoto S, Kajii E (1996) Identification of a novel exon and spliced form of Duffy mRNA that is the predominant transcript in both erythroid and postcapillary venule endothelium. Blood 87:378–385.
3. Chaudhuri A, Polyakova J, Zbrzezna V, Pogo AO (1995) The coding sequence of Duffy blood group gene in humans and simians: restriction fragment length polymorphism, antibody and malarial parasite specificities, and expression in nonerythroid tissues in Duffy-negative individuals. Blood 85: 615–621.
4. Pogo AO, Chaudhuri A (2000) The Duffy protein: a malarial and chemokine receptor. Semin Hematol 37: 122–129.
5. Hamblin MT, Di Rienzo A (2000) Detection of the signature of natural selection in humans: evidence from the Duffy blood group locus. Am J Hum Genet 66: 1669–1679.
6. Hamblin MT, Thompson EE, Di Rienzo A (2002) Complex signatures of natural selection at the Duffy blood group locus. Am J Hum Genet 70: 369–383.
7. Miller LH, Mason SJ, Clyde DF, McGinniss MH (1976) The resistance factor to Plasmodium vivax in blacks — The Duffy blood group genotype, *FyFy*. New Engl J Med 295: 302–304.
8. Mendes C, Dias F, Figueiredo J, Mora VG, Cano J, et al. (2011) Duffy negative antigen is no longer a barrier to Plasmodium vivax – molecular evidences from the




Figure 2. Neighbor-Joining (NJ) phylogenetic tree showing genetic interrelationship among seven different population samples based on pairwise population genetic distance matrix (Nei's *D*, Table S3). The colour codes of the three clades are same as that of the Figure 1, corresponding to differential incidences of vivax malaria (high, medium, low). doi:10.1371/journal.pone.0045219.g002

- African West Coast (Angola and Equatorial Guinea). *PLoS Negl Trop Dis* 5: e1192.
9. King CL, Adams JH, Xianli J, Xianli J, Grimberg BT, et al. (2011) Fy^a/Fy^b antigen polymorphism in human erythrocyte Duffy antigen affects susceptibility to *Plasmodium vivax* malaria. *Proc Natl Acad Sci USA* 108: 20113–20118.
 10. Howes RE, Patil AP, Piel FB, Nyangiri OA, Kabaria CW, et al. (2011) The global distribution of the Duffy blood group. *Nat Commun* 2: 266.
 11. Das A, Anvikar AR, Cator LJ, Dhiman RC, Eapen A, et al. (2012) Malaria in India: The center for the study of complex malaria in India. *Acta Tropica* 121: 267–273.
 12. Das A (1991) DNA fingerprinting in India. *Nature* 350: 387.
 13. Bamshad M, Kivisild T, Watkins WS, Dixon ME, Ricker CE, et al. (2001) Genetic evidence on the origins of Indian caste populations. *Genome Res* 11: 994–1004.
 14. Khan N, Chittoria A, Pande V, Jaiswal YK, Das A (2012) Development of multilocus putatively neutral DNA markers in the X-chromosome for population genetic studies in humans. *Ann Hum Biol* 39: 281–289.
 15. Mohanty SS, Singh KV, Fotedar R, Lakshminarayana J, Parihar RS (2011) Prevalence of Duffy blood groups among the population of the desert region of India. *J Rural Trop Public Health* 10: 53–56.
 16. Das MK, Singh SS, Adak T, Vasantha K, Mohanty D (2005) The Duffy blood groups of Jarawas - the primitive and vanishing tribe of Andaman and Nicobar Islands of India. *Trans Med* 15: 237–240.
 17. Verma IC, Thakur A (1993) Duffy blood group determinants and malaria in India. *J Genet* 72: 15–19.
 18. Kar S, Seth S, Seth PK (1991) Duffy blood groups and malaria in the Ao Nagas in Nagaland, India. *Hum Hered* 41: 231–235.
 19. Singh V, Mishra N, Awasthi G, Dash AP, Das A (2009) Why is it important to study malaria epidemiology in India? *Trends Parasitol* 25: 452–457.
 20. Nei M (1972) Genetic distance between populations. *Am Nat* 106: 283–292.
 21. Das A, Mohanty S, Stephan W (2004) Inferring the population structure and demography of *Drosophila ananassae* from multilocus data. *Genetics* 168: 1975–1985.
 22. Gupta B, Srivastava N, Das A (2012) Inferring the evolutionary history of Indian *Plasmodium vivax* from population genetic analyses of multilocus nuclear DNA fragments. *Mol Ecol* 21: 1597–1616.
 23. Penrose LS (1955) Evidence of heterosis in man. *Proc Royal Soc Lond Series B* 144: 203–213.
 24. Rudan I, Carothers AD, Polasek O, Hayward C, Vitart V, et al. (2008) Quantifying the increase in average human heterozygosity due to urbanisation. *Eur J Hum Genet* 16: 1097–1102.
 25. Perna SJQ, Cardoso GL, Guerrero JF (2007) Duffy blood group genotypes among African-Brazilian communities of the Amazon region. *Genet Mol Res* 6: 166–172.
 26. Tournamille C, Blancher A, Le Van Kim C, Gane P, Apoil PA, et al. (2004) Sequence, evolution and ligand binding properties of mammalian Duffy antigen/receptor for chemokines. *Immunogenet* 55: 682–694.
 27. Gupta B, Gupta P, Sharma A, Singh V, Dash AP, et al. (2010) High proportion of mixed-species *Plasmodium* infections in India revealed by PCR diagnostic assay. *Trop Med Intl Health* 15: 819–824.
 28. Sucheston L, David B, Witonsky DB, Hastings D, Yildiz O, et al. (2011) Natural selection and functional genetic variation in the p53 pathway. *Hum Mol Gen* 20: 1502–1508.
 29. Eanes WF (2010) Molecular population genetics and selection in the glycolytic pathway. *J Exp Biol* 214: 165–171.
 30. Tishkoff SA, Varkonyi R, Cahinhinan N, Abbes S, Argyropoulos G, et al. (2001) Haplotype diversity and linkage disequilibrium at human G6PD: recent origin of alleles that confer malarial resistance. *Science* 293: 455–462.
 31. Currat M, Trabuchet G, Rees GD, Perrin P, Harding RM, et al. (2002) Molecular analysis of the β -Globin gene cluster in the Niokholo Mandenka population reveals a recent origin of the β^S Senegal mutation. *Am J Hum Genet* 70: 207–223.