

RESEARCH

Open Access

Enzymatic and toxigenic ability of opportunistic fungi contaminating intensive care units and operation rooms at Assiut University Hospitals, Egypt

Mohamed Bassam Aboul-Nasr^{1*}, Abdel-Naser Ahmed Zohri² and Enas Mahmoud Amer²

Abstract

Total of 110 isolates belonging to 8 fungal species collected from intensive care units (ICUs) and operation rooms (ORs) at Assiut University hospitals were examined for their ability to produce some extracellular enzymes and mycotoxins which are considered as important factors involved in for fungal pathogenicity. The results revealed that 73, 92 and 78 out of the 110 tested isolates produced protease, lipase and urease respectively; meanwhile, 77 of the tested isolates exhibited some hemolytic activities. Chromatographic analysis (TLC) of the crude extract of the fungal isolates tested revealed that 79 isolates of them had the ability to produce at least one of these mycotoxic compounds (aflatoxins B₁, B₂, G₁, gliotoxin, fumigillin, T-2, zearalenone, roridin A & E, verrucarins A & J, trichoveroids, satratoxin H & E). These results demonstrate that the opportunistic fungal species isolated from (ICUs) and (ORs) and tested exhibited some enzymatic and mycotoxic ability which are the most effective virulence factors contributing to fungal pathogenicity indicating that the management of infection control unit at Assiut University hospitals must be aware of not only bacterial but also fungal contamination.

Keywords: Airborne fungi; Enzymes; Mycotoxins and pathogenicity

Introduction

Members belonging to the *Aspergillus*, *Fusarium* and *Mucorales* genera are regarded as the main cause of fungal infections at hospitals (Alberti et al., 2001; Faure et al., 2000; Perdelli et al., 2006). Indeed fungi contaminating hospital rooms may grow on organic matter including various building materials and develop microcolonies. Prior to possible dissemination spores emanating from these colonies could be inhaled by immuno-suppressed patients resulting in local infections (Singh & Paterson, 2005). The ability of fungi to cause human diseases (mycoses) as pathogens appears to be accidental and such diseases are primarily related to the immunological status of the host and environmental exposure, rather than to the infecting organism (Rippon, 1988; Kwon-Chung & Bennett, 1992; Ellis, 1994). A relationship between fungal contamination in hospital environments and the incidence

of invasive aspergillosis has been demonstrated (Alberti et al., 2001) and more than 500 cases of post-operative aspergillosis in immuno-competent individuals have been reported (Pasqualotto and Denning, 2006). *Aspergillus fumigatus* and *A. flavus* are the leading species of the genus *Aspergillus* causing invasive aspergillosis (Pasqualotto, 2008). Outbreaks of nosocomial aspergillosis are attributed to airborne sources and even small concentrations of spores have been associated with outbreaks (Vonberg and Gastmeier, 2006).

Fungi are known to elaborate extracellular enzymes based on the substrate they utilize for growth. Extracellularly produced enzymes have been described in certain fungi such as *Candida* (Borg & Ruchel, 1988) and *Aspergillus* species (Hanzi et al., 1993). Production and secretion of hydrolytic enzymes, such as proteases, lipases and phospholipases are very important virulence factors. These enzymes play a role in nutrition, tissue damage, fungal dissemination within the human body, iron acquisition and overcoming the host immune system

* Correspondence: mbassamaboulnasr@yahoo.com

¹Faculty of Science, Department of Botany, University of Sohag, Sohag, Egypt
Full list of author information is available at the end of the article

which strongly affects fungal pathogenicity (Ibrahim *et al.*, 1995). Secretion of enzymes into extracellular environments might be an important adaptive mechanism during the life cycle of fungi (Monod *et al.*, 2002). Earlier studies on fungal enzymatic activities aimed at establishing the role of enzymes in fungal pathogenicity, as well as their capacity to induce inflammatory reactions in the host (Rippon, 1982). It is logical to suppose that these enzymes could act by enabling tissue invasion easier, but they could also participate in causing infection by impairing some mechanisms of the immune system and/or assist in obtaining of nutrients, thus causing injury to the host (Birch *et al.*, 2004; Da Silva *et al.*, 2005). Hussein and Brasel *et al.*, (2001) examined the ability of 80 fungal isolates from keratitis patients to produce extracellular enzymes in growth medium and found that most of these isolates could produce protease, lipase, urease and catalase enzymes but at varying levels.

Mycotoxins are an extremely adverse group of low molecular weight fungal secondary metabolites which when ingested, inhaled or absorbed through the skin; can cause lowered performance, sickness or even death in man and animals including birds (Pitt, 1996 and Van Egmond & Speijers, 1999). They may affect the reproductive, immune, gastrointestinal systems, specific target organs, in addition they may exhibit hormonal activity, developmental defects including those relating to birth (tetragenic and neurotoxic) (Richard, 1991; Sharma, 1993; Kuiper-Goodman, 2004). Although fungal spores may have the highest concentrations of mycotoxins, the vegetative part of the fungus, the mycelium or the substrate upon which the fungus grows can also contain these toxins. Viability of spores is not essential to toxicity. In other words, a dead spore can still be a source of toxin (Keller, *et al.*, 2005). Infants and children are considered more susceptible to different toxins than adults, because of their lower body weight, higher metabolic rate, incomplete development of some organs and tissues such as those in the central nervous system (WHO, 1986; NAS, 1993). Nonetheless, human illnesses caused by mycotoxins may be a public health problem than one realizes because some of these cases (chronic) may go un-noticed for an extended period of time unless large amounts of mycotoxins are consumed resulting in acute symptomology (Hesseltine, 1985). Like the case may be for other hospital infection control units elsewhere, Assiut University hospitals only focus on bacterial infection. This study was directed to examine the potential of the most common fungal species isolated from the atmospheric air and dust samples collected from air conditioners of ICUs and ORs to produce extracellular hydrolytic enzymes and mycotoxic compounds which are important virulence factors involved in fungal pathogenicity.

Materials and methods

Screening of fungal isolates for extracellular enzyme production

A total of 110 filamentous fungi isolated from air as well as filters of air conditioning systems in intensive care units and operation rooms (noted in our laboratory) were screened for their ability to produce extracellular enzymes in solid media. The following fungal species were tested: i.e. *Aspergillus flavus* (20 isolates), *A. fumigatus* (16), *A. niger* (19), *Cladosporium cladosporioides* (5), *Fusarium solani* (17), *F. oxysporum* (6), *Myrothecium roridum* (1) and *Stachybotrys elegans* (26).

Protease activity was determined using a Casein hydrolysis medium in which skim milk gives an opaque final appearance and hydrolysis of the casein resulted in a clear zone around the fungal colony. (Paterson & Bridge, 1994). Lipase activity was measured using the method of Ullman & Blasins (1974) with some modification this time using Tween 80 instead of Tween 20. The lipolytic producing ability was observed as a visible precipitate due to the formation of crystals of calcium salt of the oleic acid liberated by the enzyme. Urease activity was determined using urease medium described by Paterson & Bridge (1994). Isolates capable of producing urease turned the yellow color of the acidic medium to purple-red or deep pink color., meanwhile hemolytic activity of fungal isolates was measured using human blood agar medium (Ronald, 2000).

Screening of fungal isolates for mycotoxin production

Cultivation of fungal isolates and extraction of their mycotoxins

One-hundred and ten isolates were cultivated in a 15 cm Petri dish containing solid Czapek's glucose agar under aseptic conditions and incubated at $25 \pm 2^\circ\text{C}$ for 10 days. The plates containing *Fusarium*, *Stachybotrys* and *Myrothecium* isolates were transferred into a refrigerator for another 10 days. At the end of incubation periods, an agar plug technique was employed for the extraction of mycotoxins in culture materials, whereby, all the agar medium with fungal mycelia were cut into small pieces, transferred into a 250 ml Erlenmeyer flask containing 50 ml 96% methanol. The content was shaken on a rotary shaker (200 r.p.m., 24 h) and filtered through filter paper (Aboul-Nasr; Obied-Allah, 2013).

The extracted material was then washed using 25 ml of the same extractin solvent. The methanol extracts were combined, dried over anhydrous sodium sulphate, and concentrated under a vacuum. The residue was transferred to a dram vial and further evaporated to near dryness.

Thin layer chromatographic analysis

For the screening of mycotoxins in extracts, a thin-layer chromatographic technique adopted by El-kady and Moubasher (1982) was employed.

Results and discussion

The ability of clinical fungal strains from various ICUs and ORs at Assiut University hospital units were tested for their ability to produce extracellular enzymes in solid media and data presented in Table 1. Data indicated that more than 66% of tested isolates (73 out of 110) of five fungal genera; 3 *Aspergillus spp.* (55 isolates), one *Cladosporium sp.* (5 isolates), 2 *Fusarium spp.* (23 isolates), one *Stachybotrys sp.* (26 isolates) and one isolate of *Myrothecium sp.* had the ability to produce protease. Nearly similar percent (70.9%) of the tested fungal isolates (78 out of 110) were recorded as lipase producers in this study. Salyers and Witt (1994) reported that microbial cells secrete hydrolytic enzymes that destroy the constituents of host cell membranes leading to membrane dysfunction, physical disruption as well as aid in the invasion of host tissues. Proteolytic degradation of lung tissues has been suggested as one of the key events involved in the physiopathology of *A. fumigatus* (Kothary *et al.*, 1984). Also, several species of *Aspergillus* such as *A. fumigatus*, *A. flavus*, *A. oryzae* and *A. sojae* are known to secrete protease as reported by Monod *et al.*, (1993). Stehr *et al.*, (2003) found that extracellular lipases play a role during microbial infections and suggested their role is to digest lipids for nutrient acquisition by pathogenic microbe and that these enzymes help the microbe (bacteria or fungi) to grow in environments where lipids are the sole carbon source.

A majority of the fungal isolates under study (92 out of 110: 83.6%) were able to produce urease. Urease catalyses the hydrolysis of urea to ammonia and carbamate, in which the latter by-product is further hydrolyzed to ammonia and carbonic acid resulting in an increase in pH (Zimmer, 2000). Urease activity has been found in several bacteria and fungi and has been shown to be an important pathogenic factor (Eaton *et al.*, 1991; Cox *et al.*, 2000). It has also been postulated that much of the tissue

damage induced by *Helicobacter pylori* is as a result of ammonium hydroxide produced through the actions of urease. Studies have shown that the actions of urease may alter the function of white blood cells (Mai *et al.*, 1992; Mobley, 1996). In this study, about 70.9% of the tested fungal isolates (78 out of 110) exhibited a lysis activity (hemolysis) on human blood. Vesper *et al.* (1999; 2001) isolated stachylysin a hemolytic agent from *Stachybotrys chartarum*. Donohue *et al.*, (2004) isolated chrysolysin another hemolytic agent from *P. chrysogenum*.

The toxic potentials of these fungal isolates previously tested herein for their enzymatic activity were also evaluated and the data presented in Table 2 revealed that 79 out of the 110 tested fungal isolates (71.82%) were recorded as mycotoxin producers. Several mycotoxins including aflatoxins, gliotoxin, fumigillin, cladosporin, T-2 toxin, zearalenone, rosidins, verrucarins, trichoveroides and satratoxins were produced by different fungal isolates in this study. Aflatoxins B₁ & B₂ were recorded in extracts obtained from cultures of 11 *A. flavus* with one positive isolates being able to produce additionally aflatoxins G₁ & G₂. Aflatoxins are produced by many strains of *A. flavus*. They are toxic, having carcinogenic, mutagenic and teratogenic effects in laboratory animals (Abdel-Wahhab *et al.*, 1998, 2006). Aflatoxin B₁ is the most potent carcinogenic substance naturally produced mainly by *A. flavus* and *A. parasiticus* (Squire, 1981) and is classified by the International Agency of Research on Cancer as a group 1 human carcinogen (IARC, 1982). Gliotoxin and fumigillin were produced by all the 16 tested isolates of *A. fumigatus*. Also, 7 out of 19 tested isolates of *A. niger* had the ability to produce gliotoxin. *Aspergillus* members especially *A. fumigatus* and *A. niger* are the most common causal agents of aspergillosis. Invasive aspergillosis is very common among immunocompromised patients, with reported incidence rates of 17-26% among lung transplants patients, 5-15% allogenic

Table 1 Extracellular enzymes produced by the tested fungal isolates collected from ICUs and ORs at Assiut University hospitals

Fungal isolates	No. of tested isolates	Protease		Lipase		Urease		Hemolysis	
		+ Ve	- Ve	+ Ve	-Ve	+Ve	-Ve	+ Ve	- Ve
<i>Aspergillus flavus</i>	20	16	4	19	1	20	0	19	1
<i>Aspergillus fumigatus</i>	16	10	6	15	1	15	1	12	4
<i>Aspergillus niger</i>	19	15	4	16	3	8	11	13	6
<i>Cladosporium cladosporioides</i>	5	1	4	4	1	0	5	2	3
<i>Fusarium solani</i>	17	11	6	9	8	10	7	7	10
<i>Fusarium oxysporum</i>	6	3	3	5	1	5	1	4	2
<i>Myrothecium roridum</i>	1	1	0	1	0	1	0	1	0
<i>Stachybotrys elegans</i>	26	22	4	25	1	19	7	19	7
Total	110	79	31	94	16	78	32	77	33

Table 2 Mycotoxigenicity of fungal isolates collected from ICUs and ORs at Assiut University hospital

Toxins produced	No. of strains positive	No. of strains tested	Fungal isolates
Aflatoxins B1, B2	10	20	<i>Aspergillus flavus</i>
Aflatoxins B1, B2, G1, G2	1		
Gliotoxin, fumigillin	16	16	<i>Aspergillus fumigatus</i>
Gliotoxin	7	19	<i>Aspergillus niger</i>
Cladosporin	5	5	<i>Cladosporium cladosporioides</i>
T-2 toxin	9	17	<i>Fusarium solani</i>
T-2 toxin, Zearalenone	2		
Zearalenone,	1		
Zearalenone	6	6	<i>Fusarium oxysporum</i>
Roridin A & E, Verrucarins A & J	1	1	<i>Myrothecium roridum</i>
Trichoveroides	10	26	<i>Stachybotrys elegans</i>
Trichoveroides & verrucarins J	6		
Satratoxins H & E, Trichoveroides, Verrucarins J	5		

bone marrow transplants patients, 5-24% of those with acute leukaemia and 2-13% heart transplants patients (Curtis *et al.*, 2004). Kupfahl *et al.*, (2008) investigated the presence of gliotoxin-producing *Aspergillus* strains among clinical isolates collected from different parts of Germany and Austria. In that light, we collected and examined 158 different *Aspergillus* isolates consisting of 100 *A. fumigatus*, 27 *A. terreus*, 15 *A. flavus* and 16 *A. niger* strains from different medical centers and other environmental samples. According to that study, gliotoxin was recovered from 98, 56, 37 and 13% of *A. fumigatus*, *A. niger*, *A. terreus* and *A. flavus*, respectively. Nielsen (2003) found that *A. fumigatus* and *A. niger* are frequently isolated as indoor moulds, with the former isolate having the ability to produce fumitoxins, fumitremergens, gliotoxin and other mycotoxins, while the later produced only ochratoxin A. Cladosporin was produced by five tested isolates of *C. cladosporioides* in this study. This species is one of the causative agents of skin lesions, keratitis, nail fungus and pulmonary infections. Also, acute symptoms of exposure to this fungus are edema and bronchospasms which may lead to pulmonary emphysema. Zearalenone was produced by all six tested isolates of *F. oxysporum* and three out of 17 isolates of *F. solani*. This toxin has an estrogenic activity and this estrogenic property enables exposure to its products a concern for human health (Niyono *et al.*, 1988, Richard, 1991). In humans, the major effect of zearalenone bears on the reproductive system affecting reproductive organs' structure and function that may lead to hyperestrogenism (Kuiper-Goodman *et al.*, 1993). T-2 toxin productivity by 11 of the 17 tested *F. solani* isolate was recorded in this study. This toxin is a representative of a large group of non-macrocytic trichothecenes whose major effect and that of other trichothecenes is their

ability to inhibit protein synthesis which is followed by a secondary disruption of DNA and RNA synthesis (Niyono *et al.*, 1988; Richard, 1991). Verrucarins A & J as well as roridin A & E are mycotoxins produced by *M. roridum* found in this study. Also, verrucarins J was produced by 11 out of 26 tested isolates of *S. elegans* with 5 of this species found to produce satratoxins H & E. Meanwhile 19 of the same species were able to synthesize trichoveroides. Verrucarins, roridins, trichoveroides and satratoxins. Different types of macrocyclic trichothecenes produced mainly by *Stachybotrys* (Eppley, 1977; Eppley and Bailey, 1973; Eppley *et al.*, 1980). The involvement of macrocyclic trichothecenes in stachybotryotoxicosis were suggested by isolation of satratoxins from straws fed to sheep and cattle with stachybotryotoxicosis (Harrach and Bata, 1983; Harrach *et al.*, 1983). Bata *et al.* (1985) found that all 17 strains of *Stachybotrys atra* isolated in Middle Europe and found to produce each of the five macrocyclic trichothecenes that included satratoxins H & G, verrucarins A, E & J, trichoveroides and roridin A & E. Islam *et al.*, (2006) suggested that neurotoxicity and inflammation in the nose and brain are potential adverse health effects of exposure to satratoxins.

Conclusion

In conclusion, hydrolytic enzymes and mycotoxic compounds which are considered the most important virulence factors influencing the pathogenicity of opportunistic fungal infections were detected in most of the cultures of fungal isolates tested herein. Thus the personnel managing the infection control unit of Assiut University hospitals must be aware of not only bacterial contamination, but there is a potential for the distribution of fungal infection as well.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MBA designed the research plan, followed the research plan execution point by point, helped in drafting the manuscript, revised it and was the corresponding author. AAZ helped in identifying the toxic profile and planned the enzyme profile experiments. EMA carried out the research plan point by point, identifying fungi and toxins under the supervision of MBA and helped in drafting the manuscript. All authors read and approved the final manuscript.

Author details

¹Faculty of Science, Department of Botany, University of Sohag, Sohag, Egypt. ²Faculty of Science, Department of Botany, University of Assiut, Assiut, Egypt.

Received: 16 April 2013 Accepted: 26 July 2013

Published: 29 July 2013

References

- Abdel-Wahhab MA, Nada SA, Farag IM, Abbas NF, Amra HA (1998) Potential protective effect of HSCAS and bentonite against dietary aflatoxicosis in rat: with special reference to chromosomal aberrations. *Nat J Toxins* 6:211–218
- Abdel-Wahhab MA, Ahmed HH, Hagazi MM (2006) Prevention of aflatoxin B₁-initiated hepatotoxicity in rat by marine algae extracts. *J Appl Toxicol* 26:229–238
- Aboul-Nasr MB, Obied-Allah MR (2013) Biological and Chemical Detection of Fumonisin Produced on Agar Medium by *Fusarium verticillioides* Isolates Collected from Corn in Sohag-Egypt. *J Microbiol*, Epub ahead of print
- Alberti C, Bouakline A, Ribaud P, Lacroix C, Rousselot P, Leblanc T (2001) Relationship between environmental fungal contamination and the incidence of invasive aspergillosis in haematology patients. *J Hosp Infect* 48:198–206
- Bata A, Balazs H, Ujszasz K, Kis-Tamas A, Laszity R (1985) Macrocyclic trichothecene toxins produced by *Stachybotrys atra* strains isolated in Middle Europe. *J Appl & Environ Microbiol* 49(3):678–681
- Birch M, Denning DW, Robson GD (2004) Comparison of extracellular phospholipase activities in clinical and environmental *Aspergillus fumigatus* isolates. *J Med Mycol* 42:81–86
- Borg M, Ruchel R (1988) Expression of extracellular acid proteinase by proteolytic *Candida* species during experimental infection of oral mucosa. *J Infect Immun* 56:626–663
- Cox GM, Mukherjee J, Cole GT, Casadevall A, Perfect JR (2000) Urease as a virulence factor in experimental cryptococcosis. *J Infect Immun* 68:443–448
- Curtis MS, Allen CIH, Lieberman MD, Martha SMD, William R, Marsha VMD (2004) Adverse Health Effects of Indoor Mould. *Aust College of Nutri & Environ Medicine* 23(1):3–8
- Da Silva BCM, Auler ME, Ruiz LS (2005) *Trichophyton rubrum* isolated from AIDS and human immunodeficiency virus-infected patients in São Paulo, Brazil: antifungal susceptibility and extracellular enzyme production. *J Chemotherapy* 51:21–26
- Donohue M, ChungY MML, Ward M, Selgrade MJ, Vesper SJ (2004) Hemolysin Chrysolysin from *Penicillium chrysogenum* promotes inflammatory response. *Int J Hyg Environ Health* 208(4):279–85
- Eaton KA, Brooks CL, Morgan DR, Krakowka S (1991) Essential role of urease in pathogenesis of gastritis induced by *Helicobacter pylori* in gnotobiotic piglets. *J Infect Immun* 59:2470–2475
- El-Kady, Moubasher MH (1982) Toxicogenicity and toxin of *Stachybotrys chartarum* isolates from wheat straw samples in Egypt. *J Exp Mycol* 6:25–31
- Ellis D (1994) The Human Opportunistic Mycoses. *J Clin Mycol* 36(1):166
- Eppley RM (1977) Mycotoxins in human and animal health. *Pathotox Publishers*, Park Forest South, Ill. V. Rodricks, pp 285–293
- Eppley RM, Bailey WJ (1973) 12, 13 Epoxy-A9-trichothecenes as the probable mycotoxins responsible for stachybotryotoxicosis. *J Sci* 191:758–760
- Eppley RM, Highet RJ, Bailey WJ (1980) Structure of satratoxin H, a metabolite of *Stachybotrys atra* application of proton and carbon – 13 nuclear magnetic resonance. *J Org Chem* 42:240–243
- Faure O, Fricker-Hidalgo H, Lebeau B, Mallaret MR, Ambroise-Thomas P, Grillot R (2000) Eight year surveillance of environmental fungal contamination in hospital operating rooms and haematologic units. *J Hosp Infect* 50:155–160
- Hanzi M, Shimizu M, Hearn VM, Monod M (1993) A study of alkaline protease secreted by different *Aspergillus* species. *J Myco* 36:351–356
- Harrach B, Bata A, Bajmoczy E, Benko M (1983) Isolation of satratoxins from the bedding straw of a sheep-flock with fatal stachybotryotoxicosis. *J Appl Environ Microbiol* 45:1419–1422
- Hesseltine CW (1985) Global significance of mycotoxins. In: Steyn PS, Vlegaar R (eds) *J Mycotox and Phyco- toxins*. Elsevier Scientific Publishing Co., Amsterdam, pp 1–18
- Hussein HS, Brasel JM (2001) Toxicity, metabolism, and impact of mycotoxins on humans and animals. *J Toxicol* 167(2):101–134
- Ibrahim AS, Mirbod F, Filler SG, Banno Y, Cole GT, Kitajima Y, Edwards JE Jr, Nozawa Y, Ghannoum MA (1995) Evidence implicating phospholipase as a virulence factor of *Candida albicans*. *J Infect Immun* 63:1993–1998
- International Agency for Research on Cancer (IARC) (1982) The evaluation of the carcinogenic risk of chemicals to humans. Monograph Supplement 4. International Agency for Research on Cancer, Lyon, France
- Islam J, Harkema R, Pestka JJ (2006) Satratoxin G from the Black Mold *Stachybotrys chartarum* Evokes Olfactory Sensory Neuron Loss and Inflammation in the Murine Nose and Brain. *J Environ Health Perspect* 114:1099–1107
- Keller NP, Turner G, Bennett JW (2005) Fungal secondary metabolism – from biochemistry to genomics. *J Nat Rev Microbiol* 3(12):937–947
- Kothary MH, Chase T Jr, MacMillan JD (1984) Correlation of elastase production by some strains of *Aspergillus fumigatus* with ability to cause pulmonary invasive Aspergillosis in mice. *J Infect Immun* 43:320–325
- Kuiper-Goodman T (2004) Chapter 1. Risk assessment and risk management of mycotoxins in food. In: Mogan N, Olsen M (eds) *Mycotoxins in Food, Detection and Control*. CRC Press, New York; Wood head Publishing Limited, Cam-bridge, England, pp 3–31
- Kuiper-Goodman T, Ominiski K, Marquardt RR, Malcolm S, McMullen E, Lombaert GAE, Morten T (1993) Estimating human exposure to ochratoxin A in Canada. In: Creppy E, Dirheimer G (eds) *Human Ochrotoxicosis and Its Pathologies*, Montrouge. John Libbey, Eurotech, London, pp 167–74
- Kupfahl AM, Cornelia LF, Guido F, Gerhard H, Thomas R, Gernot G, Herbert H (2008) Gliotoxin production by clinical and environmental *Aspergillus fumigatus* strains. *Med Microbiol* 298(3–4):319–327
- Kwon-Chung KJ, Bennett JE (1992) Mycosis. *J Med Mycol*, Lea & Febiger, Philadelphia 31(1):560–593
- Mai UEH, Perez-Perez GI, Allen JB, Wahl SM, Blaser MJ, Smith PD (1992) Surface proteins from *Helicobacter pylori* exhibit chemotactic activity for human leukocytes and are present in gastric mucosa. *J Exp Med* 175:517–525
- Marc Z (2000) Molecular mechanics evaluation of the proposed mechanisms for the degradation of urea by urease. *J Biomol Struct Dyn* 17(5):787–97
- Mobley HL (1996) The role of *Helicobacter pylori* urease in the pathogenesis of gastritis and peptic ulceration. *J Aliment Pharmacol Ther* 10:57–64
- Monod M, Paris S, Sarfati J, Jaton-Ogay P, Ave P, Latge' JP (1993) Virulence of alkaline protease-deficient mutants of *Aspergillus fumigatus*. *FEMS Microbiol Letters* 106:39–46
- Monod M, Capoccia S, Lechenne B (2002) Secreted proteases from pathogenic fungi. *Int Med J Microbiol* 292:405–419
- NAS (1993) Pesticides in the Diets of Infants and Children. National Academy Press, Washington, DC, pp 253–258
- Nielsen KF (2003) Mycotoxin production by indoor molds. *Fungal Genet Biol* 39:103–117
- Niyo KA, Richard JL, Tiffany LH (1988) Effect of T-2 mycotoxin ingestion on phagocytosis of *Aspergillus fumigatus* conidia by rabbit alveolar macrophages and on hematologic, serum biochemical, and pathologic changes in rabbits. *J Vet Res* 49:1766–1773
- Pasqualotto AC (2008) Differences in pathogenicity and clinical syndromes due to *Aspergillus fumigatus* and *Aspergillus flavus*. *J Med Mycol* 47(1):1–10
- Pasqualotto AC, Denning DW (2006) Post-operative aspergillosis. *Clinic Microbiol Infect* 12:1060–1076
- Paterson RRM, Bridge PD (1994) Biochemical techniques for filamentous fungi. *Int J Mycol Instit* 1:21
- Perdelli F, Cristina ML, Sartini M, Spagnolo AM, Dallera M, Ottria G (2006) Fungal contamination in hospital environments. *J Infect Cont Hosp Epidemiol* 27:44–47
- Pitt JI (1996) What are mycotoxins? *Austral. Mycotoxin Newsletter* 7(4):1–3
- Richard J (1991) Mycotoxins as immunomodulators in animal systems. In: Bray GA, Ryan DH (eds) *Mycotoxins, Cancer, and Health*. Pennington Center Nutrition Series. Louisiana State University Press, Baton Rouge, LA, pp 197–220

- Rippon JW (1988) *The Pathogenic Fungi and the Pathogenic Actinomycetes*, 3rd ed., WB Saunders, Philadelphia, 3rd ed. *J Med Mycol*:797
- Ronald MA (2000) *Hand book of Microbiological Media*, 10th edn. CRC press, Inc, USA, p 137,333,785,279
- Salyers A, Witt D (1994) Virulence factors that damage the host. *Bacterial pathogenesis: a molecular approach*. ASM Press, Washington, D.C, pp 47–62
- Sharma RP (1993) Immunotoxicity of mycotoxins. *J Dairy Sci* 76:892–897
- Singh N, Paterson DL (2005) *Aspergillus* infections in transplant recipients, Review. *J Clin Microbiol* 18:44–69
- Squire RA (1981) Ranking animal carcinogens: a proposed regulatory approach. *Sci* 214:877–880
- Stehr F, Kretschmar M, Kroger C, Hube B, Schafer W (2003) Microbial lipases as virulence factors. *Molecular Catalysis, B: Enzymatic* 22:347–355
- Ullman V, Blasins G (1974) A simple medium for the detection of different lipolytic activity of microorganisms. *Zentrabl Bakteriell J Hyg II Abt A* 229:264–267
- van Egmond HP, Speijers GA (1999) Natural toxins I. Mycotoxins. In: van der Heijden K, Younes M, Fishbone L, Miller S (eds) *International Food Safety Handbook, Science, International Regulation, and Control*. Marcel Dekker, Inc, New York, Basel, pp 342–355
- Vesper SJ, Dearborn DG, Yike I, Sorenson WG, Haugland RA (1999) Hemolysis, toxicity, and randomly amplified polymorphic DNA analysis of *Stachybotrys chartarum* strains. *J Appl Environ Microbiol* 65:3175–3181
- Vesper SJ, Magnuson S, Dearborn D, Yike I, Haugland RA (2001) Initial characterization of the hemolysin from *Stachybotrys chartarum*. *J Infect Immun* 69:912–916
- Vonberg RP, Gastmeier P (2006) Nosocomial aspergillosis in outbreak settings. *J Hosp Infect* 63:246–254
- WHO (Environmental Health Criteria) (1986) Principles for evaluating health risks from chemicals during infancy and early childhood; the need for a special approach. IPCS, Geneva

doi:10.1186/2193-1801-2-347

Cite this article as: Aboul-Nasr *et al.*: Enzymatic and toxigenic ability of opportunistic fungi contaminating intensive care units and operation rooms at Assiut University Hospitals, Egypt. *SpringerPlus* 2013 **2**:347.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
