

POSTER PRESENTATION

Open Access

Phase I study of safety and immunogenicity of ADU-623, a live-attenuated *Listeria monocytogenes* vaccine (Δ actA/ Δ inlB) expressing EGFRVIII and NY-ESO-1, in patients with WHO grade III/IV astrocytomas

Marka Crittenden^{1*}, Keith S Bahjat², Rui Li³, Pankaj Gore⁴, Chris Fountain³, Bill Hanson⁵, Justin Skoble⁵, Peter Lauer⁵, Aimee L Murphy⁵, Thomas Dubensky⁵, Dirk G Brockstedt⁵, Walter Urba⁶

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015)
National Harbor, MD, USA. 4-8 November 2015

Background

The neo-antigen EGFRVIII is expressed in multiple tumor types, including high-grade astrocytomas. It is associated with a poor prognosis and resistance to conventional therapies such as chemotherapy and radiation that are part of the standard treatment. We propose that immunization with a live-attenuated *Listeria*-based vaccine, ADU-623, expressing EGFRVIII and NY-ESO-1 will elicit robust tumor-specific immune responses capable of killing EGFRVIII and/or NY-ESO-1-expressing tumor cells and improve survival of the patients. In addition, ADU-623 induces a potent innate immune response that can kill transformed cells even in the absence of neo-antigens. We designed a translational vaccine study to evaluate the safety and immunogenicity of this vaccine in patients with high-grade astrocytomas after standard of care therapy or at progression.

Methods

Patients with a pathologic diagnosis of WHO Grade III/IV astrocytic tumors that have completed standard of care external beam radiation therapy and concurrent temozolomide followed by adjuvant temozolomide or with radiographic evidence of progression following standard of care radiation and chemotherapy treatment, including those who have gone on to a second surgical

resection are eligible. Patients are enrolled and assigned consecutively to one of the following ADU-623 dose level cohorts: Cohort 1 3×10^7 cfu, Cohort 2 3×10^8 cfu, or Cohort 3 1×10^9 cfu, each administered IV on Days 0, 21, 42 and 63. Adverse events are monitored throughout the treatment and patients are followed for up to 24 months. Patients are currently accruing to Cohort 3. The primary objective is to determine the maximum tolerated dose and characterize the safety profile of ADU-623 in patients with treated and recurrent WHO Grade III/IV astrocytomas. Secondary objectives include progression free survival, time to progression and overall survival rates in patients vaccinated with ADU-623. Exploratory studies of EGFRVIII-, NY-ESO-1-, vector-specific and innate immune responses will be performed.

Trial registration

ClinicalTrials.gov identifier NCT01967758.

Authors' details

¹Earle A. Chiles Research Institute, Providence Cancer Center; The Oregon Clinic, Portland, OR, USA. ²Earle A. Chiles Research Institute, Robert W. Franz Cancer Research Center, Providence Cancer Center, Portland, OR, USA.

³Providence Cancer Center, Portland, OR, USA. ⁴Microneurosurgical Consultants, Portland, OR, USA. ⁵Aduro Biotech Inc., Berkeley, CA, USA. ⁶Earle A. Chiles Research Institute, Portland, OR, USA.

Published: 4 November 2015

¹Earle A. Chiles Research Institute, Providence Cancer Center; The Oregon Clinic, Portland, OR, USA

Full list of author information is available at the end of the article

doi:10.1186/2051-1426-3-S2-P162

Cite this article as: Crittenden *et al.*: Phase I study of safety and immunogenicity of ADU-623, a live-attenuated listeria monocytogenes vaccine (Δ actA/ Δ inlB) expressing EGFRVIII and NY-ESO-1, in patients with who grade III/IV astrocytomas. *Journal for ImmunoTherapy of Cancer* 2015 **3** (Suppl 2):P162.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

