

A New Technical Approach to Monitor the Cellular Physiology by Atomic Force Microscopy

Kyung Hwan Jeong, M.D.,
Sang Ho Lee, M.D.

Department of Internal Medicine, College of
Medicine, Kyung Hee University, Seoul, Korea

Received: August 26, 2012

Accepted: December 12, 2012

Corresponding Author: Sang Ho Lee, M.D.
Department of Nephrology, Kyung Hee University
Hospital at Gangdong, Sangil-dong 149,
Gangdong-gu, Seoul 134-727, Korea
Tel: +82-2-440-6882, Fax: +82-2-440-8150
E-mail: rulale@dreamwiz.com

Atomic force microscopy (AFM) has become an important medical and biological tool for non-invasive imaging and measuring the mechanical changes of cells since its invention by Binnig et al. AFM can be used to investigate the mechanical properties of cellular events in individual living cells on a nanoscale level. In addition, the dynamic cellular movements induced by biochemical activation of specific materials can be detected in real time with three dimensional resolution. Force measurement with the use of AFM has become the tool of choice to monitor the mechanical changes of variable cellular events. In addition, the AFM approach can be applied to measure cellular adhesion properties. Moreover, the information gathered from AFM is important to understanding the mechanisms related to cellular movement and mechanical regulation. This review will discuss recent contributions of AFM to cellular physiology with a focus on monitoring the effects of antihypertensive agents in kidney cells.

Key Words: Atomic force microscopy; Mesangial cell; Antihypertensive agents

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Since the first invention of atomic force microscopy (AFM) by Binnig et al., it was recognized as an important imaging tool in biological research¹. AFM differs from conventional optical and electron microscopy, because it is performed by sensing the interacting force between its probe tip and sample surface. Furthermore, measuring the force acting between its probe tip and the sample, by means of force-distance curves, is important in defining the sample's physical property. Thus, the AFM force curve can be used to evaluate the stiffness and adhesive properties of cell membranes as the tip moves to and from the sample surface². Early on, AFM was recognized for its potential to observe living animal cells³. As a result of technical innovations, it is now also possible to perform imaging in any environment including aqueous solutions using AFM.

In renal physiology, AFM is mainly used to image intact renal cells, or to study sub-cellular structures. However, workers interested in renal cells and their membrane transporter function, can now use AFM since it is an ideal method for examining membrane structures. These experiments are still essential and continue but they are now complemented by studies in other aspects of living cells' function, such as the measurement of cellular stiffness and cell adhesiveness^{4,5}.

This review will discuss some recent contributions of AFM to renal physiology with a focus on monitoring the effect of BP (blood pressure) lowering agents on cellular biological and mechanical responses.

Principle of AFM

AFM consists of a cantilever with a sharp tip which is used to scan the specimen surface. When the tip is placed near the sample surface, the force between the tip and

the sample lead to a deflection of the cantilever (Fig. 1). Laser light is reflected off the back of the deflected cantilever and collected by photodiodes. This interaction between the probe tip and sample surface is then translated into an appropriate image.

In contrast, when the AFM tip is pressed against the cell, the membrane is indented. This mechanical change evokes the distortion of the AFM cantilever, which serves as a spring constant. The cantilever deflection permits force-distance curves of the sample surface (Fig. 2). The slope of the approach curve, the first half of force-distance curve where the tip approaches the sample surface, is used to measure the stiffness of the surface. After the

AFM tip has indented the sample surface, the cantilever bends upward (Fig. 2A).

The retraction curve, the second half of force-distance curve where the tip pulls away from the sample surface, is used to determine the adhesion force. Adhesion force between the tip and the surface causes the cantilever to bend downward (Fig. 2B).

A new technical approach to monitor the effects of anti-hypertensive agent by AFM

AFM has become an important biological tool for non-invasive imaging and measuring the mechanical property




Fig. 1. Atomic force microscopy schematic diagram.




Fig. 2. Force-distance curves obtained in a single mesangial cell. The slope of approach curve (A). The slope of retraction curve (B) (adopted from Ref 15).

of cells and materials in renal physiology.

By using AFM, Oberleithner et al.^{6,7)} recently demonstrated that stimulation of the aldosterone receptor changes the endothelial cell stiffness and this phenomenon is related to the endothelial dysfunction observed in high blood pressure. Aldosterone increases plasma membrane roughness, which is detectable only with AFM in cardiomyocytes scanned under physiological conditions⁸⁾.

Angiotensin II (Ang II) is a potent systemic vasoconstrictor and plays an important role in renal injury by promoting cell apoptosis and inflammation^{9,10)}. We recently used AFM to observe Ang II induced dynamic contraction and elastic changes in live and fixed mesangial cells¹¹⁾. We observed structural changes and the stiffening of mesangial cells induced by Ang II. To measure changes in live cell stiffness, force-distance curve measurements were performed on mesangial cells before, and after Ang II application. Ang II activates Ang II type 1 (AT₁) receptor at the surface of endothelial, mesangial cell and smooth muscle cells. Activation of the AT₁ receptor induces various signaling pathways leading to actin cytoskeletal remodeling and contraction of the cell¹²⁾. Cuerrier CM et al. recently extended the use of AFM to monitor AT₁ receptor activation-induced mechanical responses in renal cells¹³⁾. The main feature of the AT₁ receptor activation is the large upward displacement of the cell membrane with contraction. It is important to mention that these nanoscaled cell contractions and membrane displacements cannot be detected using conventional optical techniques.

Several studies have documented the role of BP lowering agent on cells including endothelial cell and mesangial cell. Recently, AFM was performed to measure the effect of BP lowering agent on cellular stiffness, cell volume and apical surface. Specific β_1 receptor blocker, nebivolol, decreases endothelial cell stiffness, which is associated with a significant increase in endothelial cell size¹⁴⁾. These morphological and functional modulations observed in endothelial cells may explain the improved endothelial function with nebivolol treatment. A nanoscopy of the

cell surface revealed that aldosterone-induced roughness and increase of cell surface are inhibited by spironolactone⁸⁾. These nano-scaled observations obtained through experiments on individual cells, improve our understanding of the regulation of spironolactone in physiological processes. It could be explained that aldosterone-induced rearrangements of the cytoskeleton may influence the membrane surface of cells. These rearrangements were effectively prevented by spironolactone.

In addition, AFM has been introduced to directly measure the interaction forces between living cells and the AFM tip. We demonstrated for the first time the use of AFM force-distance curves on live mesangial cells to directly monitor changes in surface adhesion and stiffness of cells upon treatment with telmisartan in real time¹⁵⁾. The main finding of our study was the observation of an inhibition of the contractile response to Ang II in mesangial cells pre-treated with telmisartan, which was associated with alterations in cell stiffness¹⁵⁾. We also demonstrated that telmisartan-pretreated cells showed smaller pull-off forces than did the mesangial cells before treatment¹⁵⁾. Measuring the stiffness and adhesive properties of mesangial cells after telmisartan treatment allowed us to explore the roles of Ang II receptor blocker in structural and dynamic changes of mesangial cells. This approach could implicate several advantages in the *in vitro* experiments using cultured kidney cells.

The ability to image and study the surface of living cells is one of the important tools of AFM (Fig. 3). Fig. 3 shows a time series of topography and three dimensional images of live mesangial cells, showing the movement of mesangial cells at a 60×60 μm scan range. Images are taken before and 20 min after Ang II treatment. Cells moved toward the center. This reflects that Ang II induced contractions might lead to movement of mesangial cell. We could obtain the high quality images of live cells with AFM. However, it is difficult to demonstrate delicate cytoskeletal structures of the cell. With respect to AFM image, we found that the cytoskeletal elements of fixed cells




Fig. 3. Images of AFM of live mesangial cell (A) Deflation image before treatment and (B) at 20 min after Ang II treatment (C) Three dimensional image before treatment and (D) at 20 min after Ang II treatment.

were more prominent than live cells. We performed AFM imaging of fixed cells in a liquid solution to prevent dehydration and to maintain the mesangial cell structure (Fig. 4). The AFM images of fixed mesangial cells show the morphologic characteristics of mesangial cells, including the cytoskeletal structures of actin filaments, other filamentous elements and lamellar structures (Fig. 4). The morphologic characteristics of mesangial cells were well preserved without dehydration. No fixative artifacts were detected. The nucleus height of fixed MCs was $2.51 \pm 0.42 \mu\text{m}$ ($n=5$) from the bottom of the culture dish. A significant increase in height was observed with Ang II stimulation ($n=5$, $3.12 \pm 0.57 \mu\text{m}$). Pretreatment with telmisartan abolished the Ang II effects ($n=5$, $2.21 \pm 0.50 \mu\text{m}$).

Conclusion

Mechanical properties, stiffness, and adhesive force, which could be directly measured simultaneously using AFM in live cells can provide important information for elucidating the biological and physical mechanism of renal injury.




Fig. 4. Representative topography images (left panels) and deflection images (right panels) of fixed mesangial cells in liquid conditions: before angiotensin II (A), 20 min after Ang II (B) and 20 min after Ang II stimulation under telmisartan treatment (C).

References

1. Binnig G, Quate CF, Gerber C: Atomic force microscope. *Physical Review Letters* 56:930-933, 1986
2. Volle CB, Ferguson MA, Aidala KE, et al.: Quantitative changes in the elasticity and adhesive properties of escherichia coli zk1056 prey cells during predation by bdellovibrio bacteriovorus 109. *J Langmuir* 24:8102-8110, 2008
3. Radmacher M, Tillamnn RW, Fritz M, et al.: From molecules to cells: Imaging soft samples with the atomic force microscope. *Science (New York, NY)* 257:1900-1905, 1992
4. Radmacher M, Fritz M, Kacher CM, et al.: Measuring the viscoelastic properties of human platelets with the atomic force microscope. *Biophysical Journal* 70:556-567, 1996
5. Schneider SW, Yano Y, Sumpio BE, et al.: Rapid aldoster-

- one-induced cell volume increase of endothelial cells measured by the atomic force microscope. *Cell Biology International* 21:759-768, 1997
6. Oberleithner H: Aldosterone makes human endothelium stiff and vulnerable. *Kidney International* 67:1680-1682, 2005
 7. Oberleithner H, Riethmuller C, Ludwig T, et al.: Aldosterone remodels human endothelium. *Acta physiologica (Oxford, England)* 187:305-312, 2006
 8. Kliche K, Kuhn M, Hillebrand U, et al.: Direct aldosterone action on mouse cardiomyocytes detected with atomic force microscopy. *Cell Physiol Biochem* 18:265-274, 2006
 9. Klahr S, Morrissey J: Angiotensin ii and gene expression in the kidney. *Am J Kidney Dis* 31:171-176, 1998
 10. Mezzano SA, Ruiz-Ortega M, Egido J: Angiotensin ii and renal fibrosis. *Hypertension* 38:635-638, 2001
 11. Lee GJ, Park EJ, Choi S, et al.: Observation of angiotensin ii-induced changes in fixed and live mesangial cells by atomic force microscopy. *Micron* 41:220-226, 2010
 12. Auger-Messier M, Turgeon ES, Leduc R, et al.: The constitutively active n111g-at1 receptor for angiotensin ii modifies the morphology and cytoskeletal organization of hek-293 cells. *Experimental Cell Research* 308:188-195, 2005
 13. Cuerrier CM, Benoit M, Guillemette G, et al.: Real-time monitoring of angiotensin ii-induced contractile response and cytoskeleton remodeling in individual cells by atomic force microscopy. *Pflugers Arch* 457:1361-1372, 2009
 14. Hillebrand U, Lang D, Telgmann RG, et al.: Nebivolol decreases endothelial cell stiffness via the estrogen receptor beta: A nanoimaging study. *Journal of Hypertension* 27:517-526, 2009
 15. Jeong KH, Lee TW, Ihm CG, et al.: Real-Time Monitoring of the Effects of Telmisartan on Angiotensin II-Induced Mechanical Changes in Live Mesangial Cells Using Atomic Force Microscopy. *Kidney Blood Press Res* 35:573-582, 2012