

{4,4'-Dibromo-2,2'-[2,2-dimethylpropane-1,3-diylbis(nitrilomethylidyne)]-diphenolato- $\kappa^4 O, N, N', O'$ }copper(II)

Hadi Kargar,^a Hoong-Kun Fun^{b*} and Reza Kia^b

^aDepartment of Chemistry, School of Science, Payame Noor University (PNU), Ardakan, Yazd, Iran, and ^bX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia
Correspondence e-mail: hkfun@usm.my


Received 20 October 2008; accepted 7 November 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.033; wR factor = 0.093; data-to-parameter ratio = 23.0.

In the title compound, $[\text{Cu}(\text{C}_{19}\text{H}_{18}\text{Br}_2\text{N}_2\text{O}_2)]$, the Cu^{II} ion is in a tetrahedrally distorted planar geometry, involving two N and two O atoms from the tetradentate Schiff base ligand. Intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds form an eight-membered $R_2^2(8)$ motif. The dihedral angle between two benzene rings is $36.34(9)^\circ$. There are intermolecular $\text{Cu}\cdots\text{Br}$ [$3.4566(5)$ Å] and $\text{Cu}\cdots\text{N}$ [$3.569(3)$ Å] contacts, which are significantly shorter than the sum of van der Waals radii of the relevant atoms. These interactions, along with the intermolecular $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ [centroid-centroid distances of $3.709(1)$ and $3.968(2)$ Å] interactions, link neighbouring molecules into a one-dimensional infinite chain along the c axis.

Related literature

For bond-length data, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For values of van der Waals radii, see: Bondi (1964). For related structures, see: Arıcı *et al.* (2001); Elmali *et al.* (2000); Hodgson (1975); Granovski *et al.* (1993). For the application of transition-metal complexes with Schiff base ligands, see: Blower (1998); Shahrokhian *et al.* (2000).



Experimental

Crystal data

$[\text{Cu}(\text{C}_{19}\text{H}_{18}\text{Br}_2\text{N}_2\text{O}_2)]$
 $M_r = 529.71$
 Triclinic, $P\bar{1}$
 $a = 9.1416(3)$ Å
 $b = 9.6398(3)$ Å
 $c = 11.5382(3)$ Å
 $\alpha = 75.210(2)^\circ$
 $\beta = 78.913(2)^\circ$
 $\gamma = 73.435(2)^\circ$
 $V = 934.42(5)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 5.46$ mm⁻¹
 $T = 100.0(1)$ K
 $0.41 \times 0.21 \times 0.15$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\text{min}} = 0.195$, $T_{\text{max}} = 0.443$
 29164 measured reflections
 5410 independent reflections
 4345 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.093$
 $S = 1.07$
 5410 reflections
 235 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.27$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.61$ e Å⁻³

Table 1

Selected bond lengths (Å).

Cu1—O2	1.9027 (19)	Cu1—N1	1.948 (2)
Cu1—O1	1.9146 (18)	Cu1—N2	1.955 (2)

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C16}-\text{H16A}\cdots\text{O2}^{\text{i}}$	0.93	2.44	3.342 (3)	163
$\text{C10}-\text{H10B}\cdots\text{Cg1}^{\text{ii}}$	0.97	2.50	3.324 (3)	142

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, -y, -z + 1$. Cg1 is the centroid of the Cu1, N2, O2, C11, C12, C17 ring.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2003).

HKF and RK thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/PFIZIK/613312. RK thanks Universiti Sains Malaysia for a post-doctoral research fellowship. HK thanks PNU for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2161).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Arci, C., Ercan, F., Kurtaran, R. & Atakol, O. (2001). *Acta Cryst. C* **57**, 812–814.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Blower, P. J. (1998). *Transition Met. Chem.* **23**, 109–112.
- Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Elmali, A., Zeyrek, C. T., Elerman, Y. & Svoboda, I. (2000). *Acta Cryst. C* **56**, 1302–1304.
- Granovski, A. D., Nivorozhkin, A. L. & Minkin, V. I. (1993). *Coord. Chem. Rev.* **126**, 1–69.
- Hodgson, D. J. (1975). *Prog. Inorg. Chem.* **19**, 173–202.
- Shahrokhian, S., Amini, M. K., Kia, R. & Tangestaninejad, S. (2000). *Anal. Chem.* **72**, 956–962.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, m1541-m1542 [doi:10.1107/S1600536808036635]

{4,4'-Dibromo-2,2'-[2,2-dimethylpropane-1,3-diylbis(nitrilomethylidyne)]diphenolato- κ^4 O,N,N',O'}copper(II)

H. Kargar, H.-K. Fun and R. Kia

Comment

Schiff base complexes are some of the most important stereochemical models in transition metal coordination chemistry, with their ease of preparation and structural variations (Elmali *et al.*, 2000; Granovski *et al.*, 1993). Transition metal complexes of Schiff base ligands are always of interest since they exhibit a marked tendency to oligomerize, thus leading to novel structural types, and also display a wide variety of magnetic properties (Blower, 1998; Shahrokhian *et al.*, 2000). Many of the reported structural investigations of these complexes are discussed in some details in a review (Hodgson, 1975). Tetradentate Schiff base metal complexes may form *trans* or *cis* planar or tetrahedral structures (Elmali *et al.*, 2000).

In the title compound (Fig. 1), the Cu^{II} ion shows a planar geometry distorted towards tetrahedral, which is defined by two imine N atoms and two phenolate O atoms of the tetradentate Schiff base ligand (Table 1). Intermolecular C—H \cdots O hydrogen bonds form an eight-membered ring $R_2^2(8)$ motif (Fig. 2) (Bernstein *et al.*, 1995). The bond lengths are within the normal ranges (Allen *et al.*, 1987) and are comparable with the related structure (Arici *et al.*, 2001). The dihedral angle between two benzene rings is 36.34 (9)°. The chelate ring composed of Cu1, N1, C8, C9, C10 and N2 atoms has a distorted boat conformation with puckering parameters of $Q = 0.807(3)$ Å, $\Theta = 91.1(2)$ ° and $\Phi = 264.58(17)$ °. The interesting feature of the crystal structure is short intermolecular Cu1 \cdots Br1ⁱⁱⁱ [3.4566(5) Å] and Cu1 \cdots N2ⁱⁱ [3.569(3) Å] interactions [symmetry codes: (ii) 1 - x, -y, 1 - z; (iii) 1 - x, -y, 2 - z], which are significantly shorter than the sum of van der Waals radii of the relevant atoms [Cu: 2.32; Br: 1.85; N: 1.55 Å (Bondi, 1964; Spek, 2003)]. These interactions along with the intermolecular C—H \cdots π (Table 2) and π - π interactions [centroid-centroid distances: Cg2 \cdots Cg3ⁱⁱⁱ = 3.709(1) and Cg2 \cdots Cg2ⁱⁱⁱ = 3.968(2) Å; Cg2 = centroid of the C1–C6 ring and Cg3 = centroid of the Cu1, N1, O1, C1, C6, C7 ring] link the neighbouring molecules into one-dimensional infinite chains along the *c* axis (Fig. 3).

Experimental

The title compound was prepared based on the reported method (Arici *et al.*, 2001). Single crystals suitable for X-ray analysis were obtained from an ethanol solution at room temperature.

Refinement

H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93 (aromatic), 0.97 (CH₂) and 0.96 (CH₃) Å and $U_{iso}(H) = 1.2$ (1.5 for methyl groups) $U_{eq}(C)$. The highest difference peak is located 0.81 Å from Br2 and the deepest hole is located 0.76 Å from Cu1.

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids.




Fig. 2. The crystal packing of the title compound, viewed down the *a* axis, showing the hydrogen-bond motif $R_2^2(8)$.




Fig. 3. The crystal packing of the title compound, viewed down the *b* axis, showing one-dimensional infinite chains along the *c* axis. Intermolecular $\text{Cu}\cdots\text{Br}$ and $\text{Cu}\cdots\text{N}$ interactions are shown as dashed lines.

{4,4'-Dibromo-2,2'-[2,2-dimethylpropane-1,3-diylbis(nitrilomethylidene)]diphenolato- $\kappa^4\text{O},\text{N},\text{N}',\text{O}'$ }copper(II)

Crystal data

$[\text{Cu}(\text{C}_{19}\text{H}_{18}\text{Br}_2\text{N}_2\text{O}_2)]$

$M_r = 529.71$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.1416(3) \text{ \AA}$

$b = 9.6398(3) \text{ \AA}$

$c = 11.5382(3) \text{ \AA}$

$\alpha = 75.210(2)^\circ$

$\beta = 78.913(2)^\circ$

$\gamma = 73.435(2)^\circ$

$V = 934.42(5) \text{ \AA}^3$

$Z = 2$

$F_{000} = 522$

$D_x = 1.883 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9921 reflections

$\theta = 2.2\text{--}33.8^\circ$

$\mu = 5.46 \text{ mm}^{-1}$

$T = 100.0(1) \text{ K}$

Block, red

$0.41 \times 0.21 \times 0.15 \text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100.0(1) \text{ K}$

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.195$, $T_{\max} = 0.443$

29164 measured reflections

5410 independent reflections

4345 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

$\theta_{\max} = 30.0^\circ$

$\theta_{\min} = 1.8^\circ$

$h = -12 \rightarrow 12$

$k = -13 \rightarrow 13$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.033$	H-atom parameters constrained
$wR(F^2) = 0.093$	$w = 1/[\sigma^2(F_o^2) + (0.0524P)^2 + 0.5947P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
5410 reflections	$(\Delta/\sigma)_{\max} = 0.002$
235 parameters	$\Delta\rho_{\max} = 1.27 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.61 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Experimental. The low-temperature data was collected with the Oxford Cyrosystem Cobra low-temperature attachment.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.60215 (4)	0.07295 (3)	0.63222 (3)	0.01866 (8)
Br1	0.15151 (3)	-0.15996 (3)	1.22034 (2)	0.02498 (8)
Br2	0.80441 (4)	0.43866 (3)	-0.00230 (2)	0.02942 (9)
O1	0.4262 (2)	0.1481 (2)	0.73927 (16)	0.0210 (4)
O2	0.5974 (2)	0.2636 (2)	0.53114 (17)	0.0226 (4)
N1	0.6688 (3)	-0.1090 (2)	0.7503 (2)	0.0195 (4)
N2	0.6996 (3)	-0.0301 (2)	0.5008 (2)	0.0193 (4)
C1	0.3696 (3)	0.0739 (3)	0.8409 (2)	0.0187 (5)
C2	0.2233 (3)	0.1404 (3)	0.8980 (2)	0.0216 (5)
H2A	0.1701	0.2336	0.8604	0.026*
C3	0.1577 (3)	0.0712 (3)	1.0073 (2)	0.0222 (5)
H3A	0.0617	0.1176	1.0423	0.027*
C4	0.2363 (3)	-0.0692 (3)	1.0654 (2)	0.0209 (5)
C5	0.3765 (3)	-0.1391 (3)	1.0130 (2)	0.0202 (5)
H5A	0.4274	-0.2325	1.0520	0.024*
C6	0.4441 (3)	-0.0708 (3)	0.9004 (2)	0.0182 (5)
C7	0.5935 (3)	-0.1509 (3)	0.8535 (2)	0.0195 (5)
H7A	0.6390	-0.2402	0.9019	0.023*
C8	0.8231 (3)	-0.1986 (3)	0.7167 (3)	0.0220 (5)
H8A	0.8583	-0.2707	0.7877	0.026*
H8B	0.8935	-0.1347	0.6889	0.026*
C9	0.8276 (3)	-0.2804 (3)	0.6167 (2)	0.0215 (5)
C10	0.7088 (3)	-0.1897 (3)	0.5308 (2)	0.0208 (5)
H10A	0.7350	-0.2262	0.4566	0.025*
H10B	0.6084	-0.2049	0.5677	0.025*
C11	0.7379 (3)	0.0315 (3)	0.3905 (2)	0.0191 (5)

supplementary materials

H11A	0.7818	-0.0304	0.3364	0.023*
C12	0.7183 (3)	0.1880 (3)	0.3441 (2)	0.0184 (5)
C13	0.7662 (3)	0.2337 (3)	0.2195 (2)	0.0200 (5)
H13A	0.8138	0.1630	0.1734	0.024*
C14	0.7432 (3)	0.3807 (3)	0.1663 (2)	0.0206 (5)
C15	0.6701 (3)	0.4888 (3)	0.2346 (2)	0.0221 (5)
H15A	0.6534	0.5887	0.1975	0.026*
C16	0.6231 (3)	0.4473 (3)	0.3559 (2)	0.0221 (5)
H16A	0.5755	0.5202	0.3999	0.027*
C17	0.6454 (3)	0.2960 (3)	0.4160 (2)	0.0193 (5)
C18	0.7856 (3)	-0.4283 (3)	0.6727 (3)	0.0255 (6)
H18A	0.7886	-0.4782	0.6098	0.038*
H18B	0.6840	-0.4106	0.7165	0.038*
H18C	0.8580	-0.4889	0.7268	0.038*
C19	0.9898 (3)	-0.3061 (3)	0.5469 (3)	0.0267 (6)
H19A	0.9944	-0.3565	0.4841	0.040*
H19B	1.0627	-0.3653	0.6011	0.040*
H19C	1.0142	-0.2124	0.5118	0.040*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.02588 (17)	0.01331 (15)	0.01503 (15)	-0.00383 (12)	-0.00011 (12)	-0.00300 (12)
Br1	0.03447 (15)	0.02289 (14)	0.01814 (13)	-0.01260 (11)	0.00257 (10)	-0.00364 (10)
Br2	0.04858 (18)	0.01898 (14)	0.01683 (13)	-0.00907 (12)	0.00283 (11)	-0.00148 (10)
O1	0.0275 (9)	0.0161 (9)	0.0166 (9)	-0.0040 (7)	0.0010 (7)	-0.0029 (7)
O2	0.0316 (10)	0.0154 (9)	0.0182 (9)	-0.0037 (8)	0.0001 (7)	-0.0036 (7)
N1	0.0240 (11)	0.0159 (10)	0.0180 (10)	-0.0030 (8)	-0.0029 (8)	-0.0047 (9)
N2	0.0282 (11)	0.0131 (10)	0.0166 (10)	-0.0057 (8)	-0.0023 (8)	-0.0032 (8)
C1	0.0268 (13)	0.0156 (11)	0.0162 (11)	-0.0069 (10)	-0.0036 (9)	-0.0055 (9)
C2	0.0268 (13)	0.0164 (12)	0.0211 (13)	-0.0055 (10)	-0.0031 (10)	-0.0031 (10)
C3	0.0245 (13)	0.0219 (13)	0.0217 (13)	-0.0070 (10)	-0.0001 (10)	-0.0080 (11)
C4	0.0287 (13)	0.0198 (12)	0.0161 (12)	-0.0103 (10)	-0.0011 (10)	-0.0037 (10)
C5	0.0286 (13)	0.0170 (12)	0.0160 (12)	-0.0079 (10)	-0.0029 (10)	-0.0027 (10)
C6	0.0251 (12)	0.0154 (11)	0.0155 (11)	-0.0062 (9)	-0.0023 (9)	-0.0045 (10)
C7	0.0258 (13)	0.0154 (11)	0.0171 (12)	-0.0039 (10)	-0.0046 (10)	-0.0028 (10)
C8	0.0236 (12)	0.0184 (12)	0.0230 (13)	-0.0033 (10)	-0.0027 (10)	-0.0048 (11)
C9	0.0262 (13)	0.0162 (12)	0.0213 (13)	-0.0040 (10)	-0.0029 (10)	-0.0042 (10)
C10	0.0302 (13)	0.0137 (11)	0.0185 (12)	-0.0066 (10)	-0.0011 (10)	-0.0036 (10)
C11	0.0256 (12)	0.0147 (11)	0.0170 (12)	-0.0049 (9)	-0.0026 (9)	-0.0037 (9)
C12	0.0227 (12)	0.0152 (11)	0.0175 (12)	-0.0057 (9)	-0.0010 (9)	-0.0040 (10)
C13	0.0246 (12)	0.0187 (12)	0.0173 (12)	-0.0057 (10)	-0.0012 (10)	-0.0056 (10)
C14	0.0272 (13)	0.0188 (12)	0.0158 (12)	-0.0082 (10)	-0.0014 (10)	-0.0021 (10)
C15	0.0303 (14)	0.0137 (11)	0.0212 (13)	-0.0059 (10)	-0.0039 (10)	-0.0011 (10)
C16	0.0289 (13)	0.0147 (12)	0.0217 (13)	-0.0043 (10)	0.0001 (10)	-0.0056 (10)
C17	0.0232 (12)	0.0162 (11)	0.0181 (12)	-0.0052 (9)	-0.0013 (9)	-0.0038 (10)
C18	0.0361 (15)	0.0163 (12)	0.0230 (13)	-0.0064 (11)	-0.0042 (11)	-0.0020 (11)
C19	0.0269 (14)	0.0234 (14)	0.0287 (15)	-0.0048 (11)	0.0008 (11)	-0.0083 (12)

Geometric parameters (Å, °)

Cu1—O2	1.9027 (19)	C8—H8A	0.9700
Cu1—O1	1.9146 (18)	C8—H8B	0.9700
Cu1—N1	1.948 (2)	C9—C18	1.530 (4)
Cu1—N2	1.955 (2)	C9—C19	1.531 (4)
Br1—C4	1.902 (3)	C9—C10	1.535 (4)
Br2—C14	1.901 (3)	C10—H10A	0.9700
O1—C1	1.305 (3)	C10—H10B	0.9700
O2—C17	1.303 (3)	C11—C12	1.437 (4)
N1—C7	1.286 (3)	C11—H11A	0.9300
N1—C8	1.467 (3)	C12—C13	1.413 (4)
N2—C11	1.287 (3)	C12—C17	1.432 (3)
N2—C10	1.470 (3)	C13—C14	1.366 (4)
C1—C2	1.422 (4)	C13—H13A	0.9300
C1—C6	1.425 (4)	C14—C15	1.405 (4)
C2—C3	1.379 (4)	C15—C16	1.373 (4)
C2—H2A	0.9300	C15—H15A	0.9300
C3—C4	1.402 (4)	C16—C17	1.420 (4)
C3—H3A	0.9300	C16—H16A	0.9300
C4—C5	1.371 (4)	C18—H18A	0.9600
C5—C6	1.411 (4)	C18—H18B	0.9600
C5—H5A	0.9300	C18—H18C	0.9600
C6—C7	1.442 (4)	C19—H19A	0.9600
C7—H7A	0.9300	C19—H19B	0.9600
C8—C9	1.544 (4)	C19—H19C	0.9600
O2—Cu1—O1	92.77 (8)	C19—C9—C10	110.3 (2)
O2—Cu1—N1	160.11 (9)	C18—C9—C8	110.1 (2)
O1—Cu1—N1	93.32 (9)	C19—C9—C8	108.4 (2)
O2—Cu1—N2	93.40 (8)	C10—C9—C8	110.7 (2)
O1—Cu1—N2	151.78 (9)	N2—C10—C9	113.6 (2)
N1—Cu1—N2	90.14 (9)	N2—C10—H10A	108.8
C1—O1—Cu1	126.57 (17)	C9—C10—H10A	108.8
C17—O2—Cu1	128.01 (16)	N2—C10—H10B	108.8
C7—N1—C8	119.4 (2)	C9—C10—H10B	108.8
C7—N1—Cu1	125.97 (19)	H10A—C10—H10B	107.7
C8—N1—Cu1	114.58 (17)	N2—C11—C12	125.4 (2)
C11—N2—C10	118.7 (2)	N2—C11—H11A	117.3
C11—N2—Cu1	125.90 (18)	C12—C11—H11A	117.3
C10—N2—Cu1	114.85 (16)	C13—C12—C17	120.1 (2)
O1—C1—C2	118.6 (2)	C13—C12—C11	116.7 (2)
O1—C1—C6	124.7 (2)	C17—C12—C11	123.1 (2)
C2—C1—C6	116.7 (2)	C14—C13—C12	120.7 (2)
C3—C2—C1	122.2 (3)	C14—C13—H13A	119.7
C3—C2—H2A	118.9	C12—C13—H13A	119.7
C1—C2—H2A	118.9	C13—C14—C15	120.3 (2)
C2—C3—C4	119.7 (2)	C13—C14—Br2	119.65 (19)
C2—C3—H3A	120.1	C15—C14—Br2	120.0 (2)

supplementary materials

C4—C3—H3A	120.1	C16—C15—C14	120.2 (2)
C5—C4—C3	120.3 (2)	C16—C15—H15A	119.9
C5—C4—Br1	119.8 (2)	C14—C15—H15A	119.9
C3—C4—Br1	119.8 (2)	C15—C16—C17	121.8 (2)
C4—C5—C6	120.7 (3)	C15—C16—H16A	119.1
C4—C5—H5A	119.7	C17—C16—H16A	119.1
C6—C5—H5A	119.7	O2—C17—C16	118.9 (2)
C5—C6—C1	120.4 (2)	O2—C17—C12	124.1 (2)
C5—C6—C7	116.8 (2)	C16—C17—C12	117.0 (2)
C1—C6—C7	122.7 (2)	C9—C18—H18A	109.5
N1—C7—C6	125.3 (2)	C9—C18—H18B	109.5
N1—C7—H7A	117.3	H18A—C18—H18B	109.5
C6—C7—H7A	117.3	C9—C18—H18C	109.5
N1—C8—C9	112.7 (2)	H18A—C18—H18C	109.5
N1—C8—H8A	109.0	H18B—C18—H18C	109.5
C9—C8—H8A	109.0	C9—C19—H19A	109.5
N1—C8—H8B	109.0	C9—C19—H19B	109.5
C9—C8—H8B	109.0	H19A—C19—H19B	109.5
H8A—C8—H8B	107.8	C9—C19—H19C	109.5
C18—C9—C19	110.4 (2)	H19A—C19—H19C	109.5
C18—C9—C10	106.9 (2)	H19B—C19—H19C	109.5
O2—Cu1—O1—C1	-174.0 (2)	C8—N1—C7—C6	177.2 (2)
N1—Cu1—O1—C1	-12.9 (2)	Cu1—N1—C7—C6	0.6 (4)
N2—Cu1—O1—C1	83.6 (3)	C5—C6—C7—N1	176.6 (2)
O1—Cu1—O2—C17	-151.9 (2)	C1—C6—C7—N1	-6.7 (4)
N1—Cu1—O2—C17	100.4 (3)	C7—N1—C8—C9	108.8 (3)
N2—Cu1—O2—C17	0.6 (2)	Cu1—N1—C8—C9	-74.2 (2)
O2—Cu1—N1—C7	114.8 (3)	N1—C8—C9—C18	-86.0 (3)
O1—Cu1—N1—C7	7.2 (2)	N1—C8—C9—C19	153.1 (2)
N2—Cu1—N1—C7	-144.8 (2)	N1—C8—C9—C10	32.0 (3)
O2—Cu1—N1—C8	-61.9 (3)	C11—N2—C10—C9	115.7 (3)
O1—Cu1—N1—C8	-169.52 (17)	Cu1—N2—C10—C9	-72.2 (2)
N2—Cu1—N1—C8	38.50 (18)	C18—C9—C10—N2	160.8 (2)
O2—Cu1—N2—C11	0.1 (2)	C19—C9—C10—N2	-79.2 (3)
O1—Cu1—N2—C11	102.4 (3)	C8—C9—C10—N2	40.8 (3)
N1—Cu1—N2—C11	-160.3 (2)	C10—N2—C11—C12	172.3 (2)
O2—Cu1—N2—C10	-171.27 (18)	Cu1—N2—C11—C12	1.2 (4)
O1—Cu1—N2—C10	-69.0 (3)	N2—C11—C12—C13	-179.2 (3)
N1—Cu1—N2—C10	28.31 (19)	N2—C11—C12—C17	-3.2 (4)
Cu1—O1—C1—C2	-169.15 (18)	C17—C12—C13—C14	-0.2 (4)
Cu1—O1—C1—C6	11.1 (4)	C11—C12—C13—C14	175.9 (2)
O1—C1—C2—C3	-178.0 (2)	C12—C13—C14—C15	-0.6 (4)
C6—C1—C2—C3	1.9 (4)	C12—C13—C14—Br2	-178.50 (19)
C1—C2—C3—C4	0.1 (4)	C13—C14—C15—C16	0.9 (4)
C2—C3—C4—C5	-1.3 (4)	Br2—C14—C15—C16	178.8 (2)
C2—C3—C4—Br1	176.5 (2)	C14—C15—C16—C17	-0.4 (4)
C3—C4—C5—C6	0.4 (4)	Cu1—O2—C17—C16	176.52 (18)
Br1—C4—C5—C6	-177.39 (19)	Cu1—O2—C17—C12	-2.5 (4)
C4—C5—C6—C1	1.7 (4)	C15—C16—C17—O2	-179.5 (2)

C4—C5—C6—C7	178.5 (2)	C15—C16—C17—C12	-0.4 (4)
O1—C1—C6—C5	177.1 (2)	C13—C12—C17—O2	179.8 (2)
C2—C1—C6—C5	-2.7 (4)	C11—C12—C17—O2	3.9 (4)
O1—C1—C6—C7	0.5 (4)	C13—C12—C17—C16	0.7 (4)
C2—C1—C6—C7	-179.3 (2)	C11—C12—C17—C16	-175.2 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C16—H16A...O2 ⁱ	0.93	2.44	3.342 (3)	163
C10—H10B...Cg1 ⁱⁱ	0.97	2.50	3.324 (3)	142

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, -y, -z+1$.

Fig. 1




Fig. 2




Fig. 3

