

Factors Affecting Cardiopulmonary Resuscitation Practice Among Nurses in Mizan Tepi University Teaching Hospital, Tepi General Hospital, and Gebretsadik Shawo Hospital, Southwest Ethiopia

Mirresa Guteta 

Nursing Department, College of Medicine and Health Science, Mizan-Tepi University, Mizan-Aman, Ethiopia

Correspondence: Mirresa Guteta, Email mirresag143@gmail.com

Background: Cardiopulmonary resuscitation is one of the basic lifesaving medical procedures which is performed for a patient with an emergency such as cardiac arrest, suffocation, near-dying, or circumstance that results in cardiac or pulmonary failure or both.

Objective: To assess factors affecting the practice of cardiopulmonary resuscitation among nurses at Mizan Tepi University Teaching Hospital, Tepi General Hospital, and Gebretsadik Shawo Hospital.

Methods: Institution-based cross-sectional study which was supported by qualitative data conducted from March 20 to April 20, 2021. Nurses who answered ≥ 7 CPR practice questions correctly were considered to have good practice. Data were entered into EpiData version 4.4 and exported to SPSS version 23 for data analysis. A multivariable logistic regression model was fitted, adjusted odds ratio (AOR) at 95% confidence interval and p-value < 0.05 were estimated to determine the statistically significant association between predictors and outcome variable. Qualitative data were analyzed based on thematic content.

Results: The prevalence of good practice towards CPR was 31.8% (95% CI: 27.5–36.3) in the study area. Experience of 6–10 years (AOR = 2.27, 95% CI: 1.25–4.13) and > 10 years (AOR = 1.81, 95% CI: 1.10–2.98), rarely (AOR = 3.77, 95% CI: 1.26–11.30) or actively (AOR = 4.60, 95% CI: 1.51–13.98) involved in CPR practice, assigned to emergency department (AOR = 1.02, 95% CI: 0.55–1.90), having CPR good knowledge (AOR = 1.37, 95% CI: 0.28–2.14) and having a nursing degree (AOR = 1.54, 95% CI: 0.93–2.54) were predictors of CPR good practice.

Conclusion: Nurses' good practice towards cardiopulmonary resuscitation was low. Therefore, efforts should be taken to enhance nurses' practice towards cardiopulmonary resuscitation.

Keywords: practice, nurse, cardiopulmonary resuscitation, associated factors

Introduction

Cardiopulmonary resuscitation (CPR) is an organized medical maneuver through which chest compressions and artificial breathing are given to provide sufficient blood perfusion towards major organs including the brain and heart. Cardiopulmonary resuscitation can be a life-saving procedure performed for a patient in an emergency such as cardiac arrest, suffocation, near-drowning, electrocution injuries, or any condition in which a victim's respiration or circulation has stopped.^{1–3}

Health professionals including nurses working in the hospital can reduce mortality resulting from cardiac and respiratory-related problems when they are equipped with the knowledge and skill of cardiopulmonary resuscitation. As a result, nurses should be equipped with the knowledge and skill of basic life support to avert critical medical-surgical emergencies.^{4–6}

Appropriate CPR is crucial for the survival of cardiopulmonary arrest sufferers. The chance of survival in cardiopulmonary arrest is reduced by 7–10% for each minute of CPR delay. Additionally, CPR can decrease the length of

hospital stay. One of the indications for CPR in cardiac arrest is caused by failure of the blood circulation due to a cardiac problem.⁷⁻⁹

For the better outcome of the patients, applying CPR early is most effective. Providing frequently updated training about cardiopulmonary resuscitation for emergency care health providers can improve the survival rate of the patients.^{10,11}

Among health professionals working in a health-care setting, nurses are the ones who face problems that need CPR procedure and perform the procedure. Even though CPR is among the important of procedures for saving the life of patients, studies depict that the skill of nurses regarding the procedure is deficient. To perform CPR procedures properly the nurse is expected to have updated knowledge and skill of cardiopulmonary resuscitation.^{12,13}

Various international and country-based verified protocols for CPR are endorsed and official training programs depending on these guidelines have been implemented by certified training centers.^{14,15}

Cardiopulmonary resuscitation is a rescue intervention, executed to maintain survival of patients until additional management can be given.¹⁶⁻¹⁹ Based on different studies nurses have good knowledge and a positive attitude toward CPR, but they have a gap in practice.^{4,5}

The objective of this study was to assess factors affecting the application of CPR among nurses working in the selected three hospitals.

Materials and Methods

Study Area and Period

The study was carried out in Mizan Tepi University Teaching Hospital (MTUTH), Tepi General Hospital, and Gebretsadik Shawo Hospital located in Bench Sheko, Sheka, and Kefa Zones from March 20 to April 20, 2021. MTUTH is in Bech Sheko Zone 560 km distant from the capital city of Ethiopia, Addis Ababa; Tepi General Hospital is in Sheka Zone, 565 km away from Addis Ababa; and Gebretsadik Shawo Hospital is located in Kefa Zone which is 411 km away from Addis Ababa. The three hospitals provide surgical, gynecological and obstetric, general medical, pediatrics, minor and major operations, ophthalmologic and diagnostic services. They have different health professionals in different fields of study and qualifications.

Study design: Institution-based cross-sectional study design which included both qualitative and quantitative data collection methods was used.

Inclusion and exclusion criteria: All nurses' working in the three hospitals were incorporated in the study whereas those nurses on permitted leave or assigned to the hospital for less than 6 months were excluded from the study.

Sample size and sampling procedure: All available nurses in the three hospitals during the data collection period were incorporated in the study by using a census (Figure 1).

Operational Definition

CPR knowledge: Cardiopulmonary resuscitation knowledge questions comprise 12 questions which were coded as 1 for the right answer and 0 for the wrong answer by SPSS. The maximum and minimum knowledge scores were 12 and 0; knowledge scores were categorized whereby scores from 9–12 were grouped as good knowledge while scores from 0–8 were grouped as poor knowledge.

CPR practice: Cardiopulmonary resuscitation practice questions comprise 9 questions which were coded as 1 for right answer and 0 for the wrong answer. The maximum and minimum practice scores were 9 and 0, practice scores were categorized whereby scores from 7–9 were grouped as good practice while scores from 0–6 were grouped as poor practice.

CPR training: Nurses who have taken CPR training for the last year are counted as having regular training.

Work experience: Total years of experience after nurses graduate and start working.

CPR involvement: Those nurses who frequently participated in the CPR procedure were considered as actively involved, those involved seldom were considered as rarely involved, and those not involved at all were considered as never involved.




Figure 1 Schematic presentation of sampling procedure in MTUTH, Tepi General Hospital and Gebretsadik Shawa Hospital, Southwest Ethiopia. **Abbreviations:** CPR, cardiopulmonary resuscitation; MTUTH, Mizan Tepi University Teaching Hospital.

Data Collection instrument: The data were gathered through a pretested and organized tool which was adopted from other similar studies done before, and the English version was used. The tool consists of three parts. Part I: socio-demographic characteristics comprising 7 items. Part II: Cardiopulmonary resuscitation knowledge questions comprising 12 questions which were coded as 1 for the right answer and 0 for the wrong answer. The maximum and minimum knowledge scores were 12 and 0, knowledge scores were categorized whereby scores from 9–12 were grouped as good knowledge while scores from 0–8 were grouped as poor knowledge. Part III: Cardiopulmonary resuscitation practice questions comprising 9 questions which were coded as 1 for right answer and 0 for the wrong answer. The maximum and minimum practice scores were 9 and 0, practice scores were categorized whereby scores from 7–9 were grouped as good practice while scores from 0–6 were grouped as poor practice. Data collection was undertaken by six data collectors and three supervisors.

Data Collection technique: The data were gathered through an interviewer-administered tool. Qualitative data collection was conducted by the researcher himself using a checklist.

Data Quality Assurance

Before conducting actual data collection, the tool was pretested on 5% of samples at Chena primary hospital. The pretest aimed to assure the study participants can understand the questions and to check the sequence of the tool. Based on the result of pretest mitigation was done. The training was given for data collectors and supervisors on the intention of the study, approach to contact with study participants, and data collection techniques. Completeness, precision and coherence of data were verified by supervisors who also provided feedback accordingly throughout data gathering time. Finally, data were coded, cleaned, entered, and checked for completeness before analysis.

Processing and Analysis of the Data

EpiData version 4.4 was used for data entry and SPSS version 23 was utilized for analysis. Data exploration was conducted to examine different characteristics of the data and descriptive statistics were used to describe the data depending on its nature. After cleaning of the data, descriptive statistics such as frequencies, proportions, and percentages were done.

After checking all assumptions of logistic regression such as independent observations (multicollinearity among independent variables, linearity of independent variables, and log odds of the dependent variable), bivariable logistic regression was carried out to select a candidate for multivariate logistic regression analysis with a p-value <0.25 at 95% confidence level. Then, candidate variables were entered into a multivariable logistic regressions model using the backward method to identify the statistically significant factors for cardiopulmonary resuscitation by controlling possible confounders. The degree of association between dependent and independent variables was assessed using odds ratio and statistically significant factors were declared at 95% confidence interval and a p-value <0.05.

Computation

The multiple-choice questions which assess the knowledge were changed into yes and no which stand for correct and incorrect answers respectively. This choice question which was assessing the practice was also transformed into yes and no to represent good practice and poor practice respectively.

Results

Demographic and Socio-Economic Characteristics

In this study, 422 nurses working in the selected three hospitals participated and yielded a 100% response rate. Among these 250 (59.2%) were female and 172 (40.8%) were male. Regarding marital status and age of the study participants, 276 (65.4%) were married and 289 (68.5%) of them were aged between 26–30 years. Out of the total, 218 (51.6%) had 6–10 years of working experience. Regarding their qualifications and working unit, 318 (75.5%) had degrees and 282 (66.8%) worked on inpatient wards (Table 1).

Knowledge of Nurses Towards CPR

In this study, the level of good knowledge about cardiopulmonary resuscitation was found to be 70.2% among nurses who work in MTUTH, Tepi General Hospital, and Gebretsadik Shawo Hospital. More than half of the respondents

Table 1 Demographic and Socio-Economic Characteristics of Nurses Working in MTUTH, Tepi General Hospital and GSH, Southwest Ethiopia

Variable		Frequency	Percentage (%)
Gender	Female	250	59.2
	Male	172	40.8
Marital status	Married	276	65.4
	Single	146	34.6
Age in years	20–25	74	17.5
	26–30	289	68.5
	31–35	45	10.7
	>36	14	3.3
Service year	1–5	98	23.3
	6–10	218	51.6
	>10	106	25.1
Qualification	Degree	318	75.5
	Diploma	104	24.6
Taking recent training <3 year	Yes	146	34.6
	No	276	65.4
Setting	OPD	59	14.0
	Emergency	81	19.2
	Ward	282	66.8

Abbreviations: OPD, outpatient department; GSH, Gebretsadik Shawo Hospital; MTUTH, Mizan Tepi University Teaching Hospital.

responded with the right answer to questions regarding when CPR is most effective, the right way to open the airway before providing mouth-to-mouth respiration, the chance of survival of a victim when CPR is given properly, rescue breaths per cycle, depth of compression and correct placement of hand on the sternum (Table 2).

The Practice of Nurses Towards CPR

The prevalence of good practice towards CPR among nurses working in the selected three hospitals was 31.8% (95% CI: 27.5–36.3). The study revealed that 35.7% of the participants had performed cardiopulmonary resuscitation on a patient in cardiac emergencies, 62.9% of the participants checked a patient's pulse rate before commencing CPR, and 53.6% ensured that the patient was in a supine position on a relatively hard surface before commencing cardiopulmonary resuscitation (Table 3).

Table 2 Participants' Knowledge Response About CPR in MTUTH, TGH, and GSH, SNNPR, Southwest f Ethiopia

S.No	Variable	Categories	Frequency	Percentage (%)
1	CPR is most effective when initiated immediately after the patient's collapse	Yes	389	92.1
		No	33	7.9
2	The correct compression to respiration ratio 3:2	Yes	106	25.0
		No	316	75.0
3	Victims are to be given another rescue breath if the chest does not rise after giving the first rescue breath	Yes	111	26.4
		No	311	73.6
4	The right chest compression to be performed per minute when giving CPR is 100	Yes	135	32.1
		No	287	67.9
5	The recommended way to open the airway before giving mouth-to-mouth respiration is to tilt the head back and lift the chin up	Yes	374	88.6
		No	48	11.4
6	The chest compression landmark on an adult is at the center of the chest	Yes	78	18.6
		No	344	81.4
7	Chest compressions should be performed on an infant with two fingers of one hand while doing CPR	Yes	187	44.3
		No	235	55.7
8	The steps of CPR in the correct sequence is compression, preserve patent airway, and artificial breathing	Yes	87	20.7
		No	335	79.3
9	One should give rescue breath at a rate of 2 times per cycle	Yes	259	61.4
		No	163	38.6
10	How much depth you should compress the chest in adult up to 2 inch	Yes	226	53.6
		No	196	46.4
11	While you are placing the other hand on top of the hand on the sternum the finger should be interlocked	Yes	241	57.1
		No	181	42.9
12	Giving rescue breathing for an infant is mouth to mouth and nose	Yes	256	60.7
		No	166	39.3

Abbreviations: CPR, cardiopulmonary resuscitation; GSH, Gebretsadik Shawa Hospital; OPD, outpatient department; MTUTH, Mizan Tepi University Teaching Hospital; TGH, Tepi General Hospital.


Table 3 Participants' Practice Response About CPR in MTUTH, Tepi General Hospital and Gebretsadik Shawo Hospital, Southwest Ethiopia

S.No	Variable	Categories	Frequency	Percentage (%)
1	Have you performed CPR on a patient in cardiac emergencies	Yes	151	35.7
		No	271	64.3
2	I check for patient pulse rate before commencing CPR	Yes	266	62.9
		No	156	37.1
3	I ensure a person in a cardiac emergency is laid supine on a relatively hard surface before commencing CPR	Yes	226	53.6
		No	196	46.4
4	I wear latex gloves before commencing CPR	Yes	166	39.3
		No	256	60.7
5	I used both hands to perform chest compression during CPR	Yes	316	75.0
		No	106	25.0
6	I pinch the patient's nostril before giving mouth to mouth ventilation	Yes	208	49.3
		No	214	50.7
7	I do CPR on a patient in a hospital environment	Yes	241	57.1
		No	181	42.9
8	I use my palm on the patient forehead and gently tilt the head forward	Yes	166	39.3
		No	256	60.7
9	I use the same procedure to administer CPR to children and adult	Yes	139	32.9
		No	283	67.1

Abbreviations: CPR, cardiopulmonary resuscitation; OPD, outpatient department; MTUTH, Mizan Tepi University Teaching Hospital.

Level of Involvement in Cardiopulmonary Resuscitation

Based on this study 18, 228, and 176 nurses were never, rarely, or actively involved in CPR procedures, respectively (Figure 2).

**Figure 2** Level of involvement in CPR in MTUTH, Tepi General Hospital and Gebretsadik Shawo Hospital, Southwest Ethiopia.

Abbreviations: CPR, cardiopulmonary resuscitation; MTUTH, Mizan Tepi University Teaching Hospital.

Cardiopulmonary Resuscitation and Its Associated Factors

Those independent variables having p-values <0.25 were analyzed in multivariable logistic regression. In bivariable logistic regression analysis, sex, marital status, experience, knowledge, taking CPR training recently, assigned ward, qualification, level of CPR involvement and confidence had p-values <0.25, so were retained for multivariable logistic regression analysis.

However, in multivariable logistic regression, experience, knowledge, qualification, assigned ward, and level of CPR involvement were independent predictors of CPR good practice (Table 4).

The likelihood of good practice was higher among nurses having 6–10 years experience (AOR = 2.27, 95% CI: 1.25–4.13) and having >10 years experience (AOR = 1.81, 95% CI: 1.10–2.98) compared with those nurses having 1–5 years of work experience.

Table 4 Bi-Variable and Multi-Variable Logistic Regression Analysis Outcome of Factors Affecting CPR Practice Among Nurses in MTUTH, Tepi General Hospital and Gebretsadik Shawa Hospital, Southwest Ethiopia

Characteristics		Frequency	CPR Practice		Univariate	Multivariate	P-value
			Poor	Good	COR (95% CI)	AOR (95% CI)	
Gender	Male	250	67	183			0.071
	Female	172	67	105	1.74 (1.15, 2.64)	0.60 (0.39, 0.91)	
Marital status	Single	146	49	97			0.646
	Married	276	85	191	1.13 (0.74, 1.74)	1.12 (0.69, 1.82)	
Service year/ experience	1–5	106	27	79			0.007*
	6–10	218	65	153	2.19 (1.21, 2.97)	2.27 (1.25, 4.13)	
	>10	98	42	56	1.76 (1.08, 2.89)	1.81 (1.10, 2.98)	
Level of CPR involvement	Never	18	12	6			0.018*
	Rarely	228	73	155	4.25 (1.53, 11.76)	3.77 (1.26, 11.30)	
	Actively	176	49	127	5.18 (1.84, 14.58)	4.60 (1.51, 13.98)	
Ward	OPD	59	11	48			0.009*
	Emergency	81	23	58	0.72 (0.42, 1.24)	1.02 (0.55, 1.90)	
	Inpatient	282	100	182	1.73 (0.78, 3.91)	1.83 (0.78, 4.26)	
Knowledge	Poor	126	52	74			0.041*
	Good	296	90	206	1.61 (0.39, 2.68)	1.37 (0.28, 2.14)	
Qualification	Diploma	104	38	66			0.009*
	Degree	318	96	222	1.33 (0.84, 2.12)	1.54 (0.93, 2.54)	
Confidence	No	147	53	94			0.260
	Yes	275	81	194	1.35 (0.88, 2.07)	1.29 (0.83, 2.00)	
Taking CPR training <3 year	Yes	146	41	105			0.287
	No	276	93	183	0.77 (0.49, 1.19)	1.28 (0.81, 2.03)	

Note: P-value * < 0.05.

Abbreviations: COR, crude odds ratio; AOR, adjusted odds ratio; CPR, cardiopulmonary resuscitation; MTUTH, Mizan Tepi University Teaching Hospital.

In this study the likelihood of good practice was higher among nurses either rarely involved in CPR practice (AOR = 3.77, 95% CI: 1.26–11.30) or actively involved in CPR practice (AOR = 4.60, 95% CI: 1.51–13.98) compared with never involved in CPR practice.

The likelihood of good practice was higher among nurses assigned to the emergency department (AOR = 1.02, 95% CI: 0.55–1.90) compared with those assigned to the outpatients department. Similarly, the odds of good practice were higher among nurses having good knowledge about CPR (AOR = 1.37, 95% CI: 0.28–2.14) than in nurses having poor CPR knowledge.

The odds of good practice were higher among nurses with a nursing degree (AOR = 1.54, 95% CI: 0.93–2.54) than nurses with a diploma.

Qualitative Result

Twelve in-depth interviews were done among the heads of nurses from each unit. As a result, the qualitative results were merged into three thematic areas. Those are the provision of CPR training in the study hospital, the personal attitude of study participants toward CPR, and the confidence of nurses to perform the CPR maneuver.

Provision of CPR Training

Cardiopulmonary resuscitation is one of the pivotal lifesaving procedures and detailed knowledge and skills are required to perform it. To acquire the knowledge and skill needed for the practice of CPR procedure requires intensified education in the theoretical class as well as during practical attachment in a real environment through contact with the patients in a health setting. After graduating and beginning professional work in the health setting continuous upgrading of the knowledge and skill of cardiopulmonary resuscitation is very crucial because the maneuvers need recurrent rehearsal since its science is updated through time. Regarding training, there is a gap in providing the training about CPR which has direct or indirect effects on practice of the procedure.

Perception of Nurses About CPR

Conducting cardiopulmonary resuscitation for patients having respiratory or cardiac arrest or both has a great contribution in saving patient lives. Nurses are among the front-line health profession identifying the need for and initiating cardiopulmonary resuscitation. The procedure has a great importance but it needs knowledge, courage, experience, and skill to perform.

Nurses' Confidence to Perform CPR

Self-confidence is very important to carry out cardiopulmonary resuscitation. Nurses should have self-confidence equally with good knowledge, attitude, skill, and experience to apply CPR. Nurses may have good knowledge about CPR but hesitation in applying the procedure due to lack of confidence in carrying out the maneuver may impose harm on the patient.

Discussion

This study assessed the practice of cardiopulmonary resuscitation and its associated factors among nurses working in the three selected hospitals. The prevalence of good practice towards CPR among nurses was 31.8% (95% CI; 27.5–36.3) in the selected hospitals. The finding was lower than a study carried out in Nigeria which found 65.2% of the study participants had performed CPR while 34.80% had not practiced CPR.²⁰ Possible explanations might be because of study tool variation in using the cut points, service year variation, and due to their qualification status. The majority of the study participants in this study have not taken enough satisfactory training as they were not able to remember information important to applying CPR.

The finding of this study was higher than a study done in Amhara region referral hospital which was 28.4%.²¹ The possible explanations might be study tool variation in using the cut points and service year variation. The above-mentioned study was done on Basic Life Support which is a vast lifesaving maneuver including CPR.

In this study, the level of good knowledge about cardiopulmonary resuscitation was found to be 70.2%. The result was similar to a study done in Wolaita Sodo Teaching Hospital.¹⁸ The possible explanation might be because both studies were done in a similar setting and the study participants' profiles are similar. In addition, the tool used in both studies was similar.

Based on this study those nurses having good knowledge of CPR are one times more likely to have good practice CPR procedures than nurses having poor knowledge. This finding is supported by a study conducted in Amhara region referral hospitals.²¹ The possible explanation might be that in order to practice the procedure the grounding is the theoretical knowledge which means knowing might be the driving force to apply it into practice. However, a study done did not reveal a significant association between knowledge and CPR practice.

This study shows an association between CPR good practice and experience. Those nurses who had work experience of 6–10 years and >10 years were 2 and one times more likely to have good CPR practice than those nurses who had 1–5 years of experience, respectively.

This finding is in line with a study done in Japan.²² Experience has an impact to apply the knowledge into practice. Those experienced nurses face a lot of challenges in health settings which provide the courage, strength, and skill to apply cardiopulmonary resuscitation.

In this study, good practice had a significant association with the assigned ward, where those nurses assigned to the emergency department were more likely to have good CPR practice than those nurses assigned to the outpatients department. This finding conforms with a study done in Turkey.²³ The possible justification for this might be that most of the time patients visiting emergency departments are those who are severe cases which require cardiopulmonary resuscitation to sustain the life of the victims. As a result, nurses assigned to the emergency department have more chances to face patients who need CPR procedures and perform the maneuvers repeatedly than nurses assigned to another ward.

According to this study frequency of CPR involvement had a significant association with good practice, where nurses rarely or actively involved in CPR practice were three and four times more likely to have good practice than those nurses never involved in CPR practice. This finding conforms with a study conducted in Amhara region referral hospitals and Japan.^{21,22} This finding may be justified by the fact that when professionals frequently apply their knowledge they have a greater tendency to use their knowledge to practice the skill (behavior).

Findings from this study revealed that nurses' qualifications had a significant association with good practice, where degree nurses are one times more likely to have good CPR practice than diploma nurses. This finding is in harmony with a study done in Amhara region referral hospitals.²¹ When a qualified professional nurse is advanced the scope of theoretical as well as practical sessions they cover is increased. Due to this, they begin to apply different procedures including cardiopulmonary resuscitation. Unlike the current finding, research work in Kuwait¹⁹ did not show a significant association between good CPR knowledge and nurses' qualifications.

Limitation

This study has strengths and limitations. Using both qualitative and quantitative data methods and incorporating three different health facilities is considered a strength. Among the possible limitations of the study are recall bias and no indication of source and consequence association.

Conclusion

The prevalence of good practice towards cardiopulmonary resuscitation was 31.8%. Having experience of 6–10 and >10 years, actively or rarely involved with cardiopulmonary resuscitation, being assigned to the emergency department, having good CPR knowledge and having a nursing degree were predictors of cardiopulmonary resuscitation good practice. Nurses' practice towards cardiopulmonary resuscitation was low. Therefore, efforts should be taken to enhance nurses' practice towards cardiopulmonary resuscitation.

Abbreviations

AOR, adjusted odds ratio; COR, crude odds ratio; CPR, cardiopulmonary resuscitation; GSH, Gebretsadik Shawo Hospital; MTUTH, Mizan Tepi University Teaching Hospital; SPSS, Statistical Package for Social Sciences; TGH, Tepi General Hospital; WHO, World Health Organization.

Data Sharing Statement

The corresponding author will make data available upon reasonable request.

Ethics Approval and Consent to Participate

Institutional Review Board of Mizan Tepi University was providing ethical clearance with the (Ref. No. Nur/0040/13). An official permission letter was secured from the three hospitals. The verbal consent which was approved by Institutional Review Board was taken from study participants after explaining the objective of the study, the risk, and benefit, issues of confidentiality, and the right of participation. Assurance was given for participants to withdraw or not to participate in the study without any hesitancy. This study was conducted per the Declaration of Helsinki.

Acknowledgments

First of all, I would like to thank Mizan Tepi University Teaching Hospital, Tepi General Hospital, and Gebretsadik Shawo Hospital and staff for their support and cooperation. Next, I extend my gratitude to study participants, data collectors, and supervisors.

Author Contributions

The author made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Disclosure

The author reports no conflicts of interest in this work.

References

1. Catherine AB, Schechter J, Berzon B, Windle ML. Cardiopulmonary resuscitation (CPR). Practical essentials [homepage on the Internet]. [cited August 29, 2017]. Available from: <http://www.emedicine.medscape.com>. Accessed April 8, 2022.
2. Cass A, Bartz A. Why it remains difficult for remote cardiologist to obtain the locus of control for ambulatory health care conditions such as congestive heart failure? A tug of war between general practice, administrators and implementable research findings. *J General Pract.* 2014;02. doi: 10.4172/2329-9126.1000146
3. American Heart Association. The American Heart Association guidelines for cardiopulmonary resuscitation. *Emerg Cardiovasc Care Elvister.* 2020;4:1–5.
4. Raghava S, Nazir R. Adult Basic Life Support (BLS) awareness and knowledge among medical and dental interns completing internship from Deemed University. *Resuscitation.* 2006;70:416–422.
5. Simmes FM, Schoonhoven L, Mintjes J, Fikkers BG, Van der Hoeven JG. Incidence of cardiac arrests and unexpected deaths in surgical; patients before and after implementation of a rapid response system. *Ann Intensive Care.* 2012;2(1):141–147. doi:10.1186/2110-5820-2-20
6. Hamilton R. Nurses Knowledge and skills following cardio-pulmonary resuscitation training: a review of the literature. *J Adv Nurs.* 2005;51(3):288–97. doi:10.1111/j.1365-2648.2005.03491.x
7. Cooper S, Johnston E, Priscott D. Immediate life support training. Impact in a primary care setting? *Resuscitation.* 2007;72:92–99. doi:10.1016/j.resuscitation.2006.06.004
8. Oh S-I II, Han -S-S. A study on the sustainable effect of reeducation on CPR on nurses' knowledge and skill. *J Korean Acad Nurs.* 2008;38(3):383–392. doi:10.4040/jkan.2008.38.3.383
9. Rea Thomas D, Carol F, Linda C, Donohoe Racheal T, Cindy H, Jennifer I. CPR with chest compression alone or with rescue breathing. *Engl J Med.* 2010;36(5):423–433.
10. Ratha K, Panda S, Pradhan R. "Evaluate the effectiveness of planned teaching program regarding basic life support(BLS) among intern (BSc nursing) student at selected nursing college", Bhubaneswar, Odisha. *Nurse Health Sci.* 2014;3(1):16–19.
11. Dariush M, Benjamin Emelia J, Go Alan S, Arnett Donna K, Blaha Michael J. Heart diseases and stroke statistics-2015 update a report from the American heart association. *Circulation.* 2015;131(4):e329–e329.
12. Weisfeldt ML. Public access defibrillation: good or great? *BMJ.* 2004;328:E271–272. doi:10.1136/bmj.328.7438.E271

13. Wenzel V, Lehmkühl P, Kubilis PS, Idris AH, Pichlmayr I. Poor correlation of mouth-to-mouth ventilation skills after BLS training and 6 Month later. *Resuscitation*. 1999;35:129–134. doi:10.1016/S0300-9572(97)00044-0
14. Zaheer H, Haque Z. Awareness about BLS (CPR) among medical students: status and requirements. *J Pak Med Assoc*. 2009;59(1):57–59.
15. Gombotz H, Weh B, Mitterndorfer W, et al. In-hospital cardiac resuscitation outside the ICU by nursing staff equipped with automated external defibrillators—the first 500 cases. *Resuscitation*. 2006;70(3):416–422. doi:10.1016/j.resuscitation.2006.02.006
16. Nagwa I, Abass N, Soliman M. Effect of implementing Advanced Cardiovascular Life Support (ACLS) Guidelines 2016 on nurse's knowledge and performance. *Am J Biomed Res*. 2020;8:534–542.
17. American Heart Association. Cardiopulmonary resuscitation: statement by the Ad Hoc Committee on cardiopulmonary resuscitation of the division of medical science, national academy of sciences, national research council. *JAMA*. 1966;198:372–379. doi:10.1001/jama.1966.03110170084023
18. Marsha Assefa D, Tilahun S. Assessment of Knowledge, Attitude and Practice towards CPR among Nurses Working in Wolaita Sodo University Teaching Hospital Wolaita Sodo, Southern Ethiopia. *Int J Biotech Trends Technol*. 2017;4:54.
19. AL-Kandary S, AL-Jeheidli A, Ghayath T, et al. Perceived competence in cardiopulmonary resuscitation, knowledge and practice among qualified nurses in Kuwait. *Alexandria Bulletin*. 2007;43(2):294–295.
20. Marzooq H, Lyneham J. Cardiopulmonary resuscitation knowledge among nurses working in Bahrain. *Int J Nurs Pract*. 2009;15(4):294–302. doi:10.1111/j.1440-172X.2009.01752.x
21. Ihunanya OM, Oke Michael RN, Babcock BN. Knowledge, attitude, and practice of cardiopulmonary resuscitation among nurses in Babcock university teaching hospital in Ilishan-Remo, Ogun State, Nigeria. *Int J Caring Sci*. 2020;13:1773–1782.
22. Mengistu Mekonnen K. Knowledge, practice and associated factors towards basic life support among nurses working in Amhara region referral hospitals, northwest Ethiopia; 2016.
23. Tanigawa K, Iwami T, Nishiyama C, et al. Are trained individuals more likely to perform bystander CPR? An observational study. *Resuscitation*. 2011;82(5):523–528. doi:10.1016/j.resuscitation.2011.01.027

Open Access Emergency Medicine

Dovepress

Publish your work in this journal

The Open Access Emergency Medicine is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of emergency medicine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/open-access-emergency-medicine-journal>